

Title	The effect of calcium supplementation on fat metabolism during recreational exercise
Authors	Bradley, John L.;Woods, Trevor;Murphy, S.;Keane, L.;O'Brien, Nora M.
Publication date	2009-09
Original Citation	Bradley, J.L., Woods, T., Murphy, S., Keane, L., O'Brien, N.M. 2009. The effect of calcium supplementation on fat metabolism during recreational exercise. In UCC (University College Cork), Proceedings of the 39th annual research conference in Food, Nutrition and Consumer Sciences.
Type of publication	Conference item
Rights	© 2009. The Authors. - http://creativecommons.org/licenses/by-nc-nd/3.0/
Download date	2024-04-28 09:35:19
Item downloaded from	https://hdl.handle.net/10468/190

THE EFFECT OF CALCIUM SUPPLEMENTATION ON FAT METABOLISM DURING RECREATIONAL EXERCISE.

J. Bradley¹, T. Woods², S. Murphy³, L. Keane³ and N. O'Brien³.

¹Department of Education; ²Office of Sport and Recreation; ³Department of Food and Nutritional Sciences; University College Cork, Cork, Ireland

ABSTRACT

Evidence has emerged suggesting dietary calcium can modulate energy metabolism. Studies have linked whole body fat oxidation during free living activities to both acute and habitual calcium intake. The aim of this study was to examine the relationship between acute calcium intake and whole body fat oxidation during simulated recreational activity. Thirty min following ingestion of 800mg calcium supplement or placebo, fat metabolism was monitored during 45min of treadmill walking. Expired air was sampled throughout by online metabolic spirometry (Cosmed CPET, Italy) and analysed for metabolic fuel using indirect calorimetry. The results showed a single, acute calcium supplement of 800mg had no effect on fat metabolism compared to the placebo condition during walking. Contrary to findings from 24 hour studies monitoring calcium intake and macronutrient oxidation, this suggests single acute calcium supplementation protocols are not enough to increase energy metabolism or influence fat metabolism. Evidence suggests that calcium intake increases energy metabolism via whole body systemic changes. Increased energy metabolism and fat oxidation may only occur when recreational exercise is combined with habitual calcium supplementation.

INTRODUCTION

Analysis of the National Health and Examination Survey reported an inverse relationship between body weight and calcium intake¹. This relationship has been developed further to show higher rates of whole body fat oxidation over a 24 hour period with higher acute calcium intake^{2,3}. The aim of this study is to examine the relationship between acute calcium intake and whole body fat oxidation during simulated recreational activity using a commercially available calcium-rich supplement.

MATERIALS AND METHODS

Ten female subjects ([mean \pm SD] age: 21.6 \pm 1.0years; weight: 64.1 \pm 6.4kg; height: 166.6 \pm 5.6cm) visited the Human Performance Laboratory (Mardyke Arena, UCC) on three occasions. Before each visit the subjects completed a food frequency questionnaire to assess habitual calcium intake. Thirty min prior to the exercise protocol on each occasion subjects ingested 500ml of 800mg calcium dissolved in water/MiWadi™ or placebo (MiWadi™ drink). The first visit involved a sub-maximal incremental exercise test on a treadmill (Powerjog GX100) to determine the peak fat oxidation rate of the subject (using indirect calorimetry⁴ adapted from Nordby et al.⁵). The treadmill speed was held constant throughout at between 4-7km/h and the gradient increased by 1% each stage. On visits 2 and 3 subjects ingested either the placebo or calcium supplement 30min prior to exercise. They then carried out a 45min bout of exercise at the peak fat oxidation intensity identified in the first visit. Heart rate (Polar S610i heart rate monitor) and metabolic fuel (via indirect calorimetry using a Cosmed CPET metabolic cart) were analysed throughout.

RESULTS AND DISCUSSION

Fat oxidation was analysed over the whole 45min bout of exercise and from each 15min through the 45min exercise period. No significant difference was seen from the effect of calcium supplementation ($p > 0.05$). There was no relationship between total calcium intake (habitual intake plus the 800mg supplement) and rate of fat oxidation. Habitual calcium intake assessed by the food frequency questionnaire showed a wide range of individual calcium intakes from the diet. However this showed no relationship to the change in fat oxidation following the supplement condition.

Dietary calcium from both supplemental and dairy sources has been shown to increase fat oxidation at rest⁶, during free-living activities that included periods of exercise^{2,3} and during periods of calorie deficit⁷. In the present study the calcium supplement showed no evidence of increasing fat metabolism compared to a placebo (MiWadi™) condition. Several mechanisms have been proposed that suggest a relationship between calcium intake and energy metabolism (summarised in Harris⁸). These include a reduction in intestinal fat absorption and

suppression of 1,25-dihydroxyvitamin D (calcitrol) that inhibit fatty acid synthesis and promote lipolysis. All of these mechanisms rely on systemic changes in the body occurring over an extended duration. The present study used an extended period of exercise (45min). However, previous studies have measured fat oxidation at rest over much longer periods (2-6 hours) and those measuring fat oxidation during free-living activities correlated prandial calcium intake and fat oxidation over 24 hour periods. The mechanisms proposed above may only become apparent over these longer durations.

CONCLUSION

The calcium supplement showed no evidence of increasing fat metabolism compared to a placebo (Miwadi™) condition and suggests single acute calcium supplementation protocols are not enough to increase energy metabolism or fat oxidation. These changes may only occur when recreational exercise is combined with habitual calcium supplementation.

REFERENCES

- ¹ McCarron, D.A., Morris, C.D., Henry, H.J. & Stanton, J.L. (1984). Blood pressure and nutrient intake in the United States. *Science* 224, 1392-1398.
- ² Melanson, E.L., Sharp, T.A., Schneider, J., Donahoo, W.T., Grunwald, G.K. & Hill, J.O. (2003). Relation between calcium intake and fat oxidation in adult humans. *Int. J. Obesity*, 27, 196-203.
- ³ Melanson, E.L., Donahoo, W.T., Dong, F., Ida, T. & Zemel, M.B. (2005). Effect of low- and high-calcium dairy-based diets on macronutrient oxidation in humans. *Obesity Research*, 13(12)2102-2112.
- ⁴ Stisen, A.B., Stougaard, O., Langfort, J., Helge, J.W., Sahlin, K. & Madsen, K. (2006). Maximal fat oxidation rates in endurance trained and untrained women. *Eur. J. Appl. Physiol.* 98, 497-506.
- ⁵ Nordby, P., Saltin, B. & Helge, J.W. (2006). Whole-body fat oxidation determined by graded exercise and indirect calorimetry: a role for muscle oxidative capacity? *Scand. J. Med. Sci. Sports* 16, 209-214.
- ⁶ Cummings, N.K., Soares, M.J., James, A.P. & Chan She Ping-Delfos, W. (2004). Comparison of dairy and non-dairy sources of calcium on thermogenesis and substrate oxidation in humans. *Asia Pac. J. Clin. Nutr.* 13, S87.
- ⁷ Teegarden, D., White, K.M., Lyle, R.M., Zemel, M.B., Van Load, M.D., Matkovic, V., Craig, B.A. & Schoeller, D.A. (2008). Calcium and dairy product modulation of lipid utilization and energy expenditure. *Obesity* 16(7), 1566-1572.
- ⁸ Harris, R.B.S. (2005). Dairy protein, calcium and body weight – the need for a mechanism. *Int. J. Obesity* 29, 388-390.