

Title	Environmental factors correlated with the metabolite profile of <i>Vitis vinifera</i> cv. Pinot Noir berry skins along a European latitudinal gradient
Authors	Del-Castillo-Alonso, María Ángeles;Castagna, Antonella;Csepregi, Kristóf;Hideg, Éva;Jakab, Gabor;Jansen, Marcel A. K.;Jug, Tjaša;Llorens, Laura;Mátai, Anikó;Martínez-Lüscher, Johann;Monforte, Laura;Neugart, Susanne;Olejnickova, Julie;Ranieri, Annamaria;Schödl-Hummel, Katharina;Schreiner, Monika;Soriano, Gonzalo;Teszák, Péter;Tittmann, Susanne;Urban, Otmar;Verdaguer, Dolors;Zipoli, Gaetano;Martínez-Abaigar, Javier;Núñez-Olivera, Encarnación
Publication date	2016-10-31
Original Citation	Del-Castillo-Alonso, M. Á., Castagna, A., Csepregi, K., Hideg, É., Jakab, G., Jansen, M. A. K., Jug, T., Llorens, L., Mátai, A., Martínez-Lüscher, J., Monforte, L., Neugart, S., Olejnickova, J., Ranieri, A., Schödl-Hummel, K., Schreiner, M., Soriano, G., Teszlák, P., Tittmann, S., Urban, O., Verdaguer, D., Zipoli, G., Martínez-Abaigar, J. and Núñez-Olivera, E. (2016) 'Environmental Factors Correlated with the Metabolite Profile of <i>Vitis vinifera</i> cv. Pinot Noir Berry Skins along a European Latitudinal Gradient', <i>Journal of Agricultural and Food Chemistry</i> , 64(46), pp. 8722-8734. doi: 10.1021/acs.jafc.6b03272
Type of publication	Article (peer-reviewed)
Link to publisher's version	10.1021/acs.jafc.6b03272
Rights	© 2016 American Chemical Society. This document is the Accepted Manuscript version of a Published Work that appeared in final form in <i>Journal of Agricultural and Food Chemistry</i> , copyright © American Chemical Society after peer review and technical editing by the publisher. To access the final edited and published work see http://pubs.acs.org/doi/abs/10.1021/acs.jafc.6b03272
Download date	2024-04-27 07:53:37

Item downloaded
from

<https://hdl.handle.net/10468/3276>



UCC

University College Cork, Ireland
Coláiste na hOllscoile Corcaigh

Article

Environmental factors correlated with the metabolite profile of *Vitis vinifera* cv. Pinot Noir berry skins along a European latitudinal gradient

María-Ángeles Del-Castillo-Alonso, Antonella Castagna, Kristóf Csepregi, Éva Hideg, Gabor Jakab, Marcel A.K. Jansen, Tjaša Jug, Laura Llorens, Anikó Máтай, Johann Martínez-Lüscher, Laura Monforte, Susanne Neugart, Julie Olejnickova, Annamaria Ranieri, Katharina Schoedl-Hummel, Monika Schreiner, Gonzalo Soriano, Péter Teszlák, Susanne Tittmann, Otmar Urban, Dolores Verdaguer, Gaetano Zipoli, Javier Martínez-Abaigar, and Encarnacion Núñez-Olivera

J. Agric. Food Chem., **Just Accepted Manuscript** • DOI: 10.1021/acs.jafc.6b03272 • Publication Date (Web): 31 Oct 2016

Downloaded from <http://pubs.acs.org> on November 7, 2016

Just Accepted

“Just Accepted” manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides “Just Accepted” as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. “Just Accepted” manuscripts appear in full in PDF format accompanied by an HTML abstract. “Just Accepted” manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). “Just Accepted” is an optional service offered to authors. Therefore, the “Just Accepted” Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the “Just Accepted” Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these “Just Accepted” manuscripts.

1 **Environmental factors correlated with the metabolite profile of *Vitis***
2 ***vinifera* cv. Pinot Noir berry skins along a European latitudinal**
3 **gradient**

4
5 María Ángeles Del-Castillo-Alonso¹, Antonella Castagna², Kristóf Csepregi³, Éva
6 Hideg³, Gabor Jakab^{3,11}, Marcel A.K. Jansen⁴, Tjaša Jug⁵, Laura Llorens⁶, Anikó
7 Máta³, Johann Martínez-Lüscher⁷, Laura Monforte¹, Susanne Neugart⁸, Julie
8 Olejnickova⁹, Annamaria Ranieri², Katharina Schödl-Hummel¹⁰, Monika Schreiner⁸,
9 Gonzalo Soriano¹, Péter Teszlák¹¹, Susanne Tittmann¹², Otmar Urban⁹, Dolors
10 Verdaguer⁶, Gaetano Zipoli¹³, Javier Martínez-Abaigar¹, and Encarnación Núñez-
11 Olivera^{1,*}

12
13 ¹Faculty of Science and Technology, University of La Rioja, Madre de Dios 53, 26006
14 Logroño (La Rioja), Spain

15 ²Department of Agriculture - Food and Environment, and Interdepartmental Research
16 Center Nutrafood “Nutraceuticals and Food for Health”, University of Pisa, via del
17 Borghetto 80, 56124 Pisa, Italy

18 ³Institute of Biology, University of Pécs, Ifjúság u. 6, 7624 Pécs, Hungary

19 ⁴School of Biological, Environmental and Earth Sciences, University College Cork,
20 College Road, Cork, Ireland

21 ⁵Agricultural and Forestry Institute of Nova Gorica, Pri hrastu 18, 5270 Nova Gorica,
22 Slovenia

23 ⁶Department of Environmental Sciences, Faculty of Sciences, University of Girona,
24 Campus Montilivi, Maria Aurèlia Capmany i Farnés 69, 17003 Girona, Spain

25 ⁷UMR 1287 EGFV, Bordeaux Sciences Agro, INRA, Université de Bordeaux, ISVV,
26 33882 Villenave d’Ornon, France

27 ⁸Department Plant Quality, Leibniz-Institute of Vegetable and Ornamental Crops
28 Grossbeeren/Erfurt e.V., Theodor-Echtermeyer-Weg 1, 14979 Grossbeeren, Germany

29 ⁹Global Change Research Institute CAS, v.v.i, Bělidla 986/4a, 60300 Brno, Czech
30 Republic

31 ¹⁰Department of Crop Sciences, BOKU - University of Natural Resources and Life
32 Sciences, Konrad-Lorenz-Str. 24, 3430 Tulln, Austria

33 ¹¹Research Institute for Viticulture and Oenology, University of Pécs, Pázmány P. u. 4,
34 7634 Pécs, Hungary

35 ¹²Institute for General and Organic Viticulture, Geisenheim University, Von-Lade-
36 Strasse 1, 65366 Geisenheim, Germany

37 ¹³Institute of Biometeorology – National Research Council, Via Caproni 8, 50144
38 Firenze, Italy

39

40 Corresponding Author

41 *(Encarnación Núñez-Olivera) Phone: +34 941299755. Fax: +34 941299721. E-mail:
42 encarnacion.nunez@unirioja.es

43 **ABSTRACT**

44

45 Mature berries of Pinot Noir grapevines were sampled across a latitudinal gradient in
46 Europe, from southern Spain to central Germany. Our aim was to study the influence of
47 latitude-dependent environmental factors on the metabolite composition (mainly
48 phenolic compounds) of berry skins. Solar radiation variables were positively correlated
49 with flavonols and flavanonols and, to a lesser extent, with stilbenes and cinnamic
50 acids. The daily means of global and erythemic UV solar radiation over long periods
51 (bud break-veraison, bud break-harvest and veraison-harvest), and the doses and daily
52 means in shorter development periods (5-10 days before veraison and harvest) were the
53 variables best correlated with the phenolic profile. The ratio between trihydroxylated
54 and monohydroxylated flavonols, which was positively correlated with antioxidant
55 capacity, was the berry skin variable best correlated with those radiation variables. Total
56 flavanols and total anthocyanins did not show any correlation with radiation variables.
57 Air temperature, degree days, rainfall and aridity indices showed fewer correlations
58 with metabolite contents than radiation. Moreover, the latter correlations were restricted
59 to the period veraison-harvest, where radiation, temperature and water availability
60 variables were correlated, making it difficult to separate the possible individual effects
61 of each type of variable. The data show that managing environmental factors, in
62 particular global and UV radiation, through cultural practices during specific
63 development periods, can be useful to promote the synthesis of valuable nutraceuticals
64 and metabolites that influence wine quality.

65

66 Keywords: *Vitis vinifera* cv. Pinot Noir, latitudinal gradient, phenolic composition,
67 berry skins, solar radiation, ultraviolet radiation, hydroxylation ratios, Europe

68

69 **INTRODUCTION**

70

71 Environmental factors, such as air temperature, ambient solar radiation (including UV)
72 and photoperiod, vary with latitude. In turn, variations in these environmental factors
73 may cause changes in physiological and/or biochemical characteristics of plants. Yet,
74 this is not always the case as plant responses to latitudinal climatic conditions may be
75 masked by, for example, local climatic factors, cultivational measures, or pest and
76 diseases. Thus, there is a need for latitudinal studies that help to identify the
77 environmental factors that impact most on plants, as well as the traits most affected.
78 Such studies are important in terms of understanding ecological processes (especially in
79 the context of climate change), but also have a direct relevance for the agricultural
80 industry. A number of plant traits have been studied in relation to latitude, including
81 plant height, seed production, growth, biomass production, photosynthesis rates,
82 chlorophyll fluorescence, photosynthetic pigment composition, mineral nutrient
83 contents and ratios, water relations and secondary metabolite contents.¹⁻⁸ Most of these
84 traits have been measured in leaves, whereas only a few studies have used fruits.
85 Latitude-related environmental variables that have been hypothesized to explain
86 changes in plant traits include air temperature, degree days, rainfall, aridity indices, soil
87 moisture, total solar radiation doses, and UV radiation doses. Most latitudinal studies
88 have been carried out using wild species, while only a few studies have dealt with
89 commercially interesting species, such as juniper,³ ryegrass⁷ and currant.⁸ To our
90 knowledge, no study has dissected the effects of latitudinal gradients, and the associated
91 environmental parameters, on grapevine, although latitude is a recognized factor used,
92 for example, to predict the suitability of territories for grapevine culture.⁹

93

94 Remarkably, the effects of latitude and associated environmental parameters on the
95 phenolic composition of grapevine berries have not been studied, in spite of the fact that
96 similar studies have been conducted on other species with less commercial impact.^{3-5,7,8}
97 This omission is even more remarkable, given that the phenolic compounds synthesized
98 in grapevine berries decisively determine wine characteristics and quality, including the
99 presence of important nutraceuticals and nutritionally-desirable antioxidants.^{10,11} Berry
100 skin is the main source of many of these phenolic compounds, including anthocyanins,
101 flavonols and stilbenes.¹²⁻¹⁴

102

103 The present study was conducted on Pinot Noir grapevines. This variety is the tenth
104 most cultivated grapevine worldwide, and the seventh fastest-expanding winegrape
105 variety in the period 2000-2010.¹⁵ Pinot Noir grapevines occupy more than 86,000 ha in
106 the world (1.88% of the total grapevine acreage), especially in Europe, where it
107 occupies 3% of the total acreage. Pinot Noir is especially adapted to cold climates, thus
108 ascending to higher latitudes than other varieties. In fact, the European distribution of
109 this cultivar ranges from southern Spain to central Germany. Given this wide ranging
110 distribution, our aim was to identify the influence of latitude and associated
111 environmental parameters (air temperature, global and UV radiation, rainfall and
112 aridity) on the metabolite composition of berry skins of *Vitis vinifera* cv. Pinot Noir in
113 Europe. This study will inform management of those environmental parameters that
114 affect berry skin composition. In turn, a better understanding of the influence of these
115 parameters can help improve wine quality.

116 **MATERIALS AND METHODS**

117

118 **Collection sites and environmental variables**

119

120 Berries of Pinot Noir grapevines (*Vitis vinifera* L.) were collected in 2013 from 11
121 localities in Spain, France, Italy, Hungary, Austria, Slovenia, the Czech Republic and
122 Germany (Figure 1, Table 1). This represented a latitudinal gradient of almost 14° (36.7-
123 50.0 °N) and a linear distance of around 1,500 km, covering most of the commercial
124 Pinot Noir growing latitudes in the Northern Hemisphere (35-55°).¹⁶ Vineyard age
125 varied between 6 and 30 years, and vineyard soils were mostly calcareous and neutral-
126 alkaline (pH between 7.0 and 8.5). No fertilization or irrigation had been applied to the
127 vineyards.

128

129 In each locality, berry samples were collected from three separate plants (replicates) at
130 commercial maturity, always around noon-time, and on a sunny day. Collection dates
131 varied from 31 July to 22 October, depending on the location. Three clusters were
132 collected for each replicate. As row orientation varied between vineyards, clusters were
133 always picked from a SE-orientated shoot. In situ, every berry was separated from its
134 cluster by cutting the pedicel. Subsequently, berry density was determined as
135 floatability in a NaCl solution series, which allowed for harvesting berries of a similar
136 ripeness using a non-destructive method.^{17,18} To reduce the variability that is normally
137 found within a cluster, berries with a density between 140-160 g NaCl l⁻¹ were selected,
138 rinsed in distilled H₂O and immediately transported to the laboratory in a portable
139 icebox. In the laboratory, berries were frozen in liquid nitrogen and kept at -80°C until
140 further analyses.

141

142 Relevant environmental data were obtained for each locality. Daily values of mean
143 temperature, rainfall and ground-station global radiation (GGR) were obtained for the
144 period bud break-harvest from the nearest meteorological observatory to each vineyard.
145 For most vineyards, meteorological stations were located less than 200 m from the
146 actual vineyards. Remaining stations were located less than 20 km away, except in the
147 case of Lednice (Czech Republic) where the station for GGR measurement was located
148 50 km from the vineyard. In the latter cases, it was ascertained that meteorological
149 stations were located at a similar latitude and altitude as the respective vineyards, which

150 makes the assumption that data were homogeneous. Based on these data, two aridity
151 indices were calculated: the ratio Rainfall/ETP, where ETP is the potential
152 evapotranspiration computed according to Hargraves formula (based on solar global
153 radiation and mean air temperature), and the Gausse Index (the ratio between rainfall
154 and twice the mean daily temperature). In addition, daily values of DSSF (Downward
155 Surface Shortwave Flux) global radiation and TEMIS-derived erythemal UV radiation
156 (T UVery) were obtained for the period bud break-harvest. Daily DSSF was calculated
157 by integrating the 30 minutes of data downloaded from the LandSaf web page
158 (<http://landsaf.meteo.pt>). The data in this archive take into account the differences in the
159 day-length of the various locations. T UVery was downloaded from the ESA-TEMIS
160 web page (<http://www.temis.nl>) and estimated on the basis of Meteosat data (to assess
161 cloud cover), SCIAMACHY data (to assess O₃ column) and a radiative transfer
162 model.¹⁹ The degree days (using 10°C as base temperature) and the daily doses of GGR,
163 DSSF and T UVery were integrated over three different periods: bud break-veraison,
164 bud break-harvest, and veraison-harvest. Additionally, DSSF and T UVery doses were
165 integrated for 5 and 10 days before veraison, and for 5 and 10 days before harvest,
166 because the periods around veraison and prior to harvest are important for the synthesis
167 of phenolic compounds in grapevine berries and, thus, for their commercial quality.²⁰⁻²²

168

169 **Analysis of berries**

170

171 Frozen berries were allowed to partially thaw and the skin was carefully removed from
172 the flesh using a scalpel, and without rupturing the hypodermal cells. The content of
173 total soluble solids (TSS) was measured in °Brix in the flesh, using a digital
174 refractometer. The skins were immediately submerged in liquid nitrogen, weighed and
175 lyophilized. Lyophilized berry skins were weighed and ground to obtain a homogeneous
176 powder for each replicate. Then, all the samples were shipped to one laboratory for
177 detailed analysis of metabolites.

178

179 For each analytical sample used for the analysis of phenolic compounds, 50 mg of skin
180 powder was frozen in liquid nitrogen and ground again in a TissueLyser (Qiagen,
181 Hilden, Germany). The total content of methanol-soluble phenolic compounds
182 (MSPCs), mainly located in the vacuoles,¹⁸ was measured by spectrophotometry. For
183 this analysis, 2 ml of a mixture of methanol: water: 7M HCl (70:29:1 v:v:v) was added

184 for extraction (24 h at 4°C in the dark). The extract was centrifuged at 6000 g for 15 min
185 and the supernatant was selected for spectrophotometry. The level of MSPCs was
186 measured as the area under the absorbance curve in the wavelength intervals between
187 280-315 and 280-400 nm (AUC₂₈₀₋₃₁₅ and AUC₂₈₀₋₄₀₀ respectively) and normalised per
188 unit of dry weight (DW),²³ using a λ 35 spectrophotometer (Perkin-Elmer, Wilton, CT,
189 USA). Individual phenolic compounds were analysed by ultra-performance liquid
190 chromatography (UPLC) using a Waters Acquity UPLC system (Waters Corporation,
191 Milford, MA, USA).²³ Solvents were: A, water/formic acid (0.1%), and B, acetonitrile
192 with 0.1% formic acid. The gradient program employed was: 0-7 min, 99.5-80% A; 7-9
193 min, 80-50% A; 9-11.7 min, 50-0% A; 11.7-15 min, 0-99.5% A. The UPLC system was
194 coupled to a micrOTOF II high-resolution mass spectrometer (Bruker Daltonics,
195 Bremen, Germany) equipped with an Apollo II ESI/APCI multimode source and
196 controlled by the Bruker Daltonics DataAnalysis software. The electrospray source was
197 operated in positive or negative mode. The capillary potential was set to 4 kV; the
198 drying gas temperature was 200 °C and its flow 9 l min⁻¹; the nebulizer gas was set to
199 3.5 bar and 25 °C. Spectra were acquired between *m/z* 120 and 1505 in positive mode
200 for anthocyanins and in negative mode for the remaining phenolic compounds. The
201 different phenolic compounds analysed were identified according to their order of
202 elution and the retention times of the following pure compounds: myricetin, quercetin,
203 catechin, epicatechin, astilbin, *trans*-resveratrol, *p*-coumaric acid, caffeic acid and
204 ferulic acid (Sigma, St. Louis, MO, USA); kaempferol-3-*O*-glucoside, isorhamnetin-3-*O*-
205 *O*-glucoside, syringetin-3-*O*-glucoside, procyanidin B1 and malvidin-3-*O*-glucoside
206 (Extrasynthese, Genay, France); isorhamnetin, quercetin-3-*O*-glucoside, quercetin-3-*O*-
207 galactoside, quercetin-3-*O*-glucopyranoside, quercetin-3-*O*-glucuronide and quercetin-
208 3-rutinoside (Fluka, Buchs, Germany). Quantification of compounds that were not
209 commercially available was carried out using the calibration curves belonging to the
210 most similar compound: myricetin for its glucosides; isorhamnetin for isorhamnetin-3-*O*-
211 *O*-glucuronide; quercetin for quercetin-3-*O*-arabinoside; astilbin for taxifolin-3-*O*-
212 glucoside; *trans*-resveratrol for its glucoside; *p*-coumaric acid for *p*-coumaroyl-tartaric
213 acid; caffeic acid for *p*-caffeoyl-tartaric acid; ferulic acid for feruloyl-tartaric acid; and
214 malvidin-3-*O*-glucoside for anthocyanins. Total contents of the different phenolic
215 groups were obtained as the sum of the individual compounds. The ratios between
216 trihydroxylated and dihydroxylated (3',4',5'-OH/3',4'-OH) anthocyanins, and between

217 trihydroxylated and monohydroxylated (3',4',5'-OH/4'-OH) and trihydroxylated and
218 dihydroxylated (3',4',5'-OH/3',4'-OH) flavonols, were also calculated.

219

220 For carotenoid and chlorophyll extraction,²⁴ 6 ml of a mixture of methanol, acetone, and
221 hexane (1:1:1 v:v:v) was added to a glass tube containing 50 mg of lyophilized skin
222 powder. The mixture was vortexed for 30 s and then stirred for 30 min at 4°C in the
223 dark. After the addition of 2 ml of MilliQ water the tube was vigorously shaken for 1
224 min and then centrifuged for 1 min at 1500 g. The non-polar phase containing
225 carotenoids and chlorophylls was recovered. The extraction was repeated by adding 2
226 ml of hexane to the remaining mixture. The two extracts were pooled and the volume
227 reduced to 1 ml by vacuum evaporation. The extract was filtered through 0.2- μ m filters
228 and immediately subjected to high-performance liquid chromatography (HPLC)
229 analysis as follows. Separation was performed at room temperature by a Spectra System
230 P4000 HPLC, equipped with a UV 6000 LP photodiode array detector (Thermo Fisher
231 Scientific, Waltham, MA, USA) using a Zorbax ODS column (5 μ m particle size, 250 x
232 4.6 mm, Agilent Technologies, Santa Clara, CA, USA). HPLC separation was carried
233 out at a flow rate of 0.8 ml min⁻¹ using the following linear gradient: 0 min, 82% A
234 (CH₃CN), 18% B (methanol/hexane/CH₂Cl₂ 1:1:1 v:v:v); 20 min, 76% A, 24% B; 30
235 min, 58% A, 42% B; 40 min, 39% A, 61% B. The column was allowed to re-equilibrate
236 in the starting solution (82% A, 18% B) for 5 min before the next injection. Different
237 individual chlorophylls and carotenoids were detected by their absorbance at 445 nm.

238

239 The antioxidant capacity of berry skins was measured by generating the radical cation
240 2,2'-azinobis(3-ethylbenzothiazoline-6-sulfonic acid) (ABTS^{•+}).²⁵ The radical solution
241 was diluted in ethanol to obtain an absorbance of 0.700 \pm 0.020 at 734 nm (Perkin-
242 Elmer λ 35 spectrophotometer). After addition of 1 ml of diluted ABTS^{•+} solution to 100
243 μ l of skin extract (250 μ g of skin powder in 1 ml of a mixture of methanol: water: 7M
244 HCl 70:29:1 v:v:v), the decrease in absorbance was monitored and compared to that of
245 the Trolox standard (Sigma) exactly 4 min after initial mixing. Antioxidant capacity was
246 expressed in terms of Trolox equivalent antioxidant capacity (TEAC) per g DW of skin.

247

248 DNA isolation from lyophilized berry skins was carried out using the ZenoGene40
249 Plant DNA Purifying Kit (Zenon Bio Kft., Szeged, Hungary). Concentration of the
250 samples was measured with a Genova Nano Spectrophotometer (Jenway, Staffordshire,

251 UK). DNA content per DW of berry skin (ng mg^{-1} DW) was calculated using the
252 formula: mean of DNA concentration ($\text{ng } \mu\text{l}^{-1}$) multiplied by the volume of extraction
253 (μl) and divided by the DW of the lyophilized sample (mg). This analysis served to
254 calculate the metabolite concentrations on a DNA basis.

255

256 **Statistical analysis**

257

258 Pearson correlation coefficients (r) were used to examine the relationships between all
259 the variables studied, both the environmental-geographical parameters and the traits
260 analyzed in berry skins, including the total contents of the different groups of phenolic
261 compounds. Correlations were considered significant when $p < 0.05$. The sampling
262 localities were ordinated by Principal Components Analysis (PCA), taking into account
263 MSPCs and the total contents of the different groups of phenolic compounds. All the
264 statistical procedures were performed with SPSS 19.0 for Windows (SPSS Inc.,
265 Chicago, IL, USA).

266 **RESULTS**

267

268 **Variation in environmental variables**

269

270 The latitudinal gradient used in this study was associated with substantial differences in
271 several meteorological variables (Table 2). For the period from bud break to harvest,
272 these differences were, amongst others, around 5°C in mean daily temperature, 500
273 degree days, almost 300 mm in rainfall, almost 900 MJ m⁻² in DSSF dose, and 241 kJ
274 m⁻² in T UVery dose. Interestingly, the parameters displaying the greatest differences
275 were the DSSF and T UVery doses accumulated during the 10 days before harvest. For
276 these variables, the differences between the maximum and the minimum values along
277 the gradient were more than 80% of the maximum value. The highest and lowest values
278 of temperature variables were usually recorded in Pécs and Rioja, respectively, except
279 for the veraison-harvest period, in which they were recorded in Spanish localities (Jerez
280 or Girona) and Lednice, respectively. The highest mean values of solar radiation (GGR,
281 DSSF, T UVery) were always recorded in Jerez, and this included also the highest
282 accumulated doses in the 5 or 10 days before veraison and before harvest. The highest
283 accumulated doses over longer periods were recorded in Spanish localities (either Rioja,
284 Girona or more rarely Jerez) or in Lednice, depending on the length of the period
285 considered, because those periods were longer in Rioja, Girona or Lednice than in Jerez
286 (see Table 1 for the length of the period bud break-harvest). The lowest values of
287 radiation variables were generally recorded in Geisenheim or Lednice.

288

289 **Variation in berries variables**

290

291 Metabolite contents were obtained and normalized against both berry skin DW (Table
292 3) and DNA amount. The correlations between metabolites and environmental
293 parameters were similar irrespective of the normalization approach, given that DNA
294 amount and berry skin DW were significantly correlated ($r = 0.79$, $p < 0.01$, $n = 11$).
295 Therefore, results are only described on a per berry skin DW basis. MSPC values varied
296 between 9.7 and 40.3 (as AUC₂₈₀₋₃₁₅ mg⁻¹ DW) and between 17.1 and 74.3 (as AUC₂₈₀₋
297 ₄₀₀ mg⁻¹ DW). Absorption levels in the two wavelength regions were strongly and
298 positively correlated (Table S1). The highest and lowest MSPC values were found in
299 Girona and Lednice, respectively. We quantified 29 phenolic compounds: 24 flavonoids

300 (14 flavonols, 5 anthocyanins, 3 flavanols –monomeric or dimeric tannins-, and 2
301 flavanonols) and 5 non-flavonoids (3 cinnamic acids and 2 stilbenes). Great differences
302 in the concentrations of most groups of phenolic compounds were found between
303 localities. Anthocyanins were the most abundant group, showing values between 18.9
304 (Bilje) and 110.1 (Girona) mg g⁻¹ DW. In every locality, malvidin-3-*O*-glucoside was
305 the major anthocyanin. Flavonols were the second most abundant group of flavonoids,
306 ranging between 1.76 (Bilje) and 7.7 (Girona) mg g⁻¹ DW. The major flavonol was
307 quercetin 3-*O*-glucuronide. Flavanonols (between 0.18 and 1.14 mg g⁻¹ DW, in Bilje
308 and Jerez, respectively) and flavanols (between 0.21 and 0.99 mg g⁻¹ DW, in Lednice
309 and Bilje, respectively) were less abundant. Among non-flavonoids, cinnamic acids
310 were the most abundant group, and also the group showing the greatest variability
311 between localities, with values between 0.16 (Lednice) and 7.2 (Firenze) mg g⁻¹ DW.
312 Finally, the least abundant compounds were stilbenes, which also showed a great
313 variability (between 14 and 928 µg g⁻¹ DW, in Lednice and Girona, respectively).

314

315 The antioxidant capacity of berry skin extracts varied between 3592 (Lednice) and 9104
316 (Firenze) µM TE g⁻¹ DW. Chlorophylls and all carotenoids showed the highest values in
317 Rioja and the lowest in Pécs. β-Carotene was the most abundant carotenoid. The berry
318 fresh weight varied between 1.1 (Girona and Bordeaux) and 2.1 g (Geisenheim),
319 although most localities showed values between 1.1 and 1.3 g. TSS varied between 19.1
320 (Bilje) and 23.7 °Brix (Jerez).

321

322 **Correlations between variables**

323

324 The correlations between all the environmental and plant response variables were
325 determined (Table S1). Unless otherwise stated, the correlations mentioned in this text
326 were significant ($p < 0.05$) and positive. With respect to the correlations between berry
327 skin variables, MSPCs were correlated with the contents of most phenolic compounds
328 (except flavanols) and carotenoids. The total contents of flavonols, flavanonols,
329 stilbenes and anthocyanins were correlated with one another, whereas the total content
330 of cinnamic acids was only correlated with that of flavanonols. Total flavanol content
331 was not correlated with the total content of any other phenolic group. The antioxidant
332 capacity of berry skin extracts was correlated with anthocyanins, MSPCs, flavonols, the
333 ratio 3',4',5'-OH/3',4'-OH flavonols and, less significantly, with flavanonols, cinnamic

334 acids, the ratio 3',4',5'-OH/4'-OH flavonols, and carotenoids. There was no correlation
335 between the antioxidant capacity and contents of stilbenes or flavanols. Carotenoid and
336 chlorophyll contents were correlated with each other, and carotenoid levels were also
337 correlated with those of stilbenes.

338

339 Possible correlations between environmental-geographical parameters and berry skin
340 variables were also explored. It was found that latitude was negatively correlated with
341 MSPCs and the total contents of flavonols, flavanols and stilbenes, but not flavanols,
342 cinnamic acids, anthocyanins and carotenoids (Figure 2).

343

344 Correlations between temperature variables and berry variables were few for the periods
345 bud break-veraison and bud break-harvest. The mean daily temperature and degree days
346 in the period bud break-veraison (but not bud break-harvest) were correlated
347 (negatively) with carotenoids, chlorophylls and TSS, only. Degree days in the period
348 bud break-veraison were also correlated with flavanols. No temperature variable in
349 these two periods was correlated with the total content of any other phenolic group,
350 although there were some correlations between temperature variables and individual
351 compounds. For the period veraison-harvest, the mean daily temperature and degree
352 days were correlated with MSPCs and the total contents of flavonols and flavanols. In
353 addition, the mean daily temperature was correlated with the ratios 3',4',5'-OH/4'-OH
354 and 3',4',5'-OH/3',4'-OH flavonols, and the degree days with the total content of
355 anthocyanins.

356

357 Rainfall and aridity indices were hardly correlated with berry skin variables for the
358 periods bud break-veraison and bud break-harvest. Only quercetin showed somewhat
359 consistent (positive) correlations with rainfall, the Rainfall/ETP ratio and Gaussen Index
360 (but only in the period bud break-harvest). For the period veraison-harvest, rainfall and
361 aridity indices were negatively correlated with the total content of flavonols and
362 flavanols. In addition, Gaussen index was negatively correlated with MSPCs and the
363 ratios 3',4',5'-OH/4'-OH and 3',4',5'-OH/3',4'-OH flavonols.

364

365 Radiation variables, particularly DSSF and T UVery variables, correlated well with
366 berry skin variables for the three periods considered. The daily means of DSSF and T
367 UVery in the periods bud break-harvest and veraison-harvest, the DSSF doses in the 10

368 days before harvest, the daily mean of T UVery in the 5 and 10 days before veraison,
369 and the T UVery doses in the 5 and 10 days before veraison were all correlated with
370 MSPCs. The same variables, together with the T UVery doses in the 10 days before
371 harvest and in the period bud break-harvest (in this last case, with a lower significance
372 level), were correlated with the total contents of flavonols and flavanonols. Total
373 stilbene content was only correlated with the DSSF and T UVery doses in the period
374 bud break-harvest, and total cinnamic acid content only with the daily mean and the
375 dose of T UVery in the 10 days before veraison. Total flavanol and anthocyanin
376 contents were not correlated with any radiation variable. Regarding individual
377 compounds, the strongest correlations were found between contents of several flavonols
378 and flavanonols and the daily means of DSSF and T UVery in the periods bud break-
379 harvest and veraison-harvest, as well as with the DSSF and T UVery doses in the
380 periods of 5 or 10 days before veraison or harvest. Levels of two flavanols, one
381 anthocyanin and the three cinnamic acids analyzed were also correlated with some of
382 those T UVery expressions.

383

384 The ratio 3',4',5'-OH/3',4'-OH anthocyanins was not correlated with any radiation or
385 temperature variable. Yet, the ratios 3',4',5'-OH/4'-OH and 3',4',5'-OH/3',4'-OH
386 flavonols were the berry skin variables that displayed the strongest correlations with
387 specific radiation variables, such as the daily means of DSSF and T UVery in the
388 periods bud break-harvest and veraison-harvest, and the accumulated doses in the 10
389 days before veraison and harvest. This correlation did, however, not extend to the
390 accumulated doses in longer periods, as Figure 3 shows for the period bud break-
391 harvest. Finally, the number of days from bud break to harvest and from veraison to
392 harvest were negatively correlated with total and several individual flavanols.

393

394 **Principal Components Analysis**

395

396 The localities studied were ordinated by PCA using MSPCs and the different groups of
397 phenolic compounds. The accumulated variance by the first three axes was 94.0%
398 (67.3% for axis I, 17.3% for axis II and 9.4% for axis III). The plot using the first two
399 axes, together with the loading factors and their significance, is shown in Figure 4. The
400 total contents of all the phenolic groups, except flavanols, were significant loading
401 factors for the positive part of axis I, which broadly ordinated the localities on the basis

402 of their latitude, with southernmost localities situated towards the positive part of the
403 axis and the northernmost ones towards the negative part. Total flavanols and total
404 cinnamic acids were the only significant loading factors for the positive part of axis II,
405 which separated localities 4, 6, 9, 7 and 1 from the remaining ones. No significant
406 loading factor was found for the negative part of axes I and II.

407 DISCUSSION

408

409 Environmental-geographical gradients, such as those related to latitude, can be exploited
410 to explore and predict the physiological and/or biochemical responses of plants by using
411 a space-for-time substitution.⁶ This type of study cannot necessarily pinpoint the
412 influence of one particular environmental parameter on a plant response, as can be done
413 in controlled studies. However, the strength of latitudinal studies is that plant responses
414 are studied under realistic conditions (i.e. commercial vineyards), where plants are
415 exposed to a natural combination of ambient, environmental parameters. In this study a
416 range of metabolites were measured in skins of Pinot Noir berries, originating from 11
417 vineyards along a latitudinal gradient of nearly 14°. The levels of the various
418 metabolites measured in Pinot Noir berry skins were broadly in agreement with levels
419 measured in other studies using this, or other cultivars.^{12,18,23}

420

421 Radiation is an important determinant of berry skin metabolite profile

422

423 A key finding of this study is that the contents of MSPCs, flavonols, flavanonols and
424 stilbenes in Pinot Noir berry skins increased with decreasing latitudes. Previously,
425 similar results were found for MSPC contents in leaves of *Lolium perenne*,⁷ but no
426 comparative results existed for specific phenolic compounds nor for grapevine. It might
427 be argued that negative correlations between latitude and the abovementioned phenolic
428 groups are due to the longer berry maturation period at lower latitudes. However, we
429 consider this unlikely because (1) latitude was not significantly correlated with the
430 number of days from veraison to harvest, and (2) the latter variable was not correlated
431 with the contents of those phenolic compounds. Rather, the correlations between
432 latitude and contents of phenolic compounds were probably due to the negative
433 correlation between latitude and radiation (both global and UV) variables. Radiation
434 variables were strongly and positively correlated with the total contents of most
435 phenolic groups, mainly flavonols and flavanonols, and to a lesser extent with stilbenes
436 and cinnamic acids, together with MSPCs. The relationship between radiation levels
437 and the content of these phenolic compounds had previously been reported for berry
438 skins of several red grapevine varieties, such as Pinot Noir, Merlot, Malbec and
439 Cabernet Sauvignon,²⁶⁻²⁹ although not in relation with latitudinal gradients.

440

441 Rather than radiation in general, the means of DSSF and T UVery over long periods
442 (bud break-veraison, bud break-harvest and veraison-harvest) and the means or doses in
443 important development periods (5-10 days before veraison and harvest) were the
444 variables best correlated with phenolic compounds, particularly flavonols, flavanonols
445 and cinnamic acids. This is related to the fact that the periods around veraison and prior
446 to harvest are important for the synthesis of phenolic compounds.²⁰⁻²² The stimulation of
447 flavonol accumulation was expected because these compounds are radiation-reactive and
448 concentrations are well known to increase with increasing levels of solar radiation
449 (particularly UV-B) in grapevine berry skins.^{13,18,27,29-33}

450

451 It is not simply total flavonol levels that correlate with radiation parameters, the ratios
452 3',4',5'-OH/4'-OH and 3',4',5'-OH/3',4'-OH flavonols were the berry skin variables
453 best correlated with specific radiation variables, such as the mean values or doses of
454 DSSF and T UVery radiation in critical periods (5-10 days before veraison and harvest),
455 but not with the accumulated doses over long periods (Figure 3). Thus, higher solar
456 radiation values (both total and UV) in those critical periods might increase the B-ring
457 hydroxylation level of flavonols in Pinot Noir berry skins. Previously, it was shown that
458 the hydroxylation level depends on both the grape variety¹² and environmental factors,
459 such as the radiation level. The effect of radiation, in turn, may depend again on the
460 variety considered: the hydroxylation ratios increased with increasing total or UV
461 radiation in Pinot Noir (this study), but decreased with increasing total or UV radiation
462 in Sangiovese²² and Tempranillo.^{18,34} This complexity may be caused by the intricate
463 regulation mechanism of the genes and enzymes involved in the synthesis of flavonols
464 with different hydroxylation levels.^{21,30,31} In petunia, the highest level of B-ring
465 hydroxylation was caused by the specific effect of increased UV-B radiation.³⁵ The
466 antioxidant activity of flavonoids strongly depends on the number of hydroxyl groups
467 bound to the aromatic B-ring.³⁶ Given that the hydroxylation ratios were positively
468 correlated with the antioxidant capacity in our study, flavonols may be important as
469 both sunscreens and antioxidants in Pinot Noir berry skins, and their role as antioxidants
470 would increase in those localities with higher radiation levels.

471

472 Flavanonols (dihydroflavonols) are bioactive compounds that contribute to tolerance to
473 fungal infections and colour expression in some red wines.³⁷ Given that flavanonols
474 comprise a relatively small fraction of total wine flavonoids, their regulation by, and

475 responses to, radiation were not clear. However, the results in this paper show that
476 flavanone levels were positively correlated with radiation. This observation is
477 consistent with a previous study that reported increases in flavanones in Malbec berry
478 skins following exposure to higher solar radiation levels due to cluster thinning.³⁷
479 Similarly, flavanone levels were found to be elevated in berries exposed to ambient
480 UV-B, in comparison with berries receiving no UV-B.¹³

481

482 The reported data indicate positive correlations of cinnamic acid levels with radiation.
483 Consistently, higher values of caffeoyl-tartaric acid were found in skins of Pinot Noir
484 berries exposed to solar radiation when compared with shaded berries.²⁶ However, not
485 all studies show increases in cinnamic acids with increasing radiation. Coumaroyl-
486 tartaric acid levels showed no response to solar UV-B radiation exposure in Malbec
487 berry skins.²⁸ Probably, the synthesis of cinnamic acids in berries is more influenced by
488 the radiation received prior to veraison, because contents are highest before berry
489 maturation.¹⁴ Besides, there is some debate on whether cinnamic acids are
490 predominantly present in pulp, rather than skin. Furthermore, the response of cinnamic
491 acid levels to variations in radiation appears to be influenced by the specific year,³⁸ and
492 each specific cinnamic acid seems to react in a different way.¹⁸

493

494 In contrast to flavonol and flavanone content, the levels of total stilbenes were only
495 correlated with the global and UV radiation doses over long periods (bud break-
496 harvest). Both stilbenes and flavonoids derive from coumaroyl-coenzyme A in the
497 general phenylpropanoid metabolism, but stilbenes are synthesized by stilbene synthase
498 instead of chalcone synthase. Stilbene synthase is found in berry skins during all stages
499 of fruit development,³⁹ which could explain the correlation of total stilbene contents
500 with global and UV doses over long periods. Yet, similar to flavonols, stilbenes
501 (resveratrol) were also found to be UV-induced, as was demonstrated in studies using
502 Malbec berry skins.²⁸

503

504 It was found in this study that the total content of anthocyanins was not correlated with
505 any radiation variable. This finding is congruent with previous findings on Pinot Noir
506 berry skins, which showed that anthocyanin content was not affected by sun exposure.²⁶
507 The finding is also consistent with the fact that anthocyanin biosynthesis is controlled
508 by a different system than that controlling flavonol biosynthesis.⁴⁰ In general,

509 anthocyanins are accumulated under conditions of low temperature and high radiation
510 levels,^{8,41} but contradictory data have been reported in grape berries as a consequence of
511 differences in cultivar, site, season, sampling and analytical techniques.⁴² In addition, it
512 has often been difficult to separate the effects of light and temperature.

513

514 The ratio 3',4',5'-OH/3',4'-OH anthocyanins was also not correlated with any radiation
515 variable (unlike the hydroxylation ratio of flavonols). Previous studies had shown that
516 the hydroxylation ratio of anthocyanins may increase⁴³ or decrease^{31,44} with increasing
517 (total or UV) radiation in different grapevine varieties, and even the responses may vary
518 depending on the year of study.^{27,30} These diverse responses to radiation may be due not
519 only to a complex regulation of the synthesis of differently hydroxylated anthocyanins
520 in the different varieties (as occurred with respect to the hydroxylation ratios of
521 flavonols), but also to the specific responses of each individual anthocyanin. For
522 example, in our study the trisubstituted malvidin-3-*O*-glucoside was the only
523 anthocyanin (positively) correlating with radiation variables, thus affecting the response
524 of the ratio to radiation.

525

526 Total flavanol levels were not correlated with any radiation variable nor with levels of
527 any other phenolic group. A likely explanation for this observation is that flavanols are
528 synthesized during the early stages of berry development and that their levels remain
529 fairly stable during subsequent berry growth. Several authors have reported that flavanol
530 levels are stable, and show little responsiveness to changes in radiation or other
531 environmental parameters.^{14,44,45} Nevertheless, there is no consensus on this point, as
532 solar UV exclusion has been reported to decrease flavanol content,²⁹ and responses to
533 temperature and water availability have also been reported.⁴⁵⁻⁴⁷

534

535 Thus, it is concluded that radiation is strongly correlated with Pinot Noir berry skin
536 phenolic profile. Radiation-related changes in phenolic profile are highly specific.
537 Radiation appears to affect one class of metabolites, while other compounds are not
538 affected. Such specific regulatory interactions offer scope to precision manipulation of
539 berry skin metabolite profiles, in order to increase berry and wine quality.

540

541

542

543 Effects of temperature and water supply on berry skin metabolic profile

544

545 Along the latitudinal gradient studied, the effect of temperature on overall phenolic
546 composition of Pinot Noir berry skins was weaker than the effect of radiation, because
547 temperature variables were correlated with phenolic composition only when they were
548 calculated for the period veraison-harvest. In this case, MSPCs, flavonols, flavanonols,
549 anthocyanins, and the ratios 3',4',5'-OH/4'-OH and 3',4',5'-OH/3',4'-OH flavonols,
550 were positively correlated with the mean daily temperature and/or degree days. These
551 correlations might be due to the fact that temperature and radiation variables were also
552 correlated for that period (Table S1), and it may be difficult to differentiate radiation
553 and temperature effects.⁴² It may not be surprising that the effects of temperature were
554 more clear in the most important period for berry maturity (veraison-harvest),²⁰
555 particularly in the case of anthocyanins, which increase strongly in that period.²⁰⁻²²
556 Anthocyanins are known to be influenced by specific temperature conditions, such as
557 ambient temperatures recorded after veraison.^{27,41,47,48} Results are also congruent for
558 flavonols because, although more influenced by radiation, these compounds can also
559 respond to temperature.⁴ Flavanols are known to be influenced by specific temperature
560 conditions, but in this study effects of a limited range of temperatures were tested, and it
561 is possible that more extreme temperatures are required to impact on these phenolics.
562 With respect to cinnamic acids, their synthesis in the first stages of berry development
563 and the strong decrease in concentrations after veraison²⁰ may mask the influence of
564 temperature on their content at harvest, thus concealing any correlation between
565 temperature parameters and cinnamic acid concentrations.

566

567 Rainfall and aridity indices showed a similar behavior as temperature variables, and
568 were correlated with some phenolic compounds only when the period veraison-harvest
569 was considered. In this period, water availability variables were correlated with
570 temperature and radiation variables, and thus the individual effect of each variable could
571 not be differentiated. Water availability typically shows strong relationships with
572 different plant traits,⁴⁹ but direct effects on the contents of grape skin phenolic
573 compounds are considered to be relatively minor.^{50,51} This could be due to the fact that
574 the effects of water availability on berry skin composition are mainly mediated by
575 changes in berry size which subsequently affect the proportion of skin in relation to
576 total berry, or by changes in photosynthesis rates modifying source-sink relationships.⁴²

577 Nevertheless, changes in anthocyanins, flavonols and stilbenes caused by water deficit
578 or excess have been described, sometimes in contradictory ways,^{42,52} and drought
579 conditions have been reported to increase the expression of different genes involved in
580 the biosynthesis of phenolic compounds.^{31,52} Overall, correlations between water
581 availability and phenolic composition were not conclusive in our study.

582

583

584 **In summary**

585

586 PCA was used to summarize the results described above. Axis I mostly represented a
587 latitude gradient, and was determined by nearly all different groups of phenolic
588 compounds that are present in berry skins (flavonols, flavanols, anthocyanins,
589 stilbenes and cinnamic acids, together with MSPCs). Thus, Pinot Noir berry skins from
590 southern localities were more enriched in most phenolics than those from northern
591 latitudes. This is congruent with the general variation in phenolic compounds (except
592 anthocyanins) with latitude.⁴ Changes in phenolic composition can influence wine
593 quality and will contribute to wine genuineness in each locality. Given that, in our
594 study, latitude was more often correlated with radiation variables than with temperature
595 or water availability variables, radiation appeared to be the most important factor
596 contributing to the differentiation of berry skin composition at the localities studied.
597 Nevertheless, in the most important period for phenolic ripeness (veraison-harvest),
598 latitude and radiation, temperature and water availability variables were correlated with
599 one another, and the effect of each type of variable was difficult to separate. Thus, apart
600 from the effect of radiation in every period considered, the interaction of radiation,
601 temperature and water availability in the period veraison-harvest was strongly correlated
602 with the phenolic composition of berry skins along the latitudinal gradient considered.
603 Flavanols and cinnamic acids were the only phenolic compounds that define axis II of
604 the PCA, thus contributing to the differentiation of berry skins from some localities, in
605 particular those situated to the positive part of the axis II, such as Bilje, Firenze, Retz,
606 Potoče and Jerez.

607

608 Genetic and environmental factors (other than radiation, temperature and water
609 availability) have not been considered in our study, but may also affect the metabolite
610 composition of berry skins. In particular, a clone effect cannot be excluded. However,

611 this effect has been demonstrated to be relatively minor and/or non-significant in
612 previous studies using both Pinot Noir^{48,53} and other grapevine cultivars.⁵⁴ On the other
613 hand, additional environmental factors related to the so-called “terroir” and not analyzed
614 in detail in our study, such as soil type or mineral nutrition, could have influenced
615 metabolites composition,^{54,55} although it is doubtful whether the impacts of such
616 variables would have been correlated with latitude. Overall, in spite of having used
617 different clones, plant ages and soils, a significant relationship between metabolites
618 composition and the latitude-dependent environmental changes in radiation, temperature
619 and water availability was found. It is likely that this environmental influence masked
620 the possible effects of genetic factors and other non-considered environmental variables.

621

622 Particularly relevant is the finding that skin phenolic composition was correlated with
623 the DSSF and T UVery means and doses in relatively short development periods (5-10
624 days before veraison and harvest). Thus, increasing the total and/or UV radiation
625 received by the clusters in those periods through management practices, such as leaf
626 removal or supplemental UV exposure, could promote the synthesis of valuable
627 phenolic metabolites. This may eventually contribute to improved wine quality because
628 of the notable contribution of phenolic compounds to wine flavor and also by increasing
629 the amount of nutraceuticals and healthy antioxidants, such as flavonols, flavanols,
630 stilbenes and cinnamic acids.^{10,11} Among others, UV radiation has been demonstrated to
631 be an important factor correlated with berry skin composition in our study. Although
632 some of the effects observed, such as the increase in MSPCs, flavonols and cinnamic
633 acids, have been repeatedly attributed to UV (particularly UV-B) radiation,^{13,18,29,31}
634 more specific manipulative experiments are needed to prove the specific effects of this
635 fraction of solar radiation across the latitudinal gradient considered.

636

637 It is concluded that radiation in several development periods, and an interaction between
638 radiation, temperature and water availability in the period veraison-harvest, were the
639 environmental factors most correlated with the phenolic composition of Pinot Noir
640 berry skins along a latitudinal gradient in Europe. In addition, it was demonstrated that
641 effects of environmental variables may be different for different compounds and that
642 some compounds were more responsive (for example, flavonols) than others
643 (flavanols).

644

645 **ASSOCIATED CONTENT**

646

647 **Supporting information**

648

649 Table S1. Correlation coefficients among environmental-geographic and berry
650 variables. Significant correlations are indicated in different colours depending on the
651 significance level: purple, $p < 0.001$; fuchsia, $p < 0.01$; pink, $p < 0.05$. Bb, bud break; v,
652 veraison; h, harvest; see the remaining abbreviations in Table 2 and 3 legends.

653

654 **AUTHOR INFORMATION**

655

656 Corresponding Author

657 *(E.N.-O.) Phone: +34-941-941299755. Fax: +34-941-299721. E-mail:
658 encarnacion.nunez@unirioja.es.

659

660 **FUNDING**

661 ENO and JMA are grateful to the Ministerio de Economía y Competitividad of Spain
662 and FEDER funds (Project CGL2014-54127-P) for financial support. MADCA and GS
663 benefited from grants of the Universidad de La Rioja (Plan Propio 2014 and 2013,
664 respectively). AM and GJ were supported by the grant OTKA K101430 of the
665 Hungarian Scientific Research Foundation. This work was supported by COST
666 (European Cooperation in Science and Technology) Action FA0906 of the European
667 Union “UV-B radiation: a specific regulator of plant growth and food quality in a
668 changing climate”.

669

670 **NOTES**

671 The authors declare no competing financial interest.

672

673 **ACKNOWLEDGEMENTS**

674 This manuscript is the result of joint work that was part of the activities of COST
675 Action FA0906 “UV4growth” (http://www.cost.eu/COST_Actions/fa/FA0906) of the
676 European Union. We thank Enrique García-Escudero (ICVV, CSIC - Gobierno de La
677 Rioja – Universidad de La Rioja), M^a José Serrano and Belén Puertas (IFAPA-Centro

678 Rancho de la Merced, Junta de Andalucía) and Josep Trallero (Bodega Serrat de
679 Montsoriu) for support to collect grapes.
680

681 **REFERENCES**

682

683 (1) Núñez-Olivera, E.; Martínez-Abaigar, J.; Escudero, J. C. Adaptability of
684 leaves of *Cistus ladanifer* to widely varying environmental conditions. *Funct. Ecol.*
685 **1996**, *10*, 636-646.

686 (2) Llorens, L.; Peñuelas, J.; Beier, C.; Emmett, B.; Estiarte, M.; Tietema, A.
687 Effects of an Experimental Increase of Temperature and Drought on the Photosynthetic
688 Performance of Two Ericaceous Shrub Species Along a North–South European
689 Gradient. *Ecosystems* **2004**, *7*, 613-624.

690 (3) Martz, F.; Peltola, R.; Fontanay, S.; Duval, R. L. E.; Julkunen-Tiitto, R.; Stark,
691 S. Effect of Latitude and Altitude on the Terpenoid and Soluble Phenolic Composition
692 of Juniper (*Juniperus communis*) Needles and Evaluation of Their Antibacterial
693 Activity in the Boreal Zone. *J. Agr. Food Chem.* **2009**, *57*, 9575–9584.

694 (4) Jaakola, L.; Hohtola, A. Effect of latitude on flavonoid biosynthesis in plants.
695 *Plant Cell Environ.* **2010**, *33*, 1239-1247.

696 (5) Jansen, M. A. K.; Le Martret, B.; Koornneef, M. Variations in constitutive and
697 inducible UV-B tolerance; dissecting photosystem II protection in *Arabidopsis thaliana*
698 accessions. *Physiol. Plant.* **2010**, *138*, 22-34.

699 (6) De Frenne, P.; Brunet, J.; Shevtsova, A.; Kolb, A.; Graae, B. J.; Chabrierie, O.;
700 Cousins, S. A. O.; Decocq, G.; Schrijver, A. D.; Diekmann, M.; Gruwez, R.; Heinken,
701 T.; Hermy, M.; Nilsson, C.; Stanton, S.; Tack, W.; Willaert, J.; Verheyen, K.
702 Temperature effects on forest herbs assessed by warming and transplant experiments
703 along a latitudinal gradient. *Global Change Biol.* **2011**, *17*, 3240-3253.

704 (7) Comont, D.; Martinez-Abaigar, J.; Albert, A.; Aphalo, P.; Causton, D. R.;
705 Figueroa, F. L.; Gaberscik, A.; Llorens, L.; Hauser, M. T.; Jansen, M. A. K.; Kardefelt,
706 M.; De la Coba Luque, P.; Neubert, S.; Núñez-Olivera, E.; Olsen, J.; Robson, M.;
707 Schreiner, M.; Sommaruga, R.; Strid, A.; Torre, S.; Turunen, M.; Veljovic-Jovanovic,
708 S.; Verdaguer, D.; Vidovic, M.; Wagner, J.; Barbro Winkler, J.; Zipoli, G.; Gwynn-
709 Jones, D. UV-responses of *Lolium perenne* raised along a latitudinal gradient across
710 Europe: A filtration study. *Physiol. Plant.* **2012**, *145*, 604-618.

711 (8) Yang, B.; Zheng, J.; Laaksonen, O.; Tahvonen, R.; Kallio, H. Effects of
712 Latitude and Weather Conditions on Phenolic Compounds in Currant (*Ribes spp.*)
713 Cultivars. *J. Agr. Food Chem.* **2013**, *61*, 3517–3532.

- 714 (9) Kenny, G. J.; Shao, J. An assessment of a latitude-temperature index for
715 predicting climate suitability for grapes in Europe. *J. Hort. Sci.* **1992**, *67*, 239-246.
- 716 (10) Sun, T.; Chen, Q. Y.; Wu, L. J.; Yao, X. M.; Sun, X. J. Antitumor and
717 antimetastatic activities of grape skin polyphenols in a murine model of breast cancer.
718 *Food Chem. Toxicol.* **2012**, *50*, 3462–3467.
- 719 (11) Calabriso, N.; Scoditti, E.; Massaro, M.; Pellegrino, M.; Storelli, C.;
720 Ingrosso, I.; Giovinazzo, G.; Carluccio, M. A. Multiple anti-inflammatory and anti-
721 atherosclerotic properties of red wine polyphenolic extracts: differential role of
722 hydroxycinnamic acids, flavonols and stilbenes on endothelial inflammatory gene
723 expression. *Eur. J. Nutr.* **2016**, *55*, 477-489.
- 724 (12) Mattivi, F.; Guzzon, R.; Vrhovsek, U.; Stefanini, M.; Velasco, R. Metabolite
725 Profiling of Grape: Flavonols and Anthocyanins. *J. Agr. Food Chem.* **2006**, *54*, 7692-
726 7702.
- 727 (13) Berli, F. J.; Fanzone, M.; Piccoli, P.; Bottini, R. Solar UV-B and ABA Are
728 Involved in Phenol Metabolism of *Vitis vinifera* L. Increasing Biosynthesis of Berry
729 Skin Polyphenols. *J. Agr. Food Chem.* **2011**, *59*, 4874-4884.
- 730 (14) Sternad Lemut, M.; Sivilotti, P.; Franceschi, P.; Wehrens, R.; Vrhovsek, U.
731 Use of Metabolic Profiling To Study Grape Skin Polyphenol Behavior as a Result of
732 Canopy Microclimate Manipulation in a ‘Pinot noir’ Vineyard. *J. Agr. Food Chem.*
733 **2013**, *61*, 8976–8986.
- 734 (15) Anderson, K. *Which winegrape varieties are grown where? A Global*
735 *Empirical Picture*; University of Adelaide Press: Adelaide, **2013**.
- 736 (16) Clarke, O.; Rand, M. *Grapes & Wines: A Comprehensive Guide to Varieties*
737 *and Flavours*; Pavilion Books: London, **2015**.
- 738 (17) Rolle, L.; Segade, S. R.; Torchio, F.; Giacosa, S.; Cagnasso, E.; Marengo, F.;
739 Gerbi, V. Influence of Grape Density and Harvest Date on Changes in Phenolic
740 Composition, Phenol Extractability Indices, and Instrumental Texture Properties during
741 Ripening. *J. Agr. Food Chem.* **2011**, *59*, 8796-8805.
- 742 (18) Carbonell-Bejerano, P.; Diago, M. P.; Martínez-Abaigar, J.; Martínez-
743 Zapater, J. M.; Tardáguila, J.; Núñez-Olivera, E. Solar ultraviolet radiation is necessary
744 to enhance grapevine fruit ripening transcriptional and phenolic responses. *BMC Plant*
745 *Biol.* **2014**, *14*, 183-198.

- 746 (19) Allaart, M.; Van Weele, M.; Fortuin, P.; Kelder, H. An Empirical model to
747 predict the UV-index based on Solar Zenith Angle and Total Ozone. *Meteorol. Appl.*
748 **2004**, *11*, 59-65.
- 749 (20) Downey, M. O.; Harvey, J. S.; Robinson, S. P. Synthesis of flavonols and
750 expression of flavonol synthase genes in the developing grape berries of Shiraz and
751 Chardonnay (*Vitis vinifera* L.). *Aust. J. Grape Wine Res.* **2003**, *9*, 110-121.
- 752 (21) Bogs, J.; Ebadi, A.; McDavid, D.; Robinson, S. P. Identification of the
753 Flavonoid Hydroxylases from Grapevine and Their Regulation during Fruit
754 Development. *Plant Physiol.* **2006**, *140*, 279-291.
- 755 (22) Pastore, C.; Zenoni, S.; Fasoli, M.; Pezzotti, M.; Tornielli, G. B.; Filippetti, I.
756 Selective defoliation affects plant growth, fruit transcriptional ripening program and
757 flavonoid metabolism in grapevine. *BMC Plant Biol.* **2013**, *13*, 30-45.
- 758 (23) Del-Castillo-Alonso, M. A.; Diago, M. P.; Monforte, L.; Tardáguila, J.;
759 Martínez-Abaigar, J.; Núñez-Olivera, E. Effects of UV exclusion on the physiology and
760 phenolic composition of leaves and berries of *Vitis vinifera* cv. Graciano. *J. Sci. Food*
761 *Agr.* **2015**, *95*, 409-416.
- 762 (24) Heredia, A.; Peinado, I.; Rosa, E.; Andrés, A. Effect of osmotic pre-treatment
763 and microwave heating on lycopene degradation and isomerization in cherry tomato.
764 *Food Chem.* **2010**, *123*, 92-98.
- 765 (25) Re, R.; Pellegrini, N.; Proteggente, A.; Pannala, A.; Yang, M.; Rice-Evans,
766 C. Antioxidant activity applying an improved ABTS radical cation decolorization assay.
767 *Free Radic. Biol. Med.* **1999**, *26*, 1231-1237.
- 768 (26) Price, S. F.; Breen, P. J.; Valladao, M.; Watson, B. T. Cluster Sun Exposure
769 and Quercetin in Pinot noir Grapes and Wine. *Am. J. Enol. Vitic.* **1995**, *46*, 187-194.
- 770 (27) Spayd, S. E.; Tarara, J. M.; Mee, D. L.; Ferguson, J. C. Separation of
771 Sunlight and Temperature Effects on the Composition of *Vitis vinifera* cv. Merlot
772 Berries. *Am. J. Enol. Vitic.* **2002**, *53*, 171-182.
- 773 (28) Berli, F.; D'Angelo, J.; Cavagnaro, B.; Bottini, R.; Wuilloud, R.; Silva, M. F.
774 Phenolic composition in grape (*Vitis vinifera* L. cv. Malbec) ripened with different solar
775 UV-B radiation levels by capillary zone electrophoresis. *J. Agr. Food Chem.* **2008**, *56*,
776 2892-2898.
- 777 (29) Koyama, K.; Ikeda, H.; Poudel, P. R.; Goto-Yamamoto, N. Light quality
778 affects flavonoid biosynthesis in young berries of Cabernet Sauvignon grape.
779 *Phytochemistry* **2012**, *78*, 54-64.

- 780 (30) Downey, M. O.; Harvey, J. S.; Robinson, S. P. The effect of bunch shading
781 on berry development and flavonoid accumulation in Shiraz grapes. *Aust. J. Grape*
782 *Wine Res.* **2004**, *10*, 55-73.
- 783 (31) Martínez-Lüscher, J.; Sánchez-Díaz, M.; Delrot, S.; Aguirreolea, J.; Pascual,
784 I.; Gomès, E. Ultraviolet-B Radiation and Water Deficit Interact to Alter Flavonol and
785 Anthocyanin Profiles in Grapevine Berries through Transcriptomic Regulation. *Plant*
786 *Cell Physiol.* **2014**, *55*, 1925–1936.
- 787 (32) Liu, L. L.; Gregan, S.; Winefield, C.; Jordan, B. From UVR8 to flavonol
788 synthase: UV-B-induced gene expression in Sauvignon blanc grape berry. *Plant Cell*
789 *Environ.* **2015**, *38*, 905-919.
- 790 (33) Malacarne, G.; Costantini, L.; Colter, E.; Battilana, J.; Velasco, R.;
791 Vrhovsek, U.; Grando, M. S.; Moser, C. Regulation of flavonol content and
792 composition in (Syrah×Pinot Noir) mature grapes: integration of transcriptional
793 profiling and metabolic quantitative trait locus analyses. *J. Exp. Bot.* **2015**, *66*, 4441–
794 4453.
- 795 (34) Martínez-Lüscher, J.; Torres, N.; Hilbert, G.; Richard, T.; Sánchez-Díaz, M.;
796 Delrot, S.; Aguirreolea, J.; Pascual, I.; Gomès, E. Ultraviolet-B radiation modifies the
797 quantitative and qualitative profile of flavonoids and amino acids in grape berries.
798 *Phytochemistry* **2014**, *102*, 106-114.
- 799 (35) Ryan, K. G.; Swinny, E. E.; Markham, K. R.; Winefield, C. Flavonoid gene
800 expression and UV photoprotection in transgenic and mutant *Petunia* leaves.
801 *Phytochemistry* **2002**, *59*, 23-32.
- 802 (36) Sroka, Z. Antioxidative and antiradical properties of plant phenolics. *Z. Nat.*
803 *Sect. C J. Biosci.* **2005**, *60*, 833-843.
- 804 (37) Fanzone, M.; Zamora, F.; Jofré, V.; Assof, M.; Peña-Neira, Á. Phenolic
805 Composition of Malbec Grape Skins and Seeds from Valle de Uco (Mendoza,
806 Argentina) during Ripening. Effect of Cluster Thinning. *J. Agr. Food Chem.* **2011**, *59*,
807 6120–6136.
- 808 (38) Feng, H.; Yuan, F.; Skinkis, P. A.; Qian, M. C. Influence of cluster zone leaf
809 removal on Pinot noir grape chemical and volatile composition. *Food Chem.* **2015**, *173*,
810 414-423.
- 811 (39) Fornara, V.; Onelli, E.; Sparvoli, F.; Rossoni, M.; Aina, R.; Marino, G.;
812 Citterio, S. Localization of stilbene synthase in *Vitis vinifera* L. during berry
813 development. *Protoplasma* **2008**, *233*, 83–93.

- 814 (40) Fujita, A.; Goto-Yamamoto, N.; Aramaki, I.; Hashizume, K. Organ-Specific
815 Transcription of Putative Flavonol Synthase Genes of Grapevine and Effects of Plant
816 Hormones and Shading on Flavonol Biosynthesis in Grape Berry Skins. *Biosci.*
817 *Biotechnol. Biochem.* **2006**, *70*, 632-638.
- 818 (41) Azuma, A.; Yakushiji, H.; Koshita, Y.; Kobayashi, S. Flavonoid
819 biosynthesis-related genes in grape skin are differentially regulated by temperature and
820 light conditions. *Planta* **2012**, *236*, 1067-1080.
- 821 (42) Downey, M. O.; Dokoozlian, N. K.; Krstic, M. P. Cultural Practice and
822 Environmental Impacts on the Flavonoid Composition of Grapes and Wine: A Review
823 of Recent Research. *Am. J. Enol. Vitic.* **2006**, *57*, 257-268.
- 824 (43) Guan, L.; Dai, Z.; Wu, B. H.; Wu, J.; Merlin, I.; Hilbert, G.; Renaud, C.;
825 Gomes, E.; Edwards, E.; Li, S. H.; Delrot, S. Anthocyanin biosynthesis is differentially
826 regulated by light in the skin and flesh of white-fleshed and teinturier grape berries.
827 *Planta* **2016**, *243*, 23-41.
- 828 (44) Cortell, J. M.; Kennedy, J. A. Effect of Shading on Accumulation of
829 Flavonoid Compounds in (*Vitis vinifera* L.) Pinot Noir Fruit and Extraction in a Model
830 System. *J. Agr. Food Chem.* **2006**, *54*, 8510-8520.
- 831 (45) Hanlin, R. L.; Downey, M. O. Condensed Tannin Accumulation and
832 Composition in Skin of Shiraz and Cabernet Sauvignon Grapes during Berry
833 Development. *Am. J. Enol. Vitic.* **2009**, *60*, 13-23.
- 834 (46) Pastor del Rio, J. L.; Kennedy, J. A. Development of Proanthocyanidins in
835 *Vitis vinifera* L. cv. Pinot noir Grapes and Extraction into Wine. *Am. J. Enol. Vitic.*
836 **2006**, *57*, 125-132.
- 837 (47) Cohen, S. D.; Tarara, J. M.; Gambetta, G. A.; Matthews, M. A.; Kennedy, J.
838 A. Impact of diurnal temperature variation on grape berry development,
839 proanthocyanidin accumulation, and the expression of flavonoid pathway genes. *J. Exp.*
840 *Bot.* **2012**, *63*, 2655-2665.
- 841 (48) Nicholas, K. A.; Matthews, M. A.; Lobell, D. B.; Willits, N. H.; Field, C. B.
842 Effect of vineyard-scale climate variability on Pinot noir phenolic composition. *Agr.*
843 *Forest Meteorol.* **2011**, *151*, 1556-1567.
- 844 (49) Moles, A. T.; Perkins, S. E.; Laffan, S. W.; Flores-Moreno, H.; Awasthy, M.;
845 Tindall, M. L.; Sack, L.; Pitman, A.; Kattge, J.; Aarssen, L. W.; Anand, M.; Bahn, M.;
846 Blonder, B.; Cavender-Bares, J.; Hans, J.; Cornelissen, C.; Cornwell, W. K.; Díaz, S.;
847 Dickie, J. B.; Freschet, G. T.; Griffiths, J. G.; Gutierrez, A. G.; Hemmings, F. A.;

- 848 Hickler, T.; Hitchcock, T. D.; Keighery, M.; Kleyer, M.; Kurokawa, H.; Leishman, M.
849 R.; Liu, K.; Niinemets, U.; Onipchenko, V.; Onoda, Y.; Peñuelas, J.; Pillar, V. D.;
850 Reich, P. B.; Shiodera, S.; Siefert, A.; Sosinski Jr, E. E.; Soudzilovskaia, N. A.; Swaine,
851 E. K.; Swenson, N. G.; Van Bodegom, P. M.; Warman, L.; Weiher, E.; Wright, I. J.;
852 Zhang, H.; Zobel, M.; Bonser, S. P. Which is a better predictor of plant traits:
853 temperature or precipitation? *J. Veg. Sci.* **2014**, *25*, 1167–1180.
- 854 (50) Kennedy, J. A.; Matthews, M. A.; Waterhouse, A. L. Effect of Maturity and
855 Vine Water Status on Grape Skin and Wine Flavonoids. *Am. J. Enol. Vitic.* **2002**, *53*,
856 268-274.
- 857 (51) Cadot, Y.; Chevalier, M.; Barbeau, G. Evolution of the localisation and
858 composition of phenolics in grape skin between veraison and maturity in relation to
859 water availability and some climatic conditions. *J. Sci. Food Agr.* **2011**, *91*, 1963–1976.
- 860 (52) Kuhn, N.; Guan, L.; Dai, Z. W.; Wu, B. H.; Lauvergeat, V.; Gomès, E.; Li, S.
861 H.; Godoy, F.; Arce-Johnson, P.; Delrot, S. Berry ripening: recently heard through the
862 grapevine. *J. Exp. Bot.* **2014**, *65*, 4543–4559.
- 863 (53) Lee, J.; Skinkis, P. A. Oregon ‘Pinot noir’ grape anthocyanin enhancement
864 by early leaf removal. *Food Chem.* **2013**, *139*, 893-901.
- 865 (54) Van Leeuwen, C.; Friant, P.; Choné, X.; Tregoat, O.; Koundouras, S.;
866 Dubourdieu, D. Influence of Climate, Soil, and Cultivar on Terroir. *Am. J. Enol. Vitic.*
867 **2004**, *55*, 207-217.
- 868 (55) Schreiner, R. P.; Scagel, C. F.; Lee, J. N, P, and K Supply to Pinot noir
869 Grapevines: Impact on Berry Phenolics and Free Amino Acids. *Am. J. Enol. Vitic.* **2014**,
870 *65*, 43-49.

871 **FIGURE AND TABLE LEGENDS**

872

873 **Figure 1.** Geographic location of the 11 European sampling localities used in this study.

874 1, Jerez de la Frontera (Spain); 2, Girona (Spain); 3, La Rioja (Spain); 4, Firenze (Italy);

875 5, Bordeaux (France); 6, Bilje (Slovenia); 7, Potoče (Slovenia); 8, Pécs (Hungary); 9,

876 Retz (Austria); 10, Lednice (Czech Republic); 11, Geisenheim (Germany).

877

878 **Figure 2.** Regressions between selected berry variables, including carotenoids and the879 different groups of phenolic compounds, and latitude. Determination coefficients (R^2)880 and p values are shown.

881

882 **Figure 3.** Regressions between the ratio trihydroxylated / monohydroxylated flavonols

883 and selected radiation variables. DSSF, Downward Surface Shortwave Flux. T UVery,

884 TEMIS-derived erythemal UV. For both variables, the daily mean in the period bud

885 break-harvest, and the accumulated dose in the same period and in the 10 days before

886 harvest, were used for calculations. Determination coefficients (R^2) and p values are

887 shown.

888

889 **Figure 4.** Ordination, through Principal Components Analysis (PCA), of the 11

890 sampling localities used in this study, taking into account the total content of methanol-

891 soluble phenolic compounds (MSPC) and the total contents of the different groups of

892 phenolic compounds. Significant loading factors for the positive and negative parts of

893 each axis, together with their corresponding significance levels, are shown (***,

894 $p < 0.001$; **, $p < 0.01$; *, $p < 0.05$). Axis 1 is the horizontal one, and axis 2 is the vertical

895 one. Each mark on the axes represents 0.5 units.

896

897 **Table 1.** Geographic location (latitude, longitude and altitude) of the 11 European

898 sampling localities used in this study, together with the number of days from bud break

899 to harvest.

900

901 **Table 2.** Ranges of the environmental variables in the 11 European sampling localities

902 used in this study, together with the localities in which each extreme value was recorded

903 (between brackets). ETP, potential evapotranspiration. GGR, Ground-station Global

904 Radiation. DSSF, Downward Surface Shortwave Flux. T UVery, TEMIS-derived

905 erythematic UV. The different variables were calculated along three periods: bud break-
906 veraison (white background), bud break-harvest (light grey background) and veraison-
907 harvest (dark grey background). In addition, DSSF doses were calculated in the 10 days
908 before harvest, and T UVery (mean values and total doses) in different periods.

909

910 **Table 3.** Values (means \pm SE) of the variables analyzed in Pinot Noir berries in the 11
911 European sampling localities used in this study. MSPC, methanol-soluble phenolic
912 compounds. AUC, area under curve. TSS, total soluble solids.

	sampling site	country	latitude (°N)	longitude (°E)	altitude (m)	days from bud break to harvest
1	Jerez de la Frontera	Spain	36.7	-6.2	40	141
2	Girona	Spain	41.8	2.6	150	174
3	La Rioja	Spain	42.5	-2.3	342	175
4	Firenze	Italy	43.9	11.2	280	131
5	Bordeaux	France	44.8	-0.6	22	176
6	Bilje	Slovenia	45.9	13.6	70	143
7	Potoče	Slovenia	45.9	13.8	120	140
8	Pécs	Hungary	46.1	18.1	200	152
9	Retz	Austria	48.8	15.9	324	172
10	Lednice	Czech Republic	48.8	16.8	176	183
11	Geisenheim	Germany	50.0	8.0	95	170

Table 1. Geographic location (latitude, longitude and altitude) of the 11 European sampling localities used in this study, together with the number of days from bud break to harvest.

	min	max
mean daily temperature (°C)	16.4 (3)	21.2 (8)
mean daily temperature (°C)	16.6 (10)	21.1 (8)
mean daily temperature (°C)	13.1 (10)	24.4 (1)
degree days (°C)	936 (3)	1367 (8)
degree days (°C)	1197 (3)	1703 (8)
degree days (°C)	113 (10)	381 (2)
rainfall (mm)	155 (4)	439 (5)
rainfall (mm)	196 (4)	481 (5)
rainfall (mm)	0 (1)	103 (10)
rainfall/ETP	0.31 (4)	0.80 (5)
rainfall/ETP	0.28 (1)	0.82 (9)
rainfall/ETP	0 (1)	0.9 (9,10)
Gaussen Index	4.0 (4)	12.8 (5)
Gaussen Index	4.9 (4)	13.7 (5)
Gaussen Index	0 (1)	4.7 (10)
GGR (mean) (MJ m ⁻² d ⁻¹)	12.7 (9)	24.2 (1)
GGR (mean) (MJ m ⁻² d ⁻¹)	11.2 (9)	24.9 (1)
GGR (mean) (MJ m ⁻² d ⁻¹)	8.1 (9)	28.6 (1)
GGR (dose) (MJ m ⁻²)	1487 (9)	3035 (3)
GGR (dose) (MJ m ⁻²)	1939 (9)	3718 (2)
GGR (dose) (MJ m ⁻²)	370 (4)	759 (10)
DSSF (mean) (MJ m ⁻² d ⁻¹)	18.3 (11)	23.8 (1)
DSSF mean (MJ m ⁻² d ⁻¹)	15.9 (11)	24.5 (1)
DSSF mean (MJ m ⁻² d ⁻¹)	10.1 (11)	28.4 (1)
DSSF (dose) (MJ m ⁻²)	2201 (11)	2908 (2)
DSSF (dose) (MJ m ⁻²)	2684 (11)	3542 (2)
DSSF (dose) (MJ m ⁻²)	384 (4)	695 (10)
T UVery (mean) (kJ m ⁻² d ⁻¹)	3.0 (11)	3.8 (1)
T UVery (mean) (kJ m ⁻² d ⁻¹)	2.4 (11)	4.0 (1)
T UVery (mean) (kJ m ⁻² d ⁻¹)	1.5 (11)	4.8 (1)
T UVery (dose) (kJ m ⁻²)	254 (11)	483 (3)
T UVery (dose) (kJ m ⁻²)	329 (11)	570 (3)
T UVery (dose) (kJ m ⁻²)	49 (4)	114 (1)
DSSF (10-days-before-harvest dose) (MJ m ⁻²)	56.6 (11)	284 (1)
T UVery (5-days-before-veraison mean) (kJ m ⁻² d ⁻¹)	2.0 (10,11)	5.1 (1)
T UVery (10-days-before-veraison mean) (kJ m ⁻² d ⁻¹)	2.4 (10,11)	5.0 (1)
T UVery (5-days-before-veraison dose) (kJ m ⁻²)	9.9 (10)	25.3 (1)
T UVery (10-days-before-veraison dose) (kJ m ⁻²)	23.8 (10)	50.2 (1)
T UVery (10-days-before-harvest dose) (kJ m ⁻²)	6.9 (11)	47.4 (1)

Table 2. Ranges of the environmental variables in the 11 European sampling localities used in this study, together with the localities in which each extreme value was recorded (between brackets). ETP, potential evapotranspiration. GGR, Ground-station Global Radiation. DSSF, Downward Surface Shortwave Flux. T UVery, TEMIS-derived erythematic UV. The different variables were calculated along three periods: bud break-veraison (white background), bud break-harvest (light grey background) and veraison-harvest (dark grey background). In addition, DSSF doses were calculated in the 10 days before harvest, and T UVery (mean values and total doses) in different periods.

	Jerez	Girona	La Rioja	Firenze	Bordeaux	Bilje	Potoče	Pécs	Retz	Lednice	Geisenheim
total content of MSPC											
AUC ₂₈₀₋₃₁₅ mg ⁻¹ DW	39.1 ± 1.5	40.3 ± 1.2	31.0 ± 3.0	32.3 ± 0.7	22.2 ± 1.3	14.7 ± 0.2	13.2 ± 0.4	32.3 ± 0.2	32.1 ± 5.3	9.7 ± 0.1	24.3 ± 1.2
AUC ₂₈₀₋₄₀₀ mg ⁻¹ DW	71.2 ± 3.5	74.3 ± 3.0	54.5 ± 5.4	58.4 ± 1.7	41.0 ± 2.4	24.5 ± 0.1	22.7 ± 0.4	56.9 ± 0.6	56.1 ± 9.9	17.1 ± 0.4	40.7 ± 2.2
flavonols (µg g⁻¹ DW)											
myricetin	139 ± 20	153 ± 8	112 ± 24	234 ± 27	38.7 ± 5.6	7.3 ± 2.8	13.2 ± 3.4	74.1 ± 9.2	164 ± 31	2.5 ± 0.8	15.3 ± 1.8
myricetin-3- <i>O</i> -glucoside	1066 ± 137	1041 ± 62	864 ± 86	918 ± 112	487 ± 37	157 ± 17	277 ± 45	473 ± 38	535 ± 92	61.2 ± 16.1	272 ± 30
myricetin-3- <i>O</i> -glucuronide	391 ± 50	355 ± 54	183 ± 32	368 ± 21	117 ± 11	62.5 ± 6.8	86.1 ± 7.5	267 ± 23	68.5 ± 9.1	22.2 ± 6.0	47.4 ± 8.6
kaempferol-3- <i>O</i> -glucoside	177 ± 37	273 ± 61	78.5 ± 9.9	109 ± 30	106 ± 7	21.6 ± 5.0	43.9 ± 8.2	40.7 ± 5.2	145 ± 36	48.1 ± 20.9	106 ± 36
isorhamnetin 3- <i>O</i> -glucoside	319 ± 31	433 ± 49	324 ± 33	274 ± 25	252 ± 16	84.4 ± 8.2	109 ± 11	234 ± 6	252 ± 27	138 ± 39	283 ± 21
isorhamnetin 3- <i>O</i> -glucuronide	72.9 ± 8.1	92.2 ± 6.5	41.8 ± 3.3	79.5 ± 6.3	50.4 ± 3.4	22.3 ± 4.6	28.2 ± 1.3	66.5 ± 1.8	27.3 ± 5.2	77.0 ± 15.2	51.6 ± 5.3
syringetin 3- <i>O</i> -glucoside	171 ± 26	130 ± 15	139 ± 16	87.8 ± 12.2	132 ± 8	62.1 ± 3.7	68.5 ± 4.9	156 ± 5	66.3 ± 8.3	57.2 ± 10.7	106 ± 7
quercetin	4.3 ± 0.4	5.6 ± 0.7	3.9 ± 0.7	2.8 ± 0.3	7.3 ± 3.2	1.3 ± 0.2	1.3 ± 0.1	3.5 ± 0.3	5.8 ± 2.4	2.3 ± 0.3	3.4 ± 0.5
quercetin 3- <i>O</i> -glucoside	105 ± 12	160 ± 21	159 ± 26	133 ± 13	50.9 ± 2.7	17.7 ± 2.2	22.9 ± 3.6	92.9 ± 9.0	181 ± 26	27.7 ± 5.0	94.3 ± 10.8
quercetin 3- <i>O</i> -galactoside	240 ± 33	400 ± 68	174 ± 11	228 ± 32	187 ± 14	39.5 ± 9.0	51.2 ± 3.1	106 ± 3	133 ± 30	50.8 ± 9.3	120 ± 24
quercetin-3- <i>O</i> -glucopyranoside	1075 ± 100	1361 ± 122	849 ± 47	973 ± 90	825 ± 45	260 ± 47	447 ± 41	629 ± 19	599 ± 107	300 ± 51	622 ± 100
quercetin-3- <i>O</i> -arabinoside	24.9 ± 3.0	22.1 ± 2.3	16.6 ± 1.6	15.3 ± 2.0	17.8 ± 2.1	3.6 ± 1.1	10.9 ± 1.7	8.6 ± 1.4	10.7 ± 2.0	5.7 ± 1.0	13.0 ± 2.2
quercetin 3- <i>O</i> -glucuronide	2726 ± 177	3121 ± 128	1951 ± 103	3014 ± 108	2119 ± 89	995 ± 132	1211 ± 19	2900 ± 44	1430 ± 253	1454 ± 259	1656 ± 156
quercetin-3- <i>O</i> -rutinoside	272 ± 35	170 ± 23	76.4 ± 9.8	279 ± 22	114 ± 10	28.3 ± 5.3	51.4 ± 3.2	144 ± 3	107 ± 38	48.7 ± 13.5	57.1 ± 5.9
flavanols (µg g⁻¹ DW)											
catechin	126 ± 9	110 ± 8.7	111 ± 14	224 ± 19	81.9 ± 7.4	355 ± 25	188 ± 48	66.4 ± 1.8	162 ± 23	77.9 ± 5.7	102 ± 5
epicatechin	8.8 ± 1.3	5.1 ± 0.6	8.4 ± 0.7	13.3 ± 1.3	5.9 ± 0.7	7.2 ± 1.2	4.5 ± 0.6	3.3 ± 0.3	9.2 ± 1.0	1.8 ± 0.2	2.7 ± 0.1
procyanidin B1	331 ± 27	324 ± 35	266 ± 23	467 ± 40	208 ± 18	633 ± 40	384 ± 59	173 ± 7	323 ± 40	130 ± 6	168 ± 10
flavanonols (µg g⁻¹ DW)											
astilbin	715 ± 61	591 ± 68	629 ± 59	511 ± 40	568 ± 45	163 ± 12	265 ± 35	476 ± 17	493 ± 43	299 ± 58	257 ± 43
taxifolin-3- <i>O</i> -glucoside	429 ± 64	114 ± 14	194 ± 37	250 ± 19	168 ± 38	21.8 ± 8.4	75.0 ± 19.1	138 ± 11	141 ± 21	10.7 ± 2.2	27.2 ± 6.0
stilbenes (µg g⁻¹ DW)											
resveratrol	54.7 ± 6.7	123 ± 28	105 ± 29	34.1 ± 12.1	31.4 ± 5.1	21.7 ± 8.5	6.4 ± 1.4	41.4 ± 4.3	57.1 ± 19.2	11.8 ± 6.5	15.4 ± 0.9
resveratrol-3- <i>O</i> -glucoside	395 ± 62	805 ± 77	385 ± 52	117 ± 32	120 ± 27	53.9 ± 27.6	17.7 ± 5.5	243 ± 32	303 ± 19	2.2 ± 0.6	29.2 ± 8.3
cinnamic Acids (µg g⁻¹ DW)											
coumaroyl-tartaric acid	876 ± 142	221 ± 14	215 ± 37	1016 ± 143	208 ± 54	72.6 ± 32.0	89.4 ± 24.9	72.0 ± 50.1	824 ± 114	14.8 ± 9.5	48.7 ± 14.9
caffeoyl-tartaric acid	4943 ± 716	2101 ± 427	1763 ± 214	6195 ± 809	1870 ± 497	894 ± 282	1047 ± 244	1597 ± 296	5855 ± 967	144 ± 108	947 ± 315
feruloyl-tartaric acid	5.7 ± 0.4	5.1 ± 0.6	2.3 ± 0.2	5.9 ± 0.7	5.8 ± 0.8	3.6 ± 0.4	5.0 ± 0.8	5.7 ± 2.4	4.0 ± 0.3	1.8 ± 0.3	2.1 ± 0.4
anthocyanins (mg g⁻¹ DW)											
delphinidin-3- <i>O</i> -glucoside	1.7 ± 0.2	2.9 ± 0.3	3.0 ± 0.5	2.9 ± 0.2	0.8 ± 0.1	0.2 ± 0.0	0.2 ± 0.0	1.4 ± 0.0	3.7 ± 0.5	0.3 ± 0.0	2.6 ± 0.3
cyanidin-3- <i>O</i> -glucoside	0.9 ± 0.1	4.4 ± 0.1	1.6 ± 0.2	1.0 ± 0.3	0.9 ± 0.0	0.2 ± 0.0	0.1 ± 0.0	1.8 ± 0.3	1.7 ± 0.2	0.3 ± 0.0	1.5 ± 0.1
petunidin-3- <i>O</i> -glucoside	5.0 ± 0.9	6.4 ± 0.8	4.3 ± 0.0	5.7 ± 0.7	1.8 ± 0.3	0.7 ± 0.1	0.9 ± 0.1	2.7 ± 0.0	4.8 ± 2.0	1.0 ± 0.1	2.8 ± 0.2
peonidin-3- <i>O</i> -glucoside	14.9 ± 1.6	34.9 ± 1.0	20.9 ± 1.3	13.8 ± 1.4	11.7 ± 1.2	5.7 ± 0.8	3.2 ± 0.2	25.9 ± 0.3	16.7 ± 2.7	5.8 ± 0.8	8.1 ± 0.8
malvidin-3- <i>O</i> -glucoside	54.6 ± 1.1	61.5 ± 0.7	39.8 ± 3.6	44.2 ± 0.3	26.4 ± 3.6	12.3 ± 0.2	17.4 ± 0.1	39.8 ± 1.4	36.2 ± 3.9	13.3 ± 0.6	27.1 ± 0.1
other variables											
antioxidant capacity (µM TE g ⁻¹ DW)	8013 ± 942	8639 ± 408	8637 ± 216	9104 ± 212	5576 ± 654	4134 ± 308	5111 ± 600	6330 ± 730	8212 ± 902	3592 ± 685	8424 ± 595
lutein (µg g ⁻¹ DW)	66.2 ± 0.8	55.5 ± 5.2	67.7 ± 1.2	32.9 ± 1.6	32.3 ± 1.2	24.1 ± 1.0	31.8 ± 1.3	16.1 ± 1.3	48.4 ± 0.6	20.2 ± 1.6	52.0 ± 10.1
zeaxanthin (µg g ⁻¹ DW)	8.6 ± 0.4	8.4 ± 0.0	9.2 ± 0.7	3.7 ± 0.3	5.5 ± 0.4	3.7 ± 0.5	4.9 ± 0.3	2.1 ± 0.0	6.7 ± 0.4	2.6 ± 0.1	9.2 ± 0.4
β-carotene (µg g ⁻¹ DW)	171 ± 7	165 ± 6	195 ± 2	96.4 ± 3.8	112 ± 11	83.1 ± 3.5	68.8 ± 7.6	56.7 ± 5.0	129 ± 9	66.8 ± 4.8	148 ± 19

chlorophylls (<i>a+b</i>) ($\mu\text{g g}^{-1}$ DW)	438 \pm 22	424 \pm 44	525 \pm 14	227 \pm 6	290 \pm 32	188 \pm 16	182 \pm 9	117 \pm 10	360 \pm 16	135 \pm 5	480 \pm 51
fresh weight per berry (g)	1.4 \pm 0.2	1.1 \pm 0.1	1.3 \pm 0.0	1.3 \pm 0.1	1.1 \pm 0.1	1.2 \pm 0.2	1.2 \pm 0.0	1.4 \pm 0.1	1.7 \pm 0.1	1.5 \pm 0.1	2.1 \pm 0.0
TSS ($^{\circ}$ Brix)	23.7 \pm 0.3	20.4 \pm 0.4	22.3 \pm 0.3	21.3 \pm 0.0	21.1 \pm 0.4	19.1 \pm 0.1	20.1 \pm 0.5	19.5 \pm 2.0	23.2 \pm 0.4	20.9 \pm 0.1	22.0 \pm 0.2

Table 3. Values (means \pm SE) of the variables analyzed in Pinot Noir berries in the 11 European sampling localities used in this study. MSPC, methanol-soluble phenolic compounds. AUC, area under curve. TSS, total soluble solids.




Figure 1. Geographic location of the 11 European sampling localities used in this study. 1, Jerez de la Frontera (Spain); 2, Girona (Spain); 3, La Rioja (Spain); 4, Firenze (Italy); 5, Bordeaux (France); 6, Bilje (Slovenia); 7, Potoče (Slovenia); 8, Pécs (Hungary); 9, Retz (Austria); 10, Lednice (Czech Republic); 11, Geisenheim (Germany).




Figure 2. Regressions between selected berry variables, including carotenoids and the different groups of phenolic compounds, and latitude. Determination coefficients (R^2) and p values are shown.




Figure 3. Regressions between the ratio trihydroxylated / monohydroxylated flavonols and selected radiation variables. DSSF, Downward Surface Shortwave Flux. T UVery, TEMIS-derived erythemal UV. For both variables, the daily mean in the period budbreak-harvest, and the accumulated dose in the same period and in the 10 days before harvest, were used for calculations. Determination coefficients (R^2) and p values are shown.




Figure 4. Ordination, through Principal Components Analysis (PCA), of the 11 sampling localities used in this study, taking into account the total content of methanol-soluble phenolic compounds (MSPC) and the total concentrations of the different groups of phenolic compounds. Significant loading factors for the positive and negative parts of each axis, together with their corresponding significance levels, are shown (***, $p < 0.001$; **, $p < 0.01$; *, $p < 0.05$). Axis 1 is the horizontal one, and axis 2 is the vertical one. Each mark on the axes represents 0.5 units.

GRAPHIC FOR TABLE OF CONTENTS

