

Title	Long-term photometric variations in the candidate white-dwarf pulsar AR Scorpii from K2 , CRTS, and ASAS-SN observations
Authors	Littlefield, Colin;Garnavich, Peter M.;Kennedy, Mark R.;Callanan, Paul J.;Shappee, Benjamin;Holoien, Thomas
Publication date	2017-08-10
Original Citation	Colin, L., Peter, G., Mark, K., Paul, C., Benjamin, S. and Thomas, H. (2017) 'Long-term Photometric Variations in the Candidate White-dwarf Pulsar AR Scorpii from K2 , CRTS, and ASAS-SN Observations', The Astrophysical Journal Letters, 845(1), pp. L7. doi: 10.3847/2041-8213/aa8300
Type of publication	Article (peer-reviewed)
Link to publisher's version	http://stacks.iop.org/2041-8205/845/i=1/a=L7 - 10.3847/2041-8213/aa8300
Rights	© 2017. The American Astronomical Society. All rights reserved.
Download date	2024-04-19 21:39:32
Item downloaded from	https://hdl.handle.net/10468/5443






UCC

University College Cork, Ireland
Coláiste na hOllscoile Corcaigh



Long-term Photometric Variations in the Candidate White-dwarf Pulsar AR Scorpii from *K2*, CRTS, and ASAS-SN Observations

Colin Littlefield¹ , Peter Garnavich¹ , Mark Kennedy^{1,2}, Paul Callanan², Benjamin Shappee³ , and Thomas Holoién⁴

¹Department of Physics, University of Notre Dame, Notre Dame, IN, USA

²Department of Physics, University College Cork, Cork, Ireland

³Carnegie Observatories, Pasadena, CA, USA

⁴Department of Astronomy, The Ohio State University, Columbus, OH, USA

Received 2017 June 30; revised 2017 July 27; accepted 2017 July 28; published 2017 August 10

Abstract

We analyze long-cadence *Kepler K2* observations of AR Sco from 2014, along with survey photometry obtained between 2005 and 2016 by the Catalina Real-Time Sky Survey and the All-Sky Automated Survey for Supernovae. The *K2* data show the orbital modulation to have been fairly stable during the 78 days of observations, but we detect aperiodic deviations from the average waveform with an amplitude of $\sim 2\%$ on a timescale of a few days. A comparison of the *K2* data with the survey photometry reveals that the orbital waveform gradually changed between 2005 and 2010, with the orbital maximum shifting to earlier phases. We compare these photometric variations with proposed models of this unusual system.

Key words: binaries: close – pulsars: general – stars: individual (AR Sco) – stars: magnetic field – white dwarfs

1. Introduction

AR Scorpii (AR Sco) is an unprecedented binary system featuring a white dwarf (WD) that generates highly periodic pulses across the electromagnetic spectrum every 1.97 minutes, even at radio wavelengths (Marsh et al. 2016). The system’s total luminosity exceeds the combined luminosity of the WD and its red dwarf companion by an average factor of ~ 4 , and its spectral energy distribution is consistent with synchrotron radiation, with the low X-ray luminosity implying minimal accretion (Marsh et al. 2016). Interpreting AR Sco as the first WD pulsar, Buckley et al. (2016) established that the optical pulses are highly linearly polarized and argued that the observed properties of AR Sco are consistent with a strongly magnetized, rapidly rotating WD whose spin-down powers the system’s luminosity. In this scenario, the WD’s magnetic axis is nearly perpendicular to its rotational axis, and the synchrotron radiation is produced when the WD’s open field lines sweep across the secondary, accelerating electrons in its wind (Geng et al. 2016). High-angular-resolution interferometric observations have shown the radio-emitting region to be smaller than ~ 0.02 au, implying the absence of a radio-bright outflow, such as a collimated jet (Marcote et al. 2017).

As an alternative to the WD-pulsar model, Katz (2017) proposed two hypotheses. In the first, the WD’s magnetic field sweeps over the face of the secondary, leading to the formation of a bow wave on the leading face of the secondary. Magnetic dissipation occurs preferentially in this bow wave, causing the observed orbital maximum at $\phi_{\text{orb}} \sim 0.4$, where superior conjunction of the secondary occurs at $\phi_{\text{orb}} = 0.5$. In Katz’s alternative hypothesis, the WD’s spin axis is misaligned with the binary’s orbital axis, and the WD’s magnetic moment is inclined with respect to its spin axis. As a result, the magnetic field experienced by the secondary varies with orbital phase, leading to a photometric modulation at the orbital period. In this second hypothesis, the misalignment causes a precession, and Katz predicts that the phase of maximum light drifts on timescales of ~ 20 – 200 years. This provides an observational test between the various models.

In the optical, the system exhibits two principal periodicities: a 3.56 hr orbital period, and the aforementioned 1.97 minute, double-peaked pulsation, the amplitude of which can be as large as ~ 1.5 mag, corresponding with the beat period between the WD spin and binary orbital periods. Curiously, the peak of the orbital modulation occurs at $\phi_{\text{orb}} \sim 0.4$ and is therefore offset from the time of superior conjunction at $\phi_{\text{orb}} = 0.5$, the phase at which maximum light from the irradiated inner hemisphere of the secondary would normally be observed.

We analyze 11 years of survey photometry of AR Sco, as well as *Kepler K2* observations, to investigate the long-term stability of the orbital modulation in order to provide additional constraints for theoretical explanations of the system.

2. Data and Analysis

2.1. *Kepler K2* Photometry

Before Marsh et al. (2016) uncovered AR Sco’s extraordinary nature, the *Kepler* satellite observed AR Sco in long-cadence mode as part of program GO2049 (PI: Andrej Prsa) during Campaign 2 of the *K2* mission between 2014 August 23 and 2014 November 10. Unfortunately, the 30 minute cadence of the *K2* data means that the 1.97 minute pulses are not temporally resolved, but the data nevertheless provide a unique opportunity to assess the stability of the orbital waveform across the 79 day *K2* run.

To extract the photometry, we downloaded the pixel file for AR Sco and performed aperture photometry on AR Sco’s centroid in each image. All data points affected by thruster firings were removed. When phased to the orbital ephemeris in Marsh et al. (2016), the *K2* observations show the orbital modulation to be remarkably consistent, with no obvious variations in the system’s overall brightness (Figure 1, left panel). The phase plot shows that the orbital modulation peaked near $\phi_{\text{orb}} \sim 0.4$, as observed in Marsh et al. (2016). The rise to maximum is steeper than the decline to minimum, with the latter showing a change in slope near $\phi_{\text{orb}} \sim 0.7$.

A power spectrum of the light curve (Figure 2) contains the fundamental orbital frequency and its next two harmonics. We




Figure 1. Left: a phase plot of the *K2* data, using the orbital ephemeris from Marsh et al. (2016). The green line is the best-fit polynomial. The data are repeated for clarity. Right: the full *K2* light curve after subtracting the polynomial model of the phase plot. The smoothed trendline highlights bumps near $T = \sim 15$ days and $T = \sim 75$ days.




Figure 2. A Lomb–Scargle power spectrum of the *K2* light curve, with the orbital frequency (Ω) and its first two harmonics labeled. There is no evidence of other periodicities within this frequency range. The highest frequency plotted is the Nyquist frequency.

did not search for frequencies above the Nyquist frequency for the data. The orbital period in the *K2* data is 0.148533(7) days, consistent with the value reported by Marsh et al. (2016). There is no evidence of additional periodicities in the power spectrum.

Based on the phase plot, we computed a polynomial that describes the orbital modulation as a function of orbital phase. We used this polynomial to subtract the orbital modulation from the full light curve to search for subtle variations in the overall brightness, and we plot these residuals in the right panel of Figure 1. There is a significant brightness variation approximately 15 days into the *K2* run, with an amplitude of 2%. Another possible rise is seen at the end of the run. The origin of these week-long brightness variations is unclear, but we viewed the images from the pixel file to ensure that they are

not artifacts caused by the passage of asteroids through the image.

2.2. CRTS and ASAS-SN Photometry

The excellent signal-to-noise ratio of the orbital modulation in the *K2* data enables us to test whether it is consistent with the orbital modulation in the Catalina Real-Time Sky Survey (CRTS; Drake et al. 2009) and the All-Sky Automated Survey for Supernovae (ASAS-SN; Shappee et al. 2014; Kochanek et al. 2017). Together, the photometry from these two surveys provides coverage of AR Sco from 2005 to 2016. Because of sparse sampling, we divide the CRTS photometry into a trio of three-year bins (2005–2007, 2008–2010, and 2011–2013) and the ASAS-SN data into three one-year bins (2014, 2015, and 2016); in none of the bins were CRTS and ASAS-SN data merged. We phased the data in each bin using the ephemeris from Marsh et al. (2016) and then fit the average *K2* waveform to each bin using χ^2 minimization to solve for two free parameters: a phase shift and a magnitude offset.

The resulting fits are shown in Figure 3. Although the three CRTS-only bins generally agree with the *K2* waveform, there is a striking trend: the shape of the rising part of the light curve changes with time. In the earliest data (2005–2007), a number of observations between $0.2 < \phi_{\text{orb}} < 0.4$ are significantly fainter than expected from the *K2* fit, some by as much as a half-magnitude. Although there are some points that are considerably brighter than the *K2* fit, these observations were probably contaminated by beat pulses, during which AR Sco can flare by a factor of four in $\lesssim 30$ s (Marsh et al. 2016). In the next bin (2008–2010), the rising part of the light curve is still fainter than the *K2* fit, but the discrepancy is not as pronounced. Finally, the most recent CRTS bin (2011–2013) shows excellent agreement with the *K2* light curve at all phases.

The *K2* fit to the ASAS-SN data underpredicts the amplitude of the orbital variation, but this is likely the result of a difference in bandpass. The Marsh et al. (2016) spectrum




Figure 3. CRTS and ASAS-SN data with the *K2* waveform superimposed in green. The *K2* waveform (green line) was fitted to each bin by applying vertical and horizontal translations to minimize χ^2 . In the earliest CRTS bin, there is significant deviation from the *K2* modulation, and the rising part of the light curve is fainter than in subsequent bins. The bright outliers in the CRTS data are consistent with beat pulses. The “CV” bandpass denotes unfiltered data with a *V* zeropoint. A bandpass difference between the unfiltered *K2* data and the *V*-band ASAS-SN photometry likely accounts for the underprediction of the amplitude of the ASAS-SN orbital modulation.

shows a strong contribution from the M-dwarf from $\sim 700\text{--}900$ nm. While an unfiltered CCD is sensitive at these wavelengths, the *V*-band has negligible transmission redward of 700 nm. As a result, the contrast between the heated inner hemisphere and the presumably unheated outer hemisphere will be greater in the *V*-band ASAS-SN data than in the unfiltered *K2* and CRTS data, giving rise to a larger orbital modulation at shorter wavelengths.

2.3. Phase of the Orbital Maximum

We fit third-order polynomials to the orbital maximum in each of the bins to estimate the phase of maximum light. The initial fits to the data used a conventional least-squares-minimization approach, but the quality of these fits is adversely affected by a number of bright outliers between orbital phases $\sim 0.1\text{--}0.4$. Marsh et al. (2016) showed that the amplitude of the

beat pulsations is largest at these orbital phases, so observations contaminated by beat pulses will drag the maximum of a conventional least-squares polynomial toward earlier phases. Thus, we also used random-sample consensus (RANSAC; Fischler & Bolles 1981), a machine-learning algorithm that identifies and masks outliers in a data set, to fit a robust polynomial to each bin. RANSAC iteratively selects random subsets of a data set, fitting them with a specified model (in this case, a third-order polynomial). The model describing the subset is compared to all points not in the subset, and if a certain number of them agree with the model to within a specified tolerance, they are considered inliers, as are the points in the original subset. The model is then fitted to the set of inliers to produce the provisional best-fit model; outliers are ignored. RANSAC repeats this process with different subsets, attempting to find the provisional best-fit model with the lowest residuals.




Figure 4. Left: the phase of the orbital maximum in each bin, as determined by fitting outlier-resistant polynomials to the phase plots in Figure 3. Right: the CRTS bin showing the largest phase shift (2005–2007), with least-squares and robust polynomials superimposed. The pure least-squares fit is skewed by a handful of bright outliers from beat pulses, causing it to peak at an earlier phase than the robust polynomial. The RANSAC algorithm used to fit the robust polynomial identifies these outliers and excludes them from the fit.

Table 1
Phase of the Orbital Maximum

Source	Band ^a	ϕ_{\max} (Robust)	Sig. ^b	ϕ_{\max} (Least Sq.)	Sig. ^c	Peak mag. ^d
CRTS (2005–2007)	CV	$0.505^{+0.031}_{-0.028}$	0.000	$0.460^{+0.022}_{-0.022}$	0.000	14.32 ± 0.07
CRTS (2008–2010)	CV	$0.415^{+0.028}_{-0.028}$	0.539	$0.395^{+0.022}_{-0.018}$	0.712	14.10 ± 0.07
CRTS (2011–2013)	CV	$0.396^{+0.030}_{-0.027}$	0.726	$0.398^{+0.020}_{-0.016}$	0.602	14.01 ± 0.09
ASAS-SN (2014)	V	$0.429^{+0.015}_{-0.015}$	0.022	$0.424^{+0.013}_{-0.014}$	0.012	14.49 ± 0.03
K2 (2014)	K_p	$0.418^{+0.004}_{-0.004}$	n/a	$0.418^{+0.003}_{-0.002}$	n/a	n/a
ASAS-SN (2015)	V	$0.404^{+0.011}_{-0.010}$	0.077	$0.401^{+0.009}_{-0.007}$	0.064	14.38 ± 0.03
ASAS-SN (2016)	V	$0.405^{+0.012}_{-0.011}$	0.164	$0.403^{+0.010}_{-0.008}$	0.158	14.41 ± 0.03

Notes.

^a CV = unfiltered with V zeropoint.

^b Fraction of simulations of robust fits in which the simulated phase shift was larger than the observed value.

^c Same as *b*, but for the least-squares fits.

^d Average maximum magnitude of the orbital modulation in robust fits.

The left panel in Figure 4 plots the phases of orbital maximum as measured by the robust polynomials, and to illustrate how the choice of fitting algorithm impacts the measured phase of orbital maximum, the right panel shows a comparison of the least-squares and RANSAC fits to the first CRTS bin. Additionally, Table 1 lists the phases of maximum light using the two fitting methods, along with 1σ uncertainties.

Both fitting techniques agree that there was a phase shift toward later phases in the 2005–2007 CRTS bin, but the robust polynomial shows it to be larger and more statistically significant. Although the contemporaneous K2 and 2014 ASAS-SN observations are weakly suggestive of a small phase shift, the data after the first CRTS bin are consistent with an essentially constant phase of orbital maximum, so any periodicity in the phase of maximum light must be much longer than the available baseline of data. In addition, as Table 1 indicates, the maximum of the orbital light curve was ~ 0.2 mag fainter in the earliest CRTS bin than in the other two CRTS bins, and the 2014 ASAS-SN bin was ~ 0.1 mag fainter than the 2015 and 2016 ASAS-SN data.

Because of a bandpass difference between CRTS and ASAS-SN, the peak magnitudes of the CRTS and ASAS-SN orbital modulation cannot be directly compared with each

other. It is for a similar reason that we did not attempt to measure the peak magnitude in the K2 data; a single *Kepler* magnitude cannot be reliably compared with magnitude estimates from other sources on the level of a tenth of a magnitude.

For the CRTS and ASAS-SN data, we used 2000 Monte Carlo simulations to test the susceptibility of the fitting procedure to false phase shifts induced by the combined effects of (1) sparse sampling, (2) propagated uncertainties from the orbital ephemeris, and (3) the difficulty of disentangling the orbital modulation from the beat pulses. The simulations made use of 22 hr of unfiltered photometry of AR Sco obtained at a typical cadence of ~ 5 s with the University of Notre Dame’s 80-cm Sarah L. Krizmanich Telescope (SLKT) during 2016 and 2017; at this cadence, the beat pulsations are reasonably well-sampled.⁵ The Monte Carlo procedure was as follows. Each simulation began by computing new orbital phases for all data, based on the uncertainties from the Marsh et al. (2016) orbital ephemeris. For each CRTS or ASAS-SN observation in a given bin, we found the SLKT observation with the most similar orbital phase. To simulate random sampling of the beat pulse, we

⁵ The SLKT data will be analyzed in a forthcoming paper.

then randomly selected an SLKT observation obtained within $\pm \frac{1}{2}P_{\text{beat}}$ of that point and calculated the average SLKT magnitude within a timespan equal to the exposure time of the survey photometry. Using this technique, each simulation created a unique synthetic light curve whose sampling and time resolution match that of the underlying survey photometry. Finally, each simulated light curve was fit with third-order least-squares and RANSAC polynomials. The phase of maximum light and the peak magnitude of the orbital modulation were extracted from the fits.

Since sparse sampling is not an issue with the *K2* photometry, we simply simulated Gaussian scatter in the flux for each *K2* observation as well as propagated uncertainties from the orbital ephemeris. While this results in comparatively small uncertainties, there is probably an unquantified systematic error stemming from the fact that the beat pulses are fully blended into the orbital modulation, thereby distorting the orbital profile. Each *K2* integration includes about 15 beat pulses, and because the beat pulses are strongest before the orbital maximum, their contamination will shift the orbital maximum toward earlier phases. The uncertainty for the *K2* phase of orbital maximum does not model this effect.

For both the least-squares and RANSAC fits, Table 1 lists the fraction of simulations for each bin in which the simulated phase shift was larger than the observed phase shift. The results suggest that the three aforementioned effects are insufficient to produce the observed phase shift in the earliest CRTS bin; not one of the simulated phase shifts for that bin was larger than the measured value. Additionally, the simulations suggest that the small phase shift in the 2014 ASAS-SN bin is of marginal significance, but given the comparatively small size of this phase shift relative to its 1σ uncertainty, it is possible that our simulations did not fully account for all possible causes of false phase shifts.

3. Conclusion

The *K2* observations from 2014 establish limits on the stability of the optical orbital modulations of AR Sco on timescales of months and show low-amplitude, apparently aperiodic fluctuations with an unknown source. Because it is

well-defined with very little scatter, the *K2* orbital modulation is a useful point of comparison for the orbital waveforms of the CRTS and ASAS-SN data sets, helping to establish that in early CRTS observations, the peak of the orbital waveform was considerably fainter than in subsequent years.

An analysis of the orbital phase of maximum light reveals a significant phase shift in the earliest CRTS data (2005–2007), but the data from 2008–2016 are consistent with an unchanged phase of orbital maximum. In the 2005–2007 CRTS data, the peak magnitude of the orbital modulation is about ~ 0.2 mag fainter than in the remaining CRTS bins. The apparent lack of a coherent trend in the phase of orbital maximum rules out sinusoidal variations on timescales of ~ 20 years, but with only 11 years of observations, there is insufficient data to discount the possibility of a longer periodicity. Katz’s misaligned-spin model predicts a precessional period of up to several centuries, so sustained long-term monitoring of the orbital modulation will be necessary to test this possibility.

We thank the referee for a thoughtful report that led to the improvement of this Letter.

ORCID iDs

Colin Littlefield  <https://orcid.org/0000-0001-7746-5795>
 Peter Garnavich  <https://orcid.org/0000-0003-4069-2817>
 Benjamin Shappee  <https://orcid.org/0000-0003-4631-1149>

References

- Buckley, D. A. H., Meintjes, P. J., Potter, S. B., Marsh, T. R., & Gänsicke, B. T. 2016, *NatAs*, 1, 29
 Drake, A. J., Djorgovski, S. G., Mahabal, A., et al. 2009, *ApJ*, 696, 870
 Fischler, M. A., & Bolles, R. C. 1981, *Commun. ACM*, 24, 381
 Geng, J.-J., Zhang, B., & Huang, Y.-F. 2016, *ApJL*, 831, L10
 Katz, J. I. 2017, *ApJ*, 835, 150
 Kochanek, C. S., Shappee, B. J., Stanek, K. Z., et al. 2017, *PASP*, submitted (arXiv:1706.07060)
 Marcote, B., Marsh, T. R., Stanway, E. R., Paragi, Z., & Blanchard, J. M. 2017, *A&A*, 601, L7
 Marsh, T. R., Gänsicke, B. T., Hümmelich, S., et al. 2016, *Natur*, 537, 374
 Shappee, B. J., Prieto, J. L., Grupe, D., et al. 2014, *ApJ*, 788, 48