

Title	Distributed optimization algorithm for discrete-time heterogeneous multi-agent systems with nonuniform stepsizes
Authors	Mo, L.;Li, J.;Huang, Jian
Publication date	2019-06-27
Original Citation	Mo, L., Li, J. and Huang, J. (2019) 'Distributed Optimization Algorithm for Discrete-Time Heterogeneous Multi-Agent Systems With Nonuniform Stepsizes', IEEE Access, 87303-87312. (7pp.) DOI: 10.1109/ACCESS.2019.2925414
Type of publication	Article (peer-reviewed)
Link to publisher's version	https://ieeexplore.ieee.org/document/8747511 - 10.1109/ ACCESS.2019.2925414
Rights	© The Author(s) 2019. This work is licensed under a Creative Commons Attribution 3.0 License. For more information, see http://creativecommons.org/licenses/by/3.0/ - http:// creativecommons.org/licenses/by/3.0/
Download date	2024-05-18 14:17:51
Item downloaded from	https://hdl.handle.net/10468/8623



University College Cork, Ireland Coláiste na hOllscoile Corcaigh



Received June 10, 2019, accepted June 20, 2019, date of publication June 27, 2019, date of current version July 17, 2019. Digital Object Identifier 10.1109/ACCESS.2019.2925414

Distributed Optimization Algorithm for Discrete-Time Heterogeneous Multi-Agent Systems With Nonuniform Stepsizes

LIPO MO^{®1}, JINGYI LI², AND JIAN HUANG^{®3} School of Mathematics and Statistics, Beijing Technology and Business University, Beijing 100048, China ²School of Mathematics and Systems Science, Beihang University, Beijing 100083, China ³School of Mathematical Sciences, University College Cork, Cork, T12 XF62 Ireland

Corresponding author: Lipo Mo (beihangmlp@126.com)

This work was supported in part by the Beijing Educational Committee Foundation under Grant KM201910011007 and Grant PXM2019_014213 _000007, in part by the Beijing Natural Science Foundation under Grant Z180005, and in part by the National Natural Science Foundation of China under Grant 61772063.

ABSTRACT This paper is devoted to the distributed optimization problem of heterogeneous multi-agent systems, where the communication topology is jointly strongly connected and the dynamics of each agent is the first-order or second-order integrator. A new distributed algorithm is first designed for each agent based on the local objective function and the local neighbors' information that each agent can access. By a model transformation, the original closed-loop system is converted into a time-varying system and the system matrix of which is a stochastic matrix at any time. Then, by the properties of the stochastic matrix, it is proven that all agents' position states can converge to the optimal solution of a team objective function provided the union communication topology is strongly connected. Finally, the simulation results are provided to verify the effectiveness of the distributed algorithm proposed in this paper.

INDEX TERMS Distributed optimization, multi-agent systems, heterogeneous, nonuniform stepsizes.

I. INTRODUCTION

Recently, distributed control theory has gained the fast progress due to the traditional centralized control methods, such as [1], [2], have been unable to meet the demands of control engineering. As a basic problem of distributed control, especially, consensus problems with or without constraints have received huge interests from many fields, such as systems and control, computer science and mathematics, and many excellent results have been reported. For example, consensus problems of first-order and second-order multiagent systems with communication time-delay were solve in [3]–[6]. Then, convex and nonconvex constrained consensus problems were considered for discrete-time and continuoustime multi-agent systems in [7]-[9], some effective distributed algorithms were designed and some consensus conditions were obtained.

As the further development of consensus, the distributed optimization problem has attracted wide attention due to its potential applications in signal processing, sensor networks, machine learning, distributed estimation, energy management problems in smart grids, event-triggered control problems, fast convergence problems and so on [10]-[20]. The main task of distributed optimization is to design proper distributed algorithm based on local information that each agent can access to force all agents cooperatively find the optimal solution of a team objective function. For firstorder multi-agent systems, the subgradient method was introduced to optimize a sum of convex functions [21], [22]. Then, these results were extended to continuous-time situations [23]-[25]. Meanwhile, the continuous-time zerogradient-sum and Newton-Raphson algorithms were proposed in [26], [27] to assure that all agents can find the optimal solution of a team objective function. Furthermore, under the balance condition, the distributed continuous-time optimization problem was solved and the corresponding convergence rate was also analyzed in [28], [29]. Taken the identical or nonidentical constraint sets into account, the distributed optimization problems were solved by a subgradient projection algorithm for general convex objective

The associate editor coordinating the review of this manuscript and approving it for publication was Bohui Wang.

functions [30], [31]. In addition, a distributed algorithm was introduced to solve the nonuniform unbounded constrained optimization problem in [32]. For second-order multi-agent systems, by introducing intermediate variables, the optimization problems were solved in [33], [34] when the objective functions are strongly convex. To reduce the cost of communication, some distributed algorithms with nonuniform gradient stepsizes were designed to minimize the general convex team objective function [35]–[37].

Most of the existing results on distributed optimization problem assumed that all agents' dynamics are the sameorder integrators. In reality, the different agents might have different dynamics due to the restriction of real environment, the corresponding system is always said to be heterogeneous systems. The consensus problems of heterogeneous multiagent systems with or without constraints were extensively studied in [38]–[42]. However, to the best of our knowledge, there is no works on the distributed optimization algorithm of heterogeneous multi-agent systems.

In this paper, we mainly study the distributed optimization problem of heterogeneous multi-agent system with jointly strongly connected communication topologies. The main contributions of this paper are as follows:

(1) In contrast to [21]–[24], [26]–[31], [33], [34], where some algorithms were proposed to solve the distributed optimization problems, but they assumed that the objective function is strictly convex or the gradient gains are constant, this paper deals with the situation of the general convex objective function and nonuniform gradient gains, which make our analysis be more complicated.

(2) In contrast to [25], [32], [35], [37], where the distributed optimization problems were studied for first-order or second-order multi-agent systems, this paper extends these results to heterogeneous multi-agent systems including first-order and second-order integrators simultaneously, which brings us more challenges and difficulties due to the inconsistency of each agent's dynamics.

(3) The proposed algorithms and analysis methods in [21]–[35], [37] are not applicable because we not only need to deal with first-order agents but also second-order agents.

(4) In contrast to [38]–[42], where only consensus problem was considered, this paper not only considers the consensus but also optimization index, which makes our closed-loop system be nonlinear essentially due to the existence of gradient terms and brings us more challenges in convergence analysis.

To overcome these difficulties, a novel distributed algorithm with nonuniform stepsizes (gradient gains) is first designed based on the local objective function and the local neighbors' information that each agent can access. By a coordination transformation, the original closed-loop system is changed into a new one, the system matrix of which is a stochastic matrix. Then, by the properties of stochastic matrix, we prove that all agents can reach an agreement on their position states and the team objective function is minimized at the same time. *Notations:* In this paper, \mathbf{R}^r represents Euclidean space with dimension r. ||x|| represents the Euclidean normal of a vector x. For a matrix A, A^T represents its transpose. I_r represents identical matrix with dimension r. \otimes represents kronecker product. **1** represent a vector with all of its entries being one.

II. PRELIMINARIES

Let $\mathcal{G}(k) = (\mathcal{I}_{m+n}, \mathcal{E}(k))$ be a directed graph at time k, where $\mathcal{I}_{m+n} = \{1, \dots, m+n\}$ is the set of nodes and $\mathcal{E}(k)$ is the set of edges at time k. $a_{ij}(k) > 0$ if and only if $(j, i) \in \mathcal{E}(k)$ and $a_{ii}(k) = 0$ for all i. Let $N_i(k) = \{j \in V : (j, i) \in \mathcal{E}(k)\}$ be the neighbor set of node i. The Laplacian L(k) of $\mathcal{G}(k)$ is defined as $[L(k)]_{ii} = \sum_{j=1}^{n} a_{ij}(k)$ and $[L(k)]_{ij} = -a_{ij}(k)$ for all $i \neq j$. The union graph of some graphs is a new graph whose node set is the same as each graph and edge set is the unions of the edge sets of these graphs. A series of edges $(v_{i_{m-1}}, v_{i_m}), (v_{i_{m-2}}, v_{i_{m-1}}), \dots, (v_{i_1}, v_{i_2})$ is called a directed path from node v_{i_1} to node v_{i_m} . A directed graph is said to be strongly connected if there exists at least one directed path between any two different nodes.

III. MAIN RESULTS

Consider a discrete-time heterogeneous multi-agent system consisting of m > 0 second-order agents and n > 0 firstorder agents. Let $\mathcal{G}(k)$ be the directed communication graph among all gents at time k and L(k) be its corresponding Laplacian. Let $L_s(k)$ and $L_f(k)$ be the Laplacian matrices of the graphs composed of second-order agents and first-order agents respectively. Thus, the Laplacian matrix of $\mathcal{G}(k)$ can be partitioned as

$$L(k) = \begin{bmatrix} L_s(k) + D_{sf}(k) & -A_{sf}(k) \\ -A_{fs}(k) & L_f(k) + D_{fs}(k) \end{bmatrix}$$

where $A_{sf}(k)$ and $A_{fs}(k)$ are the corresponding sub-blocks of L(k), $N_{i,f}(k)$ and $N_{i,s}(k)$ are the *i*th agent's first-order and second-order neighbor sets respectively, clearly, $N_i(k) =$ $N_{i,s}(k) \cup N_{i,f}(k)$,

$$D_{sf}(k) = diag\{\sum_{j \in N_{1,f}(k)} a_{1j}(k), \cdots, \sum_{j \in N_{m,f}(k)} a_{mj}(k)\},\$$
$$D_{fs}(k) = diag\{\sum_{j \in N_{m+1,s}(k)} a_{m+1,j}(k), \cdots, \sum_{j \in N_{m+n,s}(k)} a_{nj}(k)\}.$$

Suppose the dynamics of each second-order agent is

$$x_{i}(k+1) = x_{i}(k) + v_{i}(k)T$$

$$v_{i}(k+1) = v_{i}(k) + u_{i}(k)T, \quad i \in \mathcal{I}_{m}$$
(1)

where $x_i(k)$, $v_i(k)$, $u_i(k) \in \mathbf{R}^q$ are the position state, velocity state and control input of the *i*th agent at time *k* respectively; $\mathcal{I}_m = \{1, 2, \dots, m\}; T > 0$ is the sample time. Suppose the dynamics of each first-order agent is

$$x_i(k+1) = x_i(k) + u_i(k)T, \quad i \in \mathcal{I}_{m+n} - \mathcal{I}_m,$$
(2)

where $x_i(k), u_i(k) \in \mathbf{R}^q$ are the position state and control input of the *i*th agent at time k respectively.

The objective of this paper is to design a distributed algorithm based on local information each agent can access to drive all agents reach an agreement and minimize the following team objective function

$$f(s) = \sum_{i=1}^{m+n} f_i(s), \quad s \in \mathbf{R}^q,$$
(3)

where $f_i(s) : \mathbf{R}^q \to \mathbf{R}$ is the local differentiable convex objective function only accessed by the *i*th agent, $i \in \mathcal{I}_{m+n}$. To solve this problem, the following assumptions are necessary.

Assumption 1 [25]: Suppose $X_i := \{x \in \mathbb{R}^q | \nabla f_i(x) = 0\}, i \in \mathcal{I}_{m+n} \text{ are all nonempty and bounded.}$

Let X be the optimal solution set of the team objective function f(s), then all X_i and X are nonempty bounded closed convex sets (see [25] for details).

Assumption 2: There exists $\sigma \in (0, 1)$ such that $a_{ij}(k)T \ge \sigma$ and $a_{ij}(k) \le 1$ iff $a_{ij}(k) > 0$; $\sum_{i=1}^{m+n} a_{ij}(k) = \sum_{j=1}^{m+n} a_{ij}(k)$ for all $i, j \in \mathcal{I}_{m+n}$.

Assumption 3: Suppose $0 = k_0 < k_1 < \cdots$ is an infinite sequence of switching times and the union graph of all graphs over every interval $[k_m, k_{m+1})$ is strongly connected and $k_{m+1} - k_m \leq B$ for all positive integer *m*, where B > 0 is a positive constant.

Remark 1: Assumption 1 guarantees the distributed optimization problem is solvable, where the assumption is made only for the local objective functions and the global information (the team objective function) doesn't be used. Assumption 2 guarantees the interaction weights cannot be vanishing as time goes to infinity. Assumption 3 guarantees that each pair of agents can communicate directly or indirectly infinite many times and at least once in each *B* time interval.

In this paper, we design the following distributed algorithm with nonuniform stepsizes:

For $i \in \mathcal{I}_m$,

$$u_{i}(k) = -v_{i}(k) + \sum_{j \in N_{i,s}(k)} a_{ij}(k) [(x_{j}(k) - x_{i}(k)) + (v_{j}(k) - v_{i}(k))] + \sum_{j \in N_{i,f}(k)} a_{ij}(k) [(x_{j}(k) - x_{i}(k)) - v_{i}(k)] - d_{i}(k),$$
(4)

where

$$d_i(k) = \begin{cases} 0, & \sqrt{p_i(k)} \le \|\nabla f_i(w_i^{sy}(k))\|^2 \\ \frac{\nabla f_i(w_i^{sy}(k))}{\sqrt{p_i(k)}}, & \text{otherwise,} \end{cases}$$

 $p_{i}(k + 1) = p_{i}(k) + \arctan(e^{||x_{i}(k)| + v_{i}(k)||})T, p_{i}(0) > 0, \text{ and},$ $w_{i}^{sy}(k) = x_{i}(k) + v_{i}(k) + \sum_{j \in N_{i,s}(k)} a_{ij}(k)[(x_{j}(k) - x_{i}(k)) + (v_{j}(k) - v_{i}(k))]T + \sum_{j \in N_{i,f}(k)} a_{ij}(k)[(x_{j}(k) - x_{i}(k)) - v_{i}(k)]T.$ For $i \in \mathcal{I}_{m+n} - \mathcal{I}_{m},$ $u_{i}(k) = \sum_{i \in N_{i,s}(k)} a_{ij}(k)[(x_{j}(k) - x_{i}(k)) + v_{j}(k)]$

+
$$\sum_{j \in N_{i,f}(k)} a_{ij}(k) [(x_j(k) - x_i(k))] - d_i(k),$$
 (5)

where

$$d_i(k) = \begin{cases} 0, & \sqrt{p_i(k)} \le \|\nabla f_i(w_i^{fx}(k))\|^2 \\ \frac{\nabla f_i(w_i^{fx}(k))}{\sqrt{p_i(k)}}, & \text{otherwise,} \end{cases}$$

 $p_i(k + 1) = p_i(k) + \arctan(e^{\|x_i(k)\|})T, \ p_i(0) > 0, \text{ and},$ $w_i^{fs}(k) = x_i(k) + \sum_{j \in N_{i,s}(k)} a_{ij}(k)[(x_j(k) - x_i(k)) + v_j(k)]T + \sum_{j \in N_{i,f}(k)} a_{ij}(k)[(x_j(k) - x_i(k))]T.$

Remark 2: In the above algorithms, only the local information (the state information of its neighbors) is used by each agent, i.e., the corresponding data set includes all the state information of its neighbors.

Let $y_i(k) = x_i(k) + v_i(k), i \in \mathcal{I}_m$, then for $i \in \mathcal{I}_m$ and $j \in \mathcal{I}_{m+n} - \mathcal{I}_m$,

$$x_{i}(k+1) = (1-T)x_{i}(k) + Ty_{i}(k)$$

$$y_{i}(k+1) = w_{i}^{sy}(k) - d_{i}(k)T$$

$$x_{j}(k+1) = w_{j}^{fx}(k) - d_{j}(k)T.$$
(6)

Define $Z(k) = [x^{s}(k)^{T}, y^{s}(k)^{T}, x^{f}(k)^{T}]^{T}, x^{s}(k) = [x_{1}(k)^{T}, \cdots, x_{m}(k)^{T}]^{T}, y^{s}(k) = [y_{1}(k)^{T}, \cdots, y_{m}(k)^{T}]^{T}, x^{f}(k) = [x_{m+1}(k)^{T}, \cdots, x_{m+n}(k)^{T}]^{T}, D(k) = [0, D^{s}(k)^{T}, D^{f}(k)^{T}], D^{s}(k) = [d_{1}(k)^{T}, \cdots, d_{m}(k)^{T}]^{T}, D^{f}(k) = [d_{m+1}(k)^{T}, \cdots, d_{m+n}(k)^{T}]^{T}.$ Then the closed-loop system (6) can be changed into the following form:

$$Z(k+1) = [\Phi(k) \otimes I_q]Z(k) + D(k), \tag{7}$$

where

2

$$\Phi(k) = \begin{bmatrix} (1-T)I_m & TI_m & 0\\ 0 & \phi_1(k) & A_{sf}(k)T\\ 0 & A_{fs}(k)T & \phi_2(k), \end{bmatrix},$$

and $\phi_1(k) = I_m - L_s(k)T - D_{sf}(k)T$ and $\phi_2(k) = I_n - L_f(k)T - D_{fs}(k)T$.

Lemma 1: Under Assumption 2, if $T < \frac{1-\sigma}{n+m}$, then $\Phi(k)$ is a stochastic matrix and $\Phi_1(k)$ is a double stochastic matrix for all *k* with $[\Phi_1(k)]_{ij} \ge \sigma$ for all nonzero entry $[\Phi_1(k)]_{ij}$, $i, j \in \mathcal{I}_{m+n}$, where $\Phi_1(k) = \begin{bmatrix} \phi_1(k) & A_{sf}(k)T \\ A_{fs}(k)T & \phi_2(k) \end{bmatrix}$, i.e., $\Phi(k)\mathbf{1} = \mathbf{1}$, $\Phi_1(k)\mathbf{1} = \mathbf{1}$ and $\Phi_1^T(k)\mathbf{1} = \mathbf{1}$.

Proof: From the definitions of $\Phi(k)$ and $\Phi_1(k)$, it is easy to verify the conclusion by simple calculation.

Theorem 1: Under Assumptions 1, 2 and 3. If $T < \frac{1-\sigma}{m+n}$, then all agents of systems (1) and (2) can reach an agreement on their position states and the team objective function is minimized under Algorithms (4) and (5).

Proof: First, we will prove that $x_i(k)$ and $y_j(k)$ are bounded for all $k \ge 0$, $i \in \mathcal{I}_{m+n}$ and $j \in \mathcal{I}_m$. It follows from $\frac{\pi}{4} \le \arctan(e^{||y_i(k)||}) \le \frac{\pi}{2}$ and $\frac{\pi}{4} \le \arctan(e^{||x_i(k)||}) \le \frac{\pi}{2}$ that there exists $K_0 > 0$ such that $\frac{k\pi T}{4} \le p_i(k) \le k\pi T$ for any $k \ge K_0$. Note that X and X_i are all nonempty and bounded, we can choose a constant $\delta_1 > 0$ such that $X \subset$ $Y_1, X_i \subset Y_1$ for each *i*, where $Y_1 = \{s \in \mathbf{R}^q | ||s - z|| \le \delta_1\}$ for some $z \in X$, $f_i(w_i^{sy}(k)) - f_i(z) \ge 2[f_i(z) - f_i(z_i)] + T$ for all $z \in X$, $z_i \in X_i$ and $w_i^{sy}(k) \notin Y_1$, $i \in \mathcal{I}_m$, and $f_i(w_i^{fx}(k)) - f_i(z) \ge 2[f_i(z) - f_i(z_i)] + T$ for all $z \in X$, $z_i \in X_i$ and $w_i^{fx}(k) \notin Y_1$, $i \in \mathcal{I}_{n+m} - \mathcal{I}_m$. Choose Lyapunov function $V_1(k) = \max_{i \in \mathcal{I}_{m+n}, j \in \mathcal{I}_m} \{ \|x_i(k) - z\|^2, \|y_j(k) - z\|^2 \}.$

For $i \in \mathcal{I}_m$, by the convexity, we have

$$\begin{aligned} \|x_i(k+1) - z\|^2 \\ &= \|(1-T)x_i(k) + Ty_i(k) - z\|^2 \\ &\leq (1-T)\|x_i(k) - z\|^2 + T\|y_i(k) - z\|^2 \leq V_1(k). \end{aligned}$$

When $d_i(k) = 0$, we have

$$\begin{split} \|y_i(k+1) - z\|^2 \\ &= \|w_i^{sy}(k) - d_i(k)T - z\|^2 = \|y_i(k) \\ &+ \sum_{j \in N_{i,s}(k)} a_{ij}(k)[y_j(k) - y_i(k)]T \\ &+ \sum_{j \in N_{i,f}(k)} a_{ij}(k)[x_j(k) - y_i(k)]T - z\|^2 \\ &\leq (1 - \sum_{j \in N_i(k)} a_{ij}(k)T)\|y_i(k) - z\|^2 \\ &+ \sum_{j \in N_{i,s}(k)} a_{ij}(k)T\|y_j(k) - z\|^2 \\ &+ \sum_{j \in N_{i,f}(k)} a_{ij}(k)T\|x_j(k) - z\|^2 \leq V_1(k). \end{split}$$

When $d_i(k) \neq 0$, $\sqrt{p_i(k)} > \|\nabla f_i(w_i^{sy}(k))\|^2$, we have

$$\begin{split} \|y_{i}(k+1)-z\|^{2} \\ &= \|w_{i}^{sy}(k)-d_{i}(k)T-z\|^{2} = \|w_{i}^{sy}(k)-z\|^{2} \\ &+ \|d_{i}(k)T\|^{2} - 2(w_{i}^{sy}(k)-z)^{T} \frac{\nabla f_{i}(w_{i}^{sy}(k))}{\sqrt{p_{i}(k)}}T \\ &\leq (1-\sum_{j\in N_{i}(k)}a_{ij}(k)T)\|y_{i}(k)-z\|^{2} \\ &+ \sum_{j\in N_{i,s}(k)}a_{ij}(k)T\|y_{j}(k)-z\|^{2} \\ &+ \sum_{j\in N_{i,f}(k)}a_{ij}(k)T\|x_{j}(k)-z\|^{2} + \frac{T^{2}}{\sqrt{p_{i}(k)}} \\ &- 2[f_{i}(w_{i}^{sy}(k))-f_{i}(z)]\frac{T}{\sqrt{p_{i}(k)}} \\ &\leq (1-\sum_{j\in N_{i}(k)}a_{ij}(k)T)\|y_{i}(k)-z\|^{2} \\ &+ \sum_{j\in N_{i,s}(k)}a_{ij}(k)T\|y_{j}(k)-z\|^{2} \\ &+ \sum_{j\in N_{i,j}(k)}a_{ij}(k)T\|x_{j}(k)-z\|^{2} + \frac{2T^{2}}{\sqrt{k\pi T}} \\ &+ \frac{4T}{\sqrt{k\pi T}}[f_{i}(z)-f_{i}(z_{i})] \\ &- 2[f_{i}(w_{i}^{sy}(k))-f_{i}(z_{i})]\frac{T}{\sqrt{k\pi T}}, \end{split}$$

where $z_i \in X_i$. If $w_i^{sy}(k) \in Y_1$, then

$$\|y_i(k+1) - z\|^2 \le \delta_1^2 + \frac{2T^2}{\sqrt{k\pi T}} + \frac{4T}{k\pi T} [f_i(z) - f_i(z_i)],$$

which is bounded.

If $w_i^{sy}(k) \notin Y_1$, then $f_i(w_i^{sy}(k)) - f_i(z) \ge 2[f_i(z) - f_i(z_i)] + T$. Thus, $||y_i(k+1) - z||^2 \le V_1(k)$. For $i \in \mathcal{I}_{m+n} - \mathcal{I}_m$, when $d_i(k) = 0$, we have

$$\begin{aligned} \|x_i(k+1) - z\|^2 &= \|w_i^{fx}(k) - z\|^2 \\ &\leq (1 - \sum_{j \in N_i(k)} a_{ij}(k)T) \|x_i(k) - z\|^2 \\ &+ \sum_{j \in N_{i,s}(k)} a_{ij}(k)T \|y_j(k) - z\|^2 \\ &+ \sum_{j \in N_{i,f}(k)} a_{ij}(k)T \|x_j(k) - z\|^2 \\ &\leq V_1(k). \end{aligned}$$

When $d_i(k) \neq 0$, $\sqrt{p_i(k)} > \|\nabla f_i(w_i^{fx}(k))\|^2$, and

$$\begin{split} \|x_i(k+1) - z\|^2 \\ &= \|w_i^{fx}(k) - d_i(k)T - z\|^2 \\ &\leq \|w_i^{fx}(k) - z\|^2 + \|d_i(k)T\|^2 \\ &- 2(w_i^{fx}(k) - z)^T \frac{\nabla f_i(w_i^{fx}(k))}{\sqrt{p_i(k)}}T \\ &\leq (1 - \sum_{j \in N_i(k)} a_{ij}(k)T)\|x_i(k) - z\|^2 \\ &+ \sum_{j \in N_{i,s}(k)} a_{ij}(k)T\|y_j(k) - z\|^2 \\ &+ \sum_{j \in N_{i,f}(k)} a_{ij}(k)T\|x_j(k) - z\|^2 + \frac{T^2}{\sqrt{p_i(k)}} \\ &- 2[f_i(w_i^{fx}(k)) - f_i(z)]\frac{T}{\sqrt{p_i(k)}} \\ &\leq (1 - \sum_{j \in N_{i,k}(k)} a_{ij}(k)T)\|x_i(k) - z\|^2 \\ &+ \sum_{j \in N_{i,s}(k)} a_{ij}(k)T\|y_j(k) - z\|^2 \\ &+ \sum_{j \in N_{i,s}(k)} a_{ij}(k)T\|x_j(k) - z\|^2 + \frac{2T^2}{\sqrt{k\pi T}} \\ &+ \frac{4T}{\sqrt{k\pi T}}[f_i(z) - f_i(z_i)] \\ &- 2[f_i(w_i^{fx}(k)) - f_i(z_i)]\frac{T}{\sqrt{k\pi T}}, \end{split}$$

where $z_i \in X_i$. If $w_i^{fx}(k) \in Y_1$, then

$$\|x_i(k+1) - z\|^2 \le \delta_1^2 + \frac{2T^2}{\sqrt{k\pi T}} + \frac{4T}{k\pi T} [f_i(z) - f_i(z_i)].$$

which is bounded.

If $w_i^{fx}(k) \notin Y_1$, then $f_i(w_i^{fx}(k)) - f_i(z) \ge 2[f_i(z) - f_i(z_i)] + T$. Thus, $||x_i(k+1) - z||^2 \le V_1(k)$.

From the preceding analysis, we can conclude that $V_1(k)$ is bounded. Hence, $x_i(k)$ and $y_j(k)$ are bounded for all $i \in \mathcal{I}_{m+n}$ and $j \in \mathcal{I}_m$. Note that $f_i(s), i \in \mathcal{I}_{m+n}$ are differentiable, we get that $\|\nabla f_i(w_i^{sy}(k))\|$ and $\|f_j(w_j^{fx}(k))\|$ are bounded for all $i \in \mathcal{I}_m$ and $j \in \mathcal{I}_{m+n} - \mathcal{I}_m$. Together with $\lim_{t\to\infty} p_i(k) = \infty$, there exists $K_1 > K_0$ such that $d_i(k) = \frac{\nabla f_i(w_i^{sy}(k))}{\sqrt{p_i(k)}}$ and $d_j(k) = \frac{\nabla f_j(w_j^{fx}(k))}{\sqrt{p_i(k)}}$ for all $i \in \mathcal{I}_m, j \in \mathcal{I}_{m+n} - \mathcal{I}_m$ and $k > K_1$.

 $\frac{\nabla f_j(w_j^{fx}(k))}{\sqrt{p_i(k)}} \text{ for all } i \in \mathcal{I}_m, j \in \mathcal{I}_{m+n} - \mathcal{I}_m \text{ and } k \ge K_1.$ Next, let us analyze the stability of the closed-loop system. Define $z^*(k) = \frac{1}{m+n} [\sum_{i=1}^m y_i(k) + \sum_{i=m+1}^{m+n} x_i(k)], k \ge K_1,$ we will prove that $\lim_{k \to \infty} \|Z(k) - \mathbf{1}_{2m+n} \otimes z^*(k)\| = 0.$ From (7), we have

$$z^{*}(k+1) = \frac{1}{m+n} \left[\sum_{i=1}^{m} y_{i}(k+1) + \sum_{i=m+1}^{m+n} x_{i}(k+1) \right]$$

$$= \frac{1}{m+n} \left[\sum_{i=1}^{m} y_{i}(k) + \sum_{i=m+1}^{m+n} x_{i}(k) - \sum_{i=1}^{m+n} d_{i}(k)T \right]$$

$$= z^{*}(k) - \frac{1}{m+n} \left[\sum_{i=1}^{m} \frac{\nabla f_{i}(w_{i}^{sy}(k))}{\sqrt{p_{i}(k)}} T + \sum_{i=m+1}^{m+n} \frac{\nabla f_{i}(w_{i}^{fx}(k))}{\sqrt{p_{i}(k)}} T \right].$$

Take the following Lyapunov function $V_2(k) = \sum_{i=1}^{m} \|y_i(k) - z^*(k)\|^2 + \sum_{i=m+1}^{m+n} \|x_i(k) - z^*(k)\|^2$. Then

$$\begin{split} &V_{2}(k+1) \\ &= \sum_{i=1}^{m} \|y_{i}(k+1) - z^{*}(k+1)\|^{2} \\ &+ \sum_{i=m+1}^{m+n} \|x_{i}(k+1) - z^{*}(k+1)\|^{2} \\ &= \sum_{i=1}^{m} \|w_{i}^{sy}(k) - d_{i}(k)T - z^{*}(k) \\ &+ \frac{1}{m+n} [\sum_{i=1}^{m} \frac{\nabla f_{i}(w_{i}^{sy}(k))}{\sqrt{p_{i}(k)}}T \\ &+ \sum_{i=m+1}^{m+n} \frac{\nabla f_{i}(w_{i}^{fx}(k))}{\sqrt{p_{i}(k)}}T]\|^{2} \\ &+ \sum_{i=m+1}^{m+n} \|w_{i}^{fx}(k) - d_{i}(k)T - z^{*}(k) \\ &+ \frac{1}{m+n} [\sum_{i=1}^{m} \frac{\nabla f_{i}(w_{i}^{sy}(k))}{\sqrt{p_{i}(k)}}T \\ &+ \sum_{i=m+1}^{m+n} \frac{\nabla f_{i}(w_{i}^{fx}(k))}{\sqrt{p_{i}(k)}}T]\|^{2} \\ &= \sum_{i=1}^{m} \|w_{i}^{sy}(k) - d_{i}(k)T - z^{*}(k)\|^{2} \\ &+ \frac{m}{(m+n)^{2}} \|\sum_{i=1}^{m} \frac{\nabla f_{i}(w_{i}^{fx}(k))}{\sqrt{p_{i}(k)}}T \|^{2} \\ &+ 2\sum_{i=m+1}^{m} (w_{i}^{sy}(k) - d_{i}(k)T - z^{*}(k))^{T} \frac{1}{m+n} \\ &\times [\sum_{i=1}^{m} \frac{\nabla f_{i}(w_{i}^{sy}(k))}{\sqrt{p_{i}(k)}}T + \sum_{i=m+1}^{m+n} \frac{\nabla f_{i}(w_{i}^{fx}(k))}{\sqrt{p_{i}(k)}}T] \\ &+ \sum_{i=m+1}^{m+n} \frac{\nabla f_{i}(w_{i}^{fx}(k))}{\sqrt{p_{i}(k)}}T + \sum_{i=m+1}^{m+n} \frac{\nabla f_{i}(w_{i}^{fx}(k))}{\sqrt{p_{i}(k)}}T \\ &+ \sum_{i=m+1}^{m+n} \frac{\nabla f_{i}(w_{i}^{fx}(k))}{\sqrt{p_{i}(k)}}T \|^{2} \\ &+ 2\sum_{i=m+1}^{m+n} (w_{i}^{fx}(k) - d_{i}(k)T - z^{*}(k))^{T} \frac{1}{m+n} \\ &\times [\sum_{i=m+1}^{m+n} \frac{\nabla f_{i}(w_{i}^{fx}(k))}{\sqrt{p_{i}(k)}}T \|^{2} \\ &+ 2\sum_{i=m+1}^{m+n} (w_{i}^{fx}(k) - d_{i}(k)T - z^{*}(k))^{T} \frac{1}{m+n} \\ &\times [\sum_{i=1}^{m} \frac{\nabla f_{i}(w_{i}^{fx}(k))}{\sqrt{p_{i}(k)}}T + \sum_{i=m+1}^{m+n} \frac{\nabla f_{i}(w_{i}^{fx}(k))}{\sqrt{p_{i}(k)}}T] \\ \end{split}$$

$$= \sum_{i=1}^{m} \|w_i^{sy}(k) - d_i(k)T - z^*(k)\|^2 + \sum_{i=m+1}^{m+n} \|w_i^{fx}(k) - d_i(k)T - z^*(k)\|^2 - \frac{1}{m+n} \|\sum_{i=1}^{m} \frac{\nabla f_i(w_i^{sy}(k))}{\sqrt{p_i(k)}}T + \sum_{i=m+1}^{m+n} \frac{\nabla f_i(w_i^{fx}(k))}{\sqrt{p_i(k)}}T\|^2.$$

Besides,

$$\begin{split} \sum_{i=1}^{m} \|w_i^{sy}(k) - d_i(k)T - z^*(k)\|^2 \\ &= \sum_{i=1}^{m} \|w_i^{sy}(k) - z^*(k)\|^2 + \sum_{i=1}^{m} \|d_i(k)T\|^2 \\ &- 2\sum_{i=1}^{m} (w_i^{sy}(k) - z^*(k))^T d_i(k)T \\ &\leq \sum_{i=1}^{m} \|w_i^{sy}(k) - z^*(k)\|^2 + \sum_{i=1}^{m} \|d_i(k)T\|^2 \\ &- 2\sum_{i=1}^{m} [f_i(w_i^{sy}(k)) - f_i(z^*(k))] \frac{T}{\sqrt{p_i(k)}}, \\ \sum_{i=m+1}^{m+n} \|w_i^{fx}(k) - d_i(k)T - z^*(k)\|^2 \\ &= \sum_{i=m+1}^{m+n} \|w_i^{fx}(k) - z^*(k)\|^2 + \sum_{i=m+1}^{m+n} \|d_i(k)T\|^2 \\ &- 2\sum_{i=m+1}^{m+n} (w_i^{fx}(k) - z^*(k))^T d_i(k)T \\ &\leq \sum_{i=m+1}^{m+n} \|w_i^{fx}(k) - z^*(k)\|^2 + \sum_{i=m+1}^{m+n} \|d_i(k)T\|^2 \\ &- 2\sum_{i=m+1}^{m+n} \|w_i^{fx}(k) - z^*(k)\|^2 + \sum_{i=m+1}^{m+n} \|d_i(k)T\|^2 \\ &\leq \sum_{i=m+1}^{m+n} \|w_i^{fx}(k) - z^*(k)\|^2 + \sum_{i=m+1}^{m+n} \|d_i(k)T\|^2 \\ &\leq \sum_{i=m+1}^{m+n} \|w_i^{fx}(k) - z^*(k)\|^2 + \sum_{i=m+1}^{m+n} \|d_i(k)T\|^2 \end{split}$$

Since $\frac{k\pi T}{4} \leq p_i(k) \leq k\pi T$, $\nabla f_i(w_i^{sy}(k))$ and $f_i(w_i^{fx}(k))$ are bounded for all $k \geq K_1$, there must exist $c_1, c_2 > 0$ such that $V_2(k+1) \leq \sum_{i=1}^m ||w_i^{sy}(k) - z^*(k)||^2 + \sum_{i=m+1}^{m+n} ||w_i^{fx}(k) - z^*(k)||^2 + \frac{c_1}{k} + \frac{c_2}{\sqrt{k}}$.

Note that

$$\sum_{i=1}^{m} \|w_i^{sy}(k) - z^*(k)\|^2 + \sum_{i=m+1}^{m+n} \|w_i^{fx}(k) - z^*(k)\|^2$$

= $\|\Phi_1(k) \otimes I_p[y^s(k)^T, x^f(k)^T]^T - \mathbf{1}z^*(k)\|^2$
= $\|\Phi_1(k) \otimes I_p([y^s(k)^T, x^f(k)^T]^T - \mathbf{1}z^*(k))\|^2$,

we have

$$V_2(k+1) - V_2(k) \le -\phi(k) + \frac{c_1}{k} + \frac{c_2}{\sqrt{k}}$$

where $\phi(k) = ([y^s(k)^T, x^f(k)^T]^T - \mathbf{1}z^*(k))^T[I - \Phi_1(k)^T \Phi_1(k) \otimes I_p]([y^s(k)^T, x^f(k)^T]^T - \mathbf{1}z^*(k)) \ge 0.$

Summing the above inequality from k_m to $k_{m+1} - 1$, we have

$$V_2(k_{m+1}-1) - V_2(k_m) \le -\sum_{k=k_m}^{k_{m+1}-1} \phi(k) + B(\frac{c_1}{k} + \frac{c_2}{\sqrt{k}}).$$

Since $\lim_{t\to\infty} B(\frac{c_1}{k} + \frac{c_2}{\sqrt{k}}) = 0$, for any $\epsilon_1 > 0$, there exists $K_2 > K_1$ such that $B(\frac{c_1}{k} + \frac{c_2}{\sqrt{k}}) < \epsilon_1$ for all $k > K_2$. If there exists $k \in [k_m, k_{m+1}), k_m > K_2$, such that $\phi(k) > K_2$.

If there exists $k \in [k_m, k_{m+1}^*)$, $k_m > K_2$, such that $\phi(k) > 2\epsilon_1$, then $V_2(k_{m+1}-1) - V_2(k_m) < -\epsilon_1$. If $\phi(k) \le 2\epsilon_1$ for all $k \in [k_m, k_{m+1})$, $k_m > K_2$, we need to estimate $V_2(k)$. From

Assumption 2, each nonzero entry of $\Phi_1(k)$ is no smaller than σ , hence,

$$\begin{split} &([y^{s}(k)^{T}, x^{f}(k)^{T}]^{T} - 1 \ z^{*}(k))^{T}[-I + \Phi_{1}(k)^{T} \Phi_{1}(k) \\ &\otimes I_{p}]([y^{s}(k)^{T}, x^{f}(k)^{T}]^{T} - 1 \ z^{*}(k)) \\ &\leq -[\sum_{i=1}^{m} \left(\sum_{j \in N_{i,s}(k)} \sigma^{2} \ T^{2} \|y_{j}(k) - y_{i}(k)\|^{2} \\ &+ \sum_{j \in N_{i,f}(k)} \sigma^{2} \ T^{2} \|x_{j}(k) - y_{i}(k)\|^{2} \right) \\ &+ \sum_{i=m+1}^{m+n} \left(\sum_{j \in N_{i,s}(k)} \sigma^{2} \ T^{2} \|y_{j}(k) - x_{i}(k)\|^{2} \\ &+ \sum_{j \in N_{i,f}(k)} \sigma^{2} \ T^{2} \|x_{j}(k) - x_{i}(k)\|^{2} \right)]. \end{split}$$

By similar calculation as [32], it is easy to see that there exists $c_3 > 0$ such that $||y_i(k) - z^*(k)|| \le c_3(\sqrt{\epsilon_1} + \epsilon_1)$ and $||x_j(k) - z^*(k)|| \le c_3(\sqrt{\epsilon_1} + \epsilon_1)$ for all $i \in \mathcal{I}_m, j \in \mathcal{I}_{n+m} - \mathcal{I}_m$. Hence, $V_2(k) \le c_3(\sqrt{\epsilon_1} + \epsilon_1)$ for all $k \in [k_m, k_{m+1}), k_m > K_2$ when $\phi(k) \le 2\epsilon_1$.

Summarizing the preceding two cases, for any $\epsilon_2 > 0$, there exists $K_3 > K_2$ such that $V_2(k_m) < \epsilon_2$ for all $k_m > K_3$, which implies that $\lim_{m\to\infty} V_2(k_m) < \epsilon_2$. According to the arbitrariness of ϵ_2 , we can conclude that $\lim_{m\to\infty} V_2(k_m) =$ 0. Note that $V_2(k+1) - V_2(k) \le -\phi(k) + \frac{c_1}{k} + \frac{c_2}{\sqrt{k}} \le \frac{c_1}{k} + \frac{c_2}{\sqrt{k}}$ for all $k > K_2$, we have $\lim_{k\to\infty} V_2(k)$ exists by Cauchy convergence theorem. Therefore,

$$\lim_{k \to \infty} V_2(k) = \lim_{m \to \infty} V_2(k_m) = 0,$$

i.e.,

$$\lim_{k \to \infty} \|y_i(k) - z^*(k)\| = 0,$$

$$\lim_{k \to \infty} \|x_j(k) - z^*(k)\| = 0,$$

for all $i \in \mathcal{I}_m, j \in \mathcal{I}_{m+n} - \mathcal{I}_m$.

Let $l_i(k) = \sum_{j \in N_{i,s}(k)} a_{ij}(k)[y_j(k) - y_i(k)]T + \sum_{j \in N_{i,f}(k)} [x_j(k) - y_i(i)]T - d_i(k)T$ for $i \in \mathcal{I}_m$. Then $\lim_{k \to \infty} l_i(k) = 0$, $|l_i(k)| < M$ and $v_i(k+1) = (1-T)v_i(k) + l_i(k)$, where M > 0 is a constant. For any $\epsilon > 0$, there exists K > 0 such that $(1 - T)^{K+1} < \frac{T}{M}\epsilon$. Hence,

$$\begin{split} \|v_i(k+1)\| &= \|(1-T)^{k+1}v_i(0) + \sum_{s=0}^k (1-T)^s l_i(k-s)\| \\ &\leq (1-T)^{k+1} \|v_i(0)\| + \sum_{s=0}^k (1-T)^s \|l_i(k-s)\| \\ &\leq (1-T)^{k+1} \|v_i(0)\| + \sum_{s=0}^K (1-T)^s \|l_i(k-s)\| \\ &+ M \sum_{s=K+1}^k (1-T)^s \\ &= (1-T)^{k+1} \|v_i(0)\| + \sum_{s=0}^K (1-T)^s \|l_i(k-s)\| \\ &+ \frac{M}{T} (1-T)^{K+1} (1-(1-T)^{k-K}). \end{split}$$

Hence, $\lim_{k\to\infty} \|v_i(k+1)\| \leq \frac{M}{T}(1-T)^{K+1} < \epsilon$. It follows from the arbitrariness of ϵ that $\lim_{k\to\infty} \|v_i(k)\| = 0$. Therefore, $\lim_{k\to\infty} \|x_i(k) - z^*(k)\| = 0$ for all $i \in \mathcal{I}_{m+n}$, i.e., all agents reach an agreement on their position state.

At last, let us prove all agents' states could converge to the optimal solution of the team objective function. Define $V_3(k) = ||z^*(k) - P_X(z^*(k))||^2, k > K_2$. Then

$$V_{3}(k+1) = \|z^{*}(k+1) - P_{X}(z^{*}(k+1))\|^{2}$$

$$= \|z^{*}(k+1) - P_{X}(z^{*}(k)) + P_{X}(z^{*}(k))$$

$$-P_{X}(z^{*}(k+1))\|^{2}$$

$$= \|z^{*}(k+1) - P_{X}(z^{*}(k))\|^{2} + \|P_{X}(z^{*}(k))$$

$$-P_{X}(z^{*}(k+1))\|^{2} + 2[z^{*}(k+1) - P_{X}(z^{*}(k))]^{T}$$

$$\times [P_{X}(z^{*}(k)) - P_{X}(z^{*}(k+1))].$$

Note that $[z^*(k + 1) - P_X(z^*(k))]^T [P_X(z^*(k)) - P_X(z^*(k + 1))] \le 0$ and $||P_X(z^*(k)) - P_X(z^*(k + 1))|| \le ||z^*(k) - z^*(k + 1)||$, we have

$$V_{3}(k+1) \leq \|z^{*}(k+1) - P_{X}(z^{*}(k))\|^{2} + \|z^{*}(k) - z^{*}(k+1)\|^{2}$$

$$= \|z^{*}(k+1) - z^{*}(k) + z^{*}(k) - P_{X}(z^{*}(k))\|^{2}$$

$$+ \|z^{*}(k) - z^{*}(k+1)\|^{2}$$

$$= \|z^{*}(k) - P_{X}(z^{*}(k))\|^{2} + 2\|z^{*}(k) - z^{*}(k+1)\|^{2}$$

$$+ 2(z^{*}(k) - P_{X}(z^{*}(k)))^{T}(z^{*}(k+1) - z^{*}(k)).$$

Since $y_i(k), x_j(k)$ are bounded for all $i \in \mathcal{I}_m, j \in \mathcal{I}_{m+n} - \mathcal{I}_m$, there must exist $\gamma > 0$ such that $||z^*(k+1) - z^*(k)||^2 =$ $||\frac{1}{m+n} [\sum_{i=1}^m \frac{\nabla f_i(w_i^{y_i}(k))}{\sqrt{p_i(k)}} + \sum_{i=m+1}^{m+n} \frac{\nabla f_i(w_i^{f_k}(k))}{\sqrt{p_i(k)}}]||^2 \leq \frac{\gamma}{k}$. Therefore,

$$\begin{split} V_{3}(k+1) \\ &\leq \|z^{*}(k) - P_{X}(z^{*}(k))\|^{2} + \frac{2\gamma}{k} \\ &- \frac{2}{m+n} \sum_{i=1}^{m} [z^{*}(k) - P_{X}(z^{*}(k))]^{T} \frac{\nabla f_{i}(w_{i}^{sy}(k))}{\sqrt{p_{i}(k)}} T \\ &- \frac{2}{m+n} \sum_{i=m+1}^{m+n} [z^{*}(k) - P_{X}(z^{*}(k))]^{T} \frac{\nabla f_{i}(w_{i}^{fx}(k))}{\sqrt{p_{i}(k)}} T \\ &= V_{3}(k+1) + \frac{2\gamma}{k} \\ &- \frac{2}{m+n} \sum_{i=1}^{m} [z^{*}(k) - w_{i}^{sy}(k)]^{T} \frac{\nabla f_{i}(w_{i}^{sy}(k))}{\sqrt{p_{i}(k)}} T \\ &- \frac{2}{m+n} \sum_{i=m+1}^{m+n} [z^{*}(k) - w_{i}^{fx}(k)]^{T} \frac{\nabla f_{i}(w_{i}^{fx}(k))}{\sqrt{p_{i}(k)}} T \\ &- \frac{2}{m+n} \sum_{i=m+1}^{m} [w_{i}^{sy}(k) - P_{X}(z^{*}(k))]^{T} \frac{\nabla f_{i}(w_{i}^{sy}(k))}{\sqrt{p_{i}(k)}} T \\ &- \frac{2}{m+n} \sum_{i=m+1}^{m} [w_{i}^{fx}(k) - P_{X}(z^{*}(k))]^{T} \frac{\nabla f_{i}(w_{i}^{fx}(k))}{\sqrt{p_{i}(k)}} T \end{split}$$

From the definitions of $w_i^{sy}(k)$ and $w_j^{fx}(k)$, it is easy to see that $\lim_{k\to\infty} ||z^*(k) - w_i^{sy}(k)|| = 0$ and $\lim_{k\to\infty} ||z^*(k) - w_j^{fx}(k)|| = 0$ for all $i \in \mathcal{I}_m$ and $j \in \mathcal{I}_{m+n} - \mathcal{I}_m$. Note that $\nabla f_i(w^{sy}(k))$, $\nabla f_j(w_j^{fx}(k))$ are bounded and $\frac{k\pi T}{4} \leq p_i(k) \leq k\pi T$, we can conclude that for any $\epsilon_3 > 0$, there exist $\gamma_1 > 0$ and $K_3 > K_2$ such that $-\frac{2}{m+n} \sum_{i=1}^m [z^*(k) - w_i^{sy}(k)]^T \frac{\nabla f_i(w_i^{sy}(k))}{\sqrt{p_i(k)}}T -$

$$\frac{2}{m+n} \sum_{i=m+1}^{m+n} [z^*(k) - w_i^{fx}(k)]^T \frac{\nabla f_i(w_i^{fx}(k))}{\sqrt{p_i(k)}} T - \frac{2}{m+n} \sum_{i=1}^m [f_i(w_i^{sy}(k)) - f_i(z^*(k))] \frac{T}{\sqrt{p_i(k)}} - \frac{2}{m+n} \sum_{i=m+1}^{m+n} [f_i(w_i^{fx}(k)) - f_i(z^*(k))] \frac{T}{\sqrt{p_i(k)}} \le \frac{\gamma_1 \epsilon_3}{\sqrt{k}} \text{ for all } k > K_3. \text{ Hence, we have}$$

$$\begin{split} V_{3}(k+1) - V_{3}(k) &\leq \frac{2\gamma}{k} + \frac{\gamma_{1}\epsilon_{3}}{\sqrt{k}} - \frac{2T}{(m+n)\sqrt{p_{1}(k)}} \\ &\times \sum_{i=1}^{m+n} [f_{i}(z^{*}(k)) - f_{i}(P_{X}(z^{*}(k)))] \\ &- \frac{2T}{(m+n)\sqrt{p_{1}(k)}} \sum_{i=1}^{m+n} (\frac{\sqrt{p_{1}(k)}}{\sqrt{p_{i}(k)}} - 1) \\ &\times [f_{i}(z^{*}(k)) - f_{i}(P_{X}(z^{*}(k)))]. \end{split}$$

For $i \in \mathcal{I}_m$, let $\pi_i(k) = p_i(0) - p_1(0) + \sum_{s=0}^k [\arctan(e^{\|y_i(s)\|}) - \arctan(e^{\|y_1(s)\|})]T$. Then $p_i(k) = p_1(k)[1 + \frac{\pi_i(k)}{p_1(k)}]$. Since $\lim_{k\to\infty} \|y_i(k) - y_1(k)\| = 0$, for any $\epsilon' > 0$, there exists $K' > K_0$ such that $|\arctan(e^{\|y_i(s)\|}) - \arctan(e^{\|y_i(s)\|})| < \epsilon'$ for all k > K'. Note that

$$\frac{\sum_{s=K_0+1}^{k} [\arctan(e^{\|y_i(s)\|}) - \arctan(e^{\|y_1(s)\|})]T}{p_1(k)} \le \frac{4\epsilon'}{\pi T},$$

we can get $\lim_{k\to\infty} \frac{\pi_i(k)}{p_1(k)} \leq \frac{4\epsilon'}{\pi T}$. According to the arbitrariness of ϵ' , we have $\lim_{k\to\infty} \frac{\pi_i(k)}{p_1(k)} = 0$ and $\lim_{k\to\infty} \frac{p_i(k)}{p_1(k)} = 1$. Similarly, we can prove that $\lim_{k\to\infty} \frac{p_i(k)}{p_1(k)} = 1$ for all $i \in \mathcal{I}_{m+n} - \mathcal{I}_m$. Hence, there exists $K_4 > K_3$ such that $-\frac{2T}{(m+n)\sqrt{p_1(k)}} \sum_{i=1}^{m+n} (\frac{\sqrt{p_1(k)}}{\sqrt{p_i(k)}} - 1)[f_i(z^*(k)) - f_i(P_X(z^*(k)))] \leq \frac{\epsilon_3}{\sqrt{k}}$ and $\frac{2\gamma}{k} < \frac{\epsilon_3}{\sqrt{k}}$ for all $k > K_4$. Thus, for $k > K_4$,

$$\begin{aligned}
& \sqrt{3}(k+1) - \sqrt{3}(k) \\
& \leq \frac{2\epsilon_3}{\sqrt{k}} + \frac{\gamma_1\epsilon_3}{\sqrt{k}} - \frac{2T}{m+n} \frac{1}{\sqrt{p_1(k)}} \\
& \times \left[\sum_{i=1}^{m+n} f_i(z^*(k)) - \sum_{i=1}^{m+n} f_i(P_X(z^*(k)))\right] \\
& \leq -\frac{2\sqrt{T}}{(m+n)\sqrt{\pi k}} \left[\sum_{i=1}^{m+n} f_i(z^*(k)) - \sum_{i=1}^{m+n} f_i(P_X(z^*(k))) - \gamma_2\epsilon_3\right],
\end{aligned}$$

where $\gamma_2 = \frac{(m+n)\sqrt{\pi}}{2\sqrt{T}}(2+\gamma_1) > 0$. Since $f_i(s)$ are differentiable convex function, we can

Since $f_i(s)$ are differentiable convex function, we can choose $c_1 > 0$ and $C_1 = \{s | \| s - P_x(s) \| \le c_1\}$ such that $\sum_{i=1}^{m+n} f_i(s_1) - \sum_{i=1}^{m+n} f_i(s_2) > \gamma_2 \epsilon_3 + 2\epsilon_4$ for all $s_1 \notin C_1$ and $s_2 \in X$, where $\epsilon_4 > 0$. Set $c_2 = c_1 + 2\epsilon_4$ and $C_2 = \{s | \| s - P_X(s) \| \le c_2\}$. Note that $\nabla f_i(w_i^{sy}(k))$ and $\nabla f_j(w_j^{fx}(k))$ are bounded, there exists $K_5 > K_4$ such that $\| z^*(k+1) - z^*(k) \| < \epsilon_4$ for all $k > K_5$. For $k > K_5$, if $z^*(k) \in C_1$, then $z^*(k+1) \in C_2$. If $z^*(k) \notin C_1$ and $z^*(k) \in C_2$, then $V_3(k + 1) - V_3(k) < 0$, which implies $z^*(k+1) \in C_2$. If $z^*(k) \notin C_2$ for $k > K_5$, then $V_3(k+1) - V_3(k) < -\frac{2\sqrt{T}\epsilon_4}{(m+n)\sqrt{\pi k}}$. So there exists $K_6 > K_5$ such that $z^*(k) \in C_2$, i.e., $\sum_{i=1}^{m+n} [f_i(z^*(k)) - f_i(P_X(z^*(k)))] < c_1 + 2\epsilon_4$ for all $k > K_6$. Letting $\epsilon_4 \to 0$, we can get $\lim_{k\to\infty} z^*(k) \in C_1$, which suggests $\lim_{k\to\infty} \sum_{i=1}^{m+n} [f_i(z^*(k)) - f_i(P_X(z^*(k)))] \le \gamma_2\epsilon_3$. By the arbitrariness of ϵ_3 , we conclude that $\lim_{k\to\infty} \sum_{i=1}^{m+n} [f_i(z^*(k)) - f_i(P_X(z^*(k)))] = 0$, which means all agents' states could



FIGURE 1. Communication graphs.



FIGURE 2. The first component of position states.

converge to the optimal solution of team objective function as k tends to infinity.

IV. SIMULATIONS

In this section, numerical examples are given to verify the effectiveness of the algorithm proposed in this paper. Suppose there are four second-order agents (1 to 4) and four first-order agents (5 to 8), and the communication topology switches among three graphs, each of them doesn't be strongly connected, see Fig. 1. The sample time is chosen as T = 0.001s. Take initial values $x_1(0) = (5, 5)^T$, $x_2(0) = (3, 2)^T$, $x_3(0) = (1, 1.5)^T$, $x_4(0) = (-2.5, -2)^T$, $x_5(0) = (4, 3)^T$, $x_6(0) = (-3, -2)^T$, $x_7(0) = (3, 1)^T$, $x_8(0) = (-5, -3)^T$, $v_i(0) = (0, 0)^T$, i = 1, 2, 3, 4.

A. EXAMPLE 1

The local objective functions are chosen as $f_1(s) = (s_1 - 2)^2 +$ $s_2^2, f_2(s) = s_1^4 + s_2^2, f_3(s) = s_1^2 + s_2^4, f_4(s) = s_1^2 + (s_2 - 2)^2,$ $f_5(s) = (s_1 - 2)^2 + s_2^2, f_6(s) = s_1^4 + s_2^2, f_7(s) = s_1^2 + (s_2 - 2)^2, f_8(s) = s_1^2 + s_2^4.$ where $s = (s_1, s_2)^T \in \mathbf{R}^2$. By simple computation, we can obtain that the optimal solution of the team objective function f(s) is (0.5536, 0.5536). The initial values of the nonuniform stepsizes (gradient gains) are chosen as $p_i(0) = 2, i = 1, 2, \dots, 8$. By taking algorithms (4) and (5), the position trajectories of all agents are shown in Fig. 2 and Fig. 3, from which we could see that the position states of all agents can reach an agreement and converge to the optimal solution of the team objective function with an extremely small error. But they must can converge to the optimal solution of the team objective function as time tends to infinity. Fig. 4 and Fig. 5 show the velocity states of all second-order agents, from which we could see that velocities of all second-order agents converge to zero asymptotically.



FIGURE 3. The second component of position states.



FIGURE 4. The first component of velocity states.



FIGURE 5. The second component of velocity states.

B. EXAMPLE 2

In this subsection, to show the effectiveness of the nonuniform gradient gains, we make a comparison with the algorithm that has uniform gradient gains. The gradient gains in algorithms (4) and (5) are chosen very large such as $p_i(k) = 5 * 10^5, i = 1, 2, \dots, 8$ for all k (If the gradient gains are chosen to be small, then the agreement cannot be reached). Then by taking algorithms (4) and (5), the position trajectories of all agents are shown in Fig. 6 and Fig. 7, from which we could see that the position states of all agents can reach an agreement and converge to the optimal solution of the team objective function. Moreover, the velocity states of all second-order agents are shown in Fig. 8 and Fig. 9, from which we could see that velocities of all second-order agents converge to zero. However, it can see from Fig. 6, 7, 8, 9 that the distributed optimization problem of heterogeneous multi-agent systems is solved after about 50 seconds, while from Fig. 3, 4, 5, 6 we could see that the problem is solved after nearly 50 seconds with greater convergence accuracy.



FIGURE 6. The first component of position states.



FIGURE 7. The second component of position states.



FIGURE 8. The first component of velocity states.

Therefore, by taking the algorithms with nonuniform gradient gains, the distributed optimization problem of the heterogeneous multi-agent systems can be solved and consume less time, while the convergence accuracy could be greatly improved.

C. EXAMPLE 3

In this subsection, the convex function is chosen as $f_1(s)$: $\mathbf{R}^2 \rightarrow \mathbf{R}$:

$$f_1(s) = \begin{cases} 0, & \text{if } ||s|| \le 1, \\ 0.5(||s|| - 1)^2, & \text{otherwise.} \end{cases}$$

By simple calculations, we have

$$\Delta f_1(s) = \begin{cases} [0,0]^T, & \text{if } ||s|| \le 1, \\ (||s|| - 1)\frac{s}{||s||} \ne [0,0]^T, & \text{otherwise.} \end{cases}$$

 $f_2(s) = s_1^4 + s_2^2, f_3(s) = s_1^2 + s_2^4, f_4(s) = s_1^2 + (s_2 - 2)^2,$ $f_5(s) = (s_1 - 2)^2 + s_2^2, f_6(s) = s_1^4 + s_2^2, f_7(s) = s_1^2 + (s_2 - 2)^2,$



FIGURE 9. The second component of velocity states.



FIGURE 10. The position states.



FIGURE 11. The velocity states.

 $f_8(s) = s_1^2 + s_2^4$. The initial values of the nonuniform gradient gains are chosen as $p_i(0) = 2, i = 1, 2, \dots, 8$. By simple computation, we can obtain that the optimal solution of the function $\sum_{i=1}^{\infty} f_i(s)$ is s = (0.362, 0.6144). By calculation $||s|| = 0.7\overline{131} < 1$, therefore we have $f_1(s) = 0$. Since local differentiable convex objective function $f_1(s) \ge 0$, then we could conclude that the optimal solution of the team objective function f(s) is (0.362, 0.6144). By taking algorithms (4) and (5) with the nonuniform gradient gains, the position trajectories of all agents are shown in Fig. 10, from which we could see that the position states of all agents can reach an agreement and converge to the optimal solution of the team objective function with an extremely small error. Fig. 11 show the velocity states of all second-order agents, from which we could see that velocities of all second-order agents converge to zero asymptotically. Therefore, the proposed algorithms are available for non-smooth functions.

V. CONCLUSION

In this paper, we studied the distributed optimization problem of heterogeneous multi-agent systems with switching jointly strongly connected topologies, where each agent has firstorder or second-order dynamics and the team objective function is the sum of finite local convex function. To solve this problem, a distributed algorithm is proposed for each agent based on the local information accessed by each agent. Then, by a coordination transformation, the closed-loop system is converted into an equivalent system, the system matrix of which is stochastic matrix. Based on the properties of stochastic matrix, it is shown that the consensus can be achieved and the team objective function can be minimized simultaneously. Finally, simulation results were given to demonstrate the correctness of the theoretical results. In the future, the distributed optimization problems with time-delay will be considered.

ACKNOWLEDGMENT

(Jingyi Li is co-first author.)

REFERENCES

- Y. Jia, "Robust control with decoupling performance for steering and traction of 4WS vehicles under velocity-varying motion," *IEEE Trans. Control Syst. Technol.*, vol. 8, no. 3, pp. 554–569, May 2000.
- [2] Y. Jia, "Alternative proofs for improved LMI representations for the analysis and the design of continuous-time systems with polytopic type uncertainty: A predictive approach," *IEEE Trans. Autom. Control*, vol. 48, no. 8, pp. 1413–1416, Aug. 2003.
- [3] L. Moreau, "Stability of multiagent systems with time-dependent communication links," *IEEE Trans. Autom. Control*, vol. 50, no. 2, pp. 169–182, Feb. 2005.
- [4] W. Ren and R. W. Beard, "Consensus seeking in multiagent systems under dynamically changing interaction topologies," *IEEE Trans. Autom. Control*, vol. 50, no. 5, pp. 655–661, May 2005.
- [5] B. Zhang, Y. Jia, and F. Matsuno, "Finite-time observers for multi-agent systems without velocity measurements and with input saturations," *Syst. Control Lett.*, vol. 68, pp. 86–94, Jun. 2014.
- [6] L. Zhao, J. Yu, C. Lin, and H. Yu, "Distributed adaptive fixed-time consensus tracking for second-order multi-agent systems using modified terminal sliding mode," *Appl. Math. Comput.*, vol. 312, pp. 23–35, Nov. 2017.
- [7] P. Lin and W. Ren, "Constrained consensus in unbalanced networks with communication delays," *IEEE Trans. Autom. Control*, vol. 59, no. 3, pp. 775–781, Mar. 2014.
- [8] P. Lin, W. Ren, and H. Gao, "Distributed velocity-constrained consensus of discrete-time multi-agent systems with nonconvex constraints, switching topologies, and delays," *IEEE Trans. Autom. Control*, vol. 62, no. 11, pp. 5788–5794, Nov. 2017.
- [9] L. Mo and P. Lin, "Distributed consensus of second-order multiagent systems with nonconvex input constraints," *Int. J. Robust Nonlinear Control*, vol. 28, no. 11, pp. 3657–3664, Jul. 2018.
- [10] H. Fang, C. Shang, and J. Chen, "An optimization-based shared control framework with applications in multi-robot systems," *Sci. China Inf. Sci.*, vol. 61, no. 1, Jan. 2018, Art. no. 014201.
- [11] M. M. Rana, L. Li, and S. W. Su, "An adaptive-then-combine dynamic state estimation considering renewable generations in smart grids," *IEEE J. Sel. Areas Commun.*, vol. 34, no. 12, pp. 3954–3961, Dec. 2016.
- [12] M. Rana, L. Li, and S. W. Su, "Distributed state estimation over unreliable communication networks with an application to smart grids," *IEEE Trans. Green Commun. Netw.*, vol. 1, no. 1, pp. 89–96, Mar. 2017.
- [13] M. M. Rana, W. Xiang, and E. Wang, "IoT-based state estimation for microgrids," *IEEE Internet Things J.*, vol. 5, no. 2, pp. 1345–1346, Apr. 2018.
- [14] M. M. Rana, L. Li, S. W. Su, and W. Xiang, "Consensus-based smart grid state estimation algorithm," *IEEE Trans. Ind. Informat.*, vol. 14, no. 8, pp. 3368–3375, Aug. 2018.
- [15] M. M. Rana, W. Xiang, E. Wang, and X. Li, "Monitoring the smart grid incorporating turbines and vehicles," *IEEE Access*, vol. 6, pp. 45485–45492, 2018.

- [16] M. M. Rana and W. Xiang, "IoT communications network for wireless power transfer system state estimation and stabilization," *IEEE Internet Things J.*, vol. 5, no. 5, pp. 4142–4150, Oct. 2018.
- [17] M. M. Rana, W. Xiang, and E. Wang, "Smart grid state estimation and stabilisation," *Int. J. Elect. Power Energy Syst.*, vol. 102, pp. 152–159, Nov. 2018.
- [18] Q. Lü, H. Li, and D. Xia, "Distributed optimization of first-order discretetime multi-agent systems with event-triggered communication," *Neurocomputing*, vol. 235, pp. 255–263, Apr. 2017.
- [19] Q. Lü, H. Li, and D. Xia, "Geometrical convergence rate for distributed optimization with time-varying directed graphs and uncoordinated stepsizes," *Inf. Sci.*, vol. 422, pp. 516–530, Jan. 2018.
- [20] Q. Lü, H. Li, Z. Wang, Q. Han, and W. Ge, "Performing linear convergence for distributed constrained optimisation over time-varying directed unbalanced networks," *IET Control Theory Appl.*, to be published. doi: 10.1049/iet-cta.2018.6026.
- [21] A. Nedic and A. Ozdaglar, "Distributed subgradient methods for multiagent optimization," *IEEE Trans. Autom. Control*, vol. 54, no. 1, pp. 48–61, Jan. 2009.
- [22] A. Nedić and A. Olshevsky, "Distributed optimization over time-varying directed graphs," *IEEE Trans. Autom. Control*, vol. 60, no. 3, pp. 601–615, Mar. 2015.
- [23] G. Shi, K. H. Johansson, and Y. Hong, "Reaching an optimal consensus: Dynamical systems that compute intersections of convex sets," *IEEE Trans. Autom. Control*, vol. 58, no. 3, pp. 610–622, Mar. 2013.
- [24] Z. Qiu, S. Liu, and L. Xie, "Distributed constrained optimal consensus of multi-agent systems," *Automatica*, vol. 68, pp. 209–215, Jun. 2016.
- [25] P. Lin, W. Ren, and J. A. Farrell, "Distributed continuous-time optimization: Nonuniform gradient gains, finite-time convergence, and convex constraint set," *IEEE Trans. Autom. Control*, vol. 62, no. 5, pp. 2239–2253, May 2017.
- [26] J. Lu and C. Y. Tang, "Zero-gradient-sum algorithms for distributed convex optimization: The continuous-time case," *IEEE Trans. Autom. Control*, vol. 57, no. 9, pp. 2348–2354, Sep. 2012.
- [27] D. Varagnolo, F. Zanella, A. Cenedese, G. Pillonetto, and L. Schenato, "Newton-Raphson consensus for distributed convex optimization," *IEEE Trans. Autom. Control*, vol. 61, no. 4, pp. 994–1009, Apr. 2016.
- [28] B. Gharesifard and J. Cortés, "Distributed continuous-time convex optimization on weight-balanced digraphs," *IEEE Trans. Autom. Control*, vol. 59, no. 3, pp. 781–786, Mar. 2014.
- [29] S. S. Kia, J. Cortés, and S. Martínez, "Distributed convex optimization via continuous-time coordination algorithms with discrete-time communication," *Automatica*, vol. 55, pp. 254–264, May 2015.
- [30] A. Nedic, A. Ozdaglar, and P. A. Parrilo, "Constrained consensus and optimization in multi-agent networks," *IEEE Trans. Autom. Control*, vol. 55, no. 4, pp. 922–938, Apr. 2010.
- [31] P. Lin, W. Ren, and Y. Song, "Distributed multi-agent optimization subject to nonidentical constraints and communication delays," *Automatica*, vol. 65, pp. 120–131, Mar. 2016.
- [32] P. Lin, W. Ren, C. Yang, and W. Gui, "Distributed continuous-time and discrete-time optimization with nonuniform unbounded convex constraint sets and nonuniform stepsizes," *IEEE Trans. Autom. Control*, to be published. doi: 10.1109/TAC.2019.2910946.
- [33] Y. Zhang and Y. Hong, "Distributed optimization design for second-order multi-agent systems," in *Proc. 33rd Chin. Control Conf.*, Nanjing, China, Jul. 2014, pp. 1755–1760.
- [34] Q. Liu and J. Wang, "A second-order multi-agent network for boundconstrained distributed optimization," *IEEE Trans. Autom. Control*, vol. 60, no. 12, pp. 3310–3315, Dec. 2015.
- [35] P. Lin, W. Ren, C. Yang, and W. Gui, "Distributed optimization with nonconvex velocity constraints, nonuniform position constraints and nonuniform stepsizes," *IEEE Trans. Autom. Control*, vol. 64, no. 6, pp. 2575–2582, Jun. 2019.
- [36] P. Wang, P. Lin, W. Ren, and Y. Song, "Distributed subgradient-based multiagent optimization with more general step sizes," *IEEE Trans. Autom. Control*, vol. 63, no. 7, pp. 2295–2302, Jul. 2018.
- [37] L. Mo and P. Lin, "Distributed continuous-time optimization over secondorder multi-agent networks with nonuniform gains," in *Proc. 31rd Chin. Control Decis. Conf.*, Nanchang, China, Jun. 2019, pp. 35–38.

- [38] Y. Zheng, Y. Zhu, and L. Wang, "Consensus of heterogeneous multiagent systems," *IET Control Theory Appl.*, vol. 5, no. 16, pp. 1881–1888, Apr. 2011.
- [39] Y. Feng, S. Xu, F. L. Lewis, and B. Zhang, "Consensus of heterogeneous first- and second-order multi-agent systems with directed communication topologies," *Int. J. Robust Nonlinear Control*, vol. 25, no. 3, pp. 362–375, Feb. 2015.
- [40] L. Mo, G. Niu, and T. Pan, "Consensus of heterogeneous multi-agent systems with switching jointly-connected interconnection," *Phys. A, Stat. Mechanics Appl.*, vol. 427, pp. 132–140, Jun. 2015.
- [41] L. Mo, S. Guo, and Y. Yu, "Mean-square consensus of heterogeneous multi-agent systems with nonconvex constraints, Markovian switching topologies and delays," *Neurocomputing*, vol. 291, pp. 167–174, May 2018.
- [42] H. Huang, L. Mo, and X. Cao, "Nonconvex constrained consensus of discrete-time heterogeneous multi-agent systems with arbitrarily switching topologies," *IEEE Access*, vol. 7, pp. 38157–38161, 2019.
- [43] C. Godsil and G. Royle, Algebraic Graph Theory. New York, NY, USA: Springer-Verlag, 2001.



LIPO MO was born in Hebei, China, in 1980. He received the B.S. degree in mathematics and applied mathematics from Shihezi University, in 2003, and the Ph.D. degree in control theory from the School of Mathematics and Systems Science, Beihang University, in 2010. He is currently an Associate Professor with the Beijing Technology and Business University. His research interests include stochastic systems, coordination control of multi-agent systems, and distributed optimization.



JINGYI LI was born in Beijing, China, in 1998. She is currently pursuing the B.S. degree in mathematics with Beihang University. She is the cofirst author of this paper. Her research interests include statistical computing, coordination control of multi-agent systems, and stochastic processes.



JIAN HUANG was born in Sichuang, China, in 1960. He received the B.S. degree in mathematics from Chongqing University, in 1982, the M.S. degree in mathematics from North-Western Polytechnic University, in 1987, and the Ph.D. degree in statistics from University College Dublin, in 1997. Since 1999, he has been a Researcher with University College Cork, where he is currently the Chair of Risk and Actuarial Studies Committee, School of Mathematical Sci-

ence. His research interests include modeling large spatio-temporal data, time series analysis, inverse problems, medical image data analysis, distributed optimization, and machine learning.

•••