

Title	Schizophrenia patients with a history of childhood trauma have a pro-inflammatory phenotype
Authors	Dennison, Una;McKernan, Declan P.;Cryan, John F.;Dinan, Timothy G.
Publication date	2012
Original Citation	U. Dennison, D. McKernan, J. Cryan and T. Dinan (2012). Schizophrenia patients with a history of childhood trauma have a proinflammatory phenotype. <i>Psychological Medicine</i> , 42(9), pp 1865-1871 doi:10.1017/S0033291712000074
Type of publication	Article (peer-reviewed)
Link to publisher's version	http://journals.cambridge.org/action/displayAbstract?fromPage=online&aid=8656043&fulltextType=RA&fileId=S0033291712000074 - 10.1017/S0033291712000074
Rights	© Cambridge University Press, 2012
Download date	2024-04-19 13:07:40
Item downloaded from	https://hdl.handle.net/10468/1000

Psychological Medicine

<http://journals.cambridge.org/PSM>

Additional services for *Psychological Medicine*:

Email alerts: [Click here](#)

Subscriptions: [Click here](#)

Commercial reprints: [Click here](#)

Terms of use : [Click here](#)



Schizophrenia patients with a history of childhood trauma have a pro-inflammatory phenotype

U. Dennison, D. McKernan, J. Cryan and T. Dinan

Psychological Medicine / Volume 42 / Issue 09 / September 2012, pp 1865 - 1871
DOI: 10.1017/S0033291712000074, Published online: 23 February 2012

Link to this article: http://journals.cambridge.org/abstract_S0033291712000074

How to cite this article:

U. Dennison, D. McKernan, J. Cryan and T. Dinan (2012). Schizophrenia patients with a history of childhood trauma have a pro-inflammatory phenotype. *Psychological Medicine*, 42, pp 1865-1871 doi:10.1017/S0033291712000074

Request Permissions : [Click here](#)

Schizophrenia patients with a history of childhood trauma have a pro-inflammatory phenotype

U. Dennison^{1*†}, D. McKernan^{2†}, J. Cryan^{2,3} and T. Dinan^{1,2}

¹ Department of Psychiatry, University College Cork, Cork, Ireland

² Alimentary Pharmabiotic Centre, University College Cork, Cork, Ireland

³ Department of Anatomy, University College Cork, Cork, Ireland

Background. Increasing evidence indicates that childhood trauma is a risk factor for schizophrenia and patients with this syndrome have a pro-inflammatory phenotype. We tested the hypothesis that the pro-inflammatory phenotype in schizophrenia is associated with childhood trauma and that patients without a history of such trauma have a similar immune profile to healthy controls.

Method. We recruited 40 schizophrenia patients and 40 controls, all of whom completed the Childhood Trauma Questionnaire (CTQ). Using enzyme-linked immunosorbent assay (ELISA) techniques, we measured peripheral levels of interleukin (IL)-1 β , IL-6, IL-8 and tumour necrosis factor (TNF)- α . These immune parameters were compared in schizophrenia with childhood trauma, schizophrenia without childhood trauma and healthy controls.

Results. Patients with childhood trauma had higher levels of IL-6 and TNF- α than patients without trauma and healthy controls, and TNF- α levels correlated with the extent of the trauma. Patients with no trauma had similar immune profiles to controls.

Conclusions. Childhood trauma drives changes, possibly epigenetic, that generate a pro-inflammatory phenotype.

Received 20 October 2011; Revised 19 December 2011; Accepted 5 January 2012; First published online 23 February 2012

Key words: Childhood trauma, inflammation, schizophrenia.

Introduction

Schizophrenia is a chronic, debilitating mental disorder that affects about 1% of the population. The aetiology of schizophrenia is not fully understood, but several aetiological hypotheses have been put forward that focus on genetics, neurotransmitter abnormalities and impaired neurodevelopment (Tandon *et al.* 2008). A wide variety of environmental insults have been investigated, such as viral infection during pregnancy. These factors have been comprehensively reviewed by Brown (2011).

There is accumulating evidence that schizophrenia may be due to inflammatory insults, and several studies indicate alterations in the innate immune system in schizophrenia including increased chemokines (e.g. CCL 11), acute-phase proteins (e.g. C-reactive protein, CRP) and cytokines (Strous & Shoenfeld, 2006). Cytokines are a large family of intracellular signalling molecules involved in immune cell activity and the

inflammatory response. Some cytokines, such as interleukin (IL)-6, IL-8 and tumour necrosis factor (TNF)- α , are pro-inflammatory whereas others dampen the immune response, and in general there is a 'cytokine balance' between pro- and anti-inflammatory cytokines in the body (Commins *et al.* 2010). With regard to cytokine abnormalities in schizophrenia, results are often contradictory. However, increased plasma IL-2 and IL-6 are well-replicated findings. A systematic review conducted by Potvin *et al.* (2008) reported *in vivo* increases of IL-1 receptor antagonist (IL-1RA), soluble IL-2 receptor (sIL-2R) and IL-6, and *in vitro* increases of IL-2. These alterations in IL-6 and IL-1RA were not related to antipsychotic medications, suggesting that they are linked to the physiopathology of the disease, or to as yet uncharacterized phenotypic traits of the disorder (Potvin *et al.* 2008).

With regard to environmental exposures and schizophrenia, several studies suggest that children exposed to early trauma are at greater risk of later experiencing psychotic symptoms (Read *et al.* 2005; Morgan & Fisher, 2007). In addition, genetically high-risk adoptees are significantly more sensitive than genetically low-risk adoptees to adverse rearing patterns in adoptive families, highlighting the complicated

* Address for correspondence: Dr U. Dennison, c/o City General Hospital, Infirmary Road, Cork, Ireland.
(Email: una.dennison@hse.ie)

† These authors contributed equally to this work.

gene–environment interaction observed in schizophrenia (Tienari *et al.* 2004). In relation to physical and sexual abuse, there seems to be a cumulative effect whereby the more trauma a person is exposed to, the greater the risk of psychosis/schizophrenia later on (Janssen *et al.* 2004; Shevlin *et al.* 2007). A recently published, longitudinal prospective follow-up study assessed more than 800 children exposed to the 1983 Australian bushfires. Children exposed to the bushfire alone did not have increased risk of psychosis. However, exposure to multiple traumas was positively associated with the development of psychotic symptoms, and these children had higher rates of dysfunctional parenting, drug and alcohol misuse, emotional and behavioural disturbance and childhood adversity (Galletly *et al.* 2011).

Some authors relate childhood trauma to abnormal cortisol levels and a disrupted hypothalamic–pituitary–adrenal (HPA) axis, which is responsible for the body's response to stress (Braehler *et al.* 2005). More recently, studies suggest inflammatory alterations in those exposed to early trauma, including increased CRP in depressed adults with a history of childhood maltreatment and a greater inflammatory response to acute psychological stress when compared with controls (Pace *et al.* 2006; Danese *et al.* 2008). In healthy individuals with a history of childhood maltreatment, there is evidence for increased IL-6 release in response to psychological stress (Carpenter *et al.* 2010).

We hypothesize in this study that schizophrenia patients only demonstrate a pro-inflammatory phenotype if they have experienced childhood trauma.

Subject population

Following ethical approval, 40 patients with schizophrenia aged between 18 and 65 years were recruited from psychiatric out-patient clinics and the in-patient ward at Cork University Hospital. All patients satisfied DSM-IV criteria for the diagnosis of schizophrenia based on a structured clinical interview (APA, 1994). Forty healthy controls were recruited from staff members at Cork University Hospital and University College Cork. Each participant gave fully informed written consent, and there was no financial reward for participation. Participants with inflammatory diseases, coeliac disease, lactose intolerance, or immunodeficiency, and those who had undergone any abdominal surgery, with the exception of hernia repair and appendectomy, were excluded. Each subject had a structured interview with a trained psychiatrist and donated 20 ml of venous blood (between 09:00 and 12:00 h). Details recorded on all participants included current health status, body mass index (BMI),

and medical/psychiatric history, including current medication and family history.

Questionnaires and psychiatric rating scales

In the schizophrenia cohort, symptoms were rated by a psychiatrist according to the Positive and Negative Symptom Scale (PANSS). This 30-item scale measures positive and negative symptoms, their differential, which is termed the composite score, and general psychopathology symptoms. The total score of the PANSS can be used to assess illness severity (Kay *et al.* 1987; Leucht *et al.* 2005; Opler *et al.* 2007). To assess comorbid depression, the Calgary Depression Scale for Schizophrenia (CDSS) was also completed. This is an observer-rated scale, and a cut-off ≥ 6 indicates concurrent depression (Addington *et al.* 1992, 1993). All study participants completed the short version of the Childhood Trauma Questionnaire (CTQ), which is an eight-item, self-report questionnaire that identifies the presence or absence of childhood traumatic events, including (1) death of a family member or close friend in childhood, (2) parental separation, (3) exposure to physical abuse, (4) exposure to sexual abuse, (5) exposure to violence, (6) significant illness or injury in childhood, (7) traumatic events in adulthood, and (8) obstetric complications (Bernstein *et al.* 1994). According to this scale, questions 1–5 relate specifically to psychological trauma in childhood. It is worth noting that retrospective self-reports are not completely accurate and are influenced by several factors such as forgetting, biases and repression of memories. However, this leads to an under- rather than an overestimation according to Hardt & Rutter (2004), and the CTQ has proven validity and reliability for detecting childhood traumatic events (Bernstein *et al.* 1994).

Plasma isolation and enzyme-linked immunosorbent assay (ELISA)

Whole blood (15 ml of the 20 ml collected) was added to an equal volume of Histopaque 1077 (Sigma, USA) in a sterile 50 ml tube and centrifuged at 400 *g* for 30 min at room temperature. Plasma on the upper layer was transferred to a separate tube and stored at -80°C for future use. Pro-inflammatory cytokine IL-1 β , IL-6, IL-8 and TNF- α measurement in plasma was carried out using the Meso Scale Discovery (MSD) 4-plex Human Pro-inflammatory kit II. ELISA plates were analysed using the Sector 2400 imager from MSD (USA). This is an ultra-sensitive method that has a detection limit of 0.3, 0.3, 1.0 and 0.3 pg/ml for IL-1 β , IL-6, IL-8 and TNF- α respectively (O'Brien *et al.* 2006).

Table 1. Demographic data

	Controls (<i>n</i> = 40)	Patients with schizophrenia (<i>n</i> = 40)
Age (years)	36.2 ± 1.76	38.33 ± 1.7
Sex, M/F	13/27	24/16
Height (m)	1.7 ± 0.01	1.73 ± 0.01
Weight (kg)	67.74 ± 2.6	84.01 ± 2.88***
BMI (kg/m ²)	23.89 ± 0.83	27.94 ± 0.85**
CTQ score	0.67 ± 0.15	1.75 ± 0.24***
Diagnosis		
Schizophrenia	N.A.	34
Schizo-affective disorder	N.A.	6
PANSS		
(+) Symptoms	N.A.	18
(-) Symptoms	N.A.	22
History of depression		
Previous diagnosis	0	19
Current diagnosis	0	9
Family history of psychosis	0	11
Family history of mood disorder		
Unipolar depression	3	9
Bipolar affective disorder	0	2

M, Male; F, female; BMI, body mass index; N.A., not applicable; CTQ, Childhood Trauma Questionnaire; PANSS, Positive and Negative Symptom Scale.

p* < 0.01, *p* < 0.001.

Statistical analysis

Data are expressed as mean ± standard error of the mean (S.E.M.). Before analysis, the data were examined for normality using the Kolmogorov–Smirnov test. Comparisons were made using the Student *t* test and one-way analysis of variance (ANOVA), with Bonferroni *post-hoc* tests and correlation with Pearson's product-moment correlation coefficient and the χ^2 test where appropriate. Differences were considered significant at *p* < 0.05. All statistical analysis was carried out using GraphPad Prism for Windows, version 5 (GraphPad, USA).

Results

Patient demographics (see Table 1)

In the schizophrenia cohort, we recruited 40 patients (24 males, 16 females), with a mean age of 38 ± 1.7 years, weight 84.01 ± 2.88 kg and BMI 27.94 ± 0.85 kg/m². According to the DSM-IV, 34 had a diagnosis of schizophrenia and six had a diagnosis of schizo-affective disorder (APA, 1994). Using the total score on the PANSS, 20 patients had mild, 13 patients moderate and seven patients severe illness severity. According to the composite PANSS score, 22 patients had

predominantly negative and 18 patients had predominantly positive symptoms. A positive history of depression was recorded in 22 schizophrenia subjects (19 patients self-reported previous depression and nine scored positive for current depression according to the CDSS). There was a positive family history of psychosis in 11 participants, a positive family history of unipolar mood disorder in nine participants, and a positive family history of bipolar affective disorder (BPAD) in two participants. With regard to medications in the schizophrenia group, one patient was prescribed an antidepressant alone. In the remaining 39 patients, 26 were taking one antipsychotic, 12 two antipsychotics, and one was taking a combination of three antipsychotics. The total numbers of individual antipsychotics were: (18) clozapine; (nine) olanzapine; (eight) risperidone; (four) amisulpride, haloperidol; (three) fluphenazine; (two) aripiprazole, quetiapine; and (one) sulpiride, sertindole, trifluoperazine. In this group, 11 patients were also prescribed antidepressants, four patients were prescribed mood stabilizers, and 10 patients were prescribed benzodiazepines. In the control group (*n* = 40), there were 27 females and 13 males, with a mean age of 36.2 ± 1.76 years, weight 67.74 ± 2.6 kg and BMI 23.89 ± 0.83 kg/m². None of the control subjects had




Fig. 1. Plots of cytokines interleukin (IL)-1 β , IL-6, IL-8 and tumour necrosis factor (TNF)- α in schizophrenia patients with a history of childhood trauma (SCZ⁺), and in schizophrenia patients (SCZ⁻) and controls (Con⁻) with no history of childhood trauma. * $p < 0.05$, *** $p < 0.001$ SCZ⁺ versus Con⁻. # $p < 0.05$, SCZ⁺ versus SCZ⁻.

a current or previous psychiatric diagnosis, none had a positive family history of psychosis and three had a positive family history of unipolar depression.

No significant difference was found between the schizophrenia and control groups in terms of age (Table 1). Patients with schizophrenia had significantly increased weight ($p < 0.001$) and, as a result, an increased BMI ($p = 0.0026$) when compared with controls.

Cytokine results (Fig. 1)

Patients in the schizophrenia group reported on average more childhood traumatic events than control subjects (schizophrenia 1.75 ± 0.24 versus control 0.67 ± 0.15 , $p < 0.001$). As questions 1–5 of the CTQ relate specifically to psychological trauma in childhood, a comparison was performed using patient samples who responded yes to any of questions 1–5 (SCZ⁺ 24 patients) versus subjects with no history of trauma within the schizophrenia group (SCZ⁻ 16 patients) and the control group (Con⁻ 32 controls). Of the 24 schizophrenia patients with a positive history of childhood trauma, a total of 17 (more than two-thirds) had a history of depression (eight of these were currently depressed). Of the 16 non-abused schizophrenia patients, five (less than one-third) had a history of depression (of whom one was currently depressed).

Using one-way ANOVA with a Bonferroni *post-hoc* test, the results were non-significant for IL-1 β (SCZ⁺ 0.237 ± 0.043 , SCZ⁻ 0.264 ± 0.046 , Con⁻ 0.329 ± 0.103 ; $F = 0.355$, $df = 143$, $p = 0.702$) and IL-8 (SCZ⁺ 4.313 ± 0.685 , SCZ⁻ 4.817 ± 0.725 , Con⁻ 3.614 ± 0.331 ; $F = 1.177$, $df = 143$, $p = 0.311$). However, there were

significantly increased levels of both IL-6 (SCZ⁺ 2.149 ± 0.318 , SCZ⁻ 1.048 ± 0.172 , Con⁻ 1.092 ± 0.296 ; $F = 4.258$, $df = 143$, $p < 0.05$) and TNF- α (SCZ⁺ 8.248 ± 0.601 , SCZ⁻ 6.088 ± 0.465 , Con⁻ 3.614 ± 0.331 ; $F = 11.41$, $df = 143$, $p < 0.001$) in schizophrenia patients with a positive history of childhood trauma compared with control samples with no history of childhood trauma. There were also increased levels of TNF- α in schizophrenia patients positive for childhood trauma versus those negative for childhood trauma (SCZ⁺ 8.248 ± 0.601 , SCZ⁻ 6.088 ± 0.465 ; $F = 11.41$, $df = 143$, $p < 0.05$; see Fig. 1). Using the χ^2 test, there was a significant association between cytokine level and extent of childhood trauma for TNF- α , with a greater number of traumatic exposures correlating with higher cytokine response ($\chi^2 = 20.27$, $df = 10$, $p < 0.05$).

Within the schizophrenia group as a whole, using Pearson's product-moment correlation coefficient, there was no significant correlation between cytokine levels and psychopathology, based on the total score of PANSS, the composite score of PANSS and the CDSS score reflecting overall illness severity, the presence of predominant positive or negative symptoms and the presence of current depression respectively.

The effects of BMI were analysed within the schizophrenia group, specifically looking at those with a BMI < 25 kg/m² ($n = 13$) versus those with a BMI > 25 kg/m² ($n = 27$). The results were non-significant for all cytokines measured, including IL-1 β ($F = 1.189$, $df = 78$, $p = 0.813$), IL-6 ($F = 5.276$, $df = 78$, $p = 0.232$), IL-8 ($F = 1.034$, $df = 78$, $p = 0.160$) and TNF- α ($F = 1.417$, $df = 78$, $p = 0.199$).

Discussion

In this study, we found increased levels of the pro-inflammatory markers IL-6 and TNF- α only in schizophrenia patients with a positive history of childhood trauma. Despite robust data indicating childhood maltreatment or trauma as a risk factor for schizophrenia, the biological mechanisms by which early traumatic events increase the risk of subsequent psychopathology are not fully understood. We could speculate that early trauma brings about epigenetic changes that lead to a pro-inflammatory phenotype in adulthood.

To date, there is evidence to suggest that child abuse causes long-lasting alterations in the HPA axis, hippocampal function and dopaminergic neurotransmitter systems, all of which are implicated in psychosis (Cicchetti & Walker, 2001; Read *et al.* 2001). One study reported that childhood abuse diminished cortisol response, which was amplified with increasing age and independent of other variables including psychiatric illness (Carpenter *et al.* 2009). There are

several studies on childhood trauma and depression that link trauma with altered HPA axis activity, in addition to immune activation and alterations in hippocampal volume (Heim *et al.* 2008*a,b*). Studies in animals have shown that disrupted maternal care in infancy has chronic deleterious effects on the HPA axis and also the mesolimbic dopamine system, and in humans poor parental care in childhood is associated with higher levels of salivary cortisol in response to psychological stress and also increased ventral striatum dopamine release (Liu *et al.* 1997; Hall *et al.* 1999). Taken together, these results suggest that disrupted care in childhood is associated with HPA axis dysfunction and hyper-reactivity of the dopamine system. Pathological effects of stress on the hippocampus may be related to an increase in cortisol and a decrease in brain-derived neurotrophic factor (BDNF) (Bremner, 2005; Galletly *et al.* 2011). Such hippocampal abnormalities, HPA axis disruptions and low levels of BDNF have also been described in schizophrenia, which might suggest that trauma increases the risk of psychosis in vulnerable individuals by altering brain structure and function. Recent studies by Mondelli *et al.* (2009, 2011) reported that, in first-episode psychosis, childhood trauma and adult stress are partly responsible for down-regulation of BDNF expression. They also reported hyperactivity of the HPA axis in first-episode psychosis, but found that factors other than stress resulted in these abnormalities.

Other evidence also suggests increased levels of inflammatory biomarkers in those exposed to childhood trauma. In rodents, acute stress is associated with increased IL-6 activity (LeMay *et al.* 1990; Zhou *et al.* 1993). Animal studies linking early life stress with alterations in inflammation in adulthood have been published, including a study that showed that early life stress by maternal separation in rats resulted in increased pro-inflammatory cytokine release, in addition to altered HPA axis activity (O'Mahony *et al.* 2009). Studies examining depressed adults with a history of childhood maltreatment also suggest significantly increased inflammatory markers such as CRP, and greater inflammatory response compared with non-depressed controls in response to acute psychological stress (Pace *et al.* 2006; Danese *et al.* 2008). In healthy individuals, a history of childhood maltreatment correlated higher scores on the CTQ with higher IL-6 release in response to psychological stress, further supporting the evidence for alterations in inflammatory responses in those who have a history of childhood maltreatment (Carpenter *et al.* 2010). Because IL-6 is a potent activator of the HPA axis (Žarković *et al.* 2008), it is possible that the HPA axis disturbances in schizophrenia are brought about by

pro-inflammatory cytokines. Exactly how childhood trauma generates a pro-inflammatory phenotype is unclear but may be related to stress-induced priming of microglia, which, when exposed to subsequent pro-inflammatory stimuli, may become activated and result in potentiated pro-inflammatory responses (Frank *et al.* 2007; Sparkman & Johnson, 2008).

Of note, in this study, patients with schizophrenia had higher weight and consequently higher BMI than controls. Previous studies have reported an association between early-life stress and obesity, and also a relationship between BMI and stress-induced cytokine response (Brydon *et al.* 2008; Boynton-Jarrett *et al.* 2010). Obesity itself is accompanied by low-grade inflammation, and adipocytes are a source of IL-6, CRP and serum amyloid A (SAA) (Kvasnicka *et al.* 2003). However, in the current study, comparison of schizophrenia patients with a BMI <25 kg/m² with those with a BMI >25 kg/m² did not yield significant differences in levels of IL-1 β , IL-6, IL-8 or TNF- α . It is also worth noting that depression was over-represented in the subgroup of patients with childhood trauma. It is well established that depression is also associated with a pro-inflammatory state (Dantzer *et al.* 2008; Leonard & Myint, 2009), and hence may be partially responsible for the inflammatory load seen in childhood trauma patients. However, studies of non-depressed adults with a history of childhood trauma indicate greater IL-6 release in response to psychological stress (Carpenter *et al.* 2010), suggesting that childhood trauma influences pro-inflammatory cytokines independent of depression.

Important limitations in this study include the lack of medication-free patients, as psychotropic medications are known to have an influence on cytokine levels (Maes *et al.* 1997). Of the 40 control subjects recruited, eight had a positive history of childhood trauma. A larger sample size, including more controls with a history of childhood trauma, would also have been an informative comparison group. In general, more studies investigating the link between trauma and inflammation are required, particularly in patients with psychiatric illness, where epigenetic mechanisms may be generating a pro-inflammatory phenotype.

Acknowledgements

The Alimentary Pharmabiotic Centre is a Centre for Science, Engineering and Technology (CSET) funded by Science Foundation Ireland (SFI), through the Irish Government's National Development Plan. The authors and their work were supported by SFI and GlaxoSmithKline through grant 07/CE/B1368.

Declaration of Interest

None.

References

- Addington D, Addington J, Maticka-Tyndale E** (1993). Assessing depression in schizophrenia: the Calgary Depression Scale. *British Journal of Psychiatry. Supplement* 39–44.
- Addington D, Addington J, Maticka-Tyndale E, Joyce J** (1992). Reliability and validity of a depression rating scale for schizophrenics. *Schizophrenia Research* 6, 201–208.
- APA** (1994). *Diagnostic and Statistical Manual of Mental Disorders*, 4th edn. American Psychiatric Association: Washington, DC.
- Bernstein D, Fink L, Handelsman L, Foote J, Lovejoy M, Wenzel K, Sapareto E, Ruggiero J** (1994). Initial reliability and validity of a new retrospective measure of child abuse and neglect. *American Journal of Psychiatry* 151, 1132–1136.
- Boynton-Jarrett R, Fargnoli J, Suglia SF, Zuckerman B, Wright RJ** (2010). Association between maternal intimate partner violence and incident obesity in preschool-aged children: results from the Fragile Families and Child Well-being Study. *Archives of Paediatric and Adolescent Medicine* 164, 540–546.
- Braehler C, Holowka D, Brunet A, Beaulieu S, Baptista T, Debruille JB, Walker CD, King S** (2005). Diurnal cortisol in schizophrenia patients with childhood trauma. *Schizophrenia Research* 79, 353–354.
- Bremner JD** (2005). Effects of traumatic stress on brain structure and function: relevance to early responses to trauma. *Journal of Trauma and Dissociation* 6, 51–68.
- Brown AS** (2011). The environment and susceptibility to schizophrenia. *Progress in Neurobiology* 93, 23–58.
- Brydon L, Wright CE, O'Donnell K, Zachary I, Wardle J, Steptoe A** (2008). Stress-induced cytokine responses and central adiposity in young women. *International Journal of Obesity* 32, 443–450.
- Carpenter LL, Gawuga CE, Tyrka AR, Lee JK, Anderson GM, Price LH** (2010). Association between plasma IL-6 response to acute stress and early-life adversity in healthy adults. *Neuropsychopharmacology* 35, 2617–2623.
- Carpenter LL, Tyrka AR, Ross NS, Khoury L, Anderson GM, Price LH** (2009). Effect of childhood emotional abuse and age on cortisol responsivity in adulthood. *Biological Psychiatry* 66, 69–75.
- Cicchetti D, Walker EF** (2001). Stress and development: biological and psychological consequences. *Development and Psychopathology* 13, 413–418.
- Commins SP, Borish L, Steinke JW** (2010). Immunologic messenger molecules: cytokines, interferons, and chemokines. *Journal of Allergy and Clinical Immunology* 125, S53–S72.
- Danese A, Moffitt TE, Pariante CM, Ambler A, Poulton R, Caspi A** (2008). Elevated inflammation levels in depressed adults with a history of childhood maltreatment. *Archives of General Psychiatry* 65, 409–415.
- Dantzer R, O'Connor JC, Freund GG, Johnson RW, Kelley KW** (2008). From inflammation to sickness and depression: when the immune system subjugates the brain. *Nature Reviews Neuroscience* 9, 46–56.
- Frank M, Baratta M, Sprunger D, Watkins L, Maier S** (2007). Microglia serve as a neuroimmune substrate for stress induced potentiation of CNS pro-inflammatory cytokine responses. *Brain, Behavior and Immunity* 21, 47–59.
- Galletly C, van Hooff M, McFarlane A** (2011). Psychotic symptoms in young adults exposed to childhood trauma – a 20 year follow-up study. *Schizophrenia Research* 127, 76–82.
- Hall F, Wilkinson L, Humby T, Robbins T** (1999). Maternal deprivation of neonatal rats produces enduring changes in dopamine function. *Synapse* 32, 37–43.
- Hardt J, Rutter M** (2004). Validity of adult retrospective reports of adverse childhood experiences: review of the evidence. *Journal of Child Psychology and Psychiatry* 45, 260–273.
- Heim C, Mletzko T, Purselle D, Musselman DL, Nemeroff CB** (2008a). The dexamethasone/corticotropin-releasing factor test in men with major depression: role of childhood trauma. *Biological Psychiatry* 63, 398–405.
- Heim C, Newport DJ, Mletzko T, Miller AH, Nemeroff CB** (2008b). The link between childhood trauma and depression: insights from HPA axis studies in humans. *Psychoneuroendocrinology* 33, 693–710.
- Janssen I, Krabbendam L, Bak M, Hanssen M, Vollebregt W, de Graaf R, van Os J** (2004). Childhood abuse as a risk factor for psychotic experiences. *Acta Psychiatrica Scandinavica* 109, 38–45.
- Kay S, Fiszbein A, Opler L** (1987). The positive and negative syndrome scale (PANSS) for schizophrenia. *Schizophrenia Bulletin* 13, 261–276.
- Kvasnicka T, Kvasnicka J, Ceska R, Vrablik M** (2003). Increase of inflammatory state in overweight adults with combined hyperlipidemia. *Nutrition, Metabolism and Cardiovascular Diseases* 13, 227–231.
- LeMay LG, Vander AJ, Kluger MJ** (1990). The effects of psychological stress on plasma interleukin-6 activity in rats. *Physiology and Behavior* 47, 957–961.
- Leonard BE, Myint A** (2009). The psychoneuroimmunology of depression. *Human Psychopharmacology* 24, 165–175.
- Leucht S, Kane JM, Kissling W, Hamann J, Etschel E, Engel RR** (2005). What does the PANSS mean? *Schizophrenia Research* 79, 231–238.
- Liu D, Diorio J, Tannenbaum B, Caldji C, Francis D, Freedman A, Sharma S, Pearson D, Plotsky PM, Meaney MJ** (1997). Maternal care, hippocampal glucocorticoid receptors, and hypothalamic-pituitary-adrenal responses to stress. *Science* 277, 1659–1662.
- Maes M, Delange J, Ranjan R, Meltzer HY, Desnyder R, Cooremans W, Scharpe S** (1997). Acute phase proteins in schizophrenia, mania and major depression: modulation by psychotropic drugs. *Psychiatry Research* 66, 1–11.
- Mondelli V, Cattaneo A, Belvederi Murri M, Di Forti M, Handley R, Hepgul N, Miorelli A, Navari S, Papadopoulos A, Aitchinson K, Morgan C, Murray R, Dazzan P, Pariante C** (2011). Stress and inflammation

- reduce brain-derived neurotrophic factor expression in first-episode psychosis: a pathway to smaller hippocampal volumes. *Journal of Clinical Psychiatry*. Published online: 18 May 2011. doi:10.4088/JCP.10m06745.
- Mondelli V, Dazzan J, Hepgul N, Di Forti M, Aas M, D'Albenzio A, Di Nicola M, Fisher H, Handley R, Marques T, Morgan C, Navari S, Taylor H, Papadopoulos A, Aitchinson K, Murray R, Pariante C** (2009). Abnormal cortisol levels during the day and cortisol awakening response in first-episode psychosis: the role of stress and of antipsychotic treatment. *Schizophrenia Research* **116**, 234–242.
- Morgan C, Fisher H** (2007). Environmental factors in schizophrenia: childhood trauma – a critical review. *Schizophrenia Bulletin* **33**, 3–10.
- O'Brien SM, Scully P, Scott LV, Dinan TG** (2006). Cytokine profiles in bipolar affective disorder: focus on acutely ill patients. *Journal of Affective Disorders* **90**, 263–267.
- O'Mahony SM, Marchesi JR, Scully P, Codling C, Ceolho AM, Quigley EM, Cryan JF, Dinan TG** (2009). Early life stress alters behavior, immunity, and microbiota in rats: implications for irritable bowel syndrome and psychiatric illnesses. *Biological Psychiatry* **65**, 263–267.
- Opler MG, Yang LH, Caleo S, Alberti P** (2007). Statistical validation of the criteria for symptom remission in schizophrenia: preliminary findings. *BMC Psychiatry* **7**, 35.
- Pace TWW, Mletzko TC, Alagbe O, Musselman DL, Nemeroff CB, Miller AH, Heim CM** (2006). Increased stress-induced inflammatory responses in male patients with major depression and increased early life stress. *American Journal of Psychiatry* **163**, 1630–1633.
- Potvin S, Stip E, Sepehry AA, Gendron A, Bah R, Kouassi E** (2008). Inflammatory cytokine alterations in schizophrenia: a systematic quantitative review. *Biological Psychiatry* **63**, 801–808.
- Read J, Perry BD, Moskowitz A, Connolly J** (2001). The contribution of early traumatic events to schizophrenia in some patients: a traumagenic neurodevelopmental model. *Psychiatry* **64**, 319–345.
- Read J, van Os J, Morrison AP, Ross CA** (2005). Childhood trauma, psychosis and schizophrenia: a literature review with theoretical and clinical implications. *Acta Psychiatrica Scandinavica* **112**, 330–350.
- Shevlin M, Dorahy MJ, Adamson G** (2007). Trauma and psychosis: an analysis of the National Comorbidity Survey. *American Journal of Psychiatry* **164**, 166–169.
- Sparkman N, Johnson R** (2008). Neuroinflammation associated with aging sensitizes the brain to the effects of infection or stress. *Neuroimmunomodulation* **15**, 323–330.
- Strous RD, Shoenfeld Y** (2006). Schizophrenia, autoimmunity and immune system dysregulation: a comprehensive model updated and revisited. *Journal of Autoimmunity* **27**, 71–80.
- Tandon R, Keshavan MS, Nasrallah HA** (2008). Schizophrenia, 'just the facts' what we know in 2008. 2. Epidemiology and etiology. *Schizophrenia Research* **102**, 1–18.
- Tienari P, Wynne LC, Sorri A, Lahti I, Läksy K, Moring J, Naarala M, Nieminen P, Wahlberg K-E** (2004). Genotype-environment interaction in schizophrenia-spectrum disorder. Long-term follow-up study of Finnish adoptees. *British Journal of Psychiatry* **184**, 216–222.
- Žarković M, Ignjatović S, Dajak M, Ćirić J, Beleslin B, Savić S, Stojković M, Bulat P, Trbojević B** (2008). Cortisol response to ACTH stimulation correlates with blood interleukin 6 concentration in healthy humans. *European Journal of Endocrinology* **159**, 649–652.
- Zhou D, Kusnecov A, Shurin M, Depaoli M, Rabin B** (1993). Exposure to physical and psychological stressors elevates plasma interleukin 6: relationship to the activation of hypothalamic-pituitary-adrenal axis. *Endocrinology* **133**, 2523–2530.