


Title	Molecular pedigree reconstruction and estimation of evolutionary parameters in a wild Atlantic salmon river system with incomplete sampling: a power analysis
Author(s)	Aykanat, Tutku; Johnston, Susan E.; Cotter, Deirdre; Cross, Thomas F.; Poole, Russell; Prodhhl, Paulo A.; Reed, Thomas; Rogan, Ger; McGinnity, Philip; Primmer, Craig R.
Publication date	2014-03-31
Original citation	AYKANAT, T., JOHNSTON, S. E., COTTER, D., CROSS, T. F., POOLE, R., PRODÓHL, P. A., REED, T., ROGAN, G., MCGINNITY, P. & PRIMMER, C. R. 2014. Molecular pedigree reconstruction and estimation of evolutionary parameters in a wild Atlantic salmon river system with incomplete sampling: a power analysis. BMC Evolutionary Biology, 14:68, 1-17. http://dx.doi.org/10.1186/1471-2148-14-68
Type of publication	Article (peer-reviewed)
Link to publisher's version	http://dx.doi.org/10.1186/1471-2148-14-68 Access to the full text of the published version may require a subscription.
Rights	© Aykanat et al.; licensee BioMed Central Ltd. 2014. This article is published under license to BioMed Central Ltd. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/2.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly credited. The Creative Commons Public Domain Dedication waiver (http://creativecommons.org/publicdomain/zero/1.0/) applies to the data made available in this article, unless otherwise stated. http://creativecommons.org/licenses/by/2.0
Item downloaded from	http://hdl.handle.net/10468/2234

Downloaded on 2019-01-19T10:13:39Z



662

663 **Additional file 4. Assumed lifetime reproductive success of Burrishoole Atlantic**
 664 **salmon for females (a), and males (b).** The distributions are negative binomial distributions
 665 with dispersion parameters 0.25 and 0.75 for males and females, respectively. In the above
 666 example, the average reproductive success for each gender is assumed to be two (i.e. stable
 667 population size), while in the simulations average reproductive success is adjusted for
 668 empirical census population size (Table 1).