

**UCC Library and UCC researchers have made this item openly available.
Please [let us know](#) how this has helped you. Thanks!**

Title	Positive mental health and well-being among a third level student population
Author(s)	Davoren, Martin P.; Fitzgerald, Eimear; Shiely, Frances; Perry, Ivan J.
Publication date	2013
Original citation	Davoren MP, Fitzgerald E, Shiely F, Perry IJ (2013) Positive Mental Health and Well-Being among a Third Level Student Population. PLoS ONE 8(8): e74921. doi:10.1371/journal.pone.0074921
Type of publication	Article (peer-reviewed)
Link to publisher's version	http://dx.doi.org/10.1371/journal.pone.0074921 Access to the full text of the published version may require a subscription.
Rights	© 2013 Davoren et al. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited http://creativecommons.org/licenses/by/4.0/
Item downloaded from	http://hdl.handle.net/10468/2368

Downloaded on 2021-09-21T10:21:06Z

Positive Mental Health and Well-Being among a Third Level Student Population

Martin P. Davoren^{*}, Eimear Fitzgerald, Frances Shiely, Ivan J. Perry

Department of Epidemiology and Public Health, University College Cork, Cork, Ireland

Abstract

Introduction: Much research on the health and well-being of third level students is focused on poor mental health leading to a dearth of information on positive mental health and well-being. Recently, the Warwick Edinburgh Mental Well-being scale (WEMWBS) was developed as a measurement of positive mental health and well-being. The aim of this research is to investigate the distribution and determinants of positive mental health and well-being in a large, broadly representative sample of third level students using WEMWBS.

Methods: Undergraduate students from one large third level institution were sampled using probability proportional to size sampling. Questionnaires were distributed to students attending lectures in the randomly selected degrees. A total of 2,332 self-completed questionnaires were obtained, yielding a response rate of 51% based on students registered to relevant modules and 84% based on attendance. One-way ANOVAs and multivariate logistic regression were utilised to investigate factors associated with positive mental health and well-being.

Results: The sample was predominantly female (62.66%), in first year (46.9%) and living in their parents' house (42.4%) or in a rented house or flat (40.8%). In multivariate analysis adjusted for age and stratified by gender, no significant differences in WEMWBS score were observed by area of study, alcohol, smoking or drug use. WEMWBS scores were higher among male students with low levels of physical activity ($p=0.04$). Men and women reporting one or more sexual partners ($p<0.001$) were also more likely to report above average mental health and well-being.

Conclusion: This is the first study to examine positive mental health and well-being scores in a third level student sample using WEMWBS. The findings suggest that students with a relatively adverse health and lifestyle profile have higher than average mental health and well-being. To confirm these results, this work needs to be replicated across other third level institutions.

Citation: Davoren MP, Fitzgerald E, Shiely F, Perry IJ (2013) Positive Mental Health and Well-Being among a Third Level Student Population. PLoS ONE 8(8): e74921. doi:10.1371/journal.pone.0074921

Editor: Jerson Laks, Federal University of Rio de Janeiro, Brazil

Received: April 10, 2013; **Accepted:** August 7, 2013; **Published:** August 29, 2013

Copyright: © 2013 Davoren et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: The authors have no support or funding to report.

Competing interests: The authors have declared that no competing interests exist.

* E-mail: m.davoren@ucc.ie

Introduction

Suicide, deliberate self-harm and the negative aspects of mental health and well-being have previously been the primary focus of much research. Studies from around the world focusing on the mental health and well-being of third level students have investigated risky behaviours and poor mental health and well-being [1–6]. Much of this research has highlighted alcohol and drug abuse, financial pressures and academic concerns [1,2] as determinants of mental health and well-being. However, a dearth of information on positive mental health and well-being among this population has been identified.

Historically, humankind has always been fascinated with the determinants of human health, happiness and well-being [7]. Positive mental health can be defined as “the scientific study of

those positive strengths and virtues that enable people & communities to reach optimal levels of health, happiness and well-being” [8]. This adopts the view that positive mental health and well-being is a positive concept which goes beyond healthy lifestyles to overall well-being and replaces the traditional medical perspective which has mainly focused on the treatment of mental illness [9]. As highlighted by Seligman, positive mental health and well-being is grounded in the philosophy of positive psychology [8,10] and health promotion [11].

Recently, the Warwick Edinburgh Mental Well-Being Scale (WEMWBS) was developed as a measurement of positive mental health and well-being. In 2007, a report on its validity, variability and social desirability among a general and student population was published [12]. However, the determinants of positive mental health and well-being have yet to be investigated among a third level student population utilising

WEMWBS. Thus, the aim of this research is to investigate the distribution and determinants of positive mental health and well-being in a large, broadly representative sample of Irish third level students.

Materials and Methods

Sampling & Participants

Undergraduate students registered to degree programmes at one large third level institution were randomly sampled using probability proportional to size (PPS) sampling. The primary focus of this research was on hazardous alcohol consumption. The required sample size of 2,686 students was calculated based on a prevalence of hazardous alcohol consumption of approximately 70% with a precision of 1.5%. Lecturers or module coordinators were contacted to request permission to distribute and collect questionnaires during fifteen minutes of lecture time on a date convenient to them between March 12th and 23rd, 2012. Students were initially briefed on the aims and objectives of the study along with its confidential, anonymous and voluntary nature. Questionnaires were then distributed and students given fifteen minutes for completion. All questionnaires were immediately collected and a post-questionnaire information sheet distributed. As an incentive, all participants were invited to enter a draw to win a tablet computer following survey completion. As questionnaire completion was anonymous, each student was advised to send an e-mail to the lead author with his or her name and e-mail address to enter the prize draw.

In total, 2,332 students completed this face-to-face lecture theatre based, cross-sectional survey. The response rate for those registered to the specific modules visited was 51% but for those in attendance an 84% response rate was achieved. Of these, 2,044 undergraduates completed the full fourteen questions involved in the Warwick-Edinburgh Mental Wellbeing Scale and are the subject of this analysis. Age, gender and the course profiles of the sample collected was similar to those registered with the university. The sample is representative of undergraduate men and women when compared to the specific institution demographics with more women (56%) registered to undergraduate degrees than men (44%). Furthermore, it was representative of university faculty. The institution reports 33% of undergraduate students were registered to the faculty of Arts, 21% to Law and Business, 27% to Science and Engineering while 19% were registered to Health & Welfare courses in 2011-12.

Questionnaire

The questionnaire was based on previous third level surveys conducted in Ireland [3] and it addressed a range of topics including mental health and well-being, alcohol use, smoking, drug use, physical activity, sexual health, diet and self-reported height and weight. Positive mental health and well-being was estimated using the Warwick-Edinburgh Mental Well-being Scale (WEMWBS) [12]. The Cronbach's Alpha measure of internal consistency for WEMWBS in this study was 0.90.

The questionnaire also included additional validated instruments such as the Alcohol Use Disorders Identification

Test for Consumption (AUDIT-C) and the International Physical Activity Questionnaire (IPAQ) [13]. The Clinical Research Ethics Committee, University College Cork, Ireland, granted ethical approval for this research.

Data management & Statistical analysis

Data were scanned, checked and verified using TeleForm TM scanning processes. The estimated error rate for data entry was 0.06% based on manual checking of a 10% sample of all scanned questionnaires. WEMWBS, AUDIT-C and IPAQ scores were computed in accordance with instrument guidelines. WEMWBS scores were divided into categories of mental well-being as defined by Brauholtz et al [14]. Below average mental wellbeing was defined as a WEMWBS score of more than one standard deviation below the mean, average mental wellbeing was within one standard deviation of the mean and above average mental wellbeing was over one standard deviation above the mean. BMI was estimated from self-reported height and weight. Data were analysed using IBM SPSS Statistics 20. One-way ANOVAs, univariate and multivariate logistic regression analysis were utilised to investigate factors associated with above average mental health and well-being. Multivariate logistic regression, applying backward elimination, was undertaken to investigate the combination of risk factors associated with above average mental health and well-being. Risk factors appearing as statistically significant ($P < 0.20$) in the univariate logistic regressions were used to obtain an initial multivariable logistic regression fit.

Results

Baseline characteristics of the student sample are presented by gender in Table 1. Faculty, year in college and accommodation differed significantly by gender ($p < 0.05$). As expected, women were more likely to be taking Health and Welfare undergraduate degrees. Over 40% of the sample were in first year. Gender distribution varied slightly by college year with a higher proportion of women in second year. Men were less likely to be living in campus accommodation (8.93% vs. 14.9%) and more men reported living at home with their parents than women (47.74% vs. 39.2%).

No statistically significant differences were observed between men and women in relation to age, physical activity, alcohol consumption or smoking. Over 40% of students reported moderate levels of physical activity, while in excess of 65% of students reported hazardous alcohol consumption. Over a quarter of students reported having smoked 100 cigarettes in their lives. Men were significantly more likely to report six or more sexual partners compared to women (22.5% vs. 13.6%; $p < 0.001$). In addition, men were more likely to report taking at least one form of illicit drug use during the previous 12 months (40.4% vs. 26.4%; $p < 0.001$).

WEMWBS scores were normally distributed in this population (Figure 1). Mean WEMWBS scores were slightly higher in men than in women ($p = 0.015$). Distributions of men and women's WEMWBS scores, categorised as below

Table 1. Characteristics of students sociodemographic factors and WEMWBS scores by gender [N (%)/Mean (SD)].

		Males (N=755)	Females (N=1,267)	Chi-Square
Age	<=18	96 (12.9%)	163 (13.2%)	0.077
	19	248 (33.3%)	382 (30.9%)	
	20	145 (19.5%)	278 (22.5%)	
	21	93 (12.5%)	189 (15.3%)	
	22+	162 (21.8%)	225 (18.2%)	
	Faculty	Science/Engineering	235 (31.4%)	
Arts/Social Science/Education		268 (35.8%)	501 (40%)	
Law & Business		174 (23.3%)	234 (18.7%)	
Health & Welfare		57 (7.6%)	228 (18.2%)	
Other		14 (1.9%)	16 (1.3%)	
College Year		First	400 (53.1%)	548 (43.3%)
	Second	178 (23.6%)	379 (29.9%)	
	Third	130 (17.2%)	239 (18.9%)	
	Fourth	46 (6.1%)	101 (8.0%)	
	Accommodation	Campus Accommodation	67 (8.93%)	188 (14.9%)
Rented House/Flat		289 (38.53%)	531 (42.2%)	
Parents' House		358 (47.74%)	494 (39.2%)	
House Owner		36 (4.8%)	46 (3.7%)	
Physical Activity	Low	216 (29.2%)	386 (31%)	0.693
	Moderate	324 (43.8%)	529 (42.5%)	
	High	200 (27%)	330 (26.5%)	
Smoking	Yes	201 (27%)	320 (25.6%)	0.5
	No	544 (73%)	929 (74.4%)	
Alcohol Consumption	Hazardous consumption	492 (65.2%)	855 (67.5%)	0.285
	Non-hazardous consumption	263 (34.8%)	412 (32.5%)	
Drug use	Yes	305 (40.4%)	334 (26.4%)	<0.001
	No	450 (59.6%)	933 (73.6%)	

Table 1 (continued).

		Males (N=755)	Females (N=1,267)	Chi-Square
No. of sexual partners	None	168 (22.8%)	237 (19.8%)	<0.001
	1-3	324 (44%)	623 (51.9%)	
	4-5	79 (10.7%)	177 (14.8%)	
	6+	166 (22.5%)	163 (13.6%)	
Mental Health & Well-being	Below average	91 (12.1%)	182 (14.4%)	0.311
	Average	545 (72.3%)	901 (71.1%)	
	Above average	118 (15.6%)	184 (14.5%)	
	WEMWBS mean Score	50.2 (8.4)	49.3 (8.4)	0.015

doi: 10.1371/journal.pone.0074921.t001

average, average and above average mental well-being, are displayed in Table 1.

Analysis of variance was undertaken to investigate whether significant differences in WEMWBS scores were evident among different socio-demographic, social and lifestyle factors reported by students. Analysis was conducted separately for men and women (Table 2).

Among men, mean WEMWBS scores were similar in relation to age group, area of study, accommodation, smoking status, drug use, BMI and number of sexual partners. Mean WEMWBS scores differed significantly by year of study, alcohol use and physical activity. Third year students reported higher average WEMWBS scores than their first year (52.36 vs. 49.91; $p=0.012$), second year (52.36 vs. 49.39; $p=0.012$) and fourth year (52.36 vs. 50.2; $p=0.012$) counterparts. Mean WEMWBS scores were higher among hazardous alcohol consumers versus non-hazardous consumers (51.07 vs. 48.65; $p<0.001$). Evidence of a non-linear relationship between physical activity and mental health and well-being was recorded. Men reporting moderate physical activity have significantly lower WEMWBS scores than men reporting low levels of physical activity (49.37 vs. 51.53; $p=0.013$).

Among women, mean WEMWBS scores differed by age with those aged 22 and older reporting better mental health and well-being than their female counterparts aged 20. However, there was no difference in WEMWBS scores for lifestyle or socio-demographic characteristics among women.

Logistic regression

Univariate logistic regression, stratified by gender, was undertaken to uncover risk factors associated with above average mental health and well-being. Among men, those who reported hazardous alcohol consumption and one or more lifetime sexual partners were more likely to report above

Distribution of WEMWBS scores among the third level student population




Figure 1. Distribution curve for WEMWBS scores among a third level student population.

doi: 10.1371/journal.pone.0074921.g001

average mental health and well-being. Those reporting low or moderate levels of physical activity were less likely to report above average mental health and well-being. Among women those reporting 1-3 or 6 or more lifetime sexual partners were more likely to report above average mental health and well-being as were those reporting illegal drug use. Results of univariate regression can be seen in Table 3. Covariates with a p-value less than 0.2 were included in the multivariable model and included illegal drug use, level of physical activity, alcohol use, BMI and number of lifetime sexual partners for men and year of study, illegal drug use and number of sexual partners for women.

Multivariate logistic regression, stratified by gender, was utilised to investigate the variables associated with positive mental health (above average mental well-being score). Controlling for age, there was an apparent inverse association between physical activity levels and the WEMWBS score among men. Men reporting both moderate and high levels of physical activity were significantly less likely to have an above average WEMWBS score relative to those with low physical activity. Similarly, men reporting 1-3, 4-5 or 6 or more sexual partners in their lifetime were significantly more likely to have above average mental health and well-being scores relative to those with no sexual partners. Among women, those with 1-3 or six or more sexual partners in their lifetime were over twice as likely to report above average mental health and well-being. However, no association between those reporting 4-5 sexual partners and mental health and well-being was observed.

Results for multivariate logistic regression are presented in Table 4.

Discussion

This is one of the first studies to examine mental health and well-being scores in a broad, largely representative sample of third level students and the first to report on the factors associated with positive mental health and well-being using WEMWBS. The findings from this study suggest that students with a relatively adverse health and lifestyle profile have higher than average mental health and well-being. Significant, but relatively small differences in mental health and well-being scores were observed between men and women. However, no association between alcohol consumption, obesity, drug use or smoking status and positive mental health and well-being were detected in this research.

As highlighted by Cranford et al, the effect of alcohol use on mental health and well-being is unclear [15]. The current research did not uncover an association between alcohol use and positive mental health and well-being adding to the conflicting results of previous research. The Harvard college alcohol study found that poor mental health was associated with an increased risk of alcohol use. However, other research has reported positive associations between mental health and alcohol abuse with one study indicating "that associations

Table 2. WEMWBS scores by Gender & Socio-demographic factors.

			Males		Females		
			Mean WEMWBS score (95% CI)	p-value (ANOVA)	Mean WEMWBS score (95% CI)	p-value (ANOVA)	
Socio-demographic Indicators	Age group	= < 18	49.58 (48.07-51.1)	p=0.45	49.4 (48.17-50.63)	p=0.04	
		19	50.56 (49.53-51.6)		48.93 (48.11-49.75)		
		20	50.55 (49.05-52.04)		48.46 (47.47-49.45)		
		21	51.17 (49.71-52.64)		49.52 (48.27-50.78)		
		22+	49.46 (48.08-50.85)		50.7 (49.6-51.8)		
	Year of study	1 st year	49.91 (49.11-50.7)	p=0.012	49.5 (48.81-50.2)	p=0.13	
		2 nd year	49.39 (48.07-50.71)		48.89 (48.04-49.73)		
		3 rd year	52.36 (50.96-53.77)		48.74 (47.64-49.84)		
		4 th year	50.2 (49.62-50.82)		50.83 (49.2-52.46)		
	Area of study	Science/Engineering	50.34 (49.26-51.41)	p=0.44	49.32 (48.27-50.38)	p=0.32	
		Arts/ Social Science/ Education	49.61 (48.63-50.58)		48.87 (48.13-49.6)		
		Law/Business	50.61 (49.31-51.9)		49.65 (48.58-50.72)		
		Health & Welfare	51.33 (49.13-53.54)		49.83 (48.81-50.86)		
		Other	52.36 (47.19-57.52)		46.25 (40.83-51.67)		
	Accommodation	House Owner	47.58 (45.41-49.76)	p=0.15	48.85 (46.28-51.41)	p=0.52	
		Parents House	50.07 (49.17-50.98)		49.26 (48.51-50.01)		
		Rented House	50.45 (49.49-51.42)		49.05 (48.34-49.76)		
		Campus Accommodation	51.45 (49.65-53.25)		50.09 (48.86-51.31)		
	Lifestyle indicators	Smoking Status	Smoker	50.27 (49.08-51.46)	p=0.89	48.54 (47.56-49.52)	p=0.06
			Non -smoker	50.18 (49.47-50.88)		49.59 (49.06-50.12)	
Drug Use-Ever		Yes	50.16 (49.18-51.13)	p=0.86	49.23 (48.26-50.2)	p=0.9	
		No	50.27 (49.51-51.03)		49.3 (48.77-49.83)		
Alcohol Use		Hazardous	51.07 (50.36-51.77)	p<0.001	49.37 (48.82-49.93)	p=0.58	
		Non-Hazardous	48.65 (47.56-49.73)		49.09 (48.25-49.94)		
Physical Activity		Low	51.53 (50.39-52.66)	p=0.013	49.02 (48.17-49.87)	p=0.76	
		Moderate	49.37 (48.46-50.29)		49.43 (48.73-50.14)		
		High	50.07 (49.58-50.8)		49.29 (48.39-50.19)		
BMI		Underweight	49.14 (45.47-52.81)	p=0.89	48.79 (47.2-50.38)	p=0.76	
		Normal Weight	50.12 (49.38-50.85)		49.4 (48.8-50.0)		
		Overweight	50.47 (49.00-51.93)		48.82 (47.46-50.19)		
		Obese	50.6 (49.67-53.53)		49.95 (46.94-52.96)		
No. of sexual partners		None	48.77 (47.52-50.02)	p=0.07	48.15 (47.13-49.17)	p=0.17	
		1-3	50.69 (49.8-51.57)		49.35 (48.7-49.99)		
	4-5	51.05 (49.22-52.88)		49.81 (48.56-51.06)			
	6 or more	50.51 (49.09-51.94)		49.43 (47.97-50.89)			

doi: 10.1371/journal.pone.0074921.t002

between mental disorders and alcohol involvement among college students emerged only for alcohol dependence" [15].

When Needham et al investigated the effect of weight status on mental health and well-being among adolescents, the association between weight and mental health was context specific. Young girls and adolescents in lower grades experienced being overweight more negatively. Needham concludes that the rising rates of obesity among adolescents may act as a buffer to the stigma associated with being overweight and obese in this population [16]. This study found no association between BMI status and mental health and well-being in either men or women. This may be due to rates of overweight and obesity continuing to rise in Ireland. As more

individuals become overweight stigma reduces, impacting less on the individuals mental health and well-being.

Research investigating the effect of smoking on mental health and well-being were more unified with strong associations between nicotine dependence and mood and anxiety disorders being evidenced [15]. In 2004, Patterson et al conducted a review of current research in the area concluding that symptoms of anxiety and depression are associated with cigarette smoking among American college students [17]. However, the current research is at odds with pre-existing knowledge in the area. No association between cigarette smoking and mental health and well-being in the current study

Table 3. Univariate Logistic Regression: Risk factors associated with above average mental health and well-being.

		Males			Females		
		OR	95%CI	p-value	OR	95%CI	p-value
Age group	= < 18	1.00		0.06	1.00		0.31
	19	2.82	1.22-6.49		0.86	0.51-1.49	
	20	3.29	1.38-7.83		0.88	0.5-1.54	
	21	1.89	0.71-5.02		1.24	0.69-2.22	
	22+	2.1	0.87-5.11		1.32	0.75-2.3	
Year of study	1 st year	1.00		0.33	1.00		0.16
	2 nd year	1.16	0.71-1.91		0.88	0.6-1.28	
	3 rd year	1.57	0.94-2.63		0.86	0.55-1.34	
	4 th year	1.53	0.7-3.34		1.61	0.95-2.72	
Area of study	Science/Engineering	1.00		0.31	1.00		0.99
	Arts/ Social Science/ Education	0.68	0.41-1.13		0.92	0.61-1.39	
	Law/Business	1.13	0.67-1.88		0.91	0.57-1.49	
	Health & Welfare	1.2	0.57-2.52		0.97	0.6-1.59	
	Other	1.37	0.37-5.14		0.79	0.17-3.62	
Accommodation	House Owner	0.38	0.1-1.43		0.76	0.3-1.95	
	Parents House	0.72	0.37-1.41		0.86	0.55-1.37	
	Rented House	0.83	0.42-1.63		0.82	0.52-1.3	
	Campus Accommodation	1.00		0.47	1.00		0.85
Smoking Status	Smoker	1.01	0.65-1.59		0.91	0.63-1.32	
	Non -smoker	1.00		0.96	1.00		0.63
Drug Use-Ever	Yes	1.35	0.91-2.01		1.47	1.05-2.06	
	No	1.00		0.14	1.00		0.02
Alcohol Use	Hazardous	1.69	1.09-2.64		1.02	0.73-1.43	
	Non-Hazardous	1.00		0.02	1.00		0.89
Physical Activity	Low	1.00		0.009	1.00		0.76
	Moderate	0.49	0.31-0.78		1.11	0.77-1.62	
	High	0.61	0.37-1.02		0.97	0.63-1.49	
BMI	Underweight	1.00		0.11	1.00		0.4
	Normal Weight	1.65	0.38-7.22		1.33	0.7-2.51	
	Overweight	2.82	0.63-12.69		0.95	0.43-2.11	
	Obese	2.31	0.46-11.71		1.85	0.7-4.95	
No. of sexual partners	None	1.00		0.02	1.00		0.009
	1-3	2.25	1.21-4.18		2.04	1.22-3.42	
	4-5	2.79	1.29-6.06		1.8	0.95-3.4	
	6 or more	2.73	1.4-5.32		2.8	1.53-5.15	

doi: 10.1371/journal.pone.0074921.t003

was observed. Smoking habits are regularly underreported [18], an issue which may be skewing the results of this study.

Previous research investigating the effect of recreational drug use on mental health and well-being has provided conflicting results. In 2009, Cranford et al reviewed the literature surrounding mental health and well-being and recreational drug use [15]. This review provided differing results as to the relationship between recreational drug use and mental health and well-being. Some research reported a positive association between drug use and mental health and anxiety while others report a negative effect [15,19]. This study finds that recreational drug use is not associated with mental health and well-being among a third level student population.

Low levels of physical activity and one or more sexual partners in a lifetime were all associated with positive mental health and well-being in the current research. In 2005, Penedo

undertook a review investigating the benefits of physical activity. This review identified a reduction in anxiety and depression among college students who completed physical activity [20]. Specifically, physical activity was associated with improved mood and a reduction in the symptoms of depression or anxiety. Several research studies have solidified this result [21–25]. This is, however, at odds with the current research where low levels of physical activity are associated with increased positive mental health and well-being among men. Levels of physical activity reported among our student population were relatively high, measured using the International Physical Activity Questionnaire (IPAQ). Despite the fact that the IPAQ has been extensively validated and recommended as an effective method of assessing physical activity participation [26], there are some limitations associated with its use. According to Saelens and Sallis (2003) one of the

Table 4. Multivariate Logistic Regression: Risk factors associated with above average mental health and well-being (controlling for age).

Males			
	OR	95% CI	p-value
Physical Activity (IPAQ)			
<i>Low</i>	1.00		p=0.008
<i>Moderate</i>	0.48	0.3-0.77	
<i>High</i>	0.59	0.35-0.99	
No. of sexual partners			
<i>None</i>	1.00		p=0.004
<i>1-3</i>	2.29	1.22-4.28	
<i>4-5</i>	2.98	1.32-6.72	
<i>6 or more</i>	3.64	1.78-7.43	
Females			
Year of Study			
<i>1st year</i>	1.00		p=0.03
<i>2nd year</i>	0.7	0.46-1.06	
<i>3rd year</i>	0.75	0.46-1.21	
<i>4th year</i>	1.57	0.90-2.74	
No. of sexual partners			
<i>None</i>	1.00		p=0.008
<i>1-3</i>	2.18	1.28-3.72	
<i>4-5</i>	1.86	0.96-3.61	
<i>6 or more</i>	2.96	1.56-5.62	

*. The initial model included: Age, BMI, physical activity, alcohol use, no. of sexual partners and drug use for men and age, year of study, drug use and number of sexual partners for females

doi: 10.1371/journal.pone.0074921.t004

main limitations associated with IPAQ arises due to the fact that self-reported measures can often result in the over estimation of physical activity participation [27]. In fact, it has been suggested that the IPAQ may over estimate physical activity participation to a greater extent than other physical activity surveys [28,29]. Another limitation of the IPAQ is that survey participants can often experience difficulties distinguishing between moderate and vigorous intensity activities [30], a factor which can also result in the over estimation of physical activity. It is possible that these limitations affected the findings in our study.

In 2006, Martens et al reported on the differences between actual and perceived student norms in relation to sexual behaviour. Martens discovered that "students overestimated the sexual activity of the typical student" [31]. Sexual health is regularly viewed as a risk factor but is also a positive aspect of life. Some studies have reported high numbers of sexual partners as a normative occurrence among the third level

References

- Adlaf EM, Gliksman L, Demers A, Newton-Taylor B (2001) The prevalence of elevated psychological distress among Canadian undergraduates: findings from the 1998 Canadian Campus Survey. *J Am Coll Health* 50: 67-72. doi:10.1080/07448480109596009. PubMed: 11590985.
- Bewick B, Koutsopoulou G, Miles J, Slaa E, Barkham M (2010) Changes in undergraduate students' psychological well-being as they progress through university. *Stud Higher Educ* 35: 633-645. doi: 10.1080/03075070903216643.

student population [32,33]. Although risky behaviour, adherence to contraception and protection against sexually transmitted diseases should be of concern in this group, students may feel happier as they align themselves to the perceived sexual norms of their peers. Our finding of multiple lifetime sexual partners being associated with higher than average mental health supports this view.

Strengths & Weaknesses

A strength of the current research is the rigorous sampling strategy utilised so that each degree programme, regardless of size, would have an equal opportunity of being included in the study. Also, student demographics from the overall institution where the study took place are similar to those sampled in relation to year in college, faculty and gender.

There are a number of potential sources of bias in cross-sectional studies including response bias and self-report bias, which may have affected the outcome of the study. Students who attend lectures may differ significantly from their non-attending peers in relation to their mental health and risk-taking behaviours, a concern which should be considered when interpreting results. When interpreting these results, the response rate of 51% should be taken into consideration. Though the response rate is moderate, it is in line with previous surveys undertaken both in Ireland [3] and abroad [34].

Conclusion

The findings of this study are somewhat unusual in that students with an adverse lifestyle pattern have a higher than average mental health and well-being. Students represent a unique subgroup that is relatively privileged when compared to the general population [35]. Findings from this study may not be applicable to the general population. Previous research suggests that hazardous alcohol consumption and a high number of lifetime partners are social norms in this culture, meaning those closer aligned with the norms for their culture may report more positive mental health and well-being. This is evidenced in the current study. There is a need to replicate this research yielding a higher response rate to confirm the factors associated with positive mental health and well-being among third level students. A study of the general population is also warranted.

Author Contributions

Conceived and designed the experiments: MPD EF FS IJP. Performed the experiments: MPD EF IJP. Analyzed the data: MPD EF FS IJP. Contributed reagents/materials/analysis tools: MPD EF FS IJP. Wrote the manuscript: MPD EF FS IJP.

3. Hope A, Dring C, Dring J (2005) College lifestyle and attitudinal national (CLAN) survey. The health of Irish students. Dublin: Health Service Executive.
4. Stallman HM (2010) Psychological distress in university students: A comparison with general population data. *Aust Psychol* 45: 249-257. doi:10.1080/00050067.2010.482109.
5. Tosevski DL, Milovancevic MP, Gajic SD (2010) Personality and psychopathology of university students. *Curr Opin Psychiatry* 23: 48–52. doi:10.1097/YCO.0b013e328333d625. PubMed: 19890212.
6. Zivin K, Eisenberg D, Gollust SE, Golberstein E (2009) Persistence of mental health problems and needs in a college student population. *J Affect Disord* 117: 180-185. doi:10.1016/j.jad.2009.01.001. PubMed: 19178949.
7. Nagel T (1972) Aristotle on eudaimonia. *Phronesis* 17: 252-259. doi: 10.1163/156852872X00079.
8. Seligman MEP, Csikszentmihalyi M (2000) Positive psychology: an introduction. *American Psychol* 55: 5.
9. World Health Organization (2001) The world health report 2001 - Mental Health: New Understanding, New Hope. Geneva: World Health Organization.
10. Seligman M (2002) Authentic happiness: Using the new positive psychology to realize your potential for lasting fulfillment. Free Press.
11. World Health Organization (1986) World Health Organization, Ottawa charter for health promotion. Geneva. pp. 1-4.
12. Tennant R, Hiller L, Fishwick R, Platt S, Joseph S et al. (2007) The Warwick-Edinburgh mental well-being scale (WEMWBS): development and UK validation. *Health Qual Life Outcomes* 5: 63. doi: 10.1186/1477-7525-5-63. PubMed: 18042300.
13. IPAQ Research Committee (2005) Guidelines for data processing and analysis of the International Physical Activity Questionnaire (IPAQ)-short and long forms. Available., September 17, 2008
14. Brauhnoltz S, Davidson S, Myant K, O'Connor R (2007) Well? What do you think?(2006): The third national Scottish survey of public attitudes to mental health, mental wellbeing and mental health problems: Scottish Government.
15. Cranford JA, Eisenberg D, Serras AM (2009) Substance use behaviors, mental health problems, and use of mental health services in a probability sample of college students. *Addict Behav* 34: 134-145. doi: 10.1016/j.addbeh.2008.09.004. PubMed: 18851897.
16. Needham BL, Crosnoe R (2005) Overweight status and depressive symptoms during adolescence. *J Adolesc Health* 36: 48-55. doi: 10.1016/j.jadohealth.2003.12.015. PubMed: 15661596.
17. Patterson F, Lerman C, Kaufmann VG, Neuner GA, Audrain-McGovern J (2004) Cigarette smoking practices among American college students: review and future directions. *J Am Coll Health* 52: 203-212. doi:10.3200/JACH.52.5.203-212. PubMed: 15029942.
18. Han M-Y, Chen W-Q, Wen X-Z, Liang C-H, Ling W-H (2012) Differences of Smoking Knowledge, Attitudes, and Behaviors Between Medical and Non-medical Students. *Int J Behav Med* 19: 104-110. doi: 10.1007/s12529-010-9140-7. PubMed: 21246317.
19. Stewart SH, Karp J, Pihl RO, Peterson RA (1997) Anxiety sensitivity and self-reported reasons for drug use. *J Subst Abuse* 9: 223-240. doi: 10.1016/S0899-3289(97)90018-3. PubMed: 9494951.
20. Penedo FJ, Dahn JR (2005) Exercise and well-being: a review of mental and physical health benefits associated with physical activity. *Curr Opin Psychiatry* 18: 189–193. doi: 10.1097/00001504-200503000-00013. PubMed: 16639173.
21. Brosnahan J, Steffen LM, Lytle L, Patterson J, Boostrom A (2004) The relation between physical activity and mental health among Hispanic and non-Hispanic white adolescents. *Arch Pediatr Adolesc Med* 158: 818–823. doi:10.1001/archpedi.158.8.818. PubMed: 15289257.
22. Cramer SR, Nieman DC, Lee JW (1991) The effects of moderate exercise training on psychological well-being and mood state in women. *J Psychosom Res* 35: 437-449. doi: 10.1016/0022-3999(91)90039-Q. PubMed: 1920175.
23. Hayes D, Ross CE (1986) Body and mind: The effect of exercise, overweight, and physical health on psychological well-being. *J Health Soc Behav*, 27: 387-400. PubMed: 3559130.
24. Kirkcaldy BD, Shephard RJ, Siefen RG (2002) The relationship between physical activity and self-image and problem behaviour among adolescents. *Soc Psychiatry Psychiatr Epidemiol* 37: 544-550. doi: 10.1007/s00127-002-0554-7. PubMed: 12395145.
25. Scully D, Kremer J, Meade MM, Graham R, Dudgeon K (1998) Physical exercise and psychological well being: a critical review. *Br J Sports Med* 32: 111-120. doi:10.1136/bjism.32.2.111. PubMed: 9631216.
26. Lee PH, Macfarlane DJ, Lam TH, Stewart SM (2011) Validity of the international physical activity questionnaire short form (IPAQ-SF): A systematic review. *Int J Behav Nutr Phys Act* 8: 115. doi: 10.1186/1479-5868-8-115. PubMed: 22018588.
27. Saelens BE, Sallis JF, Black JB, Chen D (2003) Neighborhood-based differences in physical activity: an environment scale evaluation. *J Info* 93: 1552–8. PubMed: 12948979.
28. Ainsworth BE, Bassett DR Jr, Strath SJ, Swartz AM, O'Brien WL et al. (2000) Comparison of three methods for measuring the time spent in physical activity. *Med Sci Sports Exerc* 32: S457–S464. doi: 10.1097/00005768-200009001-00004. PubMed: 10993415.
29. Rzewnicki R, Auweele YV, De Bourdeaudhuij I (2003) Addressing overreporting on the International Physical Activity Questionnaire (IPAQ) telephone survey with a population sample. *Public Health Nutr* 6: 299-306. PubMed: 12740079.
30. Bauman A, Bull F, Chey T, Craig CL, Ainsworth BE et al. (2009). *Int J Behav Nutr Physical Act International J Behav Nutr Physical Act* 6: 21. doi:10.1186/1479-5868-6-21.
31. Martens MP, Page JC, Mowry ES, Damann KM, Taylor KK et al. (2006) Differences between actual and perceived student norms: An examination of alcohol use, drug use, and sexual behavior. *J Am Coll Health* 54: 295-300. doi:10.3200/JACH.54.5.295-300. PubMed: 16539222.
32. Desiderato LL, Crawford HJ (1995) Risky sexual behavior in college students: Relationships between number of sexual partners, disclosure of previous risky behavior, and alcohol use. *J Youth Adolesc* 24: 55-68. doi:10.1007/BF01537560.
33. Buhi ER, Marhefka SL, Hoban MT (2010) The state of the union: Sexual health disparities in a national sample of US college students. *J Am Coll Health* 58: 337-346. doi:10.1080/07448480903501780. PubMed: 20159757.
34. Kuo M, Adlaf EM, Lee H, Gliksman L, Demers A et al. (2002) More Canadian students drink but American students drink more: comparing college alcohol use in two countries. *Addiction* 97: 1583-1592. doi: 10.1046/j.1360-0443.2002.00240.x. PubMed: 12472642.
35. Hunt J, Eisenberg D (2010) Mental health problems and help-seeking behavior among college students. *J Adolesc Health* 46: 3-10. doi: 10.1016/S1054-139X(09)00612-0. PubMed: 20123251.