

|                             |   |
|-----------------------------|---|
| Title                       | Young students' use of the internet for mental health information and support   |
| Authors                     | Horgan, Aine M.;Sweeney, John F.  |
| Publication date            | 2009-09-24  |
| Original Citation           | Horgan, A. and Sweeney, J. F. (2010) 'Young students' use of the internet for mental health information and support', Journal of Psychiatric and Mental Health Nursing, 17(2), pp. 117-123. <a href="http://dx.doi.org/10.1111/j.1365-2850.2009.01497.x">http://dx.doi.org/10.1111/j.1365-2850.2009.01497.x</a>   |
| Type of publication         | Article (peer-reviewed)   |
| Link to publisher's version | <a href="http://dx.doi.org/10.1111/j.1365-2850.2009.01497.x">10.1111/j.1365-2850.2009.01497.x</a>   |
| Rights                      | © 2009 Blackwell Publishing. This is the peer reviewed version of the following article: Horgan, A. and Sweeney, J. F., Journal of Psychiatric and Mental Health Nursing, 17: 117–123, which has been published in final form at <a href="http://dx.doi.org/10.1111/j.1365-2850.2009.01497.x">http://dx.doi.org/10.1111/j.1365-2850.2009.01497.x</a> . This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Self-Archiving. |
| Download date               | 2024-04-24 22:20:21   |
| Item downloaded from        | <a href="https://hdl.handle.net/10468/3006">https://hdl.handle.net/10468/3006</a>   |

## **Title**

Young students' use of the internet for mental health information and support

## **Abstract**

It is recognised that young people experience difficulties in accessing mainstream mental health services particularly due to the stigma that remains associated with mental health problems. One potential solution is to use the many websites available offering information and support for mental health problems, such support and information could be offered by Psychiatric Nurses. However young peoples' usage and views on using the internet for this purpose has yet to be examined. This quantitative descriptive study aimed to elicit the views of 922 University students, aged between 18 and 24 years, on using the internet for mental health information and support. Data was collected using a 30 item self-designed questionnaire and analysed using descriptive statistics. The findings indicated that 72.4% of participants used the internet several times a day. In addition 30.8% had previously searched for mental health information online, predominantly on depression. Whilst it was found that 68% of participants indicated that they would use the internet for mental health support if they needed to, 79.4% would still prefer face to face support. It is concluded that young people are willing to use the internet for mental health information and that it represents a viable source of support for this age group.

## **Introduction and Background**

Using the internet for health information has become a popular choice amongst the general population in recent times. The Central Statistics Office (2008) in Ireland reported that 62% of Irish people have access to the internet in their own homes. This is similar to international trends, for example a Parisian study found that 70% of people had internet access (Renahy et al. 2008). The Health Service Executive (2007) reported that 18% of Irish people have used the internet for health information, a much lower figure than that reported by Renahy et al. (2008) who found that 49% had used the internet previously for health information. Gallagher *et al.* (2008) reported that those seeking health information online in Ireland were most likely to be female and in employment, those least likely were males of older age and the unemployed. A high percentage reported willingness to use the internet for health

information, this however decreased with age possibly due to access to the internet. Rice (2006) analysed the results of seven American databases, two in detail, to determine what influences people to search for health information online. Similar to Gallagher's *et al.* (2008) results, it was found that the most likely people to search for health information online were female and in employment, those who have received a new diagnosis and those who want to help others cope with health related problems.

With the growth in health related websites, numerous studies have been conducted in the United States, Europe and Canada, on how people use the internet for general health information (Health on the Net Foundation 2001, eUSER 2005, Skinner *et al.* 2003), however few have focussed on young people. Their findings indicated that large numbers of people use the internet for information on illnesses, treatment and support. Indeed the use of the internet for support is well documented, particularly in the area of cancer care (Lieberman *et al.* 2003, Lieberman 2005, Klemm 2005, Hoybye *et al.* 2005).

Concerns have been raised as to the quality of health information available online and as such it has been quite extensively studied. Eysenbach *et al.* (2002) conducted a systematic review of 79 research studies that evaluated the quality of health information online. While no conclusive evidence was found due to the quality of the studies, it was reported that 70% of the studies concluded that the quality of health information online was a problem.

The use of the internet for mental health information is less certain. It is estimated that one fifth of young adults have used the internet as a source of mental health information (Gould *et al.* 2002). In the UK, a study by Powell & Clarke (2006) reported that 24% of their total sample identified the internet as one of the top three sources of information on mental health. Additionally, Powell *et al.* (2003) identified that 49% of people who use the internet for mental health information had not sought formal support from the health services. Indeed, Gallagher *et al.* (2008) reported that there is an increase in the number of people searching for mental health information online. A survey in the UK found that more than one in four people would rather use the internet for advice and counselling about depression than visit their family doctor (Graham *et al.* 2000). Indeed, Ybarra & Eaton (2005) suggested that people with depressive symptoms and suicidal ideation prefer computer based assessments and interventions. The reasons for this are multifaceted but include an ability to remain anonymous and explore and share feelings on a more intimate level (Shaw *et al.* 2000) while feeling less stigmatised for gaining mental health information and support (Johnsen *et al.* 2002).

There is a high prevalence of mental health problems amongst 18-24 year olds (Andrews *et al.* 1999). This is problematic given that, some research suggests that young people demonstrate low levels of help seeking behaviour in relation to their mental health (Sullivan *et al.* 2004, Rickwood *et al.* 2007). Reasons for this are complex. For example, Donald *et al.* (2000) reported that in young adults aged 15-24 years, 39% of males and 22% of females would not seek help from formal services for personal, emotional or distressing problems. There is further support for this perspective since Berger *et al.* (2005) found that help seeking for mental health problems is often delayed because of stigma and that the internet is a means of encouraging people to seek help. It is also argued that engaging in help for mental health problems at a young age is regarded as a protective factor and early treatment during adolescence and young adulthood are essential to aid long term prognosis (Rickwood *et al.* 2007).

Gallagher *et al.* 's (2008) findings also reported that a significant number of people who seek health information online have experienced mental health problems; they argue that this raises questions as to whether the internet should be used for mental health care delivery. Moreover, Gallagher *et al.* (2008) contend that further research is needed to ensure the needs of people who seek health information online are met.

Many websites exist offering information and support for mental health problems for young people, for example [www.spunout.ie](http://www.spunout.ie) ; however studies of young peoples' usage and views on these services are not readily available. In summary, the fact that young people demonstrate low levels of help seeking behaviour for mental health problems, while being active users of the internet where services exist to offer them mental health information, suggests that it is a potentially viable alternative to face-to-face support. Nevertheless, little is known about what types of mental health information young people search for and how young people view using the internet for mental health support.

## **Methodology**

This study aimed to determine how young people use the internet for mental health information and support using a quantitative descriptive approach as a prelude to a more in-depth study. As this study was explorative and aimed to gain descriptive information on how young people use the internet without addressing causality, this was deemed an appropriate approach.

### *Data collection and analysis*

Data was collected through the use of an online questionnaire. This questionnaire was designed by the researcher based on themes identified from the literature. An extensive review of the literature was undertaken, this information allowed for the formation of the questions used in this study. Measures taken to enhance the validity of the questionnaire included the review of the questionnaire by a statistician to determine if any questions needed to be rephrased to help with statistical analysis following data collection, by two nurse researchers and three young people in the target age group.

The questionnaire consisted of 30 items. The first four questions elicited demographic details of age, gender, subject of academic study and type of accommodation. These questions were deemed important as the University where the study was undertaken has four colleges, it was deemed important to ascertain in which college the students were studying as if the majority of students were studying in the College of Medicine and Health it may have skewed the findings. It was also deemed important to ascertain the living accommodation of the participants to provide insight into where the participants used the internet. The next 13 questions focused on how they used the internet, including frequency, use of search engines, use of social networking sites and information on how they use the internet to search for general health information. The remaining questions focussed on how often they used the internet for mental health information, what types of mental health information they searched for and their views on using the internet for mental health support. The questionnaire contained a variety of different types of questions from multiple choice questions to likert scales, with a number of questions allowing for qualitative data to be entered.

The questionnaire was emailed to all students in one University in Ireland. The email contained information about the study and a link to the survey for those interested in participating. Once the participant clicked on the link they were first asked if they met the inclusion criteria, which were, being between the ages of 18 and 24 years and not a nursing or midwifery student. Nursing and Midwifery students were excluded from the study as the researcher was directly involved in their education. An online consent form was posted and the potential participants were asked to consent to participate in the study by ticking a 'yes' or 'no' button. Only those who ticked 'yes' were able to access the questionnaire. The

questionnaire was designed in electronic format using Survey Monkey (surveymonkey.com, Oregon) and was left open for a period of one week. This software also provided descriptive statistics. Given the number of respondents, the qualitative comments were analysed using content analysis (Burns & Grove 2001), whereby phrases and words that were repeated were counted to provide a numerical value for each statement made.

### ***Ethical issues***

Ethical approval to conduct this study was granted by the University's Social Research Ethics Committee. The participants were informed that participation was voluntary, that data would be kept confidential to the researcher and that their anonymity would be protected. An online consent form was posted and only those who gave consent were able to complete the questionnaire, participants were not required to provide their name. Computer IP addresses were not stored by the researcher to protect their anonymity.

## **Results**

### **The sample**

The sample consisted of University students from one University in Ireland. One thousand, two hundred and five people commenced the survey; fifty one people who did not meet the inclusion criteria were excluded from the study. Of the eligible respondents a number of people failed to complete the survey, resulting in the final sample being nine hundred and twenty-two people. However, there were fewer numbers in the data producing sample as not all participants answered every question; the number who answered each question is provided in the description of the findings. Details of the participant's age, gender, area of study and accommodation are provided in Table 1.

### **Use of the internet**

Of the 883 participants who completed this question, the findings indicated that 72.4% (n=639) of participants used the internet several times a day, 16.2% (143) once a day and the remainder several times a week (11.1%, n=98), once a week (0.2%, n=2) or less than once a week (0.1%, n=1). The participants (n=884) reported that the majority of them accessed the internet at college (94.9%, n=839) or at home (91.7%, n=811). They also accessed the

internet at work (8.9%, n=79), at friends and family's house (24.7%, n=218), on their phone (n=4) and in internet cafes (n=4).

### **Use of the internet for mental health information and support**

Of the 867 participants who responded to this question, the findings indicated that 30.8% (n=267) of participants used the internet for mental health information. Out of those who responded, 8.2% (n=22) used it once a week or more, 15.4% (n=41) once a month, 36.7% (n=98) a few times a year and 39.7% (106) less than a few times a year.

Reasons for using the internet for mental health information were provided by 220 participants; these are reported in Table 2.

Reasons for not using the internet for mental health information were provided by 472 participants. The majority (n=316, 66.9%) reported that they did not use the internet for information on mental health as they had no need. In addition, 6.1% (n=29) participants believed the information would be unreliable, 4% (n=19) never thought about it, 8.3% (n=39) had no interest, 1.9% (n=9) would prefer to discuss mental health issues with a qualified person such as a doctor, and 1.3% (n=6) reported that they would use other resources.

The majority of the 872 respondents (68%, n=593) indicated that they would use the internet for mental health support if they needed to. Reasons for this are provided in table 3.

Those who reported they would not use the internet for mental health support (n=279) predominantly cited that they would prefer to interact with a person, whether that be a professional, a friend or a family member (n=79, 28.3%). A number of participants also reported that they believed it would be unreliable (n=42, 15.1%), untrustworthy (n=14, 5%), it lacks privacy (n=7, 2.5%), is too impersonal (n=21, 7.5%) and that insufficient support would be found (n=11, 3.9%).

### **Face to face versus internet support**

Of the 868 respondents, the majority (79.4%, n=689) reported that they would prefer face to face support rather than internet support for a mental health problem.

Those who reported they would prefer face to face support for mental health problems (n=689) suggested it was because it is more personal (n=145, 21%) and that it is easier to relate, engage and communicate with a person face to face, particularly because body language can be observed (N=67, 9.7%). They also reported that this approach allows for greater development of trust (n=25, 3.6%) and is more reliable (n=28, 4.1%). They also felt that they could verify the qualifications of the person (n=22, 3.2%) and that it would be easier to judge the feedback given to them (n=4, 0.6%). Further participants believed that the support offered to them would be tailored to their specific needs (n=27, 3.9%), they could ask specific questions (n=15, 2.2%), that better support would be provided (n=18, 2.6%), that it would be more reassuring (n=13, 1.9%), real (n=11, 1.6%), comforting (n=9, 1.3%), safe (n=5, 0.7%), and direct (n=4, 0.6%).

Those participants (n=179) who indicated a preference for internet support gave a variety of reasons. One theme centred on anonymity (n=18, 10.1%), privacy (n=14, 7.8%), and confidentiality (n=9, 5%). A second theme centred on accessibility (n=10, 5.6%), speed (n=4, 2.2%) and cost (n=4, 2.2%). A third focussed on integrity, 7.3% (n=13) believed that they would not be judged, 3.4% (n=6) believed it would be a good place to get initial information. In addition they also reported that they would be more inclined to open up on the internet (n=3, 1.7%) and that it would be easier to express themselves (n=2, 1.1%), that they could communicate with others in similar situations to find out how they are coping (n=3, 1.7%), they are less likely to lie (n=3, 1.7%) and it is less intimidating (n=3, 1.7%).

### **Website design and content for mental health problems**

The participants reported what they deemed to be important in terms of website design and content, if they were to use the internet for mental health support. Most issues were identified as being at minimum somewhat important. The most important issues identified were confidentiality, anonymity, being able to ask questions, usability and knowing who produced the information. Those deemed least important were animation, colour and chat room functions. A small number of participants provided individual comments. It was suggested that a basic layout with access to chat rooms and a good search facility would be very

important. That links to self help groups should be provided and that the website should not be for profit.

## **Discussion**

The findings suggest that young people aged between 18 and 24 years are frequent users of the internet, with 72.4% of the participants using it several times a day. This is slightly lower than the findings of Rogers (2008) study with 21 to 30 year olds in the US, where it was found that 92% access the internet several times a day.

It was also found that 91.7% of participants accessed the computer at home indicating that most households have a computer. According to the last figures available the Central Statistics office in Ireland (2008) reported that 70% of all households have computers, 62% have internet access, while 43% have broadband connection. In a recent Irish study it was reported that 58.3% of participants had used the internet, this figure rose to 88.9% among the 16 to 29 years age group (Gallagher et al. 2008). While the results of this study indicate slightly higher usage of the internet, this study focussed on University students only and thus is not representative of the entire population. Additionally, 94.9% of participants accessed the internet at college indicating a high usage on campus. Over the past number of years communication within this University has changed from paper based communication to electronic communication. Each student is provided with an email account, and lecture notes and timetables are posted online through Blackboard, this could explain the high usage on campus.

Little information is currently available on how young people use the internet for mental health information and support. The findings indicated that 30.8% of participants had used the internet for mental health information. This is higher than studies by Gould et al (2002), who found that 18.2% of 13-19 year olds used the internet as a source of mental health information and by Powell & Clarke (2006), who found that 18% of the general population used the internet for this purpose. This could indicate that 18-24 year olds are more likely to use the internet or that there has been an increase in its use over the past few years. The main information searched for was information on depression, information for course work, general information on mental health problems and information on specific problems they

themselves or family or friends experienced. Indeed, in addition to depression being the most frequently searched topic, many related topics were also searched, such as suicide, self harm, stress and confidence and self esteem building. Depression is one of the most common mental health problems in young people (WHO 2003) and thus these findings are unsurprising. They do however highlight that young people are willing to source information and support for depression on the internet. Indeed, Kessler & Walters (1998) reported that while 25% of young people will experience depression before the age of 24, only 20% of those will receive good quality care. Various reasons were provided as to why they do not use the internet for mental health information, it predominantly related to lack of perceived need. In addition a number of participants were concerned with the reliability of the information, highlighting that young people may be experiencing difficulty in determining the quality of information online. It was found that 68% of participants would use the internet for mental health support if they needed to. Similar to findings of other studies (Shaw et al. 2000), the reasons for this related to the vast amount of information available, anonymity, confidentiality, privacy, easy access and speed. It is recognised that young people are reluctant to access main stream mental health services due to fear of judgement and due to the stigma that still exists in relation to mental health problems. This study highlights that the majority of young people are willing to look for support from the internet which has implications for future service delivery for this age group.

The majority of participants reported that they would prefer face to face support over internet support for mental health problems. It was found that they believed they could relate better to people when they were in front of them. Those who reported they would prefer internet support suggested that this was due to anonymity and fear of being judged by a person when face to face. Similar findings were reported by Rogers (2008) where it was found that 80% of 21 to 30 year olds in the study would prefer face-to-face therapy. In a study by Oh et al. (2008) it was reported that certain mental health information websites have improved mental health literacy and reduced symptoms. This study explored young people's preference for a website with self help books and face to face services (counselling and mental health services). A telephone survey was conducted with 3,746 young people (12-25 years) and 2,005 parents, each participant was read a vignette based on depression, depression with alcohol use, social phobia or psychosis, they were then asked their views on their beliefs and attitudes about help seeking and interventions. It was found that 71% believed the website

would be helpful, 65% believed face to face visits with mental health professionals would be helpful and 92% believed counselling would be the most helpful. This again highlights that young people are willing to use a variety of resources for support with mental health problems.

It is important to highlight that while the majority of participants reported they would prefer face to face approaches, 20.6% reported that they would prefer internet interventions; this represents a considerable number and suggests that there is a need to develop interventions via the internet for young people experiencing mental health difficulties.

The findings indicated that in the development of web based interventions for mental health problems, confidentiality, anonymity, usability, being able to ask questions and knowing who produced the information are important to young people. This information could help in the development of web based interventions.

### **Limitations**

This study was conducted with University students and thus may not be representative of all young peoples' views. It is possible that young people in continuing education are more likely to use the internet and possess their own computer due to necessity. In addition, as the questionnaire was posted online, all participants had to use the internet to complete it.

Therefore the sample consists of a computer literate population only. While this may be considered a sampling bias, it is necessary to acknowledge that all students at this particular University are required to use the internet to keep updated on course work and timetables.

While the questionnaire used in this study was self designed and its validity not tested, the large sample size increases the reliability of the findings. In addition measures were taken, as discussed in the methodology section to enhance the validity of the questionnaire. Finally, the questionnaire did not ask participants to distinguish between whether they searched for mental health information for themselves or friends and family and as such the findings in this area are unclear.

### **Conclusion**

This study has clearly indicated that young people attending University are active users of the internet and are willing to use the internet for mental health information and support. It also

indicated that they would prefer to use face to face support instead of internet support. This study can help inform the design of mental health information websites as it demonstrates what topics are frequently searched for, the difficulties they have with using the internet and the issues they deem to be important in terms of website design and content. In addition it can help inform psychiatric nurses on the types of information and support young people are looking for. Support via the World Wide Web is being offered to young people, particularly in the developing world and thus this information can help inform practice, it can also help in further developing this aspect of psychiatric nursing care. Further studies in this area are needed, particularly on young people's views of mental health information and support websites, their search strategies and filtering strategies when looking for mental health information, their assessment of the quality of information online and their needs in relation to internet support. In addition, there is a need to develop and evaluate web based interventions for this population and psychiatric nurses need to develop their knowledge of the services currently available so that they can provide accurate and reliable information on web resources for young people.

## References

- Andrews, G., Hall, W., Teesson, M., Henderson, S. (1999). *The Mental Health of Australians*. Canberra, Mental Health Branch, Commonwealth Department of Health and Aged Care
- Berger, M., Wagner, T. & Baker, L. (2005). Internet use and stigmatised illness. *Social Science and Medicine* 61, 1821-1827.
- Burns, N. & Grove, S. (2001). *The Practice of Nursing Research. Conduct, Critique and Utilization*. 4<sup>th</sup> Edition. Philadelphia, W.B. Saunders.
- Central Statistics Office (2008). *Information Society Statistics. First results 2008*. Dublin, Central Statistics Office.
- Donald, M., Dower, J., Lucke, J. & Raphael, B. (2000). *The Queensland Young Peoples' Mental Health Survey Report* Brisbane, Centre for Primary Health Care, University of Queensland.
- eUSER (2005). *Public Online Services and User Orientation* [www.euser-eu.org](http://www.euser-eu.org) (18<sup>th</sup> June 2008)
- Eysenbach, G., Powell, J., Kuss, O. & Sa, E. (2002). Empirical studies assessing the quality of health information for consumers on the World Wide Web. *JAMA* 287(20), 2691-2700.
- Gallagher, S., Doherty, T., Moran, R. & Kartalova-O'Doherty, Y. (2008). *Internet use and seeking health information online in Ireland: Demographic characteristics and mental health characteristics of users and non-users* Dublin, Health Research Board
- Gould, M., Munfakh, J., Lubell, K., Kleinmann, M. & Parker, S. (2002). Seeking help from the internet during adolescence. *Journal of the American Academy of Child and Adolescent Psychiatry* 41, 1182-1190.
- Graham, C., Franes, A., Kenwright, M. & Marks, I. (2000). Psychotherapy by computer: a postal survey of responders to a teletext article *Psychiatric Bulletin* 24, 331-332

Health on the Net Foundation (2001). Evolution of Internet use for Health Purposes – Feb/Mar 2001. <http://www.hon.ch/Survey/FebMar2001/survey.html> 29/01/09

Health Service Executive (2007) *Insight 07. Health and Social Services in Ireland – a survey of consumer satisfaction* Dublin, Health Service Executive.

Hoybye, M., Johansen, C. & Tjornhoj-Thomsen, T. (2005). Online interaction. Effects of storytelling in an internet breast cancer support group *Psycho-Oncology* 14, 211-220.

Klemm, P. (2005). Cancer caregivers online: hope, emotional roller coaster and physical/emotional/psychological responses *Computers, Informatics, Nursing* 23(1), 38-45.

Lieberman, M. (2005). Self help on-line: an outcome evaluation of breast cancer bulletin boards *Journal of Health Psychology* 10(6), 855-862.

Lieberman, M., Golant, M., Giese-Davis, J., Winzlenberg, A., Benjamin, H., Humphreys, K., Kronenwetter, C., Russo, S. & Spiegel, D. (2003). Electronic support groups for breast carcinoma. A clinical trial of effectiveness *Cancer* 97(4), 920-925.

Johnsen, J., Rosenvinge, J. & Gammon, D. (2002). Online group interactions and mental health: an analysis of three online discussion forums *Scandinavian Journal of Psychology* 43, 445-449.

Kessler, R. & Walters, E. (1998). Epidemiology of DSM-III-R major depression and minor depression among adolescents and young adults in the National Comorbidity Survey. *Depression and Anxiety* 7, 3-14.

Oh, E., Jorm, A. & Wright, A. (2008). Perceived helpfulness of websites for mental health information. *Social Psychiatry and Psychiatric Epidemiology* 10.1007/s00127-008-0443-9

Powell, J. & Clarke, A. (2006). Internet information seeking in mental health: population survey. *British Journal of Psychiatry* 189(3), 273-277.

Powell, J. & Clarke, A. (2006). Internet information seeking in mental health: population survey. *British Journal of Psychiatry* 189(3), 273-277.

Powell, J., McCarthy, N. & Eysenbach, G. (2003). Cross-sectional survey of users of internet depression communities. *BMC Psychiatry* 3(9)

<http://www.biomedcentral.com/content/pdf/1471-244X-3-19.pdf> (18th June 2008)

Renahy, E., Parizot, I. & Chauvin, P. (2008). Health information seeking on the internet: a double divide? Results from a representative survey in the Paris metropolitan area, France, 2005-2006. *BMC Public Health* 8(69), 10.1186/1471-2458-8-69.

Rice, R. (2006). Influences, usage, and outcomes of internet health information searching: multivariate results from the Pew surveys. *International Journal of Medical Informatics* 75, 8-28.

Rickwood, D., Deane, F. & Wilson, C. (2007). When and how do young people seek professional help for mental health problems? *MJA* 187(7 Supplement), S35-S39.

Rogers, V. (2008). *Emotional self disclosure: differences between young adults who prefer face-to-face therapy and those who prefer internet therapy*. Unpublished Doctoral Dissertation. Ohio, Case Western Reserve University.

Shaw, B., McTavish, F., Hawkins, R., Gustafson, D. & Pingree, S. (2000). Experiences of women with breast cancer: exchanging social support over the CHES computer network *Journal of Health Communication* 5, 135-159.

Skinner, H., Biscope, S., Poland, B. & Goldberg, E. (2003). How adolescents use technology for health information: implications for health professionals from focus group studies. *Journal of Medical Internet Research* 5(4), e32.

Sullivan, C., Arensman, E., Keeley, H., Corcoran, P. & Perry, I. (2004). *Young people's mental health. A report of the findings from the lifestyle and coping survey* Cork: National Suicide Research Foundation.

World Health Organisation (2003). *Caring for children and adolescents with mental disorders*. Geneva, WHO

Ybarra, M. & Eaton, W. (2005). Internet based mental health interventions *Mental Health Services research* 7(2), 75-87.

Table 1: Participant characteristics

|               |   |  |  |                                  |                                       |                           |  |
|---------------|---|--|--|----------------------------------|---------------------------------------|---------------------------|--|
| Age           | 18 yrs<br>(16.8%,<br>n=149)                                   | 19 yrs<br>(25.5%,<br>n=226)              | 20 yrs<br>(20.2%,<br>n=179)                                | 21 yrs<br>(17%,<br>n=151)        | 22 yrs<br>(10.3%,<br>n=91)            | 23 yrs<br>(6.4%,<br>n=57) | 24 yrs<br>(3.7%,<br>n=33)                |
| Gender        | Male (37.6%, n=333)   |  |  |                                  | Female (62.4%, n=552)                 |                           |  |
| Area of Study | Arts, Celtic<br>Studies, Social<br>Sciences<br>(33.6%, n=298) |  | Science,<br>Engineering,<br>Food Science<br>(31.7%, n=281) |                                  | Business and<br>Law (19.4%,<br>n=172) |                           | Medicine and<br>Health (15.2%,<br>n=135) |
| Accommodation | Private<br>rental<br>(41.8%,<br>n=367)                        | Parents/Family<br>home (40.2%,<br>n=353) | College<br>housing<br>(16%,<br>n=141)                      | Social<br>housing<br>(0.8%, n=7) | Own home<br>(1.3%,<br>n=11)           |                           |  |

Table 2: Reasons for using the internet for mental health information

|  |             |                       |           |
|--|-------------|-----------------------|-----------|
| Depression   | n=81, 36.8% | Stress                | n=11, 5%  |
| Information relating to college work   | n=42, 19.1% | Coping strategies     | n=9, 4.1% |
| General information on mental health<br>problems and illnesses                                 | n=35, 15.9% | Alternative therapies | n=6, 2.7% |
| Medication management  | n=16, 7.3%  | Symptoms              | n=5, 2.3% |
| Illness of family and friends  | n=15, 6.8%  | Support groups        | n=5, 2.3% |
| Bipolar disorder   | n=13, 5.9%  | Dementia              | n=4, 1.8% |
| Anxiety problems such as obsessive<br>compulsive disorder, social anxiety and<br>panic attacks | n=12, 5.5%  | Personality disorders | n=3, 1.4% |
| Eating disorders   | n=12, 5.5%  | Post-natal depression | n=3, 1.4% |
| Suicide  | n=11, 5%    | Causes of illness     | n=3, 1.4% |
| Schizophrenia  | n=11, 5%    |                       |           |

Table 3: Reasons for using the internet for mental health support

|   |              |
|---|--------------|
| Anonymous, private and confidential                                       | n=132, 22.3% |
| Vast amount of valuable information available                             | n=129, 21.8% |
| Easily accessible 24 hours a day  | n=60, 10.1%  |
| Easy to find information  | n=65, 11%    |
| Fast  | n=49, 8.3%   |
| Cheap   | n=16, 2.7%   |
| Convenient  | n=10, 1.7%   |
| Easy to communicate with other young people in the same situation         | n=22, 3.7%   |
| A good place to start and to find out where to go for further assistance  | n=28, 4.7%   |
| Less embarrassing than talking to a professional, friend or family member | n=17, 2.9%   |