

Title	A review of the design and validation of web- and computer-based 24-h dietary recall tools
Authors	van den Barg, Rinske;Blain, Richard J.;Kehoe, Laura;Evans, Katie;Walton, Janette;Timon, Claire M.;Flynn, Albert;Gibney, Eileen R.
Publication date	2016-12-13
Original Citation	Timon, C. M., van den Barg, R., Blain, R. J., Kehoe, L., Evans, K., Walton, J., Flynn, A. and Gibney, E. R. (2016) 'A review of the design and validation of web- and computer-based 24-h dietary recall tools', Nutrition Research Reviews, 29(2), pp. 268-280. doi:10.1017/S0954422416000172
Type of publication	Article (peer-reviewed)
Link to publisher's version	10.1017/S0954422416000172
Rights	© The Authors 2016. Published by Cambridge University Press (CUP) on behalf of The Nutrition Society.
Download date	2024-04-20 00:24:18
Item downloaded from	<a href="https://hdl.handle.net/10468/3716">https://hdl.handle.net/10468/3716</a>



## A review of the design and validation of web- and computer-based 24-h dietary recall tools

Claire M. Timon<sup>1</sup>, Rinske van den Barg<sup>1</sup>, Richard J. Blain<sup>1</sup>, Laura Kehoe<sup>2</sup>, Katie Evans<sup>2</sup>, Janette Walton<sup>2</sup>, Albert Flynn<sup>2</sup> and Eileen R. Gibney<sup>1\*</sup>

<sup>1</sup>*Institute of Food and Health, University College Dublin, Belfield, Dublin 4, Republic of Ireland*

<sup>2</sup>*School of Food and Nutritional Sciences, University College Cork, Cork, Republic of Ireland*

### Abstract

Technology-based dietary assessment offers solutions to many of the limitations of traditional dietary assessment methodologies including cost, participation rates and the accuracy of data collected. The 24-h dietary recall (24HDR) method is currently the most utilised method for the collection of dietary intake data at a national level. Recently there have been many developments using web-based platforms to collect food intake data using the principles of the 24HDR method. This review identifies web- and computer-based 24HDR tools that have been developed for both children and adult population groups, and examines common design features and the methods used to investigate the performance and validity of these tools. Overall, there is generally good to strong agreement between web-based 24HDR and respective reference measures for intakes of macro- and micronutrients.

**Key words:** Web-based recalls: Dietary assessment: Computer-based recalls: 24-h Dietary recall tools: Validation

### Introduction

Accurate measurements of dietary intake are fundamental to health and nutrition research; however, diet is inherently complicated to measure as it changes over time and varies across life stages amongst other factors<sup>(1)</sup>. In recent years there have been many advances in the application of technology in the area of dietary assessment<sup>(2)</sup>. Innovative dietary assessment technologies using devices such as mobile telephones<sup>(3–8)</sup>, sensors, wearable cameras<sup>(9–11)</sup> and web-based platforms are common ways of collecting dietary intake data and have become increasingly popular alternatives to traditional pen-and-paper versions of dietary assessment<sup>(12)</sup>. Web-based methodologies facilitate the collection of dietary intake across many geographic locations<sup>(13–16)</sup> and in some cases are preferred by participants compared with the traditional pen-and-paper alternatives<sup>(3,17)</sup>. Overall the application of technology in dietary intake assessment has been shown to reduce issues associated with traditional collection of dietary data, such as cost, participation rates (by reducing the burden associated with dietary assessment) and accuracy of data collected<sup>(18)</sup>. The success of these methods has been attributed to a number of factors including the ability to collect data in a remote and neutral environment, the standardised sequence of questioning, the use of digital portion size assessment aids, automated analysis of data collected and the provision of dietary feedback<sup>(12)</sup>.

The reduced participant burden and time of data collection, associated with web- and computer-based 24-h dietary recall (24HDR) tools, support the use of this method in many different populations. Web- and computer-based 24HDR tools have been used to collect data from various population groups including young children<sup>(19–22)</sup>, adolescents<sup>(5,6,23)</sup> and adults<sup>(24–27)</sup>. Two of the comparison and validation studies included in this review recruited elderly participants in their study population<sup>(28,29)</sup>; however, the validity of these tools have not been tested explicitly in an elderly population. In general, the applications are based on the multi-pass approach described by Moshfegh *et al.*<sup>(30)</sup>. The applications are either self-administered, whereby a participant completes the recall in the absence of a researcher<sup>(5,31–33)</sup>, or interviewer administered, where the interviewer uses the application to collect/analyse dietary recall data in the presence of a participant<sup>(26,28,34,35)</sup>. The software guides and prompts the participant/interviewer to recall and record food and drink consumed in the previous day (24 h). The design features, food and beverage lists, nutritional composition data, prompts and portion size assessment aids incorporated in these applications differ from model to model and are generally for target populations.

The ability of many of these tools to accurately assess dietary intakes has been investigated using different study designs, the most common of which are comparison and validation studies. A comparison study investigates the performance of a 'test measure' of dietary assessment against an alternative dietary

**Abbreviations:** 24HDR, 24-h dietary recall; ASA24, National Cancer Institute's Automated Self-Administered 24-Hour Dietary Assessment Tool; DLW, doubly labelled water.

\* **Corresponding author:** Eileen R. Gibney, fax +353 1716 7203, email eileen.gibney@ucd.ie



recall (such as interviewer-led 24-h recall) to ascertain if the data collected by the test measure are comparable with those using the existing method. A validation study investigates the accuracy of a test measure compared with an objective measure of intake, such as biological markers or direct observation of intake. Assessing the validity of dietary assessment tools is difficult due to the risk of correlated error between the test method and a reference method<sup>(36)</sup>. The cost and practicality associated with the direct observation of intake or the collection of biological samples for the analysis of biomarkers of intake can also be limiting factors. In the absence of biomarkers or direct observation, dietary records (preferably weighed food records) are often considered the 'gold standard' for the measurement of food and nutrient intake and are typically used as a reference comparison method<sup>(37)</sup>.

De Keyzer *et al.*<sup>(38)</sup> notes that the 24HDR method is currently the most commonly used tool to collect dietary intake data in national surveys, usually taking multiple recalls to assess habitual intake and, in some instances, used in conjunction with an FFQ (for example, Canada and USA)<sup>(39,40)</sup>. In Europe, efforts are being made to standardise the collection of dietary intake data and the potential of computer-assisted 24HDR tools has been recognised<sup>(26,41)</sup>. Both self-administered and interviewer-administered applications have demonstrated the ability to reach large population groups using these applications<sup>(42,43)</sup>. At present there are numerous 24HDR tools that have been developed, tested and validated worldwide. The aim of this review was to examine common design features amongst current 24HDR tools and investigate the methods used to assess the comparability and validity of these tools.

## Methods

A literature search was conducted using the online databases PubMed, EMBASE and Web of Science to collect information on all 24HDR tools. The inclusion criteria were as follows: English-language publications from 2000 until 2016 reporting the development, comparison or validation of a web- or computer-based 24-h recall tool for any population group. The following search terms were used alone and in combination: '24h recall', '24-HR', 'nutrition', 'food', 'intake', 'dietary assessment', 'validity', 'validation', 'comparison' and 'reliability'. In a similar approach described by Gemming *et al.*<sup>(44)</sup>, abstracts were initially screened by one author (C. M. T.) for relevance and then collected and distributed to three other authors. The literature search yielded a total of forty-four relevant papers. All papers were reviewed by three investigators (C. M. T., R. v. d. B. and R. J. B.). Sixteen papers were omitted for the following reasons: the papers did not describe a 24HDR tool ( $n = 4$ ) or the papers described aspects of the study, other than the development, comparison or validation of web-based dietary recall tools ( $n = 12$ ). A total of twenty-eight papers reporting twenty-one individual web- or computer-based 24HDR tools were included in the present review.

The design characteristics of the twenty-one 24HDR tools are described in Tables 1 and 2 and the findings of those tools whereby validity/comparability was investigated are described in Tables 3 and 4. To obtain an overview of the different tools,

general characteristics were examined, such as participant age, number of food and drink items incorporated in the application, the use of prompt techniques to ensure the complete capture of data and the use of portion size assessment aids. For the assessment of techniques used to investigate validity and comparability, details such as number of recalls used, reference method, type of statistical analysis and markers of nutrient intake were compared to identify common features. Where the required information was not available in the included publications, the corresponding author was contacted via email to obtain/verify information when possible.

## Results

### Design characteristics

Of the twenty-one 24HDR tools identified from the literature search, six tools were developed for use specifically with children (three developed in Europe, two in the USA and one in Canada) and thirteen tools were developed (seven in Europe, three in USA, two in Korea and one in India) specifically for adult populations and two tools (both in the UK) were developed for use both with children and adults<sup>(5,31)</sup>. All of the tools developed for use with children were self-administered, whereas four of the tools developed for adults were interviewer administered. The youngest age to test a 24HDR tool was 7 years of age<sup>(22,45)</sup> and the oldest age was 80+ years<sup>(28,29)</sup> (Tables 1 and 2).

Only three 24HDR tools reported collecting intake data at a food-group level<sup>(27,29,46)</sup>, including additional choices/questions within each food group to obtain more specific information about intakes. All other tools recorded intake by presenting lists of food and beverage items. The number of food and drink items included varied for each tool, ranging from forty<sup>(45)</sup> to 45 000 food and beverage options<sup>(31)</sup>. The use of portion size images was the most popular aid to facilitate portion size estimation. The number of portion size images varied from tool to tool, with some investigators reporting as many as 17 000 images<sup>(32)</sup>.

The use of food prompts was prevalent across 24HDR tools developed for both children and adults. 'Frequently forgotten foods' was the most common prompt function reported in 24HDR tools developed for children. This prompt involves the presentation of a list of foods that are known to be frequently omitted from dietary recalls (for example, sugar in hot beverages, etc.)<sup>(31,47)</sup>. The Synchronised Nutrition and Activity Program (SNAP<sup>TM</sup>) tool<sup>(21)</sup> used images of foods to prompt for frequently forgotten food items rather than a list of foods. In the tools developed for adults, 'frequently forgotten foods' was also a common prompt, as was 'linked foods' (Table 2). Linked food prompts offer the participant food or drink options that are often eaten in combination with the primary food or drink item selected<sup>(5,24)</sup>. Automated data entry checks, such as those implemented by Vereecken *et al.*<sup>(48)</sup>, for portion size information and meal gap reviews whereby any period greater than 3 h between meals was queried<sup>(32)</sup> were also useful prompt functions incorporated into adult 24HDR tools. Of the 24HDR tools developed for children, two recorded supplement intake<sup>(19)</sup>

**Table 1.** Design characteristics of the web- and computer-based 24-h dietary recall tools developed for children

Author	Region	Tool name	Recall method	Age (years)	No. food and drink items	Portion size estimation method	Use of prompts	Report supplement intake
Vereecken <i>et al.</i> (2014) <sup>(6)</sup>	Belgium	CANAA-W (formally YANA-C)	24HR, MR, SA	11–12	800, 25 food groups	Photographs ( <i>n</i> 2100) of increasing portion sizes presented sequentially or simultaneously.	For foods eaten in combination with other items. Reliability checking: warning when extreme amounts are entered	No
Moore <i>et al.</i> (2008) <sup>(21)</sup>	UK	SNAP™	24HR, MR, SA	7–15	49	Description and entry of g weight By count (i.e. the number of times a particular food was consumed)	Pictures used as visual memory prompts	No
Baranowski <i>et al.</i> (2014) <sup>(57)</sup>	USA	FIRSSt	24HR, MR, SA	9–11	Not mentioned	Photographs ( <i>n</i> 14 000) of eight increasing portion sizes presented simultaneously	Avatar. For frequently forgotten foods	No
Carvalho <i>et al.</i> (2014) <sup>(22)</sup>	Portugal	PAC24	24HR, MR, SA	7–10	332 food and 41 drinks items	Photographs of seven increasing portion sizes to identify the amount of food served and amount of food left over.	For approximate time and for location of food/meal consumption	No
Storey <i>et al.</i> (2012) <sup>(23)</sup>	Canada	Web-SPAN	24HR, MR, SA	11–15	Not mentioned	Description and entry of g weight Food photographs and cues regarding beverage intake	For foods eaten in combination with other items	No
Foster <i>et al.</i> (2014) <sup>(5)</sup>	UK	Intake24	24HR, MR, SA	11–16	≥1600	Photographs of increasing portion sizes to identify the amount of food served and amount of food left over. Description of the amount served and left over is entered (using household measures)	Long time periods where no food is reported. For foods eaten in combination with other items. Asking for additional information (i.e. brand names). 'Same as before' option	No
Diep <i>et al.</i> (2015) <sup>(19)</sup>	USA	ASA24-Kids-2012	24HR, MR, SA	9–11	5407 (food terms)	Photographs ( <i>n</i> 9759) of increasing portion sizes presented sequentially	For frequently forgotten foods. Meal gap review. Final probe to ensure complete capture of data	Yes
Albar <i>et al.</i> (2016) <sup>(53)</sup>	UK	Myfood24	24HR, MR, SA	11–18	about 45 000 individual food/drink items	Portion size images for 5669 food items. Portion size suggestions (household measures). Free entry of g weight by participant	For foods eaten in combination with other items. Asking for additional information (i.e. brand names)	Yes

CANAA-W, Children and Adolescents' Nutrition Assessment and Advice on the Web (a web-based 24-h dietary recall program for children and adolescents); YANA-C, Young Adolescents' Nutrition Assessment on Computer (a web-based 24-h dietary recall program for children and adolescents); 24HR, 24-h recall; MR, multiple-pass recall; SA, self-administered; SNAP™, Synchronised Nutrition and Activity Program (a web-based 24-h recall program which assesses energy balance-related behaviours in children and adolescents); FIRSSt, Food Intake Recording Software System (a computer-based 24-h dietary recall program for children); PAC24, Portuguese self-administered, computerised, 24-h dietary recall (a web-based 24-h dietary recall program for children); Web-SPAN, Web-Survey of Physical Activity and Nutrition (a web-based 24-h dietary recall program for children and adolescents); Intake24, a web-based 24-h dietary recall program for children and young adults; ASA24-Kids-2012, Automated Self-Administered 24-Hour Dietary Assessment Tool, Kids-2012 version; myfood24, Measure your food on One day (a web-based 24-h dietary recall program for children and adults).

**Table 2.** Design characteristics of the web- and computer-based 24-h dietary recall tools developed for adults

Author	Region	Tool name	Recall method	Age (years)	No. food and drink items	Portion size estimation method	Use of prompts	Report supplement intake
Slimani <i>et al.</i> (2011) <sup>(26)</sup>	Europe	EPIC-Soft	24HR, MR, IA	≥ 15	1500–2200 individual food/drink items and 150–350 recipes	Photographs (in a book). Household measures. Standard units (in g, piece) and standard portions. Weight or volume method	Reliability checking. For frequently forgotten foods. For foods eaten in combination with other items	Yes
Touvier <i>et al.</i> (2011) <sup>(17)</sup>	France	NutriNet-Santé	24HR, MR, SA	48–75	Not mentioned	Photographs of seven increasing portion sizes presented. Free entry of g weight by participant	For frequently forgotten foods. For foods eaten in combination with other items	No
Daniel <i>et al.</i> (2013) <sup>(34)</sup>	India	NINA-DISH	24HR, FFQ, IA, MR	35–69	910 individual food/drink items	Household measures. Photographs of fruits (small, medium and large)	For foods eaten in combination with other items	No
Shin <i>et al.</i> (2014) <sup>(28)</sup>	Korea	CAPIS	24HR, MR, IA	24–85	3642 and 1886 recipes	Photographs presented ½, 1 and 1½ serving sizes for each item. Free entry of g weight by participant	For frequently forgotten foods	No
Foster (2014) <sup>(5)</sup>	UK	Intake24	24HR, MR, SA	17–24	≥1600	Over 2000 portion size images representing over 100 food and drink types. Description of the amount served and left over is entered (using household measures)	Long time periods where no food is reported. For foods eaten in combination with other items. Asking for additional information (i.e. brand names). ‘Same as before’ option	No
Hillier <i>et al.</i> (2012) <sup>(49)</sup>	UK	SNAPA™	24HR, MR, SA	Mean 34.4 (SD 11.1)	120 individual food/drink items	Report fruit and vegetables consumed as numbers of portions	For frequently forgotten foods	No
Comrie <i>et al.</i> (2009) <sup>(27)</sup>	UK	FoRC	24HR, MR, SA	18–49	121 individual food/drink items	Photographs to estimate portion size. Description and free entry of g weight by participant	The use of layers of questioning to prompt recall	No
Liu <i>et al.</i> (2011) <sup>(29)</sup>	UK	Oxford WebQ	24HR, MR, SA	19– 89	21 food groups	Standard units (i.e. four slices of bread). Portion sizes are specified as servings	Expanding questions to prompt recall and further detail	No
Zoellner <i>et al.</i> (2005) <sup>(60)</sup>	USA	IMM	24HR, MR, SA	18–65	167 individual food/drink items	Photographs of four increasing portion sizes presented	Audio instructions	No
Subar <i>et al.</i> (2007) <sup>(24)</sup>	USA	ASA24	24HR, MR, SA	18–77	7200 individual food/drink items	17 000 portion size images. Foods with portion size images had eight different portion sizes (in the latest version of ASA24)	An avatar gives audio instructions. For frequently forgotten foods. For foods eaten in combination with other items. Long time periods where no food is reported (meal gap)	Yes
Arab <i>et al.</i> (2010) <sup>(25)</sup>	USA	DietDay	24HR, MR, SA	21–69	9349 individual food/drink items	7000 portion size images which could be modified using command buttons	Reliability checking	Yes
Park <i>et al.</i> (2015) <sup>(35)</sup>	Korea	GloboDiet	24HR, MR, IA	24–68	1305 individual food/drink items	A picture book of foods/dishes was prepared of food portion sizes relevant to Korean diet. Volumes (directly or as household measures or as shapes and thickness). Standard unit. Free entry of g weight by participant	Adaptation of EPIC-Soft	Yes

**Table 2.** *Continued*

Author	Region	Tool name	Recall method	Age (years)	No. food and drink items	Portion size estimation method	Use of prompts	Report supplement intake
Freese <i>et al.</i> (2015) <sup>(46)</sup>	Germany	24-h food list	24HR, SA	20–70	246 individual food/drink items	Tool does not enquire about portion size and uses statistical methods to estimate usual intake distributions	Participants are presented with a list of food groups, if they had consumed items from that food group, more specific food items were then presented	Yes
Carter <i>et al.</i> (2015) <sup>(31)</sup>	UK	myfood24	24HR, MR, SA	18–65	About 45 000 individual food/drink items	Portion size images for 5669 food items. Portion size suggestions (household measures).	For foods eaten in combination with other items. Asking for additional information (i.e. brand names)	Yes
Wardenaar <i>et al.</i> (2015) <sup>(52)</sup>	The Netherlands	Compl-eat™	24HR, MR, SA	18–35	Not mentioned	Free entry of g weight by participant. Household measures	Food group quick list which expands to further subgroups and individual items on selection. For frequently forgotten foods	No (questioned separately)

EPIC-Soft, European Prospective Investigation into Cancer and Nutrition Software (a standardised computerised 24-h dietary recall method); 24HR, 24-h recall; MR, multiple-pass recall; IA, interviewer administered; NutriNet-Santé, a web-based 24-h dietary recall program for adults; SA, self-administered; NINA-DISH, New Interactive Nutrition Assistant – Diet in India Study of Health (a computer-based 24-h dietary recall program for adults); CAP/IS, Computer-Assisted Personal Interview System (a computer-based 24-h dietary recall program for adults); Intake24, a web-based 24-h dietary recall program for children and young adults; SNAPA™, Synchronised Nutrition and Activity Program for Adults (a web-based 24-h dietary recall program for adults); FoRC, Food Recall Checklist (121-item food recall list which has been used to measure diet in undergraduate students); Oxford WebQ, a web-based 24-h dietary recall program for adults; IMM, a bilingual interactive multimedia dietary assessment tool based on the 24-h dietary recall method (a web-based 24-h dietary recall program for adults); ASA24, Automated Self-Administered 24 Hour Dietary Assessment Tool (web-based 24-h dietary recall tool for adults, developed by the National Cancer Institute, USA); DieDay, a web-based 24-h dietary recall program for children and adults; Compleat™, a computer-based 24-h dietary recall method; myfood24, Measure your food on One day (a web-based 24-h dietary recall program for children and adults);

compared with six of the identified 24HDR tools developed for adults (Table 2).

*Web- and computer-based 24-h dietary recall tools comparability/validation study design*

Tables 3 and 4 present the comparison/validation study design features, and key findings for the identified 24HDR tools. For the purposes of this review, the authors considered a comparison study to be where the test measure was compared against a traditional 24-h recall or estimated/weighed food diary. A validation study was when the test measure was compared against direct observation (of eating occasions) or biological markers of nutrient intake. According to the authors' interpretation of a comparison and validation study, of the 24HDR tools developed for both children and adults (Foster considered one study across two age groups), nine investigators conducted comparison studies and nine investigators conducted validation studies (EPIC-Soft and NutriNet-Santé were investigated in both comparison and validation studies). The remaining three tools, of the twenty-one identified in the present review, were not evaluated in either a comparison or validation study. Of the studies included in the present review the most common reference method in comparison studies was an interviewer-led multiple-pass recall and for validation studies was direct observation of eating occasions, closely followed by the use of biological markers of intake (Table 4). The number of participants ranged from forty-one<sup>(22)</sup> to 459 for comparison and validation studies conducted with children<sup>(23)</sup>. There was similar variation with the numbers that participated in the adult studies with fifty-three<sup>(27)</sup> to 1072<sup>(43)</sup>. It is important to consider the reference method used when comparing participant numbers across studies; for example Touvier *et al.*<sup>(17)</sup> recruited 147 participants and used one web-based recall compared with one interviewer-led recall, whereas Comrie *et al.*<sup>(27)</sup> asked fifty-three participants to complete a 4-d estimated food diary as a reference. The lesser burden associated with the reference used by Touvier *et al.*<sup>(17)</sup> may allow for a greater number of participants to be involved compared with the burden (participant training, interview on diary and analysis of data) associated with a 4-d estimated food diary<sup>(27)</sup>.

A variety of statistical measures were used to investigate comparability/validity of the 24HDR tools against reference methods. Some investigators used descriptive statistics such as the number of 'matches', 'intrusions' and 'omissions' between the food and drink items recorded by the web-based tool compared with the reference and in some cases linear and Poisson regression analysis was used to investigate the association of matches, intrusions and omissions between the test and reference<sup>(5,19,20,47)</sup>. Correlation analysis (Pearson, Spearman,  $\kappa$  and intraclass coefficients) was the most popular statistical method used to investigate the validity of nutrient intake and in some instances food group intake<sup>(49)</sup> reported by 24HDR tools compared with the output of reference methods. A wide range of correlation coefficients was reported by investigators. De Keyzer *et al.*<sup>(50)</sup> reported low correlation coefficients ( $r$  0.16;  $P$  < 0.001) for intakes of thiamine recorded by EPIC-Soft when compared with intakes from an estimated food diary.

**Table 3.** Main findings of the validation/comparison studies of the web- and computer-based 24-h dietary recall tools developed for children

Author	Tool name	Study type	Subjects (n)	No. of recalls	Reference method(s)	Statistical analysis	Main results for food groups	Main results for nutrients
Moore <i>et al.</i> (2008) <sup>(21)</sup>	SNAP™	Comparison	121	1 d	Interviewer-based 24HR	Mean differences in frequency (defined as count) of consumption. Bland–Altman was used to evaluate agreement between reported intakes from the test and reference method	The mean difference was less than 1 count for all but three categories – cakes (1.15 counts), energy-dense foods (1.52 counts) and CHO-rich foods (0.97 counts)	
Carvalho <i>et al.</i> (2014) <sup>(22)</sup>	PAC24	Validation	41	1 d	Direct dietary observation	All food items identified as matches, intrusions and omissions. Bland–Altman was used to evaluate agreement between reported intakes from the test and reference method	An average match rate of 67.0 %. Levels of intrusions and omissions were 11.5 and 21.5 %, respectively. 32 % of the actual intake was underestimated by PAC24	
Storey <i>et al.</i> (2011) <sup>(23)</sup>	Web-SPAN	Comparison	459	2 d	3 d estimated dietary record	ICC was used to measure the strength of agreement between the test and reference method. Pearson's correlations were used to assess the association between intakes reported by the test and reference method. Paired-samples <i>t</i> tests were used to investigate differences between both measures		ICC ranged from 0.24 to 0.40. Pearson <i>r</i> values ranged from 0.33 (fat) to 0.41 (protein) and for micronutrients 0.24 (vitamin A) to 0.39 (vitamin D). No significant differences were observed for protein, fat, Fe and Zn
Foster (2014) <sup>(5)</sup>	Intake24	Comparison	52	4 d	Interviewer-based 24HR	Bland–Altman was used to evaluate agreement between reported intakes from the test and reference method. All food items identified as matches, intrusions and omissions	Of all foods listed, 79.2 % were matches, 12.5 % omissions and 6.7 % intrusions. EI underestimated on average by 3 %. LOA for energy range from 0.52 to 1.82	LOA for CHO, fat, vitamin C and Fe were 0.52–1.88, 0.43–1.96, 0.44–2.71 and 0.45–2.11, respectively
Baranowski <i>et al.</i> (2002) <sup>(20)</sup>	FIRSt	Validation	138	1 d	Direct dietary observation and interviewer-based 24HR	All food items identified as matches, intrusions and omissions. Pearson's correlations were used to assess the association between intakes reported by the test and reference method	Compared with the lunch observation, FIRSt attained 46 % matches, 24 % intrusions and 30 % omissions. FIRSt attained 60 % matches, 15 % intrusions and 24 % omissions against the interviewer-based 24HR for all meals	
Diep <i>et al.</i> (2015) <sup>(19)</sup>	ASA24-Kids-2012	Validation	69	1 d	Direct dietary observation and interviewer-based 24HR	All food items identified as matches, intrusions and omissions. Pearson's correlations were used to assess the association between intakes reported by the test and reference method	Match, intrusion and omission rates were 37, 27 and 35 %, respectively (site 1) and 53, 12 and 36 %, respectively (site 2) compared with observed intakes. Percentage matches between ASA24 kids and interviewer-led method were higher	
Albar <i>et al.</i> (2016) <sup>(53)</sup>	myfood24	Comparison	75	2 d	Interviewer-based 24HR	ICC was used to measure the strength of agreement between the test and reference method. Bland–Altman was used to evaluate agreement between reported intakes from the test and reference method		ICC ranged from 0.46 for Na to 0.88 for EI. No significant bias between the two methods for EI and macronutrients. The mean difference between the test and reference measure (EI) was –230 kJ

SNAP™, Synchronised Nutrition and Activity Program (a web-based 24-h recall program which assesses energy balance-related behaviours in children and adolescents); 24HR, 24-h recall; CHO, carbohydrate; PAC24, Portuguese self-administered, computerised, 24-h dietary recall (a web-based 24-h dietary recall program for children); Web-SPAN, Web-Survey of Physical Activity and Nutrition (a web-based 24-h dietary recall program for children and adolescents); ICC, intraclass correlation coefficient; Intake24, a web-based 24-h dietary recall program for children and young adults; EI, energy intake; LOA, limits of agreement; FIRSt, Food Intake Recording Software System (a computer-based 24-h dietary recall program for children); ASA24-Kids-2012, Automated Self-Administered 24-Hour Dietary Assessment Tool, Kids-2012 version; myfood24, Measure your food on One day (a web-based 24-h dietary recall program for children and adults).

In contrast, Touvier *et al.*<sup>(17)</sup> reported a high correlation coefficient ( $r$  0.92;  $P < 0.001$ ) for niacin intakes recorded by women using NutriNet-Santé and an interviewer-led 24HDR. Liu *et al.*<sup>(29)</sup> also reported a mean correlation coefficient ( $r$  0.6;  $P < 0.001$ ) for twenty-one nutrients examined to demonstrate the relationship between the test and reference measure across a range of nutrients. In addition to correlation analysis, Storey *et al.*<sup>(23)</sup> and De Keyzer *et al.*<sup>(50)</sup> used paired  $t$  tests and Student's  $t$  tests to investigate significant differences in the reporting of nutrient intakes between the test and reference methods.

Bland & Altman<sup>(51)</sup> analysis was only used by six investigators, four of which investigated the agreement and bias in nutrient intakes recorded by test and reference methods<sup>(5,27,52,53)</sup>. Moore *et al.*<sup>(21)</sup> investigated agreements between methods in the recording of food and drink items and Carvalho *et al.*<sup>(22)</sup> investigated estimated *v.* observed amounts consumed using Bland–Altman analysis.

The investigators of the 24HDR tools developed for children, included in this review, did not use biomarkers of nutrient intake to investigate validity. Table 4 shows the 24HDR tools developed for adults including four 24HDR tools that were validated using biological markers for nutrient intake. DietDay<sup>(33)</sup> is the only 24HDR tool that used doubly labelled water (DLW) in validation as a biomarker of usual energy intake<sup>(54)</sup>. Results indicated that the validity of DietDay was more superior than a paper-based FFQ or diet history questionnaire. Three investigators (Wardenaar *et al.*<sup>(52)</sup>, Ferrari *et al.*<sup>(43)</sup> and Lassale *et al.*<sup>(55)</sup>) used urinary biomarkers (urinary N, urinary N and K and urinary N, K and Na, respectively) to investigate the relationship between reported and true intakes. The investigators reported moderate to strong correlations between reported nutrient intakes and their respective urinary biomarkers:  $r$  0.37 (95 % CI 0.03, 0.70) for Na intakes reported by women<sup>(55)</sup>,  $r$  0.65 (95 % CI 0.45, 0.79) for protein intakes reported by athletes<sup>(52)</sup> and  $r$  0.83 (95 % CI 0.637, 0.932) for reported N intakes<sup>(43)</sup>.

## Discussion

### Tool design characteristics

The range of ages involved in the testing of the 24HDR tools demonstrates the applicability of web- and computer-based dietary assessment methods to the majority of the population. Investigators did note age-specific adaptations to some of the 24HDR tools including the use of cartoon avatars<sup>(56,57)</sup> and portion size pictures specifically developed for children<sup>(56,57)</sup>. Baranowski *et al.*<sup>(57)</sup> and subsequently Diep *et al.*<sup>(19)</sup> used adapted versions of the National Cancer Institute's Automated Self-Administered 24-Hour Dietary Assessment Tool (ASA24<sup>®</sup>) which had been modified for children and excluded food preparation questions deemed too difficult for children to answer<sup>(58)</sup>. However, none of the investigators noted changes to the 24HDR tools that would potentially enhance the user experience for older adults (for example, larger format or font). At present, there is no single 24HDR tool suitable for all age groups; however, some investigators such as Baranowski *et al.*<sup>(57)</sup>, Subar *et al.*<sup>(32)</sup>, Hillier *et al.*<sup>(49)</sup> and Moore *et al.*<sup>(45)</sup>

have worked to develop separate age-appropriate versions of the same tool. Foster<sup>(5)</sup> also tested the same tool with children and with young adults and Albar *et al.*<sup>(53)</sup> proposes to test myfood24 with adults aged 18 years and over. Intuitive design features such as 'same as before', 'meal gap review' and spell check function were unique to some 24HDR tools but could potentially be useful in all web- and computer-based dietary assessment designs (whether it be diary or FFQ based).

### Performance

For the majority of 24HDR tools included in this review, the investigators concluded that the performance of the respective tool was acceptable when compared with reference methods (comparison and/or validation studies). There was a variety of concluding remarks from the investigators: 'The test measure was in good agreement with the reference method'<sup>(5,17,23,27,29)</sup>; 'The test method performed well but in some instances the reference method was more accurate'<sup>(19,20,47)</sup>; 'The test measure was more superior to the reference measure'<sup>(33)</sup>; and 'Although the findings were promising the test method required further modifications to improve accuracy'<sup>(22,45,49)</sup>.

Overall, the 24HDR tools whose performance was investigated, either in a comparison and/or validation study, demonstrated promising potential. Investigators that reported the reference method as being more accurate used direct observation as the reference method<sup>(19,47,59)</sup>. Whilst this objectively investigates the accuracy of the test measure, for example in the case of Kirkpatrick *et al.*<sup>(47)</sup>, it may lead to the underestimation of the performance of ASA24 if compared with how other 24HDR tools were assessed for accuracy, for example, Zoellner *et al.*<sup>(60)</sup>. The performance of the ASA24 may be underestimated as the bilingual interactive multimedia (IMM) dietary assessment tool was compared against an interviewer-led 24-h recall, the findings of which may agree more favourably compared with direct observation. As eating is an observable behaviour<sup>(61)</sup>, direct observation of eating occasions is an expensive and invasive measure and is usually only feasible for short periods of time (for example, lunch or dinner times) which may not accurately represent the performance of the tool for recording longer periods of intake (for example, 24-h period)<sup>(19)</sup>. Recovery biomarkers measure dietary intake with little error and have been successfully used to validate some of the studies included in this review; however, these biomarkers are limited to certain nutrients and cannot validate all nutrient estimates from a test dietary assessment measure. Based on the findings of this review, the most appropriate use of reference measures for investigating the validity of a 24HDR tool is the combination of an estimated dietary record in conjunction with biological markers of intake or direct observation of eating occasions as undertaken by Arab *et al.*<sup>(33)</sup> and Baranowski *et al.*<sup>(20)</sup>, respectively.

In the assessment of the performance of 24HDR tools, acceptability of the tool is an aspect that is not always considered. In the comparison of ASA24 with an interviewer-led 24-h recall, Thompson *et al.*<sup>(62)</sup> noted that a substantial proportion of study participants preferred the self-administered ASA24 approach compared with the interviewer-led recall.

**Table 4.** Main findings of the validation/comparison studies of the web- and computer-based 24-h dietary recall tools developed for adults

Author	Tool name	Study type	Subjects (n)	No. of recalls	Reference method	Summary of statistical analysis	Main results for food groups	Main results for nutrients
De Keyzer <i>et al.</i> (2011) <sup>(50)</sup>	EPIC-Soft	Comparison	127	2 d	5 d estimated dietary record	Student's <i>t</i> test was used to investigate whether intakes reported by the test and reference measure differed significantly. Spearman and $\kappa$ correlations were used to assess the association between intakes reported by the test and reference method		Macronutrient correlation coefficients ranged from 0.47 (protein) to 0.62 (CHO) and for micronutrients 0.16 (thiamine) to 0.56 (Fe). Significant differences between methods for the reporting of ten nutrients, for example, fat
Ferrari <i>et al.</i> (2009) <sup>(43)</sup>	EPIC-Soft	Validation	1072	1 d	24-h urinary biomarkers	Pearson correlations were used to assess the association between intakes reported by the test and the respective biological value		Correlation coefficients between means of urinary and 24HR measurements were 0.838 and 0.756 for N and K intakes, respectively
Touvier <i>et al.</i> (2011) <sup>(17)</sup>	NutriNet-Santé	Comparison	147	1 d	Interviewer-based 24HR	ICC was used to measure the strength of agreement between the test and reference method. Pearson's correlations were used to assess the association between intakes reported by the test and reference method	ICC ranged from 0.5 for fats/sauces (both sexes), breakfast cereals, cakes/biscuits/pastries and dairy food (women only) to 0.9 for fruits, pulses (both sexes), breakfast cereals, alcoholic drinks and meat (men only)	The range for energy-adjusted Pearson's coefficients for macronutrients was 0.80 (protein intake recorded by men) to 0.88 (protein intake recorded by women) and for micronutrients was 0.54 (retinol recorded by men) to 0.92 (niacin recorded by women)
Lassale <i>et al.</i> (2015) <sup>(55)</sup>	NutriNet-Santé	Validation	199	3 d	24-h urinary biomarkers	ICC was used to measure the strength of agreement between the test and reference method. Pearson's correlations were used to assess the association between intakes reported by the test and reference method		Correlations between reported and true intakes were 0.61, 0.78 and 0.47 for men and 0.64, 0.42 and 0.37 for women for protein, K and Na, respectively. Attenuation factors ranged from 0.23 (Na, women) to 0.60 (K, men)
Foster (2014) <sup>(5)</sup>	Intake24	Comparison	167	4 d	Interviewer-based 24HR	Bland-Altman was used to evaluate agreement between reported intakes from the test and reference method. All food items identified as matches, intrusions and omissions	For all foods listed, 82.6 % were matches, 9 % omissions and 7.5 % intrusions. Energy intake under-estimated on average by 1 %. LOA for energy range from 0.50 to 1.97	LOA for CHO, fat, vitamin C, Fe and alcohol were 0.51–1.99, 0.43–2.31, 0.18–5.75, 0.43–2.24 and 0.09–10.91, respectively
Hillier <i>et al.</i> (2012) <sup>(49)</sup>	SNAPA™	Validation	77	5 d	Direct dietary observation on 4 d	All food items identified as matches, intrusions and omissions. Pearson's correlation was used to assess the association between intakes reported by the test and reference method	The mean match rate was 81.7 %, with an intrusion rate of 5.6 %. Pearson's correlations ranged from 0.39 to 0.56 for percentage fat and fruit and vegetable intake respectively	
Comrie <i>et al.</i> (2009) <sup>(27)</sup>	FoRC	Comparison	53	4 d	4 d estimated dietary record	Spearman's correlation was used to assess the association between intakes reported by the test and reference method. Wilcoxon signed-rank test was used to investigate whether intakes reported by the test and reference measure differed significantly. Bland-Altman was used to evaluate agreement between reported intakes from the test and reference method	Intakes of fat, NSP and bread were similar between the two methods. FoRC recorded significantly lower intakes of energy and alcohol and higher intakes of fruit and vegetables and cereals	Correlation coefficients at a food-group level ranged from 0.4 for alcohol to 0.76 for bread. For the few nutrients investigated, all <i>r</i> values were statistically significant and were greater than 0.5 (except for percentage fat)
Liu <i>et al.</i> (2011) <sup>(29)</sup>	Oxford WebQ	Comparison	116	1 d	Interviewer-based 24HR	The percentage differences in energy and nutrient intake between the two methods were calculated. Spearman's correlation was used to investigate agreement between estimates for each nutrient. Comparison of tertile intakes from the test and reference measure to identify misreporters		Mean Spearman's correlation for the 21 nutrients was 0.6. Macronutrient <i>r</i> values ranged from 0.57 (fat) to 0.66 (CHO) and for micronutrients 0.37 (vitamin E) to 0.72 (Mg). The mean differences in intake were less than 10 % for all nutrients except carotene and vitamins B <sub>12</sub> and D

**Table 4.** *Continued*

Author	Tool name	Study type	Subjects (n)	No. of recalls	Reference method	Summary of statistical analysis	Main results for food groups	Main results for nutrients
Zoellner <i>et al.</i> (2005) <sup>(60)</sup>	IMM	Comparison	80	1 d	Interviewer-based 24 HR	ANOVA tests were performed to determine the effect of substituting standardised portion sizes for reported portion sizes. Unadjusted and energy-adjusted correlations were calculated to measure the strength of relationship between the IMM recall and the interview-administered recall		Unadjusted correlation coefficients were 0.6 and energy-adjusted correlations were lower. Macronutrient unadjusted <i>r</i> values ranged from 0.44 (fat) to 0.78 (CHO) and for micronutrients 0.29 (folate) to 0.7 (Ca)
Kirkpatrick <i>et al.</i> (2014) <sup>(47)</sup>	ASA24	Validation	81	3 d	Feeding study, in which the true intake for three meals was known	Linear regression models were used to examine the association between recall mode and the proportion of items truly consumed for which a match was reported. Poisson regression was used to assess the association between recall mode and the number of items reported but not truly consumed	Average match rate of 80 % between the recall and the reference method. Mean number of intrusions for all eating occasions combined was 2.6 for the ASA24	
Arab <i>et al.</i> (2011) <sup>(33)</sup>	DietDay	Validation	261	6 d	Double-labelled water and diet history questionnaire	The ratio of reported intakes to TEE measurements was calculated to identify reporting bias and misreporters. Pearson correlation for agreement between recall and biomarker. Attenuation factor was estimated by using the regression coefficient from the regression model	Attenuation factors were 0.30 for blacks and 0.26 for whites. Adjusted correlations between true energy intake and the recalls were 0.50 and 0.47 for blacks and whites, respectively	
Wardenaar <i>et al.</i> (2015) <sup>(52)</sup>	Compl-eat™	Validation	47	3 d	24-h urinary N	Paired <i>t</i> tests were used to determine difference between biomarker value and reported intake. Pearson correlations were used to rank individuals according to intake. Attenuation factors and CI (calculated using slope of the linear regression of log transformed biomarker values and reported intakes). Bland–Altman was used to evaluate agreement between reported intakes and biological value		Estimated mean dietary protein intake was 109.6 (SD 33.0) g/d according to the test measure compared with 141.3 (SD 38.2) g/d as determined by N excretion. Reasonably good association of intakes for protein estimation (0.65, 95 % CI 0.45, 0.79). Under-reporting of protein intakes was larger with higher intakes of protein <i>v.</i> lower intakes

EPIC-Soft, European Prospective Investigation into Cancer and Nutrition Software (a standardised computerised 24-h dietary recall method); CHO, carbohydrate; 24HR, 24-h recall; NutriNet-Santé, a web-based 24-h dietary recall program for adults; ICC, intraclass correlation coefficient; Intake24, a web-based 24-h dietary recall program for children and young adults; LOA, limits of agreement; SNAPA™, Synchronised Nutrition and Activity Program for Adults (a web-based 24-h dietary recall program for adults); FoRC, Food Recall Checklist (121-item food recall list which has been used to measure diet in undergraduate students); Oxford WebQ, a web-based 24-h dietary recall program for adults; IMM, a bilingual interactive multimedia dietary assessment tool based on the 24-h dietary recall method (a web-based 24-h dietary recall program for adults); ASA24, Automated Self-Administered 24 Hour Dietary Assessment Tool (web-based 24-h dietary recall tool for adults, developed by the National Cancer Institute, USA); DietDay, a web-based 24-h dietary recall program for adults; TEE, total energy expenditure; Compl-eat™: a computer-based 24-h dietary recall program for adults.

However, Vereecken *et al.*<sup>(48)</sup> highlighted the opposite, when parents preferred a traditional approach to recording diet (i.e. a written record) compared with a web- or computer-based tool. Therefore, to ensure prolonged usage of web- and computer-based dietary assessment methods, research into the acceptability of these methods should be considered in the comparison/validation study design so that improvements can be made if warranted.

### Accuracy/precision

Although the accuracy of the 24HDR tools is difficult to compare across the board, one can consider the results reported within the individual studies. It is known that one statistical measure in isolation cannot determine the validity of any method of dietary assessment<sup>(36)</sup>. However, due to the variation and combination of statistical measures used across studies, for the purposes of this review statistical measures are compared in isolation across studies. Correlation coefficients are commonly reported across studies investigating the accuracy of 24HDR tools. Comparing the ranges of correlation coefficients for macronutrients and micronutrients reported across all 24HDR tools, NutriNet-Sant <sup>(17)</sup> had the strongest *r* values for intakes derived from the web-based tool compared with a reference measure, with *r* values ranging for macronutrient intake as high as 0.88 (protein intakes recorded by women) and for micronutrient intake an *r* value of 0.92 (niacin recorded by women). As with all studies investigating the comparability/validity of a test measure of dietary assessment compared with a reference, it is important to consider the study design. Touvier *et al.*<sup>(17)</sup> compared intakes of NutriNet-Sant  against intakes derived from an interviewer-led dietary recall at a single time point, whereas De Keyzer *et al.*<sup>(50)</sup> compared intakes across two separate time points and Comrie *et al.*<sup>(27)</sup> compared intake across four separate time points. It is, therefore, difficult to make conclusions about accuracy of one tool compared with another, particularly using just one type of statistical analysis. The number of recalls administered across the comparison/validation studies varied from a single dietary recall<sup>(21)</sup> to six recalls<sup>(33)</sup>. The difference in the number of recalls administered compared with the number of days of dietary recording by the reference method is important to consider when investigating the agreement between methods in their assessment of nutrient intake.

However, when comparing correlation coefficients from studies that have similar study designs (for example, Liu *et al.*<sup>(29)</sup> (*r* 0.37 to 0.72); Touvier *et al.*<sup>(17)</sup> (*r* 0.54 to 0.92); Zoellner *et al.*<sup>(60)</sup> (*r* 0.29 to 0.78)), all of whom used a single time point of data collection in their investigation and used an interviewer-led recall as a reference, Touvier *et al.*<sup>(17)</sup> performed the best based on this type of analysis. Another type of measure that was used to investigate validity of 24HDR tools was the incidence of 'Matches, Intrusions and Omissions' whereby food and drink items recorded using the test measure are compared with the reference measure. Again, there was variation across studies depending on the reference measure used; for example comparing the output of the web-based tool against direct observation, the number of percentage matches was not as high (ranging from 46 to 67%)<sup>(20,22)</sup> when compared

with percentage matches identified when comparing the web-based tool with interviewer-led recalls (79.2 and 82.6% both)<sup>(5)</sup>.

Precision (the degree to which the same method produces the same value on repeated measures) was less frequently reported. Based on the limited number of investigators that calculated the limits of agreement in their analysis, it is not possible to compare the precision of the methods included in this review. This type of analysis was included in a report on the performance of INTAKE24 (a web-based 24-h dietary recall program for children and young adults)<sup>(5)</sup>. Foster<sup>(5)</sup> assessed the precision of INTAKE24 compared with other web- and computer-based methods of dietary assessment, which had included these findings in their analysis, by calculating the width of the limits of agreement. The lack of this statistical interpretation in dietary assessment validation studies was also noted by Lombard *et al.*<sup>(36)</sup>.

Another factor affecting the design of the comparison/validation study was the proximity and sequence of the test and reference measure. Margetts & Nelson<sup>(63)</sup> suggested that the sequence (i.e. that the test measure be administered before the reference method) and the proximity of the test and reference measure need to be carefully considered to avoid raising the apparent level of agreement between the measures. The cross-over design is a popular design for testing the performance of 24HDR tools<sup>(5,24,60)</sup>. The design involves the use of the test and reference method in the same day. Although this limits the impact of variation of diet on the results, completing both methods in the same day may introduce a 'learning effect'. Having acknowledged this in the study design, Foster<sup>(5)</sup> attempted to limit this effect by weighting the order in which the test measure was administered to participants so that the impact of a 'learning effect' could be investigated retrospectively. Interestingly, although the weighed food record is considered the 'gold standard' (in the absence of unbiased measures) of dietary assessment<sup>(60)</sup>, none of the investigators included this as a reference method. Alternatives may have been used instead due to the high respondent burden and cost associated with weighed food diaries<sup>(64)</sup>.

Biological markers of nutrient intake can serve as an objective validation of dietary assessment methods as they reflect nutritional status, but are independent of dietary intake assessment<sup>(65)</sup>. Of all the tools included in this review, Arab *et al.*<sup>(33)</sup> was the only investigator to use DLW to assess the validity. Whilst DLW is considered the 'gold standard' recovery biomarker of nutrient intake, urinary N<sup>(66)</sup> and urinary K<sup>(67)</sup> are also useful biomarkers in the validation of dietary assessment tools. Biomarkers of nutrient intake from blood were not used by any of the investigators included in this review. Biomarkers of nutrient intake from plasma and serum samples (for example, plasma vitamin C and plasma carotenoids) have been used in other validation studies concerning novel dietary assessment methodologies<sup>(13,68,69)</sup>. In the studies investigating the validity of 24HDR tools in children (included in the present review), none of the investigators used biological markers of nutrient intake. A reason for this may be that biological samples are difficult to obtain from children<sup>(70)</sup>. Baranowski *et al.*<sup>(20)</sup> used other strategies such as obtaining a bogus pipeline hair sample from children using the FIRSSt tool to encourage more accurate

reporting of dietary intake. Using this method, the researchers informed participants that the hair sample provided would reveal what they had eaten through chemical analysis. Collecting this sample appeared to reduce the level of omissions when intakes from FIRSSt were compared with an interviewer-administered recall<sup>(20)</sup>.

## Conclusion

Web- and computer-based 24HDR tools are cost-effective, useful methods for assessing dietary intake from different populations. The overall accuracy of these methods is difficult to determine because, in many cases, direct comparisons cannot be made. Across-the-board agreement between these 24HDR tools for macro- and micronutrient intakes is generally strong when compared against references. Few studies used biomarkers in the validation of 24HDR tools; however, the investigators that did use biomarkers showed that these methods provide reliable estimates of protein intake, moderate estimates of energy intakes and reliable estimates for micronutrient intake such as K and Na. This review highlights some findings which may be applied when designing and investigating the performance of 24HDR tools in the future. Age-specific adaptations have been shown to be of benefit in younger populations, and may prove beneficial for older adult populations also. In the assessment of performance and validity, direct observation is a useful reference method; however, it is often limited to short periods of intake, whereas estimated records in conjunction with biological markers of intake may be more feasible methods of investigating validity. It is important to be mindful of the type of statistical analysis used to investigate validity. Lombard *et al.*<sup>(36)</sup> noted that in some cases more than three statistical measures are required to truly assess the validity of dietary assessment measures. Lastly, this review demonstrates that although a standardised dietary assessment methodology is preferential for nutrition surveillance across countries, valuable technology-based 24HDR tools exist and could be used to collect intermittent data for continual health and nutrition research purposes.

## Acknowledgements

The present review was completed as part of a Food Institutional Research Measure (FIRM) project, DietIreland, funded by the Department of Agriculture, Food and Marine (13/F/424).

The present review was completed as part of a project conceived by E. R. G. and A. F.; C. M. T. and E. R. G. devised the rationale and concept of the review; R. v. d. B., R. J. B., L. K., K. E. and J. W. were involved in the data collection for the review; C. M. T. drafted the manuscript. All authors approved the final content of the manuscript.

The authors declare no conflict of interest.

## References

1. Kirkpatrick SI & Collins CE (2016) Assessment of nutrient intakes: introduction to the special issue. *Nutrients* **8**, 184.
2. Arens-Volland AG, Spassova L & Bohn T (2015) Promising approaches of computer-supported dietary assessment and

management: current research status and available applications. *Int J Med Inform* **84**, 997–1008.

3. Six BL, Schap TE, Zhu FM, *et al.* (2010) Evidence-based development of a mobile telephone food record. *J Am Diet Assoc* **110**, 74–79.
4. Wang D-H, Kogashiwa M, Ohta S, *et al.* (2002) Validity and reliability of a dietary assessment method: the application of a digital camera with a mobile phone card attachment. *J Nutr Sci Vitaminol* **48**, 498–504.
5. Foster E (2014) Comparison study: INTAKE24 vs interviewer led recall. Final Report for Food Standards Agency. <http://www.food.gov.uk/scotland/news-updates/news/2014-13135/intake24> (accessed June 2016).
6. Vereecken C, Covents M, Maes L, *et al.* (2014) Formative evaluation of the dietary assessment component of Children's and Adolescents' Nutrition Assessment and Advice on the Web (CANAA-W). *J Hum Nutr Diet* **27**, Suppl. 1, 54–65.
7. Rollo ME, Ash S, Lyons-Wall P, *et al.* (2015) Evaluation of a mobile phone image-based dietary assessment method in adults with type 2 diabetes. *Nutrients* **7**, 4897–4910.
8. Harray AJ, Boushey CJ, Pollard CM, *et al.* (2015) A novel dietary assessment method to measure a healthy and sustainable diet using the mobile food record: protocol and methodology. *Nutrients* **7**, 5375–5395.
9. Sun M, Burke LE, Mao Z-H, *et al.* (2014) eButton: a wearable computer for health monitoring and personal assistance. In *Proceedings of the The 51st Annual Design Automation Conference on Design Automation Conference - DAC 2014*. New York: ACM.
10. Lee CD, Chae J, Schap TE, *et al.* (2012) Comparison of known food weights with image-based portion-size automated estimation and adolescents' self-reported portion size. *J Diabetes Sci Technol* **6**, 428–434.
11. Barker ME, Blain RJ & Russell JM (2015) The influence of academic examinations on energy and nutrient intake in male university students. *Nutr J* **14**, 98.
12. Illner A-K, Freisling H, Boeing H, *et al.* (2012) Review and evaluation of innovative technologies for measuring diet in nutritional epidemiology. *Int J Epidemiol* **41**, 1187–1203.
13. Fallaize R, Forster H, Macready AL, *et al.* (2014) Online dietary intake estimation: reproducibility and validity of the Food4Me food frequency questionnaire against a 4-day weighed food record. *J Med Internet Res* **16**, e190.
14. Labonté M-È, Cyr A, Baril-Gravel L, *et al.* (2012) Validity and reproducibility of a web-based, self-administered food frequency questionnaire. *Eur J Clin Nutr* **66**, 166–173.
15. Matthys C, Pynaert I, De Keyzer W, *et al.* (2007) Validity and reproducibility of an adolescent web-based food frequency questionnaire. *J Am Diet Assoc* **107**, 605–610.
16. Kristal AR, Kolar AS, Fisher JL, *et al.* (2014) Evaluation of web-based, self-administered, graphical food frequency questionnaire. *J Acad Nutr Diet* **114**, 613–621.
17. Touvier M, Kesse-Guyot E, Méjean C, *et al.* (2011) Comparison between an interactive web-based self-administered 24 h dietary record and an interview by a dietitian for large-scale epidemiological studies. *Br J Nutr* **105**, 1055–1064.
18. Foster E, Hawkins A, Simpson E, *et al.* (2014) Developing an interactive portion size assessment system (IPSAS) for use with children. *J Hum Nutr Diet* **27**, Suppl. 1, 18–25.
19. Diep CS, Hingle M, Chen T-A, *et al.* (2015) The Automated Self-Administered 24-Hour Dietary Recall for Children, 2012 Version, for youth aged 9 to 11 years: a validation study. *J Acad Nutr Diet* **115**, 1591–1598.
20. Baranowski T, Islam N, Baranowski J, *et al.* (2002) The food intake recording software system is valid among fourth-grade children. *J Am Diet Assoc* **102**, 380–385.

21. Moore HJ, Eells LJ, McLure SA, *et al.* (2008) The development and evaluation of a novel computer program to assess previous-day dietary and physical activity behaviours in school children: the Synchronised Nutrition and Activity Program (SNAP). *Br J Nutr* **99**, 1266–1274.
22. Carvalho MA, Baranowski T, Foster E, *et al.* (2014) Validation of the Portuguese self-administered computerised 24-hour dietary recall among second-, third- and fourth-grade children. *J Hum Nutr Diet* **28**, 666–674.
23. Storey KE & McCargar LJ (2012) Reliability and validity of Web-SPAN, a web-based method for assessing weight status, diet and physical activity in youth. *J Hum Nutr Diet* **25**, 59–68.
24. Subar AF, Thompson FE, Potischman N, *et al.* (2007) Formative research of a quick list for an automated self-administered 24-hour dietary recall. *J Am Diet Assoc* **107**, 1002–1007.
25. Arab L, Wesseling-Perry K, Jardack P, *et al.* (2010) Eight self-administered 24-hour dietary recalls using the Internet are feasible in African Americans and Whites: the energetics study. *J Am Diet Assoc* **110**, 857–864.
26. Slimani N, Casagrande C, Nicolas G, *et al.* (2011) The standardized computerized 24-h dietary recall method EPIC-Soft adapted for pan-European dietary monitoring. *Eur J Clin Nutr* **65**, Suppl. 1, S5–S15.
27. Comrie F, Masson LF & McNeill G (2009) A novel online food recall checklist for use in an undergraduate student population: a comparison with diet diaries. *Nutr J* **8**, 13.
28. Shin S, Park E, Sun DH, *et al.* (2014) Development and evaluation of a web-based computer-assisted personal interview system (CAPIS) for open-ended dietary assessments among Koreans. *Clin Nutr Res* **3**, 115–125.
29. Liu B, Young H, Crowe FL, *et al.* (2011) Development and evaluation of the Oxford WebQ, a low-cost, web-based method for assessment of previous 24 h dietary intakes in large-scale prospective studies. *Public Health Nutr* **14**, 1998–2005.
30. Moshfegh A, Raper N, Ingwersen L, *et al.* (2001) An improved approach to 24-hour dietary recall methodology. *Ann Nutr Metab* **45**, Suppl., 156 Abstr.
31. Carter MC, Albar SA, Morris MA, *et al.* (2015) Development of a UK online 24-h dietary assessment tool: myfood24. *Nutrients* **7**, 4016–4032.
32. Subar AF, Kirkpatrick SI, Mittl B, *et al.* (2012) The Automated Self-Administered 24-hour dietary recall (ASA24): a resource for researchers, clinicians, and educators from the National Cancer Institute. *J Acad Nutr Diet* **112**, 1134–1137.
33. Arab L, Tseng C-H, Ang A, *et al.* (2011) Validity of a multipass, web-based, 24-hour self-administered recall for assessment of total energy intake in blacks and whites. *Am J Epidemiol* **174**, 1256–1265.
34. Daniel CR, Kapur K, McAdams MJ, *et al.* (2014) Development of a field-friendly automated dietary assessment tool and nutrient database for India. *Br J Nutr* **111**, 160–171.
35. Park E, Shin S, Hwang S, *et al.* (2009) Web-based dietary assessment software for 24-hour recall interview. *FASEB J* **23**, 223.1.
36. Lombard MJ, Steyn NP, Charlton KE, *et al.* (2015) Application and interpretation of multiple statistical tests to evaluate validity of dietary intake assessment methods. *Nutr J* **14**, 40.
37. Ratz SK, Scheett AJ, Johnson LK, *et al.* (2015) Validity of electronic diet recording nutrient estimates compared to dietitian analysis of diet records: randomized controlled trial. *J Med Internet Res* **17**, e21.
38. De Keyzer W, Bracke T, McNaughton SA, *et al.* (2015) Cross-continental comparison of national food consumption survey methods – a narrative review. *Nutrients* **7**, 3587–3620.
39. Statistics Canada (2012) Inventory of Food Consumption Surveys. <http://www5.statcan.gc.ca/cansim/pick-choisir?lang=eng&p2=33&id=1052012> (accessed June 2016).
40. United States Department of Agriculture (2004) What We Eat in America, NHANES. <http://www.ars.usda.gov/Services/docs.htm?docid=13793> (accessed June 2016).
41. European Food Safety Authority (2009) General principles for the collection of national food consumption data in the view of a pan-European dietary survey. *EFSA J* **7**, 1435.
42. Subar AF, Kirkpatrick SI, Thompson FE, *et al.* (2014) The web-based automated self-administered 24-hour dietary recall performs similarly to a traditional interviewer-administered 24-hour dietary recall. *Circulation* **129**, Suppl. 1, AMP31.
43. Ferrari P, Roddam A, Fahey MT, *et al.* (2009) A bivariate measurement error model for nitrogen and potassium intakes to evaluate the performance of regression calibration in the European Prospective Investigation into Cancer and Nutrition study. *Eur J Clin Nutr* **63**, Suppl. 4, S179–S187.
44. Gemming L, Utter J & Ni Mhurchu C (2015) Image-assisted dietary assessment: a systematic review of the evidence. *J Acad Nutr Diet* **115**, 64–77.
45. Moore HJ, Hillier FC, Batterham AM, *et al.* (2014) Technology-based dietary assessment: development of the Synchronised Nutrition and Activity Program (SNAP). *J Hum Nutr Diet* **27**, Suppl. 1, 36–42.
46. Freese J, Feller S, Hartig U, *et al.* (2014) Development and evaluation of a short 24-h food list as part of a blended dietary assessment strategy in large-scale cohort studies. *Eur J Clin Nutr* **68**, 324–329.
47. Kirkpatrick SI, Subar AF, Douglass D, *et al.* (2014) Performance of the Automated Self-Administered 24-hour Recall relative to a measure of true intakes and to an interviewer-administered 24-h recall. *Am J Clin Nutr* **100**, 233–240.
48. Vereecken CA, Covents M, Haynie D, *et al.* (2009) Feasibility of the Young Children's Nutrition Assessment on the Web. *J Am Diet Assoc* **109**, 1896–1902.
49. Hillier FC, Batterham AM, Crooks S, *et al.* (2012) The development and evaluation of a novel Internet-based computer program to assess previous-day dietary and physical activity behaviours in adults: the Synchronised Nutrition and Activity Program for Adults (SNAPA<sup>TM</sup>). *Br J Nutr* **107**, 1221–1231.
50. De Keyzer W, Huybrechts I, De Vriendt V, *et al.* (2011) Repeated 24-hour recalls versus dietary records for estimating nutrient intakes in a national food consumption survey. *Food Nutr Res* **2011**, 55.
51. Bland JM & Altman DG (1986) Statistical methods for assessing agreement between two methods of clinical measurement. *Lancet* **i**, 307–310.
52. Wardenaar FC, Steennis J, Ceelen IJM, *et al.* (2015) Validation of web-based, multiple 24-h recalls combined with nutritional supplement intake questionnaires against nitrogen excretions to determine protein intake in Dutch elite athletes. *Br J Nutr* **114**, 2083–2092.
53. Albar SA, Alwan NA, Evans CEL, *et al.* (2016) Agreement between an online dietary assessment tool (myfood24) and an interviewer-administered 24-h dietary recall in British adolescents aged 11–18 years. *Br J Nutr* **115**, 1678–1686.
54. Trabulsi J, Troiano RP, Subar AF, *et al.* (2003) Precision of the doubly labeled water method in a large-scale application: evaluation of a streamlined-dosing protocol in the Observing Protein and Energy Nutrition (OPEN) study. *Eur J Clin Nutr* **57**, 1370–1377.
55. Lassale C, Castetbon K, Laporte F, *et al.* (2015) Validation of a web-based, self-administered, non-consecutive-day dietary



- record tool against urinary biomarkers. *Br J Nutr* **113**, 953–962.
56. Biloft-Jensen A, Trolle E, Christensen T, *et al.* (2014) WebDASC: a web-based dietary assessment software for 8–11-year-old Danish children. *J Hum Nutr Diet* **27**, Suppl. 1, 43–53.
57. Baranowski T, Islam N, Douglass D, *et al.* (2014) Food Intake Recording Software System, version 4 (FIRSt4): a self-completed 24-h dietary recall for children. *J Hum Nutr Diet* **27**, Suppl. 1, 66–71.
58. Douglass D, Islam N, Baranowski J, *et al.* (2013) Simulated adaptations to an adult dietary self-report tool to accommodate children: impact on nutrient estimates. *J Am Coll Nutr* **32**, 92–97.
59. Baranowski T, Islam N, Baranowski J, *et al.* (2012) Comparison of a Web-based versus traditional diet recall among children. *J Acad Nutr Diet* **112**, 527–532.
60. Zoellner J, Anderson J & Gould SM (2005) Comparative validation of a bilingual interactive multimedia dietary assessment tool. *J Am Diet Assoc* **105**, 1206–1214.
61. Sherwood N (2008) Diet assessment in children and adolescence. In *Handbook of Childhood and Adolescent Obesity*, pp. 73–89 [E Jelalian and RG Steele, editors]. New York: Springer.
62. Thompson FE, Dixit-Joshi S, Potischman N, *et al.* (2015) Comparison of interviewer-administered and automated self-administered 24-hour dietary recalls in 3 diverse integrated health systems. *Am J Epidemiol* **181**, 970–978.
63. Margetts B & Nelson M (1997) *Design Concepts in Nutritional Epidemiology*, 2nd ed. Oxford: Oxford University Press.
64. Wrieden W, Peace H, Armstrong J, *et al.* (2003) A short review of dietary assessment methods used in National and Scottish Research Studies. <http://www.food.gov.uk/sites/default/files/multimedia/pdfs/scotdietassessmethods.pdf> (accessed May 2016).
65. McKeown N, Day N, Welch A, *et al.* (2001) Use of biological markers to validate self-reported dietary intake in a random sample of the European Prospective Investigation into Cancer United Kingdom Norfolk cohort. *Am J Clin Nutr* **74**, 188–196.
66. Bingham SA, Gill C, Welch A, *et al.* (1997) Validation of dietary assessment methods in the UK arm of EPIC using weighed records, and 24-hour urinary nitrogen and potassium and serum vitamin C and carotenoids as biomarkers. *Int J Epidemiol* **26**, Suppl. 1, S137–S151.
67. Tasevska N, Runswick SA & Bingham SA (2006) Urinary potassium is as reliable as urinary nitrogen for use as a recovery biomarker in dietary studies of free living individuals. *J Nutr* **136**, 1334–1340.
68. Patterson AC, Hogg RC, Kishi DM, *et al.* (2012) Biomarker and dietary validation of a Canadian food frequency questionnaire to measure eicosapentaenoic and docosahexaenoic acid intakes from whole food, functional food, and nutraceutical sources. *J Acad Nutr Diet* **112**, 1005–1014.
69. Timon CM, Astell AJ, Hwang F, *et al.* (2015) The validation of a computer-based food record for older adults: the Novel Assessment of Nutrition and Ageing (NANA) method. *Br J Nutr* **113**, 654–664.
70. Rockett J, Lynch C & Buck GM (2005) Home-based collection of biological measurements and specimens from women and children. *Epidemiology* **16**, 131.