

**UCC Library and UCC researchers have made this item openly available. Please [let us know](#) how this has helped you. Thanks!**

<b>Title</b>	Trial of labour after caesarean section and the risk of neonatal and infant death: a nationwide cohort study
<b>Author(s)</b>	O'Neill, Sinéad M.; Agerbo, Esben; Khashan, Ali S.; Kearney, Patricia M.; Henriksen, Tine B.; Greene, Richard A.; Kenny, Louise C.
<b>Publication date</b>	2017-02-27
<b>Original citation</b>	O'Neill, S. M., Agerbo, E., Khashan, A. S., Kearney, P. M., Henriksen, T. B., Greene, R. A. and Kenny, L. C. (2017) 'Trial of labour after caesarean section and the risk of neonatal and infant death: a nationwide cohort study', <i>BMC Pregnancy and Childbirth</i> , 17(74), pp. 1-9. doi:10.1186/s12884-017-1255-2
<b>Type of publication</b>	Article (peer-reviewed)
<b>Link to publisher's version</b>	<a href="http://dx.doi.org/10.1186/s12884-017-1255-2">http://dx.doi.org/10.1186/s12884-017-1255-2</a> Access to the full text of the published version may require a subscription.
<b>Rights</b>	© 2017, the Authors. Open Access. This article is distributed under the terms of the Creative Commons Attribution 4.0 International License ( <a href="http://creativecommons.org/licenses/by/4.0/">http://creativecommons.org/licenses/by/4.0/</a> ), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The Creative Commons Public Domain Dedication waiver ( <a href="http://creativecommons.org/publicdomain/zero/1.0/">http://creativecommons.org/publicdomain/zero/1.0/</a> ) applies to the data made available in this article, unless otherwise stated. <a href="http://creativecommons.org/licenses/by/4.0/">http://creativecommons.org/licenses/by/4.0/</a>
<b>Item downloaded from</b>	<a href="http://hdl.handle.net/10468/3827">http://hdl.handle.net/10468/3827</a>

Downloaded on 2022-06-28T21:20:39Z



**UCC**

University College Cork, Ireland  
Coláiste na hOllscoile Corcaigh

**Additional File 3 Neonatal and infant death according to mode of delivery – sensitivity analyses with adjustment for smoking, co-morbidities and body mass index**

Mode of delivery	Neonatal death ( $\leq 28$ days) n=95 entire cohort		
	AOR (95% CI)		
	* Model 1 (60 deaths)	† Model 2 (49 deaths)	‡ Model 3 (23 deaths)
CS – ERCS	<i>Ref</i>	<i>Ref</i>	<i>Ref</i>
CS – TOLAC	1.34 (0.74, 2.43)	1.26 (0.64, 2.48)	1.07 (0.41, 2.76)
Mode of delivery	Early neonatal death ( $\leq 7$ days) n=86 entire cohort		
	* Model 1 (56 deaths)	† Model 2 (46 deaths)	‡ Model 3 (23 deaths)
	CS – ERCS	<i>Ref</i>	<i>Ref</i>
CS – TOLAC	1.33 (0.72, 2.47)	1.30 (0.65, 2.63)	1.07 (0.41, 2.76)
Mode of delivery	Late neonatal death ( $> 7$ days, $\leq 28$ days) n=9 entire cohort		
	* Model 1	† Model 2	‡ Model 3
	CS – ERCS	<i>Ref</i>	<i>Ref</i>
CS – TOLAC	<i>Number of deaths too few to run logistic models</i>		
Mode of delivery	Infant death ( $\leq 365$ days) n=171 entire cohort		
	* Model 1 (103 deaths)	† Model 2 (78 deaths)	‡ Model 3 (31 deaths)
	CS – ERCS	<i>Ref</i>	<i>Ref</i>
CS – TOLAC	1.02 (0.65, 1.60)	0.91 (0.54, 1.53)	0.68 (0.31, 1.50)

**Table:** Data are adjusted odds ratios with 95% confidence intervals. **AOR**= Adjusted odds ratio; **CI**=confidence interval, **ERCS**: Elective repeat caesarean section; **TOLAC**=Trial of labour after caesarean section; \* **Model 1**: adjusted for key covariates in the second birth including maternal age, maternal country of origin, educational attainment, mother and father's gross income, marital status, infant birthplace and infant birth weight, history of pregnancy loss and birth year + smoking status (data available from 1991-2010 only, cohort n=50,880)  
† **Model 2**: adjusted for model 1 + co-morbidities in the second delivery including hypertension, eclampsia and preeclampsia, fetal distress and gestational diabetes (data available from 1994-2010 only, cohort n=45,542)  
‡ **Model 3**: adjusted for model 2 + body mass index (BMI) (data available from 2004-2010 only, cohort n=22,672).