

**UCC Library and UCC researchers have made this item openly available.  
Please [let us know](#) how this has helped you. Thanks!**

<b>Title</b>	Microbial succession and flavor production in the fermented dairy beverage kefir
<b>Author(s)</b>	Walsh, Aaron M.; Crispie, Fiona; Kilcawley, Kieran N.; O'Sullivan, Orla; O'Sullivan, Maurice G.; Claesson, Marcus J.; Cotter, Paul D.
<b>Publication date</b>	2016-10-04
<b>Original citation</b>	Walsh, A. M., Crispie, F., Kilcawley, K., O'Sullivan, O., O'Sullivan, M. G., Claesson, M. J. and Cotter, P. D. (2016) 'Microbial succession and flavor production in the fermented dairy beverage kefir', <i>mSystems</i> 1(5), e00052-16 (16pp). doi:10.1128/mSystems.00052-16
<b>Type of publication</b>	Article (peer-reviewed)
<b>Link to publisher's version</b>	<a href="http://msystems.asm.org/content/2/1/e00003-17">http://msystems.asm.org/content/2/1/e00003-17</a> <a href="http://dx.doi.org/10.1128/mSystems.00052-16">http://dx.doi.org/10.1128/mSystems.00052-16</a> Access to the full text of the published version may require a subscription.
<b>Rights</b>	© 2016, the Authors. This is an open access article distributed under the terms of the Creative Commons Attribution 4.0 International license. <a href="https://creativecommons.org/licenses/by/4.0/">https://creativecommons.org/licenses/by/4.0/</a>
<b>Item downloaded from</b>	<a href="http://hdl.handle.net/10468/3850">http://hdl.handle.net/10468/3850</a>

Downloaded on 2021-11-29T18:35:20Z

**Table S1. Absolute abundances of bacteria and fungi in kefir samples after 0, 8 and 24 hours of fermentation, as determined by quantitative PCR (qPCR) measurements.**

<b>Sample</b>	<b>Total fungi (ng of fungal DNA)</b>	<b>Total bacteria (copies of 16S rRNA gene)</b>
Milk 0 h	0.0016	1.78E+05
Fr1 08 h	0.1386	7.64E+07
Fr1 24 h	0.2179	2.49E+08
Ick 08 h	0.0542	2.62E+08
Ick 24 h	0.0972	1.63E+08
UK3 08 h	0.0896	7.86E+07
Uk3 24 h	0.3933	4.22E+08