

**UCC Library and UCC researchers have made this item openly available.
Please [let us know](#) how this has helped you. Thanks!**

Title	Complete genome sequence of the gamma-aminobutyric acid-producing strain <i>Streptococcus thermophilus</i> APC151
Author(s)	Linares, Daniel M.; Arboleya, Silvia; Ross, R. Paul; Stanton, Catherine
Publication date	2017-04-27
Original citation	Linares, D. M., Arboleya, S., Ross, R.P. and Stanton, C. (2017) 'Complete genome sequence of the gamma-aminobutyric acid-producing strain <i>Streptococcus thermophilus</i> APC151', <i>Genome Announcements</i> , 5(17), e00205-17 (2pp). doi: 10.1128/genomeA.00205-17
Type of publication	Article (peer-reviewed)
Link to publisher's version	http://dx.doi.org/10.1128/genomeA.00205-17 Access to the full text of the published version may require a subscription.
Rights	© 2017, Linares et al. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International license . https://creativecommons.org/licenses/by/4.0/
Item downloaded from	http://hdl.handle.net/10468/3941

Downloaded on 2022-12-07T03:00:20Z



UCC

University College Cork, Ireland
Coláiste na hOllscoile Corcaigh



Complete Genome Sequence of the Gamma-Aminobutyric Acid-Producing Strain *Streptococcus thermophilus* APC151

Daniel M. Linares,^{a,b} Silvia Arboleya,^{a,b} R. Paul Ross,^{a,b,c} Catherine Stanton^{a,b}

Food Biosciences Department, Teagasc Food Research Centre, Moorepark, Fermoy, County Cork, Ireland^a; APC, Microbiome Institute, University College Cork, Cork, Ireland^b; School of Microbiology, University College Cork, Cork, Ireland^c

ABSTRACT Here is presented the whole-genome sequence of *Streptococcus thermophilus* APC151, isolated from a marine fish. This bacterium produces gamma-aminobutyric acid (GABA) in high yields and is biotechnologically suitable to produce naturally GABA-enriched biofunctional yogurt. Its complete genome comprises 2,097 genes and 1,839,134 nucleotides, with an average G+C content of 39.1%.

Streptococcus thermophilus is a nonpathogenic homofermentative facultative anaerobic lactic acid bacterium with a long history of use in the artisanal and modern industrial manufacture of fermented dairy products, especially yogurt (1). In addition, this commensal bacterium is one of the pioneer colonizers of oral and small intestine mucosal surfaces in newborns (2) and remains predominant in the oropharyngeal (2, 3) and gastrointestinal (4) tracts throughout the human life span. Based on the potential beneficial effects on human health, it is sometimes marketed as a probiotic (5–9).

We here report the genome of *Streptococcus thermophilus* APC151, a strain isolated from the digestive tract of a marine fish that produces large amounts of gamma-aminobutyric acid (GABA) (10). GABA has been classified as a health-promoting bioactive component in foods and pharmaceuticals (11) due to its function as an antihypertensive and antidiabetic neurotransmitter (11–15).

Genomic DNA was purified using the GenElute bacterial genomic DNA kit (Sigma-Aldrich). The nucleotide sequence of *S. thermophilus* was resolved utilizing PacBio single-molecule real-time (SMRT) technology on an RS system (GATC Biotech, Konstanz, Germany). Whole-genome sequencing yielded a total of 72,472 reads with a mean length of 12,628 bp. Subsequent *de novo* assembly utilizing the HGAP3 protocol yielded a single polished contig with 423-fold average reference coverage.

Annotation was performed by the RAST server (16). The total genome size was 1,839,134 bp, which included 2,097 predicted open reading frames (ORFs) (2,012 coding genes and 85 RNA genes) and a G+C content of 39.1%. The genome encompasses one predicted phage packaging cassette, one predicted prophage lysogenic conversion module, three clustered regularly interspaced short palindromic repeat (CRISPR) clusters, and 12 genes related to transposases. Additionally, the genes responsible for GABA biosynthesis, encoding the glutamate decarboxylase (*gadB*) and the glutamate/GABA antiporter (*gadC*), have been identified in the genome of this strain. Sequence analysis revealed that the *gadB-gadC* genes are flanked by transposases, a genetic organization similar to other GAD⁺ genomes available in GenBank.

The strain was found to be sensitive to all antibiotics proposed by European Food Safety Authority (EFSA) standards (10). In addition, no amino acid decarboxylases

Received 28 February 2017 Accepted 7 March 2017 Published 27 April 2017

Citation Linares DM, Arboleya S, Ross RP, Stanton C. 2017. Complete genome sequence of the gamma-aminobutyric acid-producing strain *Streptococcus thermophilus* APC151. *Genome Announc* 5:e00205-17. <https://doi.org/10.1128/genomeA.00205-17>.

Copyright © 2017 Linares et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Daniel M. Linares, Daniel.Linares@teagasc.ie.

associated with biogenic amines biosynthesis or virulence-related genes were identified in this genome.

The *S. thermophilus* APC151 genome showed a high degree of identity to that of strain *S. thermophilus* ND03 (GenBank accession no. NC_017563). The comparison of the two genomes revealed that strain APC151 contains an additional cluster of 13 genes (encoding a 5S RNA, a small subunit rRNA, a large subunit rRNA, and 10 tRNA genes) which are not present in the ND03 strain. In addition, strain APC151 acquired a number of mutations that disrupted or caused a frameshift in seven genes (encoding two transposases, two acyltransferases, the phosphate-transport-ATP-binding protein PstB, the β -carotene 15-15'-monooxygenase, and the substrate-specific component MtsA methionine-regulated ECF transporter).

These data are intended to increase the availability of the genomes of GABA-producing *S. thermophilus* strains in order to better understand their biodiversity and their potential technological, probiotic, and biofunctional properties.

Accession number(s). This whole-genome sequence was deposited at DDBJ/EMBL/GenBank under the accession number [CP019935](https://doi.org/10.1038/CP019935).

ACKNOWLEDGMENTS

This work is supported by the APC Microbiome Institute, which is a research center funded by Science Foundation Ireland (SFI). This publication has emanated from research supported by a research grant from Science Foundation Ireland (SFI) under grant number SFI/12/RC/2273.

REFERENCES

1. Wu Q, Tun HM, Leung FC, Shah NP. 2014. Genomic insights into high exopolysaccharide-producing dairy starter bacterium *Streptococcus thermophilus* ASCC 1275. *Sci Rep* 4:4974. <https://doi.org/10.1038/srep04974>.
2. Seow WK, Lam JH, Tsang AK, Holcombe T, Bird PS. 2009. Oral *Streptococcus* species in pre-term and full-term children: a longitudinal study. *Int J Paediatr Dent* 19:406–411. <https://doi.org/10.1111/j.1365-263X.2009.01003.x>.
3. Bik EM, Long CD, Armitage GC, Loomer P, Emerson J, Mongodin EF, Nelson KE, Gill SR, Fraser-Liggett CM, Relman DA. 2010. Bacterial diversity in the oral cavity of 10 healthy individuals. *ISME J* 4:962–974. <https://doi.org/10.1038/ismej.2010.30>.
4. Wang M, Ahrné S, Jeppsson B, Molin G. 2005. Comparison of bacterial diversity along the human intestinal tract by direct cloning and sequencing of 16S rRNA genes. *FEMS Microbiol Ecol* 54:219–231. <https://doi.org/10.1016/j.femsec.2005.03.012>.
5. Burton JP, Wescombe PA, Cadieux PA, Tagg JR. 2011. Beneficial microbes for the oral cavity: time to harness the oral streptococci? *Benef Microbes* 2:93–101. <https://doi.org/10.3920/BM2011.0002>.
6. Teughels W, Kinder Haake S, Sliopen I, Pauwels M, Van Eldere J, Cassiman JJ, Quirynen M. 2007. Bacteria interfere with *A. actinomycetemcomitans* colonization. *J Dent Res* 86:611–617. <https://doi.org/10.1177/154405910708600706>.
7. Wescombe PA, Heng NC, Burton JP, Chilcott CN, Tagg JR. 2009. Streptococcal bacteriocins and the case for *Streptococcus salivarius* as model oral probiotics. *Future Microbiol* 4:819–835. <https://doi.org/10.2217/fmb.09.61>.
8. Mignolet J, Fontaine L, Kleerebezem M, Hols P. 2016. Complete genome sequence of *Streptococcus salivarius* HSIS54, a human commensal bacterium highly prevalent in the digestive tract. *Genome Announc* 4(1): e01637-15. <https://doi.org/10.1128/genomeA.01637-15>.
9. Jandhyala SM, Talukdar R, Subramanyam C, Vuyyuru H, Sasikala M, Nageshwar Reddy D. 2015. Role of the normal gut microbiota. *World J Gastroenterol* 21:8787–8803. <https://doi.org/10.3748/wjg.v21.i29.8787>.
10. Linares DM, O'Callaghan TF, O'Connor PM, Ross RP, Stanton C. 2016. *Streptococcus thermophilus* APC151 strain is suitable for the manufacture of naturally GABA-enriched bioactive yogurt. *Front Microbiol* 7:1876. <https://doi.org/10.3389/fmicb.2016.01876>.
11. Li H, Cao Y. 2010. Lactic acid bacterial cell factories for gamma-aminobutyric acid. *Amino Acids* 39:1107–1116. <https://doi.org/10.1007/s00726-010-0582-7>.
12. Shimada M, Hasegawa T, Nishimura C, Kan H, Kanno T, Nakamura T, Matsubayashi T. 2009. Anti-hypertensive effect of gamma-aminobutyric acid (GABA)-rich *Chlorella* on high-normal blood pressure and borderline hypertension in placebo-controlled double blind study. *Clin Exp Hypertens* 31:342–354. <https://doi.org/10.1080/10641960902977908>.
13. Soltani N, Qiu H, Aleksic M, Glinka Y, Zhao F, Liu R, Li Y, Zhang N, Chakrabarti R, Ng T, Jin T, Zhang H, Lu WY, Feng ZP, Prud'homme GJ, Wang Q. 2011. GABA exerts protective and regenerative effects on islet beta cells and reverses diabetes. *Proc Natl Acad Sci U S A* 108: 11692–11697. <https://doi.org/10.1073/pnas.1102715108>.
14. Wan Y, Wang Q, Prud'homme GJ. 2015. GABAergic system in the endocrine pancreas: a new target for diabetes treatment. *Diabetes Metab Syndr Obes* 8:79–87. <https://doi.org/10.2147/DMSO.S50642>.
15. Pouliot-Mathieu K, Gardner-Fortier C, Lemieux S, St-Gelais D, Champagne CP, Vuilleumard JC. 2013. Effect of cheese containing gamma-aminobutyric acid-producing lactic acid bacteria on blood pressure in men. *PharmaNutrition* 1:141–148. <https://doi.org/10.1016/j.phanu.2013.06.003>.
16. Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: rapid annotations using subsystems technology. *BMC Genomics* 9:75. <https://doi.org/10.1186/1471-2164-9-75>.