

Title	Advance care planning within survivorship care plans for older cancer survivors: a systematic review
Author(s)	O'Caoimh, Rónán; Cornally, Nicola; O'Sullivan, Ronan; Hally, Ruth; Weathers, Elizabeth; Lavan, Amanda H.; Kearns, Tara; Coffey, Alice; McGlade, Ciara; Molloy, D. William
Publication date	2017-06-23
Original citation	O'Caoimh, R., Cornally, N., O'Sullivan, R., Hally, R., Weathers, E., Lavan, A. H., Kearns, T., Coffey, A., McGlade, C. and Molloy, D. W. 'Advance care planning within survivorship care plans for older cancer survivors: a systematic review', <i>Maturitas</i> , In Press. doi:10.1016/j.maturitas.2017.06.027
Type of publication	Article (peer-reviewed)
Link to publisher's version	http://dx.doi.org/10.1016/j.maturitas.2017.06.027 Access to the full text of the published version may require a subscription.
Rights	© 2017 Elsevier B.V. This manuscript version is made available under the CC-BY-NC-ND 4.0 license http://creativecommons.org/licenses/by-nc-nd/4.0/
Embargo information	Access to this article is restricted until 12 months after publication at the request of the publisher.
Embargo lift date	2018-06-23
Item downloaded from	http://hdl.handle.net/10468/4197

Downloaded on 2019-03-21T05:27:13Z



UCC

University College Cork, Ireland
Coláiste na hOllscoile Corcaigh

Accepted Manuscript

Title: Advance care planning within survivorship care plans for older cancer survivors: a systematic review

Authors: Rónán O’Caoimh, Nicola Cornally, Ronan O’Sullivan, Ruth Hally, Elizabeth Weathers, Amanda H. Lavan, Tara Kearns, Alice Coffey, Ciara McGlade, D. William Molloy



PII: S0378-5122(17)30624-2
DOI: <http://dx.doi.org/doi:10.1016/j.maturitas.2017.06.027>
Reference: MAT 6847

To appear in: *Maturitas*

Received date: 20-5-2017
Revised date: 17-6-2017
Accepted date: 22-6-2017

Please cite this article as: O’Caoimh Rónán, Cornally Nicola, O’Sullivan Ronan, Hally Ruth, Weathers Elizabeth, Lavan Amanda H, Kearns Tara, Coffey Alice, McGlade Ciara, Molloy D.William. Advance care planning within survivorship care plans for older cancer survivors: a systematic review. *Maturitas* <http://dx.doi.org/10.1016/j.maturitas.2017.06.027>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Advance care planning within survivorship care plans for older cancer survivors: a systematic review.

Running head: Advance care planning in older cancer survivors.

Rónán O’Caoimh^{*a, b, c}, Nicola Cornally^{b, d}, Ronan O’Sullivan^b, Ruth Hally^b, Elizabeth Weathers^{a, d}, Amanda H Lavan^b, Tara Kearns^b, Alice Coffey^d, Ciara McGlade^b, D. William Molloy^{b, c}.

^aHealth Research Board, Clinical Research Facility Galway, National University of Ireland, Galway, Geata an Eolais, University Road, Galway, Ireland.

^bCentre for Gerontology and Rehabilitation, University College Cork, St Finbarr’s Hospital, Cork, Ireland.

^cCOLLAGE (COLLaboration on AGEing), University College Cork, Cork City and Louth Age Friendly County Initiative, Co Louth, Ireland.

^dSchool of Nursing and Midwifery, University College Cork, College road, Cork City, Ireland.


Corresponding Author:

Dr. Rónán O’Caoimh, Health Research Board, Clinical Research Facility Galway, National University of Ireland, Galway, Geata an Eolais, University Road, Galway, Ireland.

Tel: +35391493187

Email: rocaoimh@hotmail.com

Graphical abstract: Advance care planning within survivorship care plans for older cancer survivors: a systematic review.



Highlights

- We reviewed the evidence for advance care planning, including advance care directives in survivorship care plans, among older cancer survivors.
- No papers were found that met inclusion criteria, with only one survivorship care plan including mention of an advance care directive.
- Despite increasing numbers of older and frail cancer survivors, there is little evidence for the use of advance care planning after cancer.
- Future studies should establish their acceptability and utility among older cancer survivors.

Abstract

Advances in the medical treatment of cancer have increased the number of survivors, particularly among older adults, who now represent the majority of these. Survivorship care plans (SCPs) are documents that cancer patients receive summarising their care, usually at the end of treatment but preferably from initial diagnosis. These may increase patient satisfaction and represent an opportunity to initiate preventative strategies and address future care needs. Advance care planning (ACP), incorporating advance healthcare decision-making, including formal written directives, increases satisfaction and end-of-life care. This paper systematically reviews evaluations of ACP within SCPs among older (≥ 65 years) cancer survivors. No studies meeting the inclusion criteria were identified by search strategies conducted in PubMed/MEDLINE and the Cochrane databases. One paper examined cancer survivors' mainly positive views of ACP. Another discussed the use of a SCP supported by a 'distress inventory' that included an advance care directive (living will) as an issue, though no formal evaluation was reported. Although ACP is important for older adults, no study was found that evaluated its role within survivorship care planning. Despite the risk of recurrence and the potential for morbidity and mortality, especially among older cancer survivors, ACP is not yet a feature of SCPs.

Keywords: advance care planning; survivorship care plan; cancer survivors; older adults; advance care directives; systematic review.

Introduction

With improved treatment more patients are surviving cancer [1]. This includes older patients, who now represent the majority of cancer survivors [1,2]. Robust post-cancer treatment should involve multi-modal, tailored supports that address individual's needs and predicted risks [3], especially in the case of older patients who have different requirements to younger cancer survivors [4]. Providing cancer survivors with information in the form of a survivorship care plan (SCP) is important to improve and enhance delivery of this care.

While useful for patients of all ages, survivorship care planning may be particularly important for older adults who because of higher levels of frailty [5] and multi-morbidity [6] have increasingly complex care needs.

SCPs, either written or delivered in an computerised (online) format (e.g. the LIVESTRONG™ SCP developed in 2007), function as a communication tool to enhance person-centred quality of care, collaborative decision making and post-treatment surveillance, and were recommended for all cancer survivors in the United States Institute of Medicine report '*From cancer patient to cancer survivor: Lost in the Transition*' [7] and more recently by the American College of Surgeons Commission on Cancer [4]. Reflecting this, centres in North America have the most experience with survivorship care planning though they are now in use in many countries [8]. SCPs improve engagement with patients' general practitioners [9, 10], facilitate discussion between patients and their families [10, 11] and are associated with higher levels of satisfaction [12], knowledge [13] and self-reported understanding by survivors [11,12].

Elements perceived as being important by patients to incorporate into a SCP include a watch list of symptoms and signs, a treatment summary and preventative healthcare recommendations [14,15], though only half of SCPs provide all these elements and often without sufficient detail [8]. Despite recommendations from several bodies, their proposed potential benefits and a near universal desire among survivors for information in the form of a SCP [15,16], few patients report having received one [15]. Use varies from as high as 35% of American women with breast cancer in one centre surveyed between 2004 and 2011 [17], to less than 20% of a sample of patients with different cancers in a study in Australia [15] to only 11% in an American sample of brain tumour survivors [18]. This may be because the evidence behind their effectiveness remains unclear [12]. In particular, while suggested, there is no definitive evidence that they increase uptake of recommended surveillance screening [19] or that they improve outcomes for cancer survivors [12]. Some studies indicate that they may even increase anxiety, perceived symptom burden and unnecessary contact with healthcare professionals [20].

Despite the need for positivity in the face of a cancer diagnosis, it is also important for healthcare professionals and patients to be cognisant that cancer survivorship, particularly for older patients, is associated with reduced life expectancy with 40% of patients living less than five years after diagnosis [21]. Further, cancer survivorship is not defined by success of treatment and SCPs are also recommended for those with a new diagnosis or those with advanced stage disease living with cancer [22]. Many of those who progress or experience recurrence will receive palliation or end-of-life care and will be required to make decisions surrounding treatment, though they may often lack the ability to do this by that stage [23]. Advance care planning (ACP), which involves communication between individuals, families and professionals, to support, discuss and plan for future healthcare decisions in the event that an individual loses capacity [24], can help reduce the burden on patients and proxy decision

makers. ACP may result in the completion of an advance care directive (ACD) reflecting the persons' preferences for their healthcare including end-of-life care in writing, if and when that time comes. Based on the limited evidence, ACP improves patient and relative satisfaction, the quality of death and dying, while respecting autonomy [24], and is generally well received by both patients [25] and healthcare professionals [26]. There is also evidence that dedicated ACP programmes such as 'Let Me Decide' are cost effective [27] and may reduce unnecessary healthcare use [28]. Yet, few patients receive ACP or complete an ACD [29].

Despite the increasing use of SCPs, there is very little evidence for the use of survivorship care planning in older patients with cancer [4]. Given this and the growing recognition of the importance of ACP in those with potentially life limiting conditions, the objective of this systematic review is to summarise the evidence for the inclusion of advance care planning decisions and ACDs within SCPs among older cancer survivors.

Methods

Search Strategy and Selection Criteria

A systematic review of the literature was conducted using PubMed/MEDLINE and the Cochrane Central Register of Controlled Trials. The search was conducted before March 2017. Databases were searched using the following search terms, which are detailed further in Table 1:

Advance directive* OR Advance care directive* OR Healthcare directive* OR Health care directive* OR Advance care plan* OR Advance care decision* OR Patient care plan* OR Care plan* OR Nursing care plan* AND Cancer* OR Oncology OR Neoplasm* AND Survivor* OR "Survivorship care plan*" AND Elderly OR Aged OR Older adult* OR Older person* OR Geriatric*.

The Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines were used [30].

Papers were included if they met all the following criteria:

- (i) An original study excluding case studies, letters to the editor or conference proceedings. Review papers or those describing the results of surveys of opinion were searched for relevant papers,
- (ii) Describing the evaluation of ACP within a multi-faceted SCP or long-term follow-up care plan that specifically includes reference to the process of ACP (i.e. active communication between patients, carers or family members and healthcare professionals that recognise, prepare for, discuss and document (i.e. written ACD) future healthcare decisions. Those presenting a survey of experience or satisfaction with SCPs without reference to ACP were excluded.
- (iii) Sampling cancer survivors (across the spectrum of a diagnosis of cancer i.e. from initial diagnosis to those who were undergoing or had completed treatment, which may have included surgery, chemotherapy or radiotherapy, and from patients living with advanced cancer to those in complete remission with ‘no evidence of disease’ [22], excluding secondary cancer survivors i.e. carers, family and friends), aged ≥ 65 years (i.e. older adults).
- (iv) Reported in English.

The selection of relevant papers is presented in a PRISMA flow diagram in Figure 1. The titles and abstracts of the selected articles were screened for inclusion criteria and the full papers of those meeting these were then reviewed. In total, 369 abstracts were reviewed. No

duplicates were found. Of these, 309 citations were excluded for different reasons, the most common being that the paper did not relate to SCPs (n=206) or address the issue of ACP if it discussed or evaluated survivorship care planning. Of the remaining citations 60 papers were reviewed.

Results

No papers met the inclusion criteria for this review i.e. no paper described the evaluation of ACP as part of the wider process of survivorship care planning with older cancer survivors. Two papers, not meeting the criteria, were relevant to the topic. One paper presented the views of ACP among a small sample (n=18) of survivors with a haematological malignancy (leukaemia or lymphoma), surveyed at least six months and up to one year after receiving a haematopoietic cell transplant, which included some older patients (≥ 65), though the median age was only 47 (range 33-67) years [31]. Of these, all but one discussed some element of ACP and half made a 'living will'. The ACP intervention was not documented as part of a formal SCP. The researchers found that cancer survivors engaging with ACP felt less anxiety. Although half who had discussed mortality with their family found it upsetting, all but one would recommend ACP to those undergoing similar treatment [31].

Another paper describing an American cancer centre's revised action plan with an emphasis on psychosocial concerns, for a small sample of 26 patients, documented the presence of an ACD, included as a 'living will' as an issue coupled with 'power of attorney' in a 'distress inventory' under the heading of 'financial/practical needs' [32]. This inventory was used to support the development and implementation of a SCP. While the SCP was considered by the authors to improve psychosocial care for patients no details were provided on outcomes nor was reference made to the process or utility of recording whether the patient had an ACD as part of survivorship care planning. It was not possible to conduct a

methodological review to assess the quality of papers, given the lack of data and suitable papers available.

Discussion

This paper systematically evaluates the evidence for a role for advance decision-making and the implementation of advance care plans or directives within SCPs for older cancer survivors. The results show that there is a paucity of data available on the utility or effectiveness of including ACP and more specifically advance care directives within a SCP. While survivorship care planning has been recommended by several organisations for more than ten years, uptake has been variable and research into the process limited. Further, none of these advocate for the inclusion of ACP discussions or the formal implementation of an ACD [4,7,8,33]. Most studies published on the topic of survivorship care planning focus on surveys of opinion and practice, possibly reflecting the early stages of experience and evidence with using SCPs. The most frequent preventative actions prompted by survivorship care planning were lifestyle changes (diet and exercise) [34] rather than long-term or anticipatory planning for future healthcare requirements.

This review found only one reference to ACP within a SCP [32], although this did not evaluate the SCP. A second paper describing ACP decisions in haematological malignancy survivors did not include reference to survivorship care planning or the use of a SCP [31] and captured the views of a young cohort whose experiences may not entirely reflect those of older patients. However, the study did suggest that ACP is acceptable to most cancer survivors, across a range of age groups from those in their early thirties to late sixties [31].

The field of survivorship care does not as yet appear to place importance on the use of advance directives with cancer survivors. Reflecting this, the validated Cancer Survivors' Unmet Needs (CaSUN) measure, the most widely used measure of unmet need in cancer

survivors, does not directly include the requirement to make anticipatory decisions or provisions for future healthcare needs [35], though the item “*I need help to try to make decisions about my life in the context of uncertainty*” [36] might suggest that this is something that patients do consider, albeit indirectly. Likewise when surveyed, most requested the inclusion of strategies to deal with the fear of cancer recurrence [37] suggesting that this is something that they are keenly aware of. Advance directives and SCPs are both forms of anticipatory care planning, which can reduce unplanned hospitalisations for older, frail patients [38] and promote autonomy and shared decision-making in those with advanced cancer [39].

The emerging subspecialty of geriatric oncology, already spearheading the use of comprehensive geriatric assessment (CGA) in older cancer patients [40] is likely to act as a catalyst for introducing the discussion and encouraging the uptake of ACDs as part of SCP. The Cancer and Ageing Research Group in collaboration with the National Cancer Institute and the National Institute on Ageing have recently advocated this approach, recommending the development of standardised SCPs for older cancer survivors based on CGA [41]. Although the evidence is yet to be established, studies examining the effectiveness of geriatric assessment are on-going and will likely help establish the role of CGA in promoting positive outcomes among these patients [41]. However, these do not as yet include ACP [41], despite the shorter life expectancy and high levels of co-morbidity [42,43,44] and reoccurrence in older patients with some cancers [45]. Indeed, few if any guidelines specifically address these issues in older cancer survivors [44]. This is despite the fact that evidence suggests that those with comorbidity had a greater desire for additional information [46].

Several challenges exist to the use of SCPs that have implications for the introduction of advance directives. The most important of these are the resource constraints related to the

manpower, time and costs required to deliver both survivorship care planning [47,48,49] and ACP [26]. A lack of training, experience and confidence among healthcare professional are also barriers to both [26,50,51]. Evidence suggests that like ACP discussions [26], SCPs may not be acceptable to all patients, particularly given that a small minority prefer not to receive any information at end of treatment [15]. The optimal timing of survivorship care planning is uncertain and should be carefully considered, particularly as many older cancer survivors experience anxiety regarding the symbolic nature of a cancer diagnosis in terms of their own mortality and likelihood of recurrence and may not wish to discuss it openly [52]. Three-six months after treatment is suggested as an appropriate timeline [48,53,54], though some advocate that it should be initiated early at diagnosis, well before end of treatment and modified throughout the course of the illness [8,55]. Further, it is unclear if cancer survivors wish to incorporate ACP within SCPs. The construct of survivorship is less clear to many older cancer survivors with some regarding it as opportunity, while others see it is a reminder of vulnerability [56] and a source of anxiety [52]. It may be that survivorship itself is better conceptualised among younger patients whose expectations of treatment and surviving cancer are different to previous generations [56]. Older cancer survivors place emphasis on issues relating to quality of life after cancer [57] with a focus on practical issues, which are included in some but not all SCPs, such as stress reduction, managing concerns regarding recurrence and the impact of cancer treatment on their partner and personal relations [58].

Where survivorship care planning is best delivered is not established with little evidence supporting the superiority of primary or secondary (clinic) care [59], though patients express more confidence in specialists suggesting that advance planning discussions may be best initiated by cancer specialists [60,61]. Community providers require additional resources to implement ACP and are unclear on their role [49]. Thus, advanced nurse practitioners may be best placed to facilitate this and encourage discussion about and uptake

of both SCPs and ACP in these older cancer survivors, particularly when frailty and multi-morbidity are recognised [54,62]. Personal (face-to-face) contact with trained staff is acceptable and preferable to many patients [63] and has been shown to improve adherence with SCPs [16]. Thus, it could be a useful format for delivering ACP where desired.

Improving healthcare professionals training and understanding of survivorship care planning and the role that ACP could have in this is also important [51]. Finally, the optimal vehicle for implementing SCPs with older adults is also unclear. Although Information technology (web-based applications) solutions already support the implementation of survivorship planning and are now replacing written SCPs [23], the use of these approaches has not been documented for older survivors. Nevertheless because of complex and fluctuating health states older survivors may benefit most from the regular updating afforded by these [23,64].

Overall, the construct of ACP is similar to survivorship care planning, focusing on discussions around patients future healthcare needs [24], and is consistent with the taxonomy of health related values identified by older multi-morbid cancer survivors, which includes the importance of a balance between quality and quantity of life, and their desire to actively participate in care decisions [57]. At a minimum the offer should be available to patients to promote choice, autonomy and informed decision-making. Thus, future research should now confirm the benefits of survivorship care planning, assess the priorities of older cancer survivors and attempt to appreciate their understanding of the need and desire for a SCPs as well as the optimal timing and setting of such discussions. This research should then investigate the effects of ACP on costs and important healthcare outcomes such as the uptake of post-treatment cancer surveillance, cancer recurrence rates and the use of healthcare resources, particularly among older cancer survivors with an advance directive in place.

Limitations of this review are that it only included articles written in English, which may have resulted in the omission of important studies. Further, only two sources were

reviewed for papers, potentially limiting the results. Despite these, the findings highlight the gaps that exist in the literature.

Conclusion

In summary, survivorship care planning incorporates many important elements that could potentially improve the care and wellbeing of older cancer survivors, who already represent the majority of cancer survivors [1,2]. While these typically include post-treatment summaries, diet, nutritional and exercise recommendations, and schedules for surveillance for recurrence, they do not yet address the important issue of ACP. This review could not find any evidence for the evaluation of a SCP that incorporated the construct on ACP or a formal written advance directive but did find some evidence that ACP is acceptable for cancer survivors [31]. At present high quality evidence for the use, benefits and risks of ACP within a SCP for older cancer survivors is lacking and further research is required, particularly given the expected increase in treatment options and numbers of frail, multi-morbid older patients receiving and benefiting from cancer therapies. However, even without supporting evidence, it would seem appropriate to consider instigating an ACP discussion with older cancer survivors to coincide with the creation of a SCP, a ‘geriatric-centred care plan’ [17]. In this sense, constructs around survivorship and ACP are not mutually exclusive and should not be treated as such for older patients. We suggest that SCPs should be tailored to the characteristics of individual older cancer survivors whose needs and openness to the concept may vary considerably [46] but are generally positive [31]. Irrespective, increasing awareness of the need for SCPs and ACP in cancer survivors, particularly older persons with advanced disease and comorbidity, is required to provide patients and their families with the option to implement these important anticipatory healthcare plans and promote independent living.

Contributors

RO'C contributed to data collection, analyzed and interpreted the data, and drafted the manuscript.

NC contributed to data collection.

RO'S contributed to data collection and review.

RH contributed to data collection and review.

EW contributed to data collection.

AHL contributed to data collection.

TK contributed to data collection.

AC contributed to data collection.

CMcG contributed to data collection.

DWM conceived and designed the review, and drafted the manuscript.

All authors saw and approved the final manuscript.

Funding

No funding was received for this review.

Ethical approval

No ethical approval or consent was required under Irish law to conduct this systematic review. No human or animal subjects were directly included in this study.

Provenance and peer review

This article has undergone peer review.

Conflict of interest

The authors declare that they have no conflict of interest.

Conflict of Interest Statement

The authors report no other conflict of interest.

References

1. Parry C, Kent EE, Mariotto AB, Alfano CM, Rowland JH. Cancer survivors: a booming population. *Cancer Epidemiology and Prevention Biomarkers* 2011;20(10):1996-2005.
2. Miller KD, Siegel RL, Lin CC, et al. Cancer treatment and survivorship statistics, 2016. *CA: a cancer journal for clinicians* 2016;66(4):271-89.
3. Jefford M, Kinnane N, Howell P, et al. Implementing novel models of posttreatment care for cancer survivors: enablers, challenges and recommendations. *Asia-Pacific J of Clin Oncol* 2015;11(4):319-27.
4. Guerard EJ, Nightingale G, Bellizzi K, et al. Survivorship care for older adults with cancer: U13 conference report. *J Geriatr Oncol* 2016;7(4):305-12.
5. Santos-Eggimann B, Cuénoud P, Spagnoli J, Junod J. Prevalence of frailty in middle-aged and older community-dwelling Europeans living in 10 countries. *The Journals of Gerontology Series A*: 2009;64(6):675-81.
6. Prince MJ, Wu F, Guo Y, et al. The burden of disease in older people and implications for health policy and practice. *Lancet* 2015; 385: 549–62.
7. Hewitt M, Greenfield S, Stovall E. From cancer patient to cancer survivor: lost in transition. Committee on cancer survivorship: improving care and quality of life, institute of medicine and national research council. Washington (DC): The National Academies Press; 2006.
8. Daudt HM, van Mossel C, Dennis DL, Leitz L, Watson HC, Tanliao JJ. Survivorship care plans: a work in progress. *Current oncology* 2014;21(3):466-79.

9. Blanch-Hartigan D, Forsythe LP, Alfano CM, et al. Provision and discussion of survivorship care plans among cancer survivors: results of a nationally representative survey of oncologists and primary care physicians. *J Clin Oncol* 2014;32:1578–1585.
10. Blinder VS, Norris VW, Peacock NW, et al. Patient perspectives on breast cancer treatment plan and summary documents in community oncology care. *Cancer* 2013;119(1):164–172.
11. Palmer SC, Stricker CT, Panzer SL, et al. Outcomes and satisfaction after delivery of a breast cancer survivorship care plan: results of a multicenter trial. *J Oncol Pract* 2015;11(2):e222-9.
12. Brennan ME, Gormally JF, Butow P, Boyle FM, Spillane AJ. Survivorship care plans in cancer: a systematic review of care plan outcomes. *Br J of Cancer* 2014;111(10):1899-908.
13. Bulloch KJ, Irwin ML, Chagpar AB, et al. Systematic approach to providing breast cancer survivors with survivorship care plans: a feasibility study. *J Oncol Pract* 2015;11(2):e170-6.
14. Mayer DK, Gerstel A, Leak AN, Smith SK. Patient and provider preferences for survivorship care plans. *J Oncol Pract* 2012;8(4):e80-6.
15. Kinnane NA, Piper AJ, Jefford M. How will cancer survivors use survivorship care plans? *Acta Oncol* 2017;56(2):183-189.
16. Rosenberg CA, Flanagan C, Brockstein B, et al. Promotion of self-management for post treatment cancer survivors: evaluation of a risk-adapted visit. *J Cancer Surviv* 2016;10(1):206-19.

17. Faul LA, Luta G, Sheppard V, et al. Associations among survivorship care plans, experiences of survivorship care, and functioning in older breast cancer survivors: CALGB/Alliance 369901. *J Cancer Surviv* 2014;8(4):627-37
18. Sloane K, Vachani C, Hampshire MK, Metz JM, Hill-Kayser CE. Late effects in survivors of central nervous system tumors: reports by patients and proxies. *J Cancer Surviv* 2016;10(2):234-40.
19. Oeffinger KC, Hudson MM, Mertens AC, et al. Increasing rates of breast cancer and cardiac surveillance among high-risk survivors of childhood Hodgkin lymphoma following a mailed, one-page survivorship care plan. *Pediatr Blood Cancer* 2010;56 (5):818–824.
20. Nicolaije KA, Ezendam NP, Vos MC, et al. Impact of an Automatically Generated Cancer Survivorship Care Plan on Patient-Reported Outcomes in Routine Clinical Practice: Longitudinal Outcomes of a Pragmatic, Cluster Randomized Trial. *J Clin Oncol* 2015;33(31):3550-9.
21. Rowland JH, Bellizzi KM. Cancer survivorship issues: life after treatment and implications for an aging population. *J Clin Oncol* 2014;32(24):2662–2668.
22. Twombly R. What's in a name: who is a cancer survivor? *J Natl Cancer Inst* 2004, 96 (19):1414-1415.
23. Silveira MJ, Kim SY, Langa KM. Advance directives and outcomes of surrogate decision making before death. *N Engl J Med* 2010; 362: 1211-1218.
24. Detering KM, Hancock AD, Reade MC, Silvester W. The impact of advance care planning on end of life care in elderly patients: randomised controlled trial. *BMJ* 2010; 340:c1345.

25. Weathers E, O’Caoimh R, Cornally N, et al. Advance care planning: a systematic review of randomized controlled trials conducted with older adults. *Maturitas* 2016, 91:101-9.
26. Cornally N, McGlade C, Weathers E et al. Evaluating the systematic implementation of the 'Let Me Decide' advance care planning programme in long term care through focus groups: A user's perspective. *BMC Palliative Care* 2015, 14:55.
27. O’Sullivan R, Murphy A, O’Caoimh R, et al. Economic Analysis of Systematically Implementing a Programme of Advance Care Planning in Three Irish Nursing Homes, *BMC Research Notes* 2016; 9:237.
28. Molloy DW, Guyatt GH, Russo R, et al. Systematic implementation of an advance directive program in nursing homes: a randomized controlled trial. *JAMA* 2000; 283(11): 1437-1444.
29. Kass-Bartelmes BL, Hughes R. Advance care planning: preferences for care at the end of life. *J Pain Palliat Care Pharmacother.* 2004;18(1):87-109.
30. Liberati A, Altman DG, Tetzlaff J, et al. The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate healthcare interventions: explanation and elaboration. *BMJ* 2009;339: b2700.
31. Loggers ET, Lee S, Chilson K, Back AL, Block S, Loberiza FR. Advance care planning among hematopoietic cell transplant patients and bereaved caregivers. *Bone Marrow Transplant* 2014;49(10):1317-22.
32. Arnett J, Henry B, Fankell A. Managing late effects of breast cancer treatment. *Clin J Oncol Nurs* 2014;18 S:32-5.


33. Runowicz CD, Leach CR, Henry NL, et al. American Cancer Society/American Society of Clinical Oncology Breast Cancer Survivorship Care Guideline. *CA Cancer J Clin*. 2016 Jan-Feb;66(1):43-73.
34. Hill-Kayser CE, Vachani CC, Hampshire MK, Di Lullo G, Jacobs LA, Metz JM. Impact of internet-based cancer survivorship care plans on health care and lifestyle behaviors. *Cancer* 2013;119(21):3854-60.
35. Hodgkinson K, Butow P, Hunt GE et al. The development and evaluation of a measure to assess cancer survivors' unmet supportive care needs: The CaSUN Survivors' Unmet Needs measure). *Psychooncology* 2007; 16: 796–804.
36. Brennan ME, Butow P, Spillane AJ, Boyle F. Patient-reported quality of life, unmet needs and care coordination outcomes: Moving toward targeted breast cancer survivorship care planning. *Asia Pac J Clin Oncol* 2016;12(2):e323-31.
37. Boyajian RN, Grose A, Grenon N, et al. Desired elements and timing of cancer survivorship care: one approach may not fit all. *J Oncol Pract* 2014;10(5):e293-8.
38. Baker A, Leak P, Ritchie LD, Lee AJ, Fielding S. Anticipatory care planning and integration: a primary care pilot study aimed at reducing unplanned hospitalisation. *Br J Gen Pract*. 2012;62(595):e113-20.
39. Boyd K, Mason B, Kendall M, et al. Advance care planning for cancer patients in primary care: a feasibility study. *Br J Gen Pract* 2010;60(581):e449-58.
40. Balducci L, Colloca G, Cesari M, Gambassi G. Assessment and treatment of elderly patients with cancer. *Surgical oncology* 2010; 19(3):117-23.

41. Mohile SG, Hurria A, Cohen HJ, et al. Improving the quality of survivorship for older adults with cancer. *Cancer* 2016;122(16):2459-568.
42. Deckx L, van den Akker M, Metsemakers J, Knottnerus A, Schellevis F, Buntinx F. Chronic diseases among older cancer survivors. *J of Cancer Epi* 2012;2012.
43. Kurnit KC, Ward KK, McHale MT, Saenz CC, Plaxe SC. Increased prevalence of comorbid conditions in women with uterine cancer. *Gynecol Oncol* 2015;138(3):731-4.
44. Meneses K, Benz R, Azuero A, Jablonski-Jaudon R, McNeese P. Multimorbidity and breast cancer. *Semin Oncol Nurs* 2015;31(2):163-9.
45. Beadle BM, Woodward WA, Buchholz TA. The impact of age on outcome in early-stage breast cancer. *Semin Oncol* 2011;21(1): 26-34.
46. O'Malley DM, Hudson SV, Ohman-Strickland PA, et al. Follow-up Care Education and Information: Identifying Cancer Survivors in Need of More Guidance. *J Cancer Educ* 2016;31(1):63-9.
47. Hewitt ME, Bamundo A, Day R, et al. Perspectives on post-treatment cancer care: Qualitative research with survivors, nurses, and physicians. *J Clin Oncol* 2007, 25: 2270-73.
48. Mayer DK, Gerstel A, Walton AL, et al. Implementing survivorship care plans for colon cancer survivors. *Oncol Nurs Forum* 2014;41(3):266-73.
49. Salz T, McCabe MS, Onstad EE, et al. Survivorship care plans: is there buy-in from community oncology providers? *Cancer* 2014;120(5):722-30.
50. Buriak SE, Potter J, Bleckley MK. Using a predictive model of clinician intention to improve continuing health professional education on cancer survivorship. *J Contin Educ Health Prof* 2015;35(1):57-64.

51. Faithfull S, Samuel C, Lemanska A, Warnock C, Greenfield D. Self-reported competence in long term care provision for adult cancer survivors: A cross sectional survey of nursing and allied health care professionals. *Int J Nurs Stud* 2016;53:85-94.
52. Deimling GT, Bowman KF, Sterns S, Wagner LJ, Kahana B. Cancer - related health worries and psychological distress among older adult, long-term cancer survivors. *Psycho - Oncology* 2006;15(4):306-20.
53. Grant M, Economou D, Ferrell BR. Oncology nurse participation in survivorship care. *Clin J Oncol Nurs* 2010; 14: 709-15.
54. Dulko D, Pace CM, Dittus KL, et al. Barriers and facilitators to implementing cancer survivorship care plans. *Oncol Nurs Forum* 2013;40(6):575-80.
55. Haq R, Heus L, Baker NA, et al. Designing a multifaceted survivorship care plan to meet the information and communication needs of breast cancer patients and their family physicians: results of a qualitative pilot study. *BMC Med Inform Decis Mak.* 2013;13:76.
56. Hannum SM, Clegg Smith K, Coa K, Klassen AC. Identity reconstruction among older cancer survivors: Age and meaning in the context of a life-altering illness. *J Psychosoc Oncol* 2016;34(6):477-92.
57. Naik AD, Martin LA, Moye J, Karel MJ. Health Values and Treatment Goals of Older, Multimorbid Adults Facing Life-Threatening Illness. *J Am Geriatr Soc* 2016;64(3):625-31.
58. Geller BM, Vacek PM, Flynn BS, Lord K, Cranmer D. What are cancer survivors' needs and how well are they being met? *J Fam Pract* 2014;63(10):E7-16.
59. Boekhout AH, Maunsell E, Pond GR, et al. A survivorship care plan for breast cancer survivors: extended results of a randomized clinical trial. *J Cancer Surviv* 2015;9(4):683-91.

60. Nyarko E, Metz JM, Nguyen GT, Hampshire MK, Jacobs LA, Mao JJ. Cancer survivors' perspectives on delivery of survivorship care by primary care physicians: an internet-based survey. *BMC family practice* 2015;16(1):143.
61. Chubak J, Aiello Bowles EJ, Tuzzio L, et al. Perspectives of cancer survivors on the role of different healthcare providers in an integrated delivery system. *J Cancer Surviv* 2014;8(2):229-38.
62. Morgan B, Tarbi E. The Role of the Advanced Practice Nurse in Geriatric Oncology Care. *Semin Oncol Nurs* 2016;32(1):33-43.
63. Wen KY, Hu A, Ma GX, Fang CY, Daly MB. Information and communication needs of Chinese American breast cancer patients: perspectives on survivorship care planning. *J Community Support Oncol* 2014;12(12):439-45.
64. Groen WG, Kuijpers W, Oldenburg HS, Wouters MW, Aaronson NK, van Harten WH. Empowerment of Cancer Survivors Through Information Technology: An Integrative Review. *J Med Internet Res* 2015;17(11):e270.

Figure 1. PRISMA flow diagram of the search strategy.



Search Details	PubMed Citations	Cochrane Citations	Total Citations
Search #1: Advance directive* OR Advance care directive* OR Healthcare directive* OR Health care directive* OR Advance care plan* OR Advance care decision* OR Patient care plan* OR Care plan* OR Nursing care plan*	52,288	6,020	58,308
Search #2: Cancer* OR Oncology OR Neoplasm*	3,567,298+	3,008	3,570,306+
Search #3: Survivor* OR “Survivorship care plan*”	88,260	702	88,962
Search #4: Elderly OR Aged OR Older adult* OR Older person* OR Geriatric*	4,672,266	8,734	4,681,702
#1 AND #2	5,974	2,106	8,080
#1 AND #3	592	542	1,134
#1 AND #4	13,536	5,840	19,376
#2 AND #3	30,079	277	30,356
#1 AND #2 AND #3	452	218	670
#1 AND #2 AND #3 AND #4	152	217	369
Number of abstracts reviewed excluded from combined searches	152	217	369
Number of abstracts excluded from combined searches	103	206	309
Number of articles reviewed from combined searches	49	11	60
Number of articles included	0	0	0