

**UCC Library and UCC researchers have made this item openly available.
 Please [let us know](#) how this has helped you. Thanks!**


Title	Reducing consumption of confectionery foods: A post-hoc segmentation analysis using a social cognition approach
Author(s)	Naughton, Paul; McCarthy, Mary; McCarthy, Sinéad N.
Publication date	2017-06-27
Original citation	Naughton, P., McCarthy, M. and McCarthy, S. (2017) 'Reducing consumption of confectionery foods: A post-hoc segmentation analysis using a social cognition approach', <i>Appetite</i> , 117, pp. 168-178. doi:10.1016/j.appet.2017.06.027
Type of publication	Article (peer-reviewed)
Link to publisher's version	http://www.sciencedirect.com/science/article/pii/S0195666316305736 http://dx.doi.org/10.1016/j.appet.2017.06.027 Access to the full text of the published version may require a subscription.
Rights	© 2017 Elsevier Ltd. This manuscript version is made available under the CC-BY-NC-ND 4.0 license http://creativecommons.org/licenses/by-nc-nd/4.0/
Embargo information	Access to this article is restricted until 24 months after publication by request of the publisher.
Embargo lift date	2019-07-27
Item downloaded from	http://hdl.handle.net/10468/4680

Downloaded on 2020-12-02T04:35:40Z

Accepted Manuscript

Reducing consumption of confectionery foods: A post-hoc segmentation analysis using a social cognition approach

Paul Naughton, Mary McCarthy, Sinéad McCarthy



PII: S0195-6663(16)30573-6

DOI: [10.1016/j.appet.2017.06.027](https://doi.org/10.1016/j.appet.2017.06.027)

Reference: APPET 3533

To appear in: *Appetite*

Received Date: 16 October 2016

Revised Date: 29 May 2017

Accepted Date: 25 June 2017

Please cite this article as: Naughton P., McCarthy M. & McCarthy Siné., Reducing consumption of confectionery foods: A post-hoc segmentation analysis using a social cognition approach, *Appetite* (2017), doi: 10.1016/j.appet.2017.06.027.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Title

Reducing Consumption of Confectionery Foods: A post-hoc segmentation analysis using a social cognition approach.

Authors

Dr Paul Naughton^{*1,2}

Dr Mary McCarthy² M.McCarthy@ucc.ie,

Dr Sinéad McCarthy¹ Sinead.McCarthy@teagasc.ie

¹Teagasc Food Research Centre, Ashtown, Dublin 15, Ireland.

²Department of Food Business & Development, University College Cork, Ireland.

* Corresponding author at: The Business School, Edinburgh Napier University, Edinburgh, EH141DJ, UK. Email: P.Naughton@napier.ac.uk

1 **Title:** Reducing Consumption of Confectionery Foods: A post-hoc segmentation analysis using a
2 social cognition approach

3 **Abstract**

4 Considering confectionery consumption behaviour this cross-sectional study used social cognition
5 variables to identify distinct segments in terms of their motivation and efforts to decrease their
6 consumption of such foods with the aim of informing targeted social marketing campaigns. Using
7 Latent Class analysis on a sample of 500 adults four segments were identified: unmotivated, triers,
8 successful actors, and thrivers. The unmotivated and triers segments reported low levels of
9 perceived need and perceived behavioural control (PBC) in addition to high levels of habit and
10 hedonic hunger with regards their consumption of confectionery foods. Being a younger adult was
11 associated with higher odds of being in the unmotivated and triers segments and being female was
12 associated with higher odds of being in the triers and successful actors segments. The findings
13 indicate that in the absence of strong commitment to eating low amounts of confectionery foods
14 (i.e. perceived need) people will continue to overconsume free sugars regardless of motivation to
15 change. It is therefore necessary to identify relevant messages or 'triggers' related to sugar
16 consumption that resonate with young adults in particular. For those motivated to change,
17 counteracting unhealthy eating habits and the effects of hedonic hunger may necessitate changes to
18 food environments in order to make the healthy choice more appealing and accessible.

19 *Keywords*

20 Healthy Eating, Sugar, Social Marketing, Social Cognition Models, Audience Segmentation

21 **1. Introduction**

22 Excessive consumption of non-milk extrinsic sugars¹ (often referred to as free or added sugars) is
23 associated with obesity and related illnesses, type 2 diabetes, and dental caries (Malik *et al.*, 2006;
24 Vartanian *et al.*, 2007; Lustig *et al.*, 2012; Te Morenga, *et al.*, 2013). Dietary guidelines across the
25 world recommend that adults and children should consume sparingly foods that contain high
26 amounts of these types of sugars (FSAI, 2011; PHE, 2014; The HHS and USDA, 2015). However,
27 studies show that on average adults are exceeding guidelines set by the WHO to limit intake of free
28 sugars to less than 10% of total energy per day (around 50grams) (WHO, 2015). Moreover, a
29 reduction in intake of free sugars to below 5% of total energy per day (around 25 grams) would have

¹ Sugars that are not incorporated in the cellular structure of foods like fruits and vegetables, whether natural or unprocessed, such as honey, or refined, such as table sugar, and, consequently, may have adverse effects on health (Department of health, 1991). Often referred to as free sugars (i.e. sugars added to foods).

30 additional health benefits (WHO, 2015). Achieving such a significant change in people's eating
31 behaviours is the challenge facing health experts and policy makers.

32 Social Marketing is concerned with voluntary behaviour change achieved through the adoption and
33 adaption of contemporary commercial marketing theory and practice (Eagle, et al., 2013). As in
34 commercial marketing theory, market segmentation and targeting is a key concept (Andreasen,
35 2002; Geier & Bryant, 2005, Dann, 2010). This entails dividing up a large heterogonous market into a
36 number of homogeneous segments and devising customised marketing programmes for one or
37 more target segments (Kotler et al., 2007). Social cognition theories of behaviour are considered a
38 core component of successful social marketing interventions (French & Blair-Stevens, 2006; Luca &
39 Suggs, 2013) as they provide useful insights for elements of the social marketing mix (i.e. product,
40 price, place and promotion) by identifying the key determinants behind overt behaviour (e.g.
41 attitudes or perceived behavioural control). Therefore, the social marketer can make an informed
42 decision on the focus of the intervention, e.g. changing attitudes or increasing behavioural control,
43 directed at behavioural change. In this paper, a latent class analysis approach to segmentation
44 analysis was conducted using social cognition variables to identify different types of people with
45 regards their consumption of confectionery foods.

46 *Segmentation & Behavioural Theory*

47 The identification of homogeneous segments is dependent on the segmentation bases and methods
48 used to divide the market (Wedel and Kamakura, 2000). According to Geier and Bryant (2005) bases
49 such as readiness to change and psychographics (e.g. lifestyle, values) have been commonly used to
50 identify distinct subgroups in social marketing campaigns. These bases are considered to be more
51 effective in identifying differentiated segments compared to demographic bases such as age and
52 ethnicity (Wedel and Kamakura, 2000; Vyncke, 2002; Weinstein, 2004), which are commonly used in
53 health behaviour research (Slater, 1996). In addition to segmentation bases, there are a variety of
54 methods that can be used to group individuals into segments and they can be broadly categorised
55 into a-priori and post-hoc methods. A-priori involves determining the type and number of segments
56 in advance whereas in a post-hoc segmentation approach the type and number of segments
57 emerges from data analysis. Kazbare et al. (2010), using the segmentation evaluation criteria
58 proposed by Kotler and Keller (2009), found that post-hoc segmentation of social cognition variables
59 was more helpful in designing healthy eating campaigns than a-priori segmentation of demographic
60 and behavioural variables as this approach provided more insight on who should be targeted and
61 what should be communicated. Segment evaluation criteria include: measurability, referring to the
62 extent to which segments can be feasibly identified and measured using segmentation variables;

63 substantiality, meaning segments must be large enough to warrant developing and maintaining a
64 special marketing mix; accessibility, considers the demographic profiling of segments in order that
65 they can be effectively reached/targeted; and differentiability, which means that segments should
66 be genuinely different on measured criteria and therefore should respond to different marketing
67 mix initiatives.

68 In social marketing interventions audience segmentation must be accompanied by a detailed study
69 of peoples' lives, behaviours, motives, and the environment in which they make choices. The
70 objective is to develop an attractive value proposition based on understanding the costs and
71 benefits associated with a new behaviour (Geier & Bryant, 2005; French & Blair Stevens, 2006).
72 Exchange theory is a fundamental principle of commercial marketing (e.g. consumer receives a
73 product or service for a cash outlay) but social marketing is more complicated as there is rarely an
74 immediate benefit for the adoption of a new behaviour and there are often immediate costs such as
75 time and emotional discomfort (Geier and Bryant, 2005). For this reason, behavioural theories,
76 including social cognition models, are considered a core component of successful social marketing
77 interventions (French & Blair-Stevens, 2006; Luca & Suggs, 2013). According to Gordon et al. (2006),
78 based on a systematic review of studies that evaluate social marketing effectiveness, social
79 marketing provides a very promising framework for improving health but issues related to research
80 design and a lack of conceptual understanding must be addressed. Luca and Suggs (2013) carried out
81 a systematic review on theory and model use in social marketing interventions between 1990 and
82 2009 and concluded that there was an ongoing lack of use of theory or an underreporting of theory
83 in social marketing campaigns.

84 *Social Cognition*

85 Social cognition models can offer value in endeavours to integrate theory into the application of
86 social marketing campaigns. These reductionistic models identify key variables that account for the
87 numerous influences on behaviour (Bagozzi, 1992) with the most frequently used theory in social
88 marketing campaigns being the transtheoretical model (TTM) (Luca and Suggs, 2013). This model is
89 built on the proposition that when addressing a problematic behaviour individuals go through
90 similar stages of change and different influencing variables are important at different stages
91 (Prochaska & DiClemente, 1983). However, a systematic review carried out by Bridle et al. (2005)
92 showed limited evidence to support the effectiveness of health behaviour change interventions
93 based on the TTM. Indeed stage models have been criticised as being too vague in explaining what
94 actually happens in each stage (Povey et al., 1999; Armitage & Conner, 2000). Other social cognition
95 models, such as the theory of planned behaviour (TPB), have been more specific in identifying the

96 variables that underlie behavioural motivation (Armitage & Conner, 2000). In these models intention
97 to perform a specific behaviour is conceptualised as the most important and most immediate
98 predictor of behaviour (Ajzen, 1991). According to Sheeran et al. (2005) intentions conclude the
99 decision making process by signalling one's commitment towards the performance of a behaviour.
100 However, meta-analysis studies show that behavioural intentions do not correspond strongly with
101 actual behaviour (Armitage & Conner, 2001; Conner and Sparks, 2005). Moreover, evidence suggests
102 that the intention-behaviour discrepancies are largely due to people having good intentions but
103 failing to act on them (Sheeran, 2002).

104 In the health behaviour literature a number of variables have been identified to explain transitions
105 from intention to action. Research indicates that having a dietary related lifestyle goal (e.g. weight
106 loss) is associated with successfully implementing and maintaining healthy dietary change as
107 individuals are more engaged in the change process and, therefore, more likely to overcome
108 potential barriers to success (Berg-Smith, 1999; Schnoll and Zimmerman, 2001; Nothwehr & Yang,
109 2006). The concept implementation intention emphasises the significance of planning in translating
110 intentions into behaviour. According to Gollwitzer (1993 pg. 152) "*The purpose of an implementation
111 intention is to lay down a specific plan that helps to promote the initiation and efficient execution of
112 goal-directed activity*". A number of studies have found that healthy dietary change is
113 significantly related to planning over and above the effects of intentions (Scholz *et al.*, 2009;
114 Osch *et al.*, 2010). In addition, perceived behavioural control (PBC) is a central concept in
115 explaining not only the actions a person is motivated to perform but also, once an activity is
116 initiated, the likelihood of maintaining effort in the face of obstacles. It reflects an individual's
117 assessment of external issues such as access to resources and internal issues such as emotions
118 that act as barriers to healthy behaviour (Ajzen, 1991). Numerous studies have demonstrated a
119 significant effect of PBC on dietary change and the concept has been incorporated into dietary
120 change interventions that have yielded favourable outcomes (Steptoe *et al.*, 2004; Linde *et al.*,
121 2006; Ahluwalia *et al.*, 2007). A less empirical examined concept but potentially important
122 determinant of dietary change is perceived need i.e. whether or not people feel the need to carry
123 out the health behaviour in question (Povey *et al.*, 2000; Payne *et al.* 2004). Evidence indicates
124 that if people perceive a problem to be associated with their diet (e.g. a feeling that one is
125 overweight) then they are more likely to make relevant dietary changes (e.g. reduce fat-intake
126 in their diet) (Glanz *et al.*, 1998; Payne *et al.* 2004). Paisley and Sparks (1998) argue that
127 people's perceptions of need may not be reflected in their attitudes and therefore should be
128 considered separately. For example, a behaviour may be seen as beneficial and wise (i.e. a

129 positive attitude) but there may be a low perceived need to perform the behaviour because the
130 outcome is not valued and/or the outcome is believed to be attainable through other means.
131 Finally, it is recognised that habits and emotions are powerful determinants of regularly performed
132 behaviours such as food consumption and often act as barriers to change (Verplanken & Aarts,
133 1999; Macht, 2008; De Bruijn, 2010). As people strive to create healthy eating habits it is likely that
134 they will have to break unhealthy eating habits as human beings are instinctively driven to foods
135 high in fat and sugar (Rozin, 2007). In addition to habits, hedonic hunger, a term used to describe a
136 person's motivation to consume tempting food even if he/she is not hungry. Thus, eating habits and
137 hedonic hunger can result in instinctive unhealthy behavioural choices that are not consistent with a
138 person's dietary intentions (Lowe and Butryn, 2007).

139 To examine dietary change this study used the following social cognitive variables: lifestyle goal,
140 dietary planning, perceived behavioural control (PBC), perceived need, confectionery habit and
141 hedonic hunger as segmentation bases. Latent class analysis was applied as a post-hoc segmentation
142 method to identify different cohorts of people with regards to confectionery consumption
143 reduction. In taking this approach this study addresses some of the concerns expressed about social
144 marketing health interventions, specifically that interventions lack theoretical foundations and are
145 designed with little appreciation of what empirical research indicates will work best and why.

146 **2. Method**

147 *Sample*

148 Data collection was carried out by a market research agency in August 2011 using a stratified
149 random sampling procedure. 500 Irish adult's representative of the population in terms of gender,
150 age, living location (i.e. rural v urban) and social class² based on the most recent Irish census data
151 were recruited for the study. The survey instrument was interviewer administered to ensure
152 accuracy and a high completion rate. Ethical approval was sought from and granted by the Social
153 Research Ethics Committee at University College Cork, Ireland. Table 1 provides an overview of the
154 sample characteristics. The distribution of self-reported body mass index (BMI) data compares
155 well with the self-reported BMI data from the SLAN 2007 study, which was a national survey of
156 lifestyle, attitudes and nutrition of the Irish population using a probability sample (Harrington *et*
157 *al.*, 2008). In the present study there were slightly more individuals classified as normal weight
158 (53% compared to 48%) and slightly less classified as overweight (34% compared to 36%) and

² The classification of social class used by the Central Statistics Office in Ireland (O'Hare *et al.*, 1991)

159 obese (12% compared to 14%). The mean BMI for the sample was 25.31 (SD = 4.21), which is
 160 similar to that of SLAN 2007 (\bar{x} = 25.52, SD = 4.03 self-reported).

161

162 **Table 1** Sample profile

	%		%		%
Gender		Social Class		Location	
Male	50	Professional (A)	7.1	Urban	55
Female	50	Managerial & technical (B)	27.5	Rural	45
Age		Non-manual (C1)	21.2	Self Reported BMI	
18-24	13.5	Skilled-manual (C2)	19.3	Underweight (<18.5)	1
25-44	43.5	Semi-skilled (D)	12	Normal Weight (18.5 – 24.9)	53
45 - 64	33.7	Unskilled (E)	4.1	Overweight (25.0 – 29.9)	34
> 65	9.2	Other	8.8	Obese (> 30.0)	12

163 *Measures*

164 Consumption of confectionery foods was measured using 12 items selected and adapted from a food
 165 frequency questionnaire (FFQ) originally designed for the EPIC-Norfolk study (UK) (EPIC-Norfolk,
 166 2012) (see appendix A for the list of items). These 12 items represent the top sources of free sugars
 167 in the British diet (PHE, 2016). Respondents indicated their consumption of these foods on a nine
 168 point response grid ranging from ‘never or less than once a month’ to ‘more than six times per day’.
 169 The Composition of Foods (1995) by the Food Standards Agency and Food Portion Sizes (1995) by
 170 Helen Crawley were used to estimate the grams of free sugar attributed to an average serving of
 171 each confectionery food item. For example, an average milk chocolate bar, 54g, contains 286
 172 calories and 31 grams of sugar. Based on these figures an estimate of an individual’s total daily
 173 consumption of sugars in grams from confectionery food was calculated.

174 The dietary behaviour of interest in this study is decreasing consumption of confectionery foods and
 175 was measured using one question: ‘Thinking back over the last six months, have you changed your
 176 average weekly intake of confectionery foods?’ Responses were classified into two categories:
 177 decreased and did not decrease. In addition, study participants were asked whether they had a
 178 lifestyle goal during the previous six months. As this was a cross-sectional study with the aim of
 179 assessing change over time, respondents were required to retrospectively reflect on their behaviour
 180 over a six-month period (Naughton *et al.*, 2015).

181 Exploratory factor analysis (EFA) using principle components (varimax) was used to test the
 182 dimensionality of the combined social cognition variables confectionery habit, hedonic hunger,
 183 perceived need, PBC, and dietary planning, as the items used to measure these variables were taken
 184 from multiple sources and some items were created for the study in terms of TACT (target, action,
 185 context and time), which is a commonly used approach (Ajzen, 2002). Table 2 shows the means,

186 standard deviations and factor loadings for each measurement item as well as the eigenvalues,
187 percentage of variance explained and reliability scores for each latent variable. Confectionery habit
188 was measured by seven items designed to represent two of the most important characteristics of
189 habitual behaviour: repetition and automaticity. The majority of these items were taken from the
190 self-report habit index (SRHI) (Verplanken & Orbell, 2003). Based on the EFA one item was removed
191 as it failed to meet the criteria of all factor loadings being > 0.4 (Stevens, 2002). Hedonic hunger
192 was measured using the Power of Food Scale (PFS), which is a validated 15 item measurement scale
193 designed to assess the appetitive aspects of eating (Lowe and Butryn, 2007; Cappelleri *et al.*, 2009;
194 Lowe *et al.*, 2009). Perceived need was measured by three items adapted from Paisley & Sparks
195 (1998) and Payne *et al.* (2004) and PBC was measured by five items adapted from Armitage and
196 Conner, (1999), and Povey *et al.* (2000). Based on the EFA one PBC item was removed as it failed to
197 meet the criteria of all factor loadings being > 0.4 (Stevens, 2002). The Cronbach's alpha for PBC was
198 0.55. While this is generally considered low, Kline (1999) notes that for psychological variables,
199 values below 0.7 can be expected because of the diversity of the variables being measured. Nunnally
200 (1967) (as cited in Peterson, 1994) suggested that a reliability score from 0.5 to 0.6 is the minimum
201 acceptable level. The items representing these four variables were measured on seven-point
202 agreement scales (1 = strongly disagree, 4 = neither agree/disagree, 7 = strongly agree). Dietary
203 planning was represented by five items adapted from Luszczynska & Schwarzer (2003), Rise *et al.*
204 (2003), Otis & Pelletier (2008) & Zandstra *et al.* (2010) measured on frequency scales (1 = never, 2 =
205 very rarely, 3 = rarely, 4 = sometimes, 5 = frequently, 6 = somewhat frequently 7 = very frequently).
206 All the variables had eigenvalues over Kaiser's criterion of 1 and in combination explained 61% of the
207 variance.

208

Table 2 Descriptive statistics, factor loadings and Cronbach's alpha for the study measures

	Mean (SD)	Factor Loading	% of variance	Eigen value	Cronbach's alpha*
Confectionery habit			13.62	4.22	0.85
When I am busy and I am hungry I am very likely to eat Confectionery foods	4.48 (1.89)	0.75			
When I am hungry and I am in a rush, I am very likely to eat Confectionery foods	4.24 (1.93)	0.78			
I would find it difficult not to eat Confectionery foods	4.21 (1.94)	0.72			
Eating Confectionery foods is something I do frequently	4.13 (1.91)	0.77			
Eating Confectionery foods is something I do without thinking	4.02 (1.92)	0.71			
Eating Confectionery foods is something I start doing before I realise I am doing it	3.78 (1.93)	0.74			
Hedonic Hunger			21.45	6.65	0.92
When I know a delicious food is available, I can't help myself from thinking about having some	4.11 (1.81)	0.76			
It's frightening to think of the power that food has over me	3.01 (1.79)	0.72			
It seems like I have food on my mind a lot	3.01 (1.71)	0.76			
Just before I taste a favourite food, I feel intense excitement	3.33 (1.74)	0.74			
If I see or smell a food I like, I get a powerful urge to have some	4.15 (1.75)	0.75			
Hearing someone describing a great meal makes me really want to have something to eat	3.61 (1.80)	0.73			
When I am around a fattening food I love, it's hard to stop myself from at least tasting it	4.22 (1.80)	0.70			
I think that I enjoy eating a lot more than most other people	3.17 (1.67)	0.70			
I love the taste of certain foods so much that I can't avoid eating them even if they are bad for me	3.98 (1.86)	0.60			
I get more pleasure from eating than I do from almost anything else	3.09 (1.72)	0.66			
I find myself thinking about food even when I am not physically hungry	3.59 (1.82)	0.63			
Before I eat a favourite food my mouth starts to water	3.61 (1.79)	0.67			
When I eat delicious food I focus a lot on how good it tastes	4.56 (1.65)	0.64			
Perceived behavioural control (PBC): Over the last six months, during the course of a typical week...			5.45	1.70	0.55
I found it easy not to overindulge on Confectionery foods.	4.50 (1.71)	0.38			
I was confident that, if I wanted to, I could avoid eating Confectionery foods.	5.38 (1.38)	0.60			

Whether I did or did not eat Confectionery foods was entirely up to me.	5.86 (1.20)	0.74			
There were plenty of opportunities for me to choose healthier alternatives to Confectionery foods.	5.37 (1.21)	0.48			
Dietary Planning: <i>Over the last six months, during the course of a typical week...</i>			13.40	4.14	0.93
I would plan ahead how often I could eat Confectionery foods in a day.	3.09 (1.73)	0.71			
I had plans in place to avoid Confectionery foods whenever I felt bored.	3.18 (1.80)	0.83			
I had plans in place to avoid Confectionery foods whenever I felt stressed.	3.01 (1.74)	0.92			
I had plans in place to avoid Confectionery foods whenever I felt in bad mood.	2.97 (1.74)	0.91			
I had plans in place to avoid Confectionery foods whenever I felt tempted.	3.41 (1.81)	0.92			
Perceived Need			7.40	2.30	0.81
Eating Confectionery foods is something I need to avoid doing as part of my diet.	5.19 (1.80)	0.81			
I need to stay away from Confectionery foods in order to have a healthy lifestyle.	5.40 (1.61)	0.87			
In order to have a healthy diet, it is important that I have a low intake of Confectionery foods.	5.79 (1.37)	0.80			

Reliability test: > 0.7 is generally considered acceptable (Kline, 1999). For psychological variables, values below

0.7 can be expected because of the diversity of the variables being measured (Kline, 1999)

Data Analysis

Latent class analysis was performed using PROC LCA in SAS 9.3 (Lanza *et al.*, 2007). This is a statistical procedure used to identify a set of discrete, mutually exclusive latent classes of individuals based on responses to categorical variables (Lanza *et al.*, 2007). In the present study two variables are categorical; confectionery consumption decrease and lifestyle goal. All of the social cognitive variables used are continuous. Therefore, in order to run the latent class analysis, the overall mean score for confectionery habit, hedonic hunger, perceived need, and PBC was calculated and each variable was recoded to create three categories representing strong, moderate and weak. Strong represents respondents that on average agreed with the statements (≥ 5 on the likert scale). Weak represents respondents who on average disagreed with the statements (< 4 on the likert scale). Moderate represents respondents neither agreed nor disagreed with the statements on average (4 – 4.9 on the likert scale). As dietary planning was measured on a frequency scale this variable was recoded into three categories: never, rarely, and sometimes/frequently. The split points were: $< 2 =$ never, rarely = 2 - 3.9; sometimes/frequently ≥ 4 .

Latent class analysis (LCA) identifies one categorical latent variable with several categories or classes to explain the relationship between the measured variables (i.e. confectionery consumption decrease, lifestyle goal, strong confectionery habit, strong hedonic hunger, strong perceived need, strong perceived behavioural control and sometimes/frequently planned). The objective is to identify a parsimonious model that uses as few latent classes as possible to explain the data (Geiser, 2010). Model selection was determined by examining Akaike information criteria (AIC) and Bayesian Information criteria (BIC), with lower values representing better suitability based on model fit and parsimony, as well as model interpretability based on theoretical knowledge (Collins & Lanza, 2010). An easy to interpret solution is characterised by the majority of conditional response probabilities being close to 1 or close to 0 with few medium sized conditional response probabilities (Geiser, 2010). Sugar consumption g/d from confectionery foods was examined across the segments identified from the latent class analysis using one-way ANOVA analysis. In addition, multi-nominal logistic regression was run to predict segment membership based on socio-demographics and BMI. This entails examining the change in odds (i.e. the odds ratio) of belonging to a particular segment relative to a reference segment from a unit change in the predictor. Data screening using box-plots identified extreme outliers in the total sugar consumption variable, which reflected unrealistic daily intake of added sugar from confectionery foods. These cases were removed resulting in a final sample of $n=477$.

3. Results

The descriptive statistics show that 38% of the total study sample ($n=477$) decreased their intake of sugar in the preceding six months and 42% of the sample had a lifestyle goal related to sugar consumption during that six-month period. The most mentioned goals were to improve health and protect against illness. On average, people were consuming 52.45 grams of free sugar per day from confectionery foods alone, which is greater than the 50g limit of total free sugar consumption specified by the WHO. A comparison of the observed means for the social cognition variables with the scale midpoints for each of these variables (i.e. 4) shows that people were moderately disposed to habitually consume confectionery foods, and hedonic hunger also had a moderate influence on their diet. PBC towards the consumption of confectionery foods was generally strong, indicating that in general people felt they had control over their consumption. Similarly, people agreed that they needed (i.e. perceived need) to regulate the amount of confectionery foods they consumed. However, people's tendency to plan consumption of confectionery was only moderate.

Table 3 Descriptive statistics for the behavioural social cognition variables N = 477

	%		\bar{X} (SD)
Decrease in confectionery consumption	37.7	Sugar from confectionery foods g/d	52.45 (40.52)
Lifestyle goal (yes)	42.3	Confectionery Habit	4.23(1.40)
Lose weight	31.0	Hedonic Hunger	3.70 (1.24)
Improve Health	81.7	Perceived Need	5.46(1.37)
Protect against illness	54.2	Perceived Behavioural Control (PBC)	5.28 (0.90)
Enhance appearance	21.8	Dietary Planning	3.20 (1.40)

Note: the scale midpoint is 4

An examination of the model fit statistics from the latent class analysis indicate that a 3 - 6 class/segmentation³ solution provides the best model fit; while the BIC values indicate that a 3 segmentation solution is optimal the AIC values indicate a 6 segmentation solution (Table 4). In LCA the interpretability of a solution is equally important as the model fit statistical criteria (Collins & Lanza, 2010). An inspection of the proportional probabilities of the 3, 4, 5 and 6 segment models suggest that the 4 segment model provides the best fit as the segments are substantial and distinguishable labels can be assigned to each (Table 5 & Figure 1). As table 5 shows the largest segment, segment 4 (unmotivated), represented 35% of the sample and was characterised by having no lifestyle goals and virtually no decrease in confectionery consumption. This segment had high probabilities for confectionery habits and hedonic hunger and low probabilities for perceived need, PBC and dietary planning. A second segment, segment 3 (thrivers), representing 28% of the sample was distinguished by virtually nobody having strong confectionery habits and hedonic hunger. The remaining two segments were characterised by high probabilities for having lifestyle goals related to sugar consumption. Segment 2 (successful actors), representing 17% of the population, was distinguished by high probabilities for perceived need, PBC and dietary planning. The majority of people in this segment (92%) decreased their sugar consumption over the previous six months and nearly everybody (96%) had a sugar related lifestyle goal. Triers (Segment 1), representing 20% of the sample, were less likely to have decreased sugar consumption (67%) despite everybody having a sugar related lifestyle goal. This segment was characterised by having a low probability for strong PBC and compared to successful actors a lower probability for strong perceived need and higher probabilities for strong habit and strong hedonic hunger. The relationship between sugar consumption in grams per day from confectionery foods and segment membership was significant ($F(3, 473) = 24.600, p < 0.001$).

³ In this paper the term segment is used rather than class

Post-hoc tests (Tukey HSD test) revealed that triers consumed the most sugar in grams per day, significantly more than thrivers and successful actors but there were no significant differences between triers and the unmotivated segment and no differences between thrivers and successful actors.

Table 4 Model Fit information for competing latent class models (n = 477)


Number of segments	G ²	df	AIC	BIC	Entropy
1	1370.96	959	1394.96	1444.97	1.0
2	973.08	946	1023.08	1127.27	0.84
3	834.29	933	910.29	1068.66	0.80
4	764.41	920	866.41	1078.95	0.81
5	710.89	907	838.89	1105.68	0.81
6	680.33	894	834.33	1155.23	0.81

Table 5 Conditional response probabilities for each social cognitive variable as a function of segment membership and sugar consumption g/d from confectionery foods across the segments (n = 477)

	Segment 1 Triers 20% (n = 97)	Segment 2 Successful Actors 17% (n = 79)	Segment 3 Thrivers 28% (n = 134)	Segment 4 Unmotivated 35% (n = 168)
Decrease confectionery consumption	0.674	0.915	0.229	0.085
Lifestyle goal (yes)	1.000	0.959	0.245	0.000
Strong Confectionery Habit	0.537	0.067	0.000	0.600
Strong Hedonic Hunger	0.685	0.174	0.021	0.449
Strong Perceived Need	0.409	0.805	0.242	0.219
Strong PBC	0.260	0.723	0.546	0.127
Planning (sometimes/freq.)	0.492	0.694	0.147	0.125
Sugar consumption g/d (M, sd)	68.04 (45.12)a	34.53 (27.33)b	36.96 (27.03)b	61.23 (43.47)a

Note: ab subscripts denote statistically significant differences between segments for sugar consumption g/d

Figure 1 Line graph of proportional probabilities



The results of the multinomial logistic regression analysis are presented in tables 6, 7 and 8. Gender, age and BMI were significant predictors of segment membership but social class was not. Table 6 shows that compared to adults over 55 years of age, 18-34 year olds and 35 – 54 year olds were less likely to be in the successful actors segment ($OR = 0.25, p < 0.05$ & $OR = 0.46, p < 0.05$ respectively) and thrivers segment ($OR = 0.33, p < 0.05$ & $OR = 0.30, p < 0.05$ respectively) than be in the unmotivated segment, indicating a negative relationship between sugar consumption levels and age i.e. sugar consumption decreases with increasing age. Similarly, Table 7 shows that compared to adults over 55 years of age, 18-34 year olds were less likely to be in the successful actors segment ($OR = 0.16, p < 0.05$) and thrivers segment ($OR = 0.22, p < 0.05$) than be in the triers segment. Regarding gender, men were half as likely to be in the successful actors segment ($OR = 0.41, p < 0.05$) than be in the unmotivated segment, reflecting a greater propensity among women to try to reduce sugar consumption (table 6). However, men were also 2.21 times more likely to be in the thrivers segment than the triers segment ($p < 0.05$) (Table 7) and three times more likely to be in the thrivers segment than the successful actors segment ($p < 0.05$) (Table 8). This reflects the dichotomy among the male population as they are either successfully consuming low intakes of sugar g/d from confectionery foods in line with the guidelines or they are consuming excessive amounts with less inclination than women to make a change. An examination of BMI suggests that the unmotivated and triers segments represent 'at risk' groups. Compared to obese adults, adults with normal BMI were 2.66 times more likely to be in the thrivers segment than the unmotivated segment ($p < 0.05$) (table 6) and 4.45 times more likely to

be in the thrivers segment than the triers segment ($p < 0.05$) (table 7). BMI may also explain the higher ratings for perceived need among successful actors as people with normal BMI compared to obese adults are 3.4 times more likely to be in the thrivers segment than the successful actors segment ($p < 0.05$) (table 8). There were no differences between the unmotivated segment and the Triers segment in terms of gender, age, social class and BMI.

Table 6 Multi-nominal logistic regression analysis for the influence of socio-demographics and BMI on segment membership with segment 4 unmotivated as the reference category

	Segment 1 Triers OR (95% CI)	Segment 2 Successful Actors OR (95% CI)	Segment 3 Thrivers OR (95% CI)	Segment 4 Unmotivated 35% (REF)	Overall pvalue ^a
Gender (Male)	0.57 (0.32, 1.01)	0.41 (0.22, 0.78)*	1.27 (0.75, 2.13)	REF.	<0.01
Age (18-34 years)	1.52 (0.65, 3.56)	0.25 (0.10, 0.60)*	0.33 (0.16, 0.66)*	REF.	<0.01
Age (35-54 years)	0.70 (0.30, 1.64)	0.46 (0.22, 0.98)*	0.30 (0.15, 0.58)*	REF.	
Age (over 55 years)	.	.	.	REF.	
Social class (AB)	1.12 (0.49, 2.54)	0.84 (0.36, 1.95)	1.50 (0.72, 3.13)	REF.	0.92
Social class (C1C2)	1.08 (0.49, 2.39)	0.84 (0.37, 1.88)	1.26 (0.61, 2.60)	REF.	
Social class (DE)	.	.	.	REF.	
BMI (Normal)	0.60 (0.25, 1.44)	0.77 (0.31, 1.93)	2.66 (1.05, 6.79)*	REF.	0.06
BMI (Overweight)	0.50 (0.20, 1.22)	0.70 (0.28, 1.76)	1.68 (0.66, 4.30)	REF.	
BMI (Obese)	.	.	.	REF.	

^a p-value reflects test of overall association between predictor and segment membership based on likelihood ratio difference * $p < 0.05$

Table 7 Multi-nominal logistic regression analysis for the influence of socio-demographics and BMI on segment membership with segment 1 triers as the reference category

	Segment 1 Triers (REF)	Segment 2 Successful Actors OR (95% CI)	Segment 3 Thrivers OR (95% CI)	Segment 4 Unmotivated 35% (95% CI)	Overall pvalue ^a
--	------------------------------	--	--------------------------------------	---	--------------------------------

Gender (Male)	REF.	0.72 (0.35, 1.46)	2.21 (1.21, 4.06)*	1.75 (0.99, 3.11)	<0.01
Age (18-34 years)	REF.	0.16 (0.06, 0.43)*	0.22 (0.09, 0.50)*	0.66 (0.28, 1.55)	<0.01
Age (35-54 years)	REF.	0.66 (0.27, 1.62)	0.43 (0.18, 0.99)*	1.43 (0.61, 3.37)	
Age (over 55 years)	REF.	.	.		
Social class (AB)	REF.	0.75 (0.29, 1.97)	1.34 (0.56, 3.24)	0.90 (0.39, 2.03)	0.92
Social class (C1C2)	REF.	0.77 (0.31, 1.97)	1.17 (0.49, 2.79)	0.93 (0.42, 2.06)	
Social class (DE)	REF.	.	.		
BMI (Normal)	REF.	1.30 (0.48, 3.46)	4.45 (1.61, 12.29)*	1.67 (0.70, 4.00)	0.06
BMI (Overweight)	REF.	1.42 (0.51, 3.91)	3.40 (1.20, 9.63)*	2.01 (0.82, 4.96)	
BMI (Obese)	REF.	.	.		

^a p-value reflects test of overall association between predictor and segment membership based on likelihood ratio difference * $p < 0.05$

Table 8 Multi-nominal logistic regression analysis for the influence of socio-demographics and BMI on segment membership with segment 2 successful actors as the reference category

	Segment 1 Triers OR (95% CI)	Segment 2 Successful Actors REF.	Segment 3 Thrivers OR (95% CI)	Segment 4 Unmotivated 35% (95% CI)	Overall pvalue ^a
Gender (Male)	1.39 (0.68, 2.84)	REF.	3.08 (1.60, 5.96)*	2.44 (1.29, 4.61)*	<0.01
Age (18-34 years)	6.19 (2.31, 16.59)*	REF.	1.33 (0.56, 3.17)	4.08 (1.67, 9.94)*	<0.01
Age (35-54 years)	1.51 (0.62, 3.70)	REF.	0.65, (0.31, 1.35)*	2.17 (1.02, 4.61)*	
Age (over 55 years)	.	REF.	.	.	
Social class (AB)		REF.			0.92
Social class (C1C2)		REF.			
Social class (DE)		REF.	.	.	
BMI (Normal)	0.77 (0.29, 2.06)	REF.	3.43 (1.22, 9.59)*	1.29 (0.52, 3.20)	0.06
BMI (Overweight)	0.71 (0.26, 1.94)	REF.	2.40 (0.85, 6.77)	1.42 (0.57, 3.56)	
BMI (Obese)	.	REF.	.		

^a p-value reflects test of overall association between predictor and latent segment membership based on likelihood ratio difference * $p < 0.05$

4. Discussion

The post-hoc segmentation analysis suggests that there are four cohorts of people with regards the probability of reducing consumption of confectionery foods; triers, successful actors, thrivers, and unmotivated. These segments can be considered operationally useful as they are measurable using social cognition variables, substantial regarding the relative size of each segment,

differentiable across sugar consumption g/d from confectionery foods, and accessible based on the demographic profiles of each segment (Kotler & Keller, 2009). The unmotivated and thrivers segments are the least likely to decrease their consumption of confectionery foods but while people in the thrivers segment are adhering to dietary guidelines people in the unmotivated segment are consuming over 50 g/d, which is the guideline limit on free sugar intake set by the WHO (WHO, 2015). People in the triers segment are more likely to decrease their consumption of confectionery foods but are less likely than people in the successful actors segment and are consuming comparable amounts of sugar to people in the unmotivated segment. The findings provide insights for social marketing interventions targeting changes in individual behaviour by addressing downstream influences (e.g. people's lack of perceived need and dietary planning) and the upstream influences (e.g. the food environment) that promote the consumption of confectionery foods.

The triers and the unmotivated segments reported low levels of perceived need compared to the successful actors segment. For the unmotivated segment this finding is consistent with no segment member having a lifestyle goal and achieving a decrease in confectionery consumption. This is a concern from a health policy perspective considering that the unmotivated segment contains significantly more obese people compared to the thrivers segment. In the triers segment 100% of the sample had a lifestyle goal related to sugar consumption but the rate of successful change was 67% compared to 92% for the successful actors segment. While it is well documented that setting goals at the outset of a behaviour change process is important in achieving the desired behavioural change (Schnoll & Zimmerman, 2001; Nothwehr & Yang, 2006; Papies *et al.*, 2007) it is possible that perceived need (in addition to PBC discussed below) may act as a boundary condition on the likelihood of successful change. According to Paisley and Sparks (1998), even if a behaviour is seen as beneficial and wise, indicating a positive attitude, there may be a low perceived need to perform the behaviour because a person perceives that it is not necessary for him-or herself to carry out the behaviour and the outcome (i.e. the lifestyle goal) is attainable through other means. Therefore, social marketing interventions will need to address the lack of perceived need by identifying triggers that move people from pre-contemplation via contemplation to action (Andreasen, 2003). This findings indicates that such interventions should be targeted predominately at men as perceived need may explain the gender differences between the segments with men more likely to be in the unmotivated segment compared to the motivated segments (i.e. successful actors and triers). This corresponds with previous studies that show women are more concerned about their diet and more motivated to make dietary changes (Wardle *et al.*, 2004; Davey *et al.* 2006; Hearty *et al.*, 2007). Regarding sugar specifically, Davey *et*

al. (2006) found that a significantly higher percentage of women agreed that they had too much sugar in their diets. However, while men are more likely to be in the unmotivated segment compared to the motivated segments (i.e. successful actors and triers) they are also more likely to be in the thrivers segment, which is characterised by low ratings on confectionery habit and hedonic hunger. This finding may be explained by research which suggests that men are less ambivalent towards nutrition and restraint eating, dieting and eating disorders are less common (Kiefer, *et al.*, 2005).

Compared to the successful actors and thrivers segments, the triers and the unmotivated segments reported high levels of confectionery habit and hedonic hunger in addition to weak levels of PBC. A key component of habitual behaviour is automaticity or a lack of conscious thought, which often leads to environmentally cued behaviour that is not consistent with ones' behavioural intentions (Verplanken and Aarts, 1999). Similarly, hedonic hunger is driven by affective rather than cognitive responses to food stimuli which results in people failing to control consumption when presented with tempting foods (Lowe and Butryn, 2007). Furthermore, PBC, confectionery habit and hedonic hunger may explain the age differences between the identified segments in this study as successful actors and thrivers were significantly older than both the triers and the unmotivated segments. Studies have consistently shown a positive correlation between stronger PBC and increasing age with younger adults more likely to list barriers to eating healthily (Kearney and McElhone, 1999; Escoto *et al.*, 2012). Therefore, in addressing the consumption of confectionery foods among younger adults it is important to consider upstream influences such as the food environment (Hasting's *et al.*, 2000). According to Swinburn *et al.* (2011 pg. 804) the increasing availability of cheap, high energy, nutrient poor foods, has resulted in a '*passive overconsumption of energy*'. Social marketers could target stakeholders interested in promoting healthier eating aimed at limiting the availability of confectionery food displays (e.g. end-of-aisle displays & island displays) within stores and workplaces as these displays may promote habitual and hedonic consumption and override self-control (Thornton *et al.*, 2012). Research has shown that situational changes (e.g. expanding the availability of healthy products in work canteens and moving healthier products to the point of purchase) have positive effects on people's food choices and eating patterns (Engbers *et al.*, 2005; Gittelsohn *et al.*, 2012).

Dietary planning has been proposed as a means to facilitate people to take more control of their diet and consequently avoid unhealthy eating practices (Armitage, 2004; Scholz *et al.*, 2009; Naughton *et al.* 2015). In this study there was a 20% difference in planning between triers and successful actors, which suggests the importance of social marketing interventions targeting

young adults who are motivated to reduce their consumption of confectionery foods with initiatives that promote dietary planning. According to Gollwitzer (1993) a specific plan helps to promote the initiation and efficient execution of goal-directed activity by laying down 'if-then' contingencies between situational cues and goal fulfilling responses. Once such contingencies are present, actions that lead to goal fulfilment gain a degree of automaticity by being under the control of relevant situational cues. Therefore, planning can help break the influence of habits on future behaviour. Verplanken and Faes (1999) found that individuals had formed implementation intentions ate healthily irrespective of their level of unhealthy eating habits.

Interestingly, social class was not associated with segment membership as research indicates that lower social class groupings are least likely to adhere to dietary guidelines (Parmenter *et al.*, 2000; Harrington *et al.*, 2008), are less likely to be concerned about their health and healthy eating (Wardle & Steptoe, 2003; Dibsall, *et al.*, 2003; Hearty *et al.*, 2007) and less likely to implement healthy lifestyle changes (NHF, 2007). However, Darmon & Dreowski (2008) in a review of the epidemiological data on the relationship between diet quality and social class, found there was less evidence that social class was related to confectionery consumption in comparison with the other food groupings (i.e. fruit and vegetables, fatty meats etc.). It may be the case that in general people are more perceptive to the message on the importance of reducing fat consumption, the predominant focus of dietary guidelines for the last three decades (Hite *et al.*, 2010), rather than the message on limiting sugar intake. Carrillo *et al.* (2011), found that consumers associated the items 'low in calories', and 'helps me control my weight' more strongly with the label 'is low in fat' than the label 'is low in sugar'.

Conclusion

In support of Lefebvre's (2000) and more recently Luca & Suggs's (2013) call to action for the increased use of theory to guide social marketing interventions, this paper provides theoretical support for the potential utility of using behavioural theory in social marketing campaigns. Using a social cognition approach, a number of important variables underlying food choice and healthy eating were identified and selected as segmentation bases. Further, in line with best practice a post-hoc segmentation method (i.e. Latent Class Analysis) was applied. This resulted in the identification of four segments that are operationally useful with respect to informing the design of effective and tailored social marketing strategies. The unmotivated segment and triers segment are the concern from a public health perspective and the defining demographic characteristic of both these segments is the disproportionately high percentage of young and middle aged adults. The social cognition findings indicate a lack of perceived need and perceived behavioural control

related to sugar intake among this cohort of adults. Therefore, a marketing strategy that can identify relevant messages or 'triggers' related to sugar consumption that resonate with the target segment will be an important step towards addressing the low ratings in perceived need. As the behaviour being promoted is likely to be unpalatable to the target market (i.e. reducing sugar consumption) it may be important to go beyond simply communicating the consequences associated with a diet high in free sugars (Peattie and Peattie, 2009). In addition, dietary change and maintenance entails improved self-control to counteract unhealthy eating habits and the effects of hedonic hunger. This may require a focus on the food environment by targeting stakeholders interested in promoting healthier eating with the aim of addressing structural barriers to healthy eating including the ease and availability of confectionery foods relative to the availability and cost of nutritious foods.

Study Limitations

While this study contributes to and builds on existing literature in the domain of health behaviour change and the use of segmentation and theory to guide social marketing campaigns there are nonetheless some limitations. Firstly, in order to examine causality the current study adopted a retrospective approach in the data collection. Retrospective studies provide cost and time efficiencies over longitudinal (prospective) studies, but there are drawbacks to this method of data collection particularly linked to the potential for recall bias and the impact of one set of answers affecting the answers to other questions in the survey. To limit this possibility the survey was designed to present the multiple questions measuring each variable in a random order (i.e. the same types of questions were not presented consecutively). Therefore, it would have required substantial cognitive effort to manipulate one's answers to all the questions. Secondly, this study used a food frequency measure of confectionery food consumption in order to calculate consumption of sugars g/d from confectionery food items. Research indicates that food frequency questionnaires tend to produce an underestimation of true dietary intake and a food diary is a more valid measure of consumption (Bedard, *et al.*, 2004; Day *et al.*, 2001). A food diary measure of consumption pre and post study would have also provided a more accurate estimate of confectionery consumption change. Finally, it should be clarified that while the measure of sugar intake used in this study relates to the main sources of free sugars, it does not represent all the identified sources of free sugars, for example free sugars from alcohol and savoury items such as sauces.

Appendix A

Items measuring confectionery food consumption
Chocolate biscuits, e.g. digestive, cookies
Plain biscuits e.g. Nice, ginger (one)
Cakes e.g. fruit, sponge, chocolate
Buns, pastries e.g. croissants, doughnuts, muffins
Fruit pies, tarts, crumbles
Milk pudding e.g. rice, custard, trifle
Ice cream and flavoured yogurts
Sweets e.g. chocolates, toffees, mints, jellies
Chocolate snack bars e.g. Mars, Crunchie (standard size)
Sugar added to Tea, Coffee, cereal (teaspoon)
Fizzy soft drinks, e.g. Coca cola, lemonade (glass)
Jam, marmalade, honey, syrup (teaspoon)

References

Ajzen, I. (1991) 'The theory of planned behaviour', *Organisational behaviour and human decision processes*, 50: 179-211.

Ajzen, I. (2002) Constructing a TPB questionnaire: conceptual and methodological considerations. [Online](Updated 2006) Available at <http://www.people.umass.edu/aizen/pdf/tpb.measurement.pdf>

Ahluwalia, J. S., Nollen, N., Kaur, H., James, A. S., Mayo, M. S. & Resnicow, K. (2007) 'Pathway to health: Cluster-randomized trial to increase fruit and vegetable consumption among smokers in public housing', *Health Psychology*, 26: 214-221.

Andreasen, A. R. (2003) 'The life trajectory of social marketing: some implications', *Marketing Theory*, 3: 293-303.

Andreasen, A. R. (2002) 'Marketing Social Marketing in the Social Change Marketplace', *Journal of Public Policy & Marketing*, 21: 3-13.

- Armitage, C. J. (2004) 'Evidence That Implementation Intentions Reduce Dietary Fat Intake: A Randomized Trial', *Health Psychology*, 23: 319-323.
- Armitage, C. J. & Conner, M. (1999) 'The theory of planned behaviour: Assessment of predictive validity and 'perceived control', *British Journal of Social Psychology*, 38: 35-54.
- Armitage, C. J. & Conner, M. (2000), 'Social Cognition Models and Health Behaviour: A Structured Review', *Psychology & Health*, 15: 173-89.
- Armitage, C. J. & Conner, M. (2001) 'Efficacy of the Theory of Planned Behaviour: A meta-analytic review', *British Journal of Social Psychology*, 40: 471.
- Bagozzi, R. P. (1992) 'The self-regulation of attitudes, intentions, and Behaviour', *Social Psychology Quarterly*, 55: 178-204.
- Bedard, D., Shatenstein, B. & Nadon, S. (2004) 'Underreporting of energy intake from a self-administered food-frequency questionnaire completed by adults in Montreal', *Public health nutrition*, 7: 675-682.
- Berg-Smith, S. M., Stevens, V. J., Brown, K. M., Van Horn, L., Gernhofer, N., Peters, E., Greenberg, R., Snetselaar, L., Ahrens, L. & Smith, K. (1999) 'A brief motivational intervention to improve dietary adherence in adolescents', *Health Education Research*, 14: 399-410.
- Bridle, C., R. P. Riemsma, J. Pattenden, A. J. Sowden, L. Mather, I. S. Watt, and A. Walker (2005) 'Systematic Review of the Effectiveness of Health Behavior Interventions Based on the Transtheoretical Model', *Psychology & Health*, 20: 283-301.
- Cappelleri, J. C., Bushmakin, A. G., Gerber, R. A., Leidy, N. K., Sexton, C. C., Karlsson, J. & Lowe, M. R. (2009) 'Evaluating the Power of Food Scale in obese subjects and a general sample of individuals: development and measurement properties', *International Journal of Obesity*, 33: 913-922.
- Carrillo, E., P. Varela, A. Salvador, and S. Fizman (2011) 'Main Factors Underlying Consumers' Food Choice: A First Step for the Understanding of Attitudes toward Healthy Eating', *Journal of Sensory Studies*, 26: 85-95.
- Collins, L. M., & Lanza, S. T. (2010) *Latent class and latent transition analysis: With applications in the social, behavioral, and health sciences*. New York: Wiley.
- Conner, M. & Sparks, P. (2005) 'Theory of Planned Behaviour and Health Behaviour', in Conner, M. & Norman, P. (Eds.) *Predicting Health Behaviour*. Berkshire: McGraw-Hill, 170-223.
- Crawley, H. (1993) *Food Portion Sizes*. London: Food Standards Agency.
- Dann, S. (2010) 'Redefining Social Marketing with Contemporary Commercial Marketing Definitions', *Journal of Business Research*, 63: 147-53.
- Darmon, N. & Drewnowski, A. (2008) 'Does Social Class Predict Diet Quality?', *Am J Clin Nutr*, 87: 1107-1117.

- Davy, S. R., Benes, B.A. and Driskell, J.A. (2006) 'Sex Differences in Dieting Trends, Eating Habits, and Nutrition Beliefs of a Group of Midwestern College Students', *J Am Diet Assoc*, 106: 1673-1677.
- Day, N. E., McKeown, N., Wong, M. Y., Welch, A. & Bingham, S. (2001) 'Epidemiological assessment of diet: a comparison of a 7-day diary with a food frequency questionnaire using urinary markers of nitrogen, potassium and sodium', *International Journal of Epidemiology*, 30: 309-317.
- De Bruijn, G. J. (2010) 'Understanding college students fruit consumption. Integrating habit strength in the theory of planned behaviour', *Appetite*, 54: 16-22.
- Dibsdall, LA, Lambert, N., Bobbin, R.F. and Frewer, L. J. (2003) 'Low-Income Consumers' Attitudes and Behaviour Towards Access, Availability and Motivation to Eat Fruit and Vegetables', *Public health nutrition*, 6: 159-168.
- Eagle, L., Stephan, D. Hill, S. , Bird, S. , Spotswood, F., and Tapp, A. (2013) *Social Marketing*. Edinburgh: Pearson.
- Engbers, L. H., van Poppel, M. N., Paw, M. J. C. A., & van Mechelen, W. (2005) 'Worksite health promotion programs with environmental changes: a systematic review', *American journal of preventive medicine*, 29: 61-70.
- Escoto, K. H. , Laska, M.N., Larson, N., Neumark-Sztainer, D. and Hannan, P.J. (2012) 'Work Hours and Perceived Time Barriers to Healthful Eating among Young Adults', *American journal of health behavior*, 36: 786-796.
- French, J. & Blair-Stevens, C. (2006) *Social Marketing National Benchmark Criteria*. London: National Social Marketing Centre.
- FSAI (2011) Scientific Recommendations for Healthy Eating Guidelines in Ireland. Dublin, Food Safety Authority of Ireland (FSAI).
- Glanz, K., Basil, M., Maibach, E., Goldberg, J. & Snyder, D. (1998) 'Why Americans eat what they do: taste, nutrition, cost, convenience, and weight control concerns as influences on food consumption', *Journal of the American Dietetic Association*, 10: 1118-1126.
- Geiser, C. (2010) *Data Analysis with Mplus*. London: Guilford Press.
- Gittelsohn, J., Rowan, M., & Gadhoke, P. (2012) 'Interventions in small food stores to change the food environment, improve diet, and reduce risk of chronic disease', *Preventing chronic disease*, 9.
- Gollwitzer, P. M. (1993) 'Goal Achievement: The Role of Intentions', *European Review of Social Psychology*, 4: 141 - 185.
- Gordon, R., McDermott, L. Stead, M. & Angus. K (2006) 'The Effectiveness of Social Marketing Interventions for Health Improvement: What's the Evidence?', *Public Health*, 120: 1133-1139.
- Grier, S. and Bryant, C. A. (2005) 'Social Marketing in Public Health', *Annual Review Public Health*, 26: 319-339.

- Harrington, J., Perry, L., Lutomski, J., Morgan, K., McGee, H., Shelley, E., Watson, D. & Barry, M. (2008) SLAN 2007: Survey of Lifestyle, Attitudes and Nutrition in Ireland. Dietary Habits of the Irish Population. Dublin: Department of Health and Children.
- Hastings, G, MacFadyen, L. and Anderson, S. (2000) 'Whose Behavior Is It Anyway? The Broader Potential of Social Marketing', *Social Marketing Quarterly*, 6: 46-58.
- Hearty, A., McCarthy, S., Kearney, J. & Gibney, M. (2007) 'Relationship between attitudes towards healthy eating and dietary behaviour, lifestyle and demographic factors in a representative sample of Irish adults', *Appetite*, 48: 1-11.
- Hite, A. H., Feinman, R.D., Guzman, G.E., Satin, M., Schoenfeld, P. & Wood, R. J. (2010) 'In the Face of Contradictory Evidence: Report of the Dietary Guidelines for Americans Committee', *Nutrition*, 26: 915-24.
- Kazbare, L., van Trijp, H. CM. and Eskildsen, J.K. (2010) 'A-Priori and Post-Hoc Segmentation in the Design of Healthy Eating Campaign', *Journal of Marketing Communications*, 16: 21-45.
- Kearney, M. J. & McElhone, S. (1999) 'Perceived barriers in trying to eat healthier-results of a pan EU consumer attitudinal survey' *British Journal of Nutrition*, 81: (suppl.2): s133-s137.
- Kiefer, I. Rathmanner, T. & Kunze, M. (2005) 'Eating and Dieting Differences in Men and Women', *The Journal of Men's Health & Gender*, 2: 194-201.
- Kline, P. (1999) *The Handbook of Psychological Testing*. London: Routledge.
- Kotler, P., Brown, L., Adam, S., Burton, S. & Armstrong, G. (2007) *Marketing*. Frenchs Forest: Pearson.
- Kotler, P. and Keller, K. L. (2009). *Marketing management* (13th ed.). Upper Saddle River, NJ: Prentice Hall.
- Lanza, S. T., Collins, L. M., Lemmon, D.R. & Schafer, J. L. (2007) 'Proc Lca: A Sas Procedure for Latent Class Analysis', *Structural Equation Modeling*, 14: 671-94.
- Lefebvre, R. C. (2000) 'Theories and Models in Social Marketing' in *Handbook of Marketing and Society*, ed. P. N. Bloom and G. T. Gundlach. Newbury Park, CA: Sage.
- Linde, J. A., Rothman, A. J., Baldwin, A. S. & Jeffery, R. W. (2006) 'The impact of self-efficacy on behaviour change and weight change among overweight participants in a weight loss trial', *Health Psychology*, 25: 282.
- Lowe, M. R. & Butryn, M. L. (2007) 'Hedonic hunger: A new dimension of appetite?' *Physiology & Behavior*, 91: 432-439.
- Lowe, M. R., Butryn, M. L., Didie, E. R., Annunziato, R. A., Thomas, J. G., Crerand, C. E., Ochner, C. N., Coletta, M. C., Bellace, D. & Wallaert, M. (2009) 'The Power of Food Scale. A new measure of the psychological influence of the food environment', *Appetite*, 53: 114-118.
- Luca, N. R. and Suggs, S. (2013) 'Theory and Model Use in Social Marketing Health Interventions', *Journal of Health Communication*, 18: 20-40.

- Lustig, Robert H., Laura A. Schmidt, and Claire D. Brindis (2012) 'Public Health: The Toxic Truth About Sugar', *Nature*, 482: 27-29.
- Luszczynska, A. & Schwarzer, R. (2003) 'Planning and self-efficacy in the adoption and maintenance of breast self-examination: a longitudinal study on self-regulatory cognitions', *Psychology & Health*, 18: 93-108.
- Macht, M. (2008) 'How emotions affect eating: A five-way model', *Appetite*, 50: 1-11.
- Malik, V. S., M. B. Schulze, and F. B. Hu (2006) 'Intake of Sugar-Sweetened Beverages and Weight Gain: A Systematic Review', *Am J Clin Nutr*, 84: 274-88.
- Naughton, P., McCarthy, M. and McCarthy, S., (2015) 'Acting to self-regulate unhealthy eating habits. An investigation into the effects of habit, hedonic hunger and self-regulation on sugar consumption from confectionery foods', *Food Quality and Preference*, 46: 173-183.
- Nothwehr, F. & Yang, J. (2006) 'Goal setting frequency and the use of behavioral strategies related to diet and physical activity', *Health Education Research*, 22: 532-538.
- O'Hare, A., Whelan, C. T., & Commins, P. (1991) 'The development of an Irish census-based social class scale', *Economic and Social Review*, 22: 135.
- Osch, L., Reubsæet, A., Lechner, L., Beenackers, M., Candel, M. & Vries, H. (2010) 'Planning health behaviour change: Comparing the behavioural influence of two types of self-regulatory planning', *British Journal of Health Psychology*, 15: 133-149.
- Otis, N. & Pelletier, L. G. (2008) 'Women's regulation styles for eating behaviours and outcomes: The mediating role of approach and avoidance food planning', *Motivation & Emotion*, 32: 55-67.
- Paisley, C. M. & Sparks, P. (1998) 'Expectations of reducing fat intake: The role of perceived need within the theory of planned behaviour', *Psychology & Health*, 13: 341 – 353.
- Papies, E. K., Stroebe, W., & Aarts, H. (2007) 'Pleasure in the mind: Restrained eating and spontaneous hedonic thoughts about food', *Journal of Experimental Social Psychology*, 43, 810-817.
- Parmenter, K., Waller, J. & Wardle, J. (2000) 'Demographic variation in nutrition knowledge in England', *Health Education Research*, 15(2): 163-174.
- Payne, N., Jones, F. & Harris, P., R. (2004) 'The role of perceived need within the theory of planned behaviour: A comparison of exercise and healthy eating', *British Journal of Health Psychology*, 9: 489-504.
- Peattie, K. & Peattie, S. (2009), 'Social Marketing: A Pathway to Consumption Reduction?,' *Journal of Business Research*, 62: 260-68.
- PHE (2014), A Quick Guide to the Governments Healthy Eating Recommendations.
- Public Health England. Available at:
[https://www.stmartins.academy/admin/userfiles/key%20information/Student%20wellbeing%20/A quick guide to govt healthy eating.pdf](https://www.stmartins.academy/admin/userfiles/key%20information/Student%20wellbeing%20/A%20quick%20guide%20to%20govt%20healthy%20eating.pdf)

PHE (2016) The National Diet and Nutrition Survey. Public Health England.

https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/216485/dh_128556.pdf

Povey, R., Conner, M., Sparks, P., James, R. & Shepherd, R. (1999) 'A critical examination of the application of the Transtheoretical Model's stages of change to dietary behaviours', *Health Education Research*, 14: 641-651.

Povey, R., Conner, M., Sparks, P., James, R. & Shepherd, R. (2000) 'The theory of planned behaviour and healthy eating: examining additive and moderating effects of social influence variables', *Psychology & Health*, 14: 991-1006.

Prochaska, J. Q. & DiClemente, C. C. (1983) 'Stages and processes of self-change of smoking: Toward an integrative model of change', *Journal of Consulting and Clinical Psychology*, 51: 390-395.

Rozin, P. (2007) 'Food choice: an introduction', in L. Frewer & H. von Trijp (ed.) *Understanding Consumers of Food Products*. Cambridge: Woodland Publishing Ltd, 3-30.

Rise, J., Thompson, M. & Verplanken, B. (2003) 'Measuring implementation intentions in the context of the theory of planned behaviour', *Scandinavian Journal of Psychology*, 44: 87-95.

Schnoll, R. & Zimmerman, B. J. (2001) 'Self-regulation training enhances dietary self-efficacy and dietary fiber consumption', *Journal of the American Dietetic Association*, 101: 1006-1011.

Scholz, U., Nagy, G., Gohner, W., Luszczynska, A. & Kliegel, M. (2009) 'Changes in self-regulatory cognitions as predictors of changes in smoking and nutrition behaviour', *Psychology & Health*, 24: 545-561.

Sheeran, P. (2002) 'Intention-Behaviour Relations: A Conceptual and Empirical Review', *European Review of Social Psychology*, 12: 1-36.

Sheeran, P., Milne, S., Webb, T. L. & Gollwitzer, P. M. (2005) 'Implementation intentions and health behaviour', in Conner, M. & Norman, P. (Eds.) *Predicting health behaviour*. 2 ed. Berkshire: McGraw-Hill Education, 276-324.

Slater, M.D., 1996. Theory and method in health audience segmentation. *Journal of health communication*, 1(3), pp.267-284.

Stephens, A., Perkins-Porras, L., Rink, E., Hilton, S. & Cappuccio, F. P. (2004) 'Psychological and social predictors of changes in fruit and vegetable consumption over 12 months following behavioral and nutrition education counseling', *Health Psychology; Health Psychology*, 23: 574.

Stevens, J. P. (2002) *Applied multivariate statistics for the social sciences*. New Jersey: Lawrence Erlbaum Associates.

Swinburn, B. A., Sacks, G. Hall, K.D., McPherson, K. Finegood, D. Moodie, M.L. & Gortmaker, S. L. (2011) 'The Global Obesity Pandemic: Shaped by Global Drivers and Local Environments', *The Lancet*, 378: 804-14.

- Thornton, L.E., Cameron, A.J., McNaughton, S.A., Worsley, A. and Crawford, D.A., 2012. The availability of snack food displays that may trigger impulse purchases in Melbourne supermarkets. *BMC Public Health*, 12(1), p.194.
- Te Morenga, L., Mallard, S. & Mann, J. (2013) 'Dietary Sugars and Body Weight: Systematic Review and Meta-Analyses of Randomised Controlled Trials and Cohort Studies', *BMJ*, 346.
- The HHS and USDA, (2015) 2015-2020 Dietary Guidelines for Americans. United States Department of Agriculture and the US Department of Health and Human Services
- Vartanian, L. R., Schwartz, M. B., & Brownell, K. D. (2007) 'Effects of Soft Drink Consumption on Nutrition and Health: A Systematic Review and Meta-Analysis', *American Journal of Public Health*, 97: 667-75.
- Verplanken, B. & Aarts, H. (1999) 'Habit, Attitude, and Planned Behaviour: Is Habit an Empty Construct or an Interesting Case of Goal-directed Automaticity?' *European Review of Social Psychology*, 10: 101 - 134.
- Verplanken, B. & Faes, S. (1999) 'Good intentions, bad habits, and effects of forming implementation intentions on healthy eating', *European Journal of Social Psychology*, 29: 591-604.
- Verplanken, B. & Orbell, S. (2003) 'Reflections on past behaviour: a self-report index of habit strength', *Journal of Applied Social Psychology*, 33: 1313-1330.
- Vyncke, P. (2002) 'Lifestyle Segmentation', *European Journal of Communication*, 17: 445.
- Wardle, J. & Steptoe, A. (2003) 'Socioeconomic Differences in Attitudes and Beliefs About Healthy Lifestyles', *Journal of Epidemiology and Community Health*, 57: 440-443.
- Wardle, J., Haase, M. A., Steptoe, A., Nillapun, M., Jonwutiwes, K. and Bellisle, F. (2004) 'Gender difference in food choice: the contribution of health beliefs and dieting', *Annals of Behavioural Medicine*, 27 (2): 107-116.
- Wedel, M. & Kamakura, W. (2000) *Market Segmentation: Conceptual and Methodological Foundations*. Boston: Springer.
- Weinstein, A. (2004) *Handbook of market segmentation: Strategic targeting for business and technology firms*. Chicago Routledge.
- WHO (2015), "Sugars Intake for Adults and Children Guideline," Geneva.
- Zandstra, E. H., den Hoed, W., van der Meer, N. & van der Maas, A. (2010) 'Improving compliance to meal-replacement food regimens: Forming implementation intentions (conscious IF-THEN plans) increases compliance', *Appetite*, 55(3): 666-670.