

**UCC Library and UCC researchers have made this item openly available.
Please [let us know](#) how this has helped you. Thanks!**

Title	Do changing levels of maternal exercise during pregnancy affect neonatal adiposity? Secondary analysis of the babies after SCOPE: evaluating the longitudinal impact using neurological and nutritional endpoints (BASELINE) birth cohort (Cork, Ireland)
Author(s)	Norris, Tom; McCarthy, Fergus P.; Khashan, Ali S.; Murray, Deirdre M.; Kiely, Mairead E.; Hourihane, Jonathan O'B.; Baker, Philip N.; Kenny, Louise C.
Publication date	2017
Original citation	Norris, T., McCarthy, F. P., Khashan, A. S., Murray, D. M., Kiely, M., Hourihane, J. O. B., Baker, P. N. and Kenny, L. C. (2017) 'Do changing levels of maternal exercise during pregnancy affect neonatal adiposity? Secondary analysis of the babies after SCOPE: evaluating the longitudinal impact using neurological and nutritional endpoints (BASELINE) birth cohort (Cork, Ireland)', <i>BMJ Open</i> , 7(11), e017987 (100pp). doi: 10.1136/bmjopen-2017-017987
Type of publication	Article (peer-reviewed)
Link to publisher's version	http://bmjopen.bmj.com/content/7/11/e017987 http://dx.doi.org/10.1136/bmjopen-2017-017987 Access to the full text of the published version may require a subscription.
Rights	© 2017, the Authors. This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial http://creativecommons.org/licenses/by-nc/4.0/
Item downloaded from	http://hdl.handle.net/10468/5407

Downloaded on 2022-01-20T19:35:09Z

Supplementary table 1: Descriptive results for those vs those without PEAPOD measurements

	With PEAPOD (n=1258)	Without PEAPOD (n=513)	Difference (Cork_with – Cork_without) (95% CI)
Sex of infant (% (n) female)	48.57 (611)	50.68 (260)	-2.11 (-7.25; 3.02)*
Birth weight (g) (median;IQR)	3500 (3180;3800)	3380 (3020;3730)	130 (80;190)**
Gestational age (weeks) (median;IQR)	40.29 (39.43;41)	40 (38.71;40.86)	0.29 (0.15;0.43)**
Maternal age (years) (mean:SD)	29.95 (4.44)	29.88 (4.61)	0.07 (-0.39;0.53)**
Maternal BMI (kg/m ²) (median:IQR)	23.9 (22;26.9)	24 (22;26.9)	-0.1 (-0.30;0.50)**
Maternal Socioeconomic status (median;IQR)	45 (29;51)	45 (29;50)	0 (-1;0)**
Maternal schooling (years) (median;IQR)	13 (13;14)	13 (13;14)	0 (0;0)**
Household income (<€21 000) (yes)(% (n))	8.47 (106)	8.22 (42)	0.25 (-2.58;3.10)*
Maternal unemployment (yes) (% (n))	5.56 (70)	3.51 (18)	2.06 (0.02;4.09)*
1 st trimester smoking (yes) (% (n))	9.30 (117)	11.70 (60)	-2.40 (-5.60;0.82)*
1 st trimester alcohol intake (units/week (median;IQR)	3 (0.6;7)	2.8 (0.62;5)	0 (-0.5;0)**

*tests for proportions for categorical variables (% difference); **t-tests/Wilcoxon rank-sum for continuous variables (difference in means/medians)

Supplementary table 2: Descriptive results for those with vs those without complete data

	Complete data group (n=1200)	Missing data group (n=58)	Difference (Complete – incomplete) (95% CI)
Sex of infant (% (n) female)	49.00 (588)	39.66 (23)	9.34 $-(3.56;22.25)^*$
Birth weight (g) (mean;SD)	3510 (465)	3322 (518)	188 (64;311)**
Gestational age (weeks) (median;IQR)	40.29 (39.43;41.00)	40.57 (39.29;42.00)	-0.43 $(-0.86;0.15)^{**}$
Maternal age (years) (mean;SD)	29.98 (4.44)	29.13 (4.54)	0.85 $(-0.33;2.02)^{**}$
Maternal BMI (kg/m ²) (median;IQR)	23.9 (22.00;26.8)	24.7 (22.00;29.00)	-0.70 $(-1.90;0.40)^{**}$
Maternal Socioeconomic status (median;IQR)	45 (29;51.00)	44.5 (29;50.00)	0 $(-3.00;4.00)^{**}$
Maternal schooling (years) (median;IQR)	13 (13;14)	13 (13;14)	0 (0;0)**
Household income <€21 000 (yes) (% (n))	8.30 (99)	12.07 (7)	-3.77 $(-12.30; 4.76)^*$
Maternal unemployment (yes) (% (n))	5.50 (66)	6.90 (4)	-1.40 $(-8.04;5.25)^*$
1 st trimester smoking (yes) (% (n))	9.08 (109)	13.79 (8)	-4.71 $(-13.73;4.31)^*$
1 st trimester alcohol intake (units/week) (median;IQR)	3 (0.6;7)	3 (0.6;10)	-0.10 $(-1.5;0.67)^{**}$

*tests for proportions for categorical variables (% difference); **t-tests/Wilcoxon rank-sum for continuous variables (difference in means/medians)

