

**UCC Library and UCC researchers have made this item openly available.
Please [let us know](#) how this has helped you. Thanks!**

| | |
|------------------------------------|---|
| Title | The natural history of degenerative cervical myelopathy and the rate of hospitalization following spinal cord injury: an updated systematic review |
| Author(s) | Tetreault, Lindsay A.; Karadimas, Spyridon; Wilson, Jefferson R.; Arnold, Paul M.; Kurpad, Shekar; Dettori, Joseph R.; Fehlings, Michael G. |
| Publication date | 2017 |
| Original citation | Tetreault, L. A., Karadimas, S., Wilson, J. R., Arnold, P. M., Kurpad, S., Dettori, J. R. and Fehlings, M. G. (2017) 'The natural history of degenerative cervical myelopathy and the rate of hospitalization following spinal cord injury: an updated systematic review', <i>Global Spine Journal</i> , 7(3S), pp. 28-34. doi: 10.1177/2192568217700396 |
| Type of publication | Article (peer-reviewed) |
| Link to publisher's version | http://journals.sagepub.com/doi/10.1177/2192568217700396 http://dx.doi.org/10.1177/2192568217700396 Access to the full text of the published version may require a subscription. |
| Rights | © 2017, the Authors. This article is distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 License (http://www.creativecommons.org/licenses/by-nc-nd/4.0/) which permits non-commercial use, reproduction and distribution of the work as published without adaptation or alteration, without further permission provided the original work is attributed as specified on the SAGE and Open Access pages (https://us.sagepub.com/en-us/nam/open-access-at-sage). http://www.creativecommons.org/licenses/by-nc-nd/4.0/ |
| Item downloaded from | http://hdl.handle.net/10468/5618 |

Downloaded on 2021-12-07T23:42:55Z

The Natural History of Degenerative Cervical Myelopathy and the Rate of Hospitalization Following Spinal Cord Injury: An Updated Systematic Review

Global Spine Journal
2017, Vol. 7(3S) 28S-34S
© The Author(s) 2017
Reprints and permission:
sagepub.com/journalsPermissions.nav
DOI: 10.1177/2192568217700396
journals.sagepub.com/home/gsj


Lindsay A. Tetreault, PhD^{1,2}, Spyridon Karadimas, MD¹,
Jefferson R. Wilson, MD, PhD³, Paul M. Arnold, MD^{4,5}, Shekar Kurpad, MD, PhD⁶,
Joseph R. Dettori, PhD, MPH⁷, and Michael G. Fehlings, MD, PhD, FRCSC, FACS^{1,3}

Abstract

Study Method: Systematic review (update).

Objective: Degenerative cervical myelopathy (DCM) is a degenerative spine disease and the most common cause of spinal cord dysfunction in adults worldwide. The objective of this study is to determine the natural history of DCM by updating the systematic review by Karadimas et al. The specific aims of this review were (1) to describe the natural history of DCM and (2) to determine potential risk factors of disease progression.

Method: An updated search based on a previous protocol was conducted in PubMed and the Cochrane Collaboration library for studies published between November 2012 and February 15, 2015.

Results: The updated search yielded 3 additional citations that met inclusion criteria and reported the incidence of spinal cord injury and severe disability in patients with DCM. Based on 2 retrospective cohort studies, the incidence rate of hospitalization for spinal cord injury is 13.9 per 1000 person-years in patients with cervical spondylotic myelopathy and 4.8 per 1000 person-years in patients with myelopathy secondary to ossification of the posterior longitudinal ligament (OPLL). In a third small prospective study, the risk of being wheelchair bound or bedridden was 66.7% in DCM patients with OPLL.

Conclusion: The overall level of evidence for these estimated rates of hospitalization following spinal cord injury was rated as low.

Keywords

degenerative cervical myelopathy, cervical spondylotic myelopathy, systematic review

Introduction

Degenerative cervical myelopathy (DCM) is a degenerative spine disease and the most common cause of spinal cord dysfunction in adults worldwide.¹ The term DCM encompasses cord compression secondary to osteoarthritic changes to the spine, disc degeneration, ligamentous aberrations, and progressive kyphosis. Patients with DCM may present with a wide range of neurological signs and symptoms, including pain, lower limb spasticity, decreased hand dexterity, hyperreflexia, and sphincter disturbance.

The pattern of progression in DCM is not well defined. Early reports of the natural history suggest that DCM is a relatively benign disorder and that patients are more likely to

¹ Toronto Western Hospital, University Health Network, Toronto, Ontario, Canada

² University College Cork, Cork, Ireland

³ University of Toronto, Toronto, Ontario, Canada

⁴ University of Kansas Medical Center, Kansas City, KS, USA

⁵ The University of Kansas, Kansas City, KS, USA

⁶ Medical College of Wisconsin, Milwaukee, WI, USA

⁷ Spectrum Research, Inc, Tacoma, WA, USA

Corresponding Author:

Michael G. Fehlings, MD, PhD, FRCSC, FACS, Division of Neurosurgery, Toronto Western Hospital, University Health Network, 399 Bathurst Street (SCI-CRU, 11th Floor McLaughlin Pavilion), Toronto, Ontario M5T 2S8, Canada.

Email: michael.fehlings@uhn.ca






Figure 1. Results of updated and originally published literature searches. KQ = key question.

remain stable over time than to deteriorate.² In the most recent systematic review of the literature, there was moderate evidence that 20% to 62% of patients with symptomatic myelopathy will decline by at least 1 point on the Japanese Orthopaedic Association scale if not managed surgically.³ The objective of this study was to update the systematic review by Karadimas et al³ that investigated (1) the natural history of DCM and (2) the potential risk factors of disease progression.

Materials and Methods

Electronic Literature Search

An updated search based on a previous protocol⁴ was conducted in PubMed and the Cochrane Collaboration library for studies published between November 2012 and February 15, 2015. Inclusion and exclusion criteria for the search were previously published, as well as methods for data abstraction, data analysis, evaluation of study quality, and assessment of the overall strength of evidence.

Results

Study Selection

The updated electronic search yielded 122 new citations (Figure 1). An additional 40 citations were identified through

Table 1. Excluded Studies and Reasons for Exclusion.

| Author (Year) | Reasons for Exclusion |
|-------------------|-------------------------------------|
| Kalb et al (2011) | Surgery study |
| Matsunaga (2008) | Wrong population: Asymptomatic OPLL |

Abbreviation: OPLL, ossification of the posterior longitudinal ligament.

directed manual search. One hundred and fifty-seven studies were excluded following title and abstract review, and 5 studies were further investigated. Following full text review, a single study was excluded because all patients underwent surgery for DCM, and another for including nonmyelopathic patients with ossification of the posterior longitudinal ligament (OPLL; Table 1).⁵ Three other studies presented new information relevant to key question 1 and reported the incidence of spinal cord injury and severe disability in patients with DCM.⁶⁻⁸

Two retrospective cohort studies used the National Health Insurance Research Database, which contains records for 23 million administered insurants in Taiwan (approximately 99% of the entire population).^{7,8} The first study included 14 140 patients hospitalized for cervical spondylotic myelopathy (CSM) with at least 1 year of follow-up (Table 2).⁸ The second study consisted of 5604 patients with myelopathy secondary to OPLL and at least 3 years of follow-up.⁷ A third prospective

Table 2. Characteristics of New Studies Addressing the Natural History of DCM.

| Author (Year)/Study Design | Patient Characteristics | Mean Follow-up; % Follow-up | Inclusion Criteria |
|---|---|--|---|
| Wu et al (2013) ⁸ / Retrospective cohort | N = 14 140; mean age = NR; % male NR | ≥ 1 year; % NR | Subjects hospitalized and discharged with the diagnostic ICD-9 code for CSM (721.1) (National Health Research Institute of Taiwan) |
| Wu et al (2012) ⁷ / Retrospective cohort | N = 5604; mean age = 60.35 ± 14 years; 70% male | ≥ 3 years; % NR | Subjects hospitalized within the study period with a first time discharge summary containing the diagnostic ICD-9 code for OPLL (723.7x) (National Health Research Institute of Taiwan) Patients hospitalized for OPLL who have not received spinal intervention within the previous 6 months (National Health Research Institute of Taiwan) |
| Matsunaga et al (2004) ⁶ / Prospective cohort | N = 36 ^a ; mean age = 61.8 years; 59% male | 17.6 years (range = 10-30 years); % NR | Patients with DCM from OPLL |

Abbreviations: DCM, degenerative cervical myelopathy; ICD-9, International Classification of Diseases, Ninth Edition; CSM, cervical spondylotic myelopathy; NR, not reported; OPLL, ossification of the posterior longitudinal ligament.

^aN = 450 in study; 36 patients were treated conservatively for myelopathy symptoms.

Table 3. Incidence Rate or Risk of Spinal Cord Injury and Disability in Patients Not Treated Surgically.

| Outcome | Study | Risk of Bias | Diagnosis | N | Person-Years or Number of Persons | Incidence Rate or Risk ^a (95% CI) |
|-------------------------|-------------------------------|-----------------|-------------------|-----|-----------------------------------|--|
| SCI ^b | Wu (2013) | Moderately low | CSM | 122 | 8776.7 | 13.9 (11.6-16.6) |
| | Wu (2012) | Moderately low | OPLL ^c | 7 | 1455.2 | 4.8 (2.3-10.1) |
| Disability ^d | Wu (2012) ⁷ | Moderately low | OPLL ^c | 5 | 1463.6 | 3.4 (1.5-8.0) |
| | Matsunaga (2004) ⁶ | Moderately high | OPLL ^c | 24 | 36 | 66.7% |

Abbreviations: CI, confidence interval; SCI, spinal cord injury; CSM, cervical spondylotic myelopathy; OPLL, ossification of the posterior longitudinal ligament; DCM, degenerative cervical myelopathy.

^aRate is per 1000 person-years; risk = percentage.

^bDefined as hospitalizations for SCI.

^cDCM secondary to OPLL

^dWu (2012) defined disability as severe neurological deficits caused by SCI such as paraplegia, tetraplegia, and incontinence; Matsunaga (2004) defined disability as becoming wheelchair bound or bedridden.

cohort study reported outcomes on 450 patients with myelopathy secondary to OPLL; however, only 36 were managed conservatively after refusing surgical treatment.

Hospitalization for Spinal Cord Injury

Based on 2 retrospective cohort studies, the incidence rate of hospitalization for spinal cord injury was 13.9 per 1000 person-years (95% confidence interval [CI] = 11.6-16.6) in patients diagnosed with CSM⁸ and 4.8 per 1000 person-years (95% CI = 2.3-10.1) in patients with myelopathy secondary to OPLL⁷ (Table 3). The rate of hospitalization for spinal cord injury in patients with DCM from OPLL was significantly higher than the rate observed in a healthy population (0.18 per 1000 person-years; hazard ratio = 32.2; 95% CI = 10.4-99.0; $P < .001$).⁷ These studies both had moderately low risk of bias (Table 4).

Disability

One large retrospective study evaluated the incidence of severe neurological deficits from spinal cord injury in patients with

Table 4. Class of Evidence for Prognostic Studies.

| Methodological Principle | Wu (2013) ⁸ | Wu (2012) ⁷ | Matsunaga (2004) ⁶ |
|---|------------------------|------------------------|-------------------------------|
| Study design | | | |
| Prospective cohort study | | | ✓ |
| Retrospective cohort study | ✓ | ✓ | |
| Case-control study | | | |
| Case series | | | |
| Patients at similar point in the course of their disease or treatment | ✓ | ✓ | |
| Patients followed long enough for outcome to occur | ✓ | ✓ | ✓ |
| Complete follow-up of ≥80% | ✓ | ✓ | |
| Controlling for extraneous prognostic factors ^a | ✓ | ✓ | |

^aAuthors must summarize baseline characteristics, and control for those that differ between treatment groups.

CSM, including paraplegia, tetraplegia, and incontinence; the reported rate was 3.4 per 1000 person-years.⁷ A second small prospective study reported that the risk of being wheelchair bound or bedridden was 66.7% (24/36) in patients with DCM

Table 5. Evidence Summary.

| | Strength of Evidence | Conclusions/Comments | Baseline ^a | Upgrade (Levels) ^b | Downgrade (Levels) ^c |
|---|----------------------|--|-----------------------|-------------------------------|---|
| What is the natural history of CSM? | | | | | |
| <i>Neurological outcome</i> | | | | | |
| • JOA change compared with baseline | Moderate | • Although mean scores tend to remain constant, there is moderate evidence (2 small prospective ^{9,10} and 4 small retrospective ¹⁰⁻¹³ observational studies) that 20% to 62% of patients will deteriorate (at least 1 point on the JOA) 3-6 years after initial assessment. Proportions vary based on definition of deterioration. | High | | Risk of bias (1) |
| • Nurick Grade change compared with baseline | Very Low | • There is very low evidence from one small retrospective observational study (N = 76) ¹⁴ that a majority of DCM patients will not experience a change on the Nurick over time with nonoperative treatment. Sixty-seven percent of patients were stable, 20% improved, and 13% deteriorated after 8 years. | Low | | Inconsistent (1) |
| • Spinal cord injury | Low | • There is low-level evidence that the rate of hospitalization for spinal cord injury is 13.9 per 1000 person-years in patients with CSM. ⁸ The rate is 4.8 per 1000 person-years in patients with DCM from OPLL and 0.18 per 1000 person-years in a healthy population (HR = 32.2; 95% CI = 10.4-99.0). ⁷ | Low | | |
| • Disability | Low | • There is low-level evidence that the rate of severe disability is 3.4 per 1000 person-years in patients with DCM secondary to OPLL. ⁷ | Low | | |
| • Conversion to surgery | Very Low | • There is very low evidence (2 small prospective ^{10,15,16} and 4 small retrospective observational studies) ^{11,17-19} that the proportion of patients undergoing surgery following worsening of symptoms increases over time. The proportion of patients converting to surgery ranges from 4% to 40% over 3 to 7 years, respectively. | Low | | Imprecise (1) |
| <i>Functional outcome</i> | | | | | |
| • Activities of daily living | Moderate | • There is moderate evidence (2 small prospective studies, N = 31 ²⁰ and N = 33 ²¹) that patients with DCM worsen in performing activities of daily living (ADL) with nonoperative treatment. One study reported 6%, 21%, 28%, and 56% worsening of ADL from baseline values at 1-, 2-, 3-, and 10-years follow-up, respectively. | High | | Risk of bias (1) |
| • Timed 10-meter walk | Very Low | • There is very low evidence (one small prospective study, N = 33 ²¹) that there is no significant difference in 10-meter walking test times between baseline and 1-, 2-, 3-, and 10-years following conservative treatment. | High | | Risk of bias (1); Imprecise (1); Inconsistent (1) |
| • Overall functional status | Very Low | • There is very low evidence (one small prospective observational study, N = 31 ²⁰) that the overall functional status improves over time in patients treated conservatively for DCM. | Low | | Inconsistent (1) |
| Are there risk factors that affect the progression of DCM? | | | | | |
| <i>Neurological outcome</i> | | | | | |
| <i>Demographic characteristics</i> | | | | | |
| • Age | Very Low | • There is very low evidence to support the association between age at diagnosis and neurological deterioration based on the JOA. One prospective study reported no association using multivariate analysis; ¹³ one prospective study reported that older age (mean 58 years) before treatment was a positive predictor for neurological improvement ($P < .05$); ⁹ and one retrospective study reported that younger age (<52 years) was a positive predictor for neurological improvement using univariate analysis. ¹⁷ | Low | | Inconsistent (1); Imprecise (1) |

(continued)

Table 5. (continued)

| | Strength of Evidence | Conclusions/Comments | Baseline ^a | Upgrade (Levels) ^b | Downgrade (Levels) ^c |
|--|----------------------|--|-----------------------|-------------------------------|---|
| • Sex | Very Low | • There is very low evidence to support the association between sex and neurological progression of myelopathy on the JOA. One prospective study reported no association using multivariate analysis ¹⁰ and one retrospective study indicated that female sex was associated with a progressively worse neurological condition ($P < .05$). ²⁰ | Low | | Inconsistent (I); Risk of bias (I) |
| • Height | Very Low | • There is very low evidence (one prospective study) that lower body height (mean 170 cm) is a positive predictor of JOA improvement ($P < .05$). ⁹ | High | | Imprecise (I); Inconsistent (I) |
| Radiographic characteristics | | | | | |
| • Circumferential spinal cord compression | Low | • There is low evidence (one prospective study using multivariate analysis) that circumferential spinal cord compression (compared with only partial cord compression) is associated with neurological deterioration (JOA) (adjusted OR = 26.6; 95% CI = 1.7-421.5). ¹⁰ | High | Large effect (I) | Risk of bias (I); Imprecise (I); Inconsistent (I) |
| • Transverse area of the spinal cord; Pavlov's Index | Very Low | • There is very low evidence (one prospective observational study) that a larger transverse area of the spinal cord (mean 76 mm ²) ($P < .05$) and a higher Pavlov Index (mean 0.9) ($P < .05$) are associated with improved neurological status (JOA). ⁹ | High | | Risk of bias (I); Imprecise (I); Inconsistent (I) |
| • Other radiological factors | Very Low | • There is very low evidence (one prospective study using multivariate analysis) that there is no significant association between developmental or dynamic canal factors, high T2WI signal intensity and neurological deterioration (JOA). ¹⁰ | High | | Risk of bias (I); Imprecise (I); Inconsistent (I) |
| Clinical characteristics | | | | | |
| • Initial level of disability | Very Low | • There is very low evidence (one prospective ⁹ and one retrospective ¹⁷ observational study) that milder disability before treatment is associated with greater neurological improvement (JOA) ($P < .05$). | Low | | Imprecise (I) |
| • Duration of disease | Very Low | • There is very low evidence (2 retrospective observational studies) that a shorter duration of symptoms is associated with neurological improvement (JOA) ($P = .001$). ^{13,17} | Low | | Inconsistent (I) |
| • Range of motion | Very Low | • There is very low evidence (one retrospective study) that greater neck range of motion (ROM) ($P < .05$), greater head ROM ($P < .01$), and difference between total head and neck ROM ($P < .01$) are associated with progressively worse neurological condition (JOA). ¹⁴ | Low | | Inconsistent (I); Imprecise (I) |
| Conversion to surgery | | | | | |
| Demographic characteristics | | | | | |
| • Age; sex | Very Low | • There is very low evidence (one retrospective study using multivariate analysis) that there is no association between age ≥ 60 years or sex and conversion to surgery. ¹⁹ | Low | | Inconsistent (I); Imprecise (I) |
| Radiographic characteristics | | | | | |
| • Cervical range of motion | Very Low | • There is very low evidence (one small retrospective study, N = 45 using multivariate analysis ¹⁹) that there is an association between increased risk of surgery and the following factors: | Low | Large effect (I) | Imprecise (I); Inconsistent (I) |
| • Segmental lordotic angle | | • Total cervical range of motion ($\geq 50^\circ$) (adjusted HR = 3.3; 95% CI = 1.03-10.25) | | | |
| • Local slip | | • Segmental lordotic angle ($< 0^\circ$) (adjusted HR = 4.5; 95% CI = 1.59-12.8) | | | |
| | | • Presence of a local slip (adjusted HR = 4.7; 95% CI = 1.67-13.0) | | | |

(continued)

Table 5. (continued)

| | Strength of Evidence | Conclusions/Comments | Baseline ^a | Upgrade (Levels) ^b | Downgrade (Levels) ^c |
|------------------------------|----------------------|--|-----------------------|-------------------------------|---------------------------------|
| • Other radiographic factors | Very Low | • There is very low evidence (one small retrospective study, N = 45 using multivariate analysis) that there is no association between increased risk of surgery and C2-7 alignment (<0°), spinal cord diameter (<50%), presence of developmental canal stenosis, and segmental range of motion (≥10°). ¹⁹ | Low | | Imprecise (1); Inconsistent (1) |

Abbreviations: CSM, cervical spondylotic myelopathy; JOA, Japanese Orthopaedic Association; DCM, degenerative cervical myelopathy; OPLL, ossification of the posterior longitudinal ligament; HR, hazard ratio; CI, confidence interval; OR, odds ratio.

^aBaseline quality: High = majority of articles low/moderately low risk of bias; Low = majority of articles moderately high/high risk of bias.

^bUpgrade: Large magnitude of effect (1 or 2 levels); dose response gradient (1 level); plausible confounding decreases magnitude of effect (1 level).

^cDowngrade: Inconsistency of results (1 or 2 levels); indirectness of evidence (1 or 2 levels); imprecision of effect estimates (1 or 2 levels); risk of bias (1 or 2 levels); failure to specify subgroup analysis a priori (1 level); reporting bias (1 level).

secondary to OPLL (Table 3).⁶ This study had moderately high risk of bias (Table 4).

Evidence Summary

The rate of hospitalization due to spinal cord injury was 4.8 per 1000 person-years in patients with DCM secondary to OPLL and 13.9 per 1000 person-years in patients with CSM. The rate of severe disability in DCM patients with OPLL was 3.4 per 1000 person-years. The strength of evidence for these estimates was Low (Table 5).

Conclusions

The results of this update indicate that the presence of OPLL or CSM may increase a patient's risk of severe disability and hospitalization for spinal cord injury. Although these findings are unlikely to directly influence management strategies, patients should be counseled of the possibility of spinal cord injury when discussing the benefits and risks of various treatment options.

Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This research was supported by AOSpine and also received funding from the Cervical Spine Research Society (CSRS). Dr Fehlings wishes to acknowledge support from the Gerald and Tootsie Halbert Chair in Neural Repair and Regeneration and the DeZwirek Family Foundation. Dr Tetreault acknowledges support from a Krembil Postdoctoral Fellowship Award.

References

- Nouri A, Tetreault L, Singh A, Karadimas SK, Fehlings MG. Degenerative cervical myelopathy: epidemiology, genetics, and pathogenesis. *Spine (Phila Pa 1976)*. 2015;40:E675-E693. doi:10.1097/BRS.0000000000000913.
- Tetreault L, Goldstein CL, Arnold P, et al. Degenerative cervical myelopathy: a spectrum of related disorders affecting the aging spine. *Neurosurgery*. 2015;77(suppl 4):S51-S67. doi:10.1227/NEU.0000000000000951.
- Karadimas S, Erwin W, Ely C, Dettori JR, Fehlings MG. Pathophysiology and natural history of cervical spondylotic myelopathy. *Spine (Phila Pa 1976)*. 2013;38:36. doi:10.1097/BRS.0b013e3182a7f2c3.
- Rhee JM, Shamji MF, Erwin WM, et al. Nonoperative management of cervical myelopathy: a systematic review. *Spine (Phila Pa 1976)*. 2013;38(22 suppl):S55-S67. doi:10.1097/BRS.0b013e3182a7f41d.
- Kalb S, Martirosyan NL, Perez-Orribo L, Kalani MY, Theodore N. Analysis of demographics, risk factors, clinical presentation, and surgical treatment modalities for the ossified posterior longitudinal ligament. *Neurosurg Focus*. 2011;30:E11. doi:10.3171/2010.12.focus10265.
- Matsunaga S, Sakou T, Taketomi E, Komiya S. Clinical course of patients with ossification of the posterior longitudinal ligament: a minimum 10-year cohort study. *J Neurosurg*. 2004;100:245-248.
- Wu JC, Chen YC, Liu L, et al. Conservatively treated ossification of the posterior longitudinal ligament increases the risk of spinal cord injury: a nationwide cohort study. *J Neurotrauma*. 2012;29:462-468. doi:10.1089/neu.2011.2095.
- Wu JC, Ko CC, Yen YS, et al. Epidemiology of cervical spondylotic myelopathy and its risk of causing spinal cord injury: a national cohort study. *Neurosurg Focus*. 2013;35:E10. doi:10.3171/2013.4.FOCUS13122.
- Kadanka Z, Mares M, Bednarik J, et al. Predictive factors for mild forms of spondylotic cervical myelopathy treated conservatively or surgically. *Eur J Neurol*. 2005;12(1):16-24.
- Shimomura T, Sumi M, Nishida K, et al. Prognostic factors for deterioration of patients with cervical spondylotic myelopathy after nonsurgical treatment. *Spine (Phila Pa 1976)*. 2007;32(22):2474-2479.
- Matsumoto M, Chiba K, Ishikawa M, Maruiwa H, Fujimura Y, Toyama Y. Relationships between outcomes of conservative treatment and magnetic resonance imaging findings in patients with mild cervical myelopathy caused by soft disc herniations. *Spine (Phila Pa 1976)*. 2001;26(14):1592-1598.
- Matsumoto M, Toyama Y, Ishikawa M, Chiba K, Suzuki N, Fujimura Y. Increased signal intensity of the spinal cord on magnetic

- resonance images in cervical compressive myelopathy. Does it predict the outcome of conservative treatment? *Spine (Phila Pa 1976)*. 2000;25(6):677-682.
13. Yoshimatsu H, Nagata K, Goto H, et al. Conservative treatment for cervical spondylotic myelopathy. prediction of treatment effects by multivariate analysis. *Spine J*. 2001;1(4):269-273.
 14. Barnes MP, Saunders M. The effect of cervical mobility on the natural history of cervical spondylotic myelopathy. *J Neurol Neurosurg Psychiatry*. 1984;47(1):17-20.
 15. Lees F, Turner JW. Natural History and Prognosis of Cervical Spondylosis. *Br Med J*. 1963;2(5373):1607-1610.
 16. Sumi M, Miyamoto H, Suzuki T, Kaneyama S, Kanatani T, Uno K. Prospective cohort study of mild cervical spondylotic myelopathy without surgical treatment. *J Neurosurg Spine*. 2012;16(1):8-14.
 17. Nakamura K, Kurokawa T, Hoshino Y, Saita K, Takeshita K, Kawaguchi H. Conservative treatment for cervical spondylotic myelopathy: achievement and sustainability of a level of “no disability”. *J Spinal Disord*. 1998;11(2):175-179.
 18. Roberts A. Myelopathy due to cervical spondylosis treated by collar immobilization. *Neurology*. 1966;16:951-954.
 19. Oshima Y, Seichi A, Takeshita K, et al. Natural course and prognostic factors in patients with mild cervical spondylotic myelopathy with increased signal intensity on T2-weighted magnetic resonance imaging. *Spine (Phila Pa 1976)*. 2012;37(22):1909-1913.
 20. Sampath P, Bendebba M, Davis JD, Ducker TB. Outcome of patients treated for cervical myelopathy. A prospective, multi-center study with independent clinical review. *Spine (Phila Pa 1976)*. 2000;25(6):670-676.
 21. Kadanka Z, Mares M, Bednanik J, et al. Approaches to spondylotic cervical myelopathy: conservative versus surgical results in a 3-year follow-up study. *Spine (Phila Pa 1976)*. 2002;27(20):2205-2210;