

**UCC Library and UCC researchers have made this item openly available.  
Please [let us know](#) how this has helped you. Thanks!**

<b>Title</b>	Prospective biomarkers in preterm preeclampsia: a review
<b>Author(s)</b>	McCarthy, Fergus P.; Ryan, Roisin M.; Chappell, Lucy C.
<b>Publication date</b>	2018-03-23
<b>Original citation</b>	McCarthy, F. P., Ryan, R. M. and Chappell, L. C. (2018) 'Prospective biomarkers in preterm preeclampsia: a review', <i>Pregnancy Hypertension</i> , In Press, doi:10.1016/j.preghy.2018.03.010
<b>Type of publication</b>	Article (peer-reviewed)
<b>Link to publisher's version</b>	<a href="http://dx.doi.org/10.1016/j.preghy.2018.03.010">http://dx.doi.org/10.1016/j.preghy.2018.03.010</a> Access to the full text of the published version may require a subscription.
<b>Rights</b>	© 2018 International Society for the Study of Hypertension in Pregnancy. Published by Elsevier B.V. All rights reserved. This manuscript version is made available under the CC-BY-NC-ND 4.0 license. <a href="http://creativecommons.org/licenses/by-nc-nd/4.0/">http://creativecommons.org/licenses/by-nc-nd/4.0/</a>
<b>Embargo information</b>	Access to this article is restricted until 12 months after publication by request of the publisher.
<b>Embargo lift date</b>	2019-03-23
<b>Item downloaded from</b>	<a href="http://hdl.handle.net/10468/5722">http://hdl.handle.net/10468/5722</a>

Downloaded on 2022-12-07T02:44:39Z



**UCC**

University College Cork, Ireland  
Coláiste na hOllscoile Corcaigh

## Accepted Manuscript

Prospective biomarkers in preterm preeclampsia: a review

Fergus P. McCarthy, Roisin M. Ryan, Lucy C. Chappell

PII: S2210-7789(17)30495-6

DOI: <https://doi.org/10.1016/j.preghy.2018.03.010>

Reference: PREGHY 411

To appear in: *Pregnancy Hypertension: An International Journal of Women's Cardiovascular Health*


Received Date: 1 December 2017

Revised Date: 21 March 2018

Accepted Date: 22 March 2018

Please cite this article as: McCarthy, F.P., Ryan, R.M., Chappell, L.C., Prospective biomarkers in preterm preeclampsia: a review, *Pregnancy Hypertension: An International Journal of Women's Cardiovascular Health* (2018), doi: <https://doi.org/10.1016/j.preghy.2018.03.010>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



**Title**

Prospective biomarkers in preterm preeclampsia: a review

**Author's names**

Fergus P. McCarthy, Roisin M. Ryan, Lucy C. Chappell

**Author Details**

Dr Fergus P McCarthy, Consultant Obstetrician and Gynaecologist/ Maternal & Fetal Medicine Subspecialist, Senior Lecturer, The Irish Centre for Fetal and Neonatal Translational Research, University College Cork, Wilton, Ireland.

Honorary Clinical Lecturer, Department of Women and Children's Health, School of Life Course Sciences, King's College London, 10th floor North Wing, St Thomas' Hospital London SE1 7EH

Dr Roisin M Ryan, London School of Hygiene and Tropical Medicine, London, UK

Professor Lucy Chappell, NIHR Research Professor in Obstetrics, King's College London. Honorary Consultant Obstetrician, St Thomas' Hospital, London SE1 7EH, UK.

**Corresponding author**

Dr Fergus McCarthy, address: as above.

Tel: +353830409737 Email: Fergus.mccarthy@ucc.ie

**Competing interests statement**

The authors have no competing interests to declare.

ACCEPTED MANUSCRIPT

**Abstract**

Preterm pre-eclampsia (prior to 37 weeks' gestation) remains a major cause of maternal and fetal morbidity and mortality particularly in low to middle income countries. Much research has focused on first and second trimester predictors of pre-eclampsia with the aim of allowing stratification of antenatal care and trialling of potential preventative and therapeutic agents. However, none have been shown to be of benefit in randomised controlled trials.

In this literature review we critically evaluate predictive and diagnostic tests for preterm pre-eclampsia and discuss their clinical use and potential value in the management of preterm pre-eclampsia. We defined preterm pre-eclampsia as pre-eclampsia occurring prior to 37 weeks' gestation. Substantial progress has been made in the development of predictive screening tests for preterm pre-eclampsia, but further research is needed prior to their introduction and integration into routine clinical practice. The performance of diagnostic tests mainly utilising angiogenic and anti-angiogenic factors for determining time to delivery in later pregnancy currently hold more promise than first trimester predictive tests, possibly reflecting the heterogeneity of pre-eclampsia.

**Keywords**

Hypertension, preterm pre-eclampsia, placental growth factor, pregnancy, screening.

## Introduction

Pre-eclampsia is defined by the International Society for the Study of Hypertension in Pregnancy (ISSHP) as hypertension and the coexistence of one or more of the following new-onset conditions; proteinuria (spot urine protein/creatinine  $>30$  mg/ mmol or  $>300$  mg/day or at least 1 g/L(2+) on dipstick testing); other maternal organ dysfunction including renal insufficiency, liver involvement, neurological complications, haematological complications; or evidence of uteroplacental dysfunction indicated by fetal growth restriction.<sup>1</sup> For the purposes of this review, preterm pre-eclampsia is defined as pre-eclampsia which occurs prior to 37 weeks gestation, though we acknowledge the current debate regarding this binary classification<sup>2 3</sup>.

This hypertensive condition usually occurs during the second half of pregnancy, complicates 2% to 8% of pregnancies and remains a major cause of maternal and perinatal mortality and morbidity worldwide, accounting for 17-24% of all maternal deaths in low income settings.<sup>4 5</sup>

Preterm pre-eclampsia is thought to arise as a result of placental ischaemia which arises from impaired placentation (trophoblast invasion of the maternal uterine spiral arteries), which in turn leads to stimulation of sustained endoplasmic reticulum and

oxidative stress.<sup>6-8</sup> This pathophysiological cascade generates the characteristic systemic symptoms of the maternal disease; it is affected by genetic, behavioural or environmental influences and is mediated by release of factors into the maternal circulation.<sup>9-12</sup> In contrast, term pre-eclampsia is generally not associated with abnormal trophoblast invasion<sup>7</sup> and it is thought that other maternal constitutive factors or the pre-existing susceptibility of the maternal vasculature is responsible for the development of term pre-eclampsia<sup>13</sup>.

There are currently few preventative strategies and no cure other than delivery of the placenta; for women presenting with disease preterm, this often results in iatrogenic delivery to prevent serious maternal or fetal adverse outcomes.

There have been wide ranging investigations into possible biomarkers for early identification of preterm pre-eclampsia<sup>14</sup>. The release of factors from the placenta into the maternal blood stream, or maternal generation of factors may precede clinical symptoms<sup>14</sup>. Therefore, there remains substantial interest in the use of these factors as potential biomarkers for subsequent disease. Initial research focused on angiogenic factors and placental-specific factors, whilst recent advances in the so called omics fields have led to further possible biomarkers<sup>15 16</sup>. Table 1 summarises the biomarkers discussed in this review. These biomarkers, alone or in combination with biophysical and sonographic findings, may allow development of a reliable and valid screening or diagnostic test for pre-eclampsia to enable risk stratification and timely use of

pharmacological interventions such as aspirin (or other novel therapies) to reduce the risk of pre-eclampsia.<sup>17,18</sup>

A difficulty when assessing studies which look at pre-eclampsia is that many do not subclassify pre-eclampsia into early onset and late onset. As noted in Wu et al, this results in a poorly defined phenotype of pre-eclampsia which may contribute to the low predictive value of studies to date.<sup>19</sup> Furthermore, as early and late onset pre-eclampsia are now believed to have pathogenetic mechanisms contributing in varying degrees, it is likely they are characterised by different biomarkers. This highlights the importance of stratifying by pre-eclampsia type in future studies of potential biomarkers.

Many proposed tests for the short and longer-term prediction of pre-eclampsia are currently under investigation, and some are now being offered in clinical practice. In this literature review we will critically evaluate predictive and diagnostic tests for pre-eclampsia and discuss their clinical use and potential value in the management of preterm pre-eclampsia. Studies are described in the text, with Table 1 providing further detail on statistical results.

### **Angiogenic factors**

The placental hypoxia resulting from the impaired trophoblast invasion that occurs in pre-eclampsia results in an imbalance between pro- and anti-angiogenic factors, in particular, the anti-angiogenic soluble fms-like tyrosine kinase-1 (sFlt-1) and the pro-angiogenic vascular endothelial growth factor (VEGF) and placental growth factor (PlGF).<sup>20-23</sup> VEGF induces endothelial proliferation and vascular permeability by



inducing the expression of integrins on endothelial cells that are involved in angiogenic invasion.<sup>24</sup> In the placenta, it is involved not only in vascularisation, but also in trophoblast syncytialisation and proliferation of extravillous trophoblasts (as assessed in primary culture of trophoblasts and a cell line of human extravillous trophoblasts).<sup>25,26</sup>

The anti-angiogenic factor sFlt-1 is a splice variant of the VEGF receptor, Flt-1 that lacks both the transmembrane and cytoplasmic domains of the cell membrane bound receptor. sFlt-1 acts as a soluble receptor that binds PlGF and also binds and deactivates circulating VEGF.<sup>27-30</sup> Measurement of these angiogenic factors has resulted in several clinically commercially available assays.<sup>29-32</sup>

Chappell and colleagues previously investigated the diagnostic accuracy of low plasma PlGF concentration (using the Alere Triage assay) in women presenting with suspected pre-eclampsia between 20 and 35 weeks' gestation (and up to 41 weeks' gestation as a secondary analysis). Of 287 women recruited prior to 35 weeks' gestation, PlGF <5th centile had high sensitivity (96%) and negative predictive value (98%) for pre-eclampsia leading to delivery within 14 days; specificity was lower (55%).<sup>33</sup>

For short term prediction of pre-eclampsia, a sFlt-1:PlGF ratio cut-off of 38 has been shown to have a negative predictive value for the development of pre-eclampsia in the subsequent week of 99.3% (95% CI, 97.9 to 99.9), with 80.0% sensitivity (95% CI, 51.9 to 95.7) and 78.3% specificity (95% CI, 74.6 to 81.7). The positive predictive value of an sFlt-1:PlGF ratio above 38 in women between 24 weeks and 36 weeks (and 6 days) of

gestation in whom pre-eclampsia was suspected, for a diagnosis of pre-eclampsia within four weeks was 36.7% (95% CI, 28.4 to 45.7), with 66.2% sensitivity (95% CI, 54.0 to 77.0) and 83.1% specificity (95% CI, 79.4 to 86.3).<sup>32</sup> A separate study demonstrated that PIGF (ROC 0.87, 95% CI 0.83–0.92), sFlt-1 (ROC 0.83, 95% CI 0.78–0.88) and endoglin (ROC 0.83, 95% CI 0.79–0.88) all have comparable results in the short term prediction for pre-eclampsia requiring delivery within 14 days in women with suspected preterm pre-eclampsia.<sup>34</sup>

Whilst the use of these tests for diagnosis is promising, predictive studies have had more mixed results. A prospective international multicentre study recruited 5121 participants between 2006 and 2009 who had risk factors for pre-eclampsia (nulliparity, diabetes, previous pre-eclampsia or chronic hypertension). Participants had their serum tested for sFlt-1, PIGF and endoglin concentrations and urine PIGF measured at fixed time-points of  $\leq 20$ , 23-27 and 32-35 weeks' gestation. Of this cohort, 3.9% (198 women) developed pre-eclampsia, with 0.9% (47 women) developing pre-term pre-eclampsia. The median maternal serum concentrations of PIGF, sFlt-1, sFlt-1/PIGF and endoglin were significantly different in women who subsequently developed pre-eclampsia compared with those who did not. However, the areas under receiver operating characteristics curves at  $\leq 20$  weeks' gestation were poor for predicting any pre-eclampsia (0.52 and 0.59) or pre-eclampsia at less than 34 weeks' gestation (0.50 to 0.63)). The corresponding sensitivity, specificity and likelihood ratios were also poor.<sup>35</sup>

A further recent longitudinal prospective cohort observational study of angiogenic markers and Angiotensin II type 1 receptor antibodies (AT1AA) was conducted. In pre-eclampsia there is an enhanced vascular sensitivity to angiotensin II and norepinephrine resulting in vasoconstriction and hypertension. The enhanced sensitivity to angiotensin II may be secondary to increased bradykinin upregulation seen in preeclamptic patients. Hence it has been proposed as a potential biomarker.<sup>36</sup> Sequential recruitment of 351 women with a singleton pregnancy was undertaken. Plasma concentrations of women with a singleton pregnancy at 12, 18, 28, 36, 40 weeks' gestation and 6 weeks post-partum were assessed for levels of AT1AA, PIGF, sFlt-1 and endoglin. Women with pre-eclampsia had higher sFlt-1 from 28 weeks onwards ( $p=0.003$ ) and lower PIGF from 18 weeks ( $p=0.004$ ). Endoglin and AT1AA concentrations did not vary over time or between groups.<sup>37</sup> PIGF compares favourably when used as a first trimester screening tool as evidenced by a recent systematic review which included a total of 103 studies and 432,621 participants. This reviewed the accuracy of serum biochemical markers PAPP-A, human Chorionic Gonadotropin, PIGF and Placental Protein 13 (PP13) in the first trimester to predict pre-eclampsia.<sup>38</sup> For pre-eclampsia, the best predictor was PIGF with a positive likelihood ratio of 4.01 (3.74, 4.28) and a negative likelihood ratio of 0.67 (0.64, 0.69).

### **Endothelial dysfunction**

Endothelial cells are important in the systemic inflammatory response as well as mediating local inflammation by upregulating the secretion of cytokines and adhesion molecules such as intracellular adhesion molecule-1 (ICAM-1), vascular cell adhesion molecule 1 (VCAM-1) and E-selectin. Plasma concentrations of these adhesion cell molecules are significantly elevated in women with pre-eclampsia.<sup>42-44</sup> Elevations in these molecules have been demonstrated before clinical symptoms develop, suggesting they may serve as predictors of pre-eclampsia. One prospective study of samples from 57 pregnant women before and after 32 weeks' gestation demonstrated that VCAM-1 was significantly elevated in women developing hypertensive diseases as compared to normotensive women (pre-eclampsia:  $p < 0.001$ ; gestational hypertension:  $p < 0.05$ ; chronic hypertension:  $p < 0.001$ ).

### **Cell free fetal DNA**

Cell free fetal DNA (cffDNA) refers to fetal DNA found in the maternal blood stream. This is now a minimally- invasive technique increasingly being used to screen for genetic anomalies. It has also been proposed as a biomarker for pre-eclampsia. A systematic review examined 13 studies of which 11 had found an increase in cffDNA among pregnant women who subsequently developed pre-eclampsia. In addition, all four studies examining early-onset or severe pre-eclampsia found significantly elevated

cffDNA levels prior to disease onset.<sup>45</sup> Additional recent case control studies examining first trimester cffDNA showed no association with first trimester cffDNA and subsequent development of pre-eclampsia.<sup>46</sup> However, these data conflict with a larger case control study supporting an association. In this study, compared with controls, the median levels and multiples of the median (MoM) values of HYP2, a cell free fetal DNA marker, were significantly higher in the pre-eclampsia and hypertensive disorders of pregnancy groups sampled at 6-14 and 15-23 weeks' gestation.<sup>47</sup> HYP2 is located on chromosome 13 and is hypermethylated in the placenta as well as in maternal blood cells. It has been studied as an epigenetic marker for total cffDNA<sup>48</sup>.

Levine *et al* conducted a nested case control study within the Calcium for Pre-eclampsia Prevention trial cohort of healthy nulliparous women. One hundred twenty pairs were randomly chosen for analysis of serum cell-free fetal DNA (cffDNA), a marker of placental debris, and C-reactive protein (CRP), a marker of inflammation. Serum specimens were obtained from participants at 8 to 20 weeks (at enrolment in the trial), at 26 to 29 weeks, at 36 weeks if still pregnant, and when hypertension or proteinuria were noted and a diagnosis of pre-eclampsia confirmed. At 29 to 41 weeks of gestation, cffDNA concentrations were significantly higher after a diagnosis of pre-eclampsia was made than before (219 vs 112 genome equivalents [GE]/mL,  $P < .001$ ). Before the diagnosis of pre-eclampsia was made, cffDNA levels in cases exceeded controls at 17 to 28 weeks (36 vs 16 GE/mL,  $P < .001$ ). However, samples taken between 29 and 41 weeks were only significantly elevated within 3 weeks prior to the diagnosis of pre-eclampsia (176 vs 75 GE/mL,  $P < .001$ ).<sup>49</sup> These data suggest that cffDNA may be useful

in the both early and late stage detection of preterm pre-eclampsia, as pre-eclampsia appears to be accompanied by a 2-stage elevation of fetal DNA.

Cotter *et al* reanalysed blood samples from women attending first antenatal clinics which were stored in a biobank. 88 women who subsequently developed pre-eclampsia (cases) and 176 matched controls were sampled at a mean gestation (+/-SD) of 15.7 +/- 3.6 weeks. The presence of fetal DNA in the maternal circulation in early pregnancy was associated with an 8-fold increased risk of developing pre-eclampsia.<sup>50</sup> This study did not differentiate between preterm and term pre-eclampsia.

Cell free RNA has also been proposed as a marker for pre-eclampsia. One study examined maternal plasma samples from 62 patients at a tertiary hospital in Indonesia from 15-20 weeks gestation who subsequently developed either early or late onset pre-eclampsia. These were compared with samples from 310 controls (unmatched subjects with a normal pregnancy) using a panel of messenger RNA markers. A receiver operating characteristic curve that was obtained with the estimated score for pre-eclampsia as a test variable yielded a detection rate of 84% (95% CI, 71.8-91.5) at a 5% false-positive rate with an area under the curve of 0.927 (P <0 .001).<sup>51</sup>

### **Immunological response, oxidative stress and inflammation**

In pre-eclampsia, the systemic inflammatory response reported in uncomplicated normal pregnancies<sup>52</sup> becomes exaggerated resulting in abnormal activation of monocytes, platelets and the endothelium.<sup>53</sup> In addition, many acute phase inflammatory proteins increase including C-reactive protein, angiotensinogen, alpha-1-

acid glycoprotein, alpha-1-antitrypsin, caeruloplasmin and fibrinogen. Moreover, as part of the inflammatory response, microparticles (plasma membrane-derived vesicles) are released and elevated levels of circulating microparticles have been shown to correlate with the endothelial dysfunction documented in pre-eclampsia.<sup>54,55</sup>

During placentation, Natural Killer (NK) cells play an important role in the decidual inflammatory response. Decidual NK cells, which are a specialised subset of NK cells, are present pre-conceptually in the endometrium of the luteal phase<sup>56</sup>. Decidual inflammation contributes to the shallow placentation in pre-eclampsia by promoting excess decidual cell IP-10 (IFN- $\gamma$ -induced protein 10), a chemokine. This has been shown to be elevated in the first trimester in a study of 90 women who subsequently developed pre-eclampsia.<sup>57</sup>

Oxidative stress is defined as an increase in the steady state levels of reactive oxygen species (ROS) and may occur as a result of increased free radical generation and/or reduced antioxidant activity. In pre-eclampsia there is an increase in the levels of placental oxidative stress and this oxidative stress may mediate endothelial cell dysfunction and contribute to the pathophysiology of pre-eclampsia.<sup>58</sup> In pre-eclampsia, oxidative stress may result from interactions between the maternal component (including pre-existing conditions such as obesity, diabetes, and hyperlipidemia) and/or the placental component that may involve secretion of lipid peroxides.<sup>59</sup> In a prospective observational study of 306 nulliparous women, markers of oxidative stress including cholesterol and caeruloplasmin levels taken at 14-16 weeks' gestation (sample 1) and

at 18-20 weeks (sample 2) were shown to correlate with the development of pre-eclampsia.<sup>60</sup>

### **Metabolomics and proteomics**

Metabolomics, the scientific study of chemical processes involving metabolites, the intermediates and products of metabolism, has more recently been applied in the prediction of pre-eclampsia. Current metabolomic technologies have allowed the establishment of metabolic signatures of pre-eclampsia in early pregnancy. A two-phase discovery/validation metabolic profiling study from samples taken at 15 weeks' gestation demonstrated an odds ratio for developing pre-eclampsia of 36 (95% CI: 12 to 108), with an area under the receiver operator characteristic curve of 0.94 using a multivariate predictive model combining 14 metabolites.<sup>61</sup>

These findings were then validated using plasma obtained at 15 weeks' gestation from 39 women who subsequently developed pre-eclampsia and 40 similarly matched controls. In an independent case-control study the same 14 metabolites produced an odds ratio of 23 (95% CI: 7 to 73) with an area under receiver operator characteristic curve of 0.92. A phase IIa clinical study of predictive testing for pre-eclampsia to evaluate whether this leads to improved pregnancy outcomes through earlier detection is currently recruiting using this metabolomic platform.<sup>16</sup> The 14 metabolites included in this model were 5-Hydroxytryptophan, Monosaccharide(s), Decanoylcarnitine, Methylglutaric acid and/or adipic acid, Oleic acid, Docosahexaenoic acid and/or docosatrienoic acid,  $\gamma$ -Butyrolactone and/or oxolan-3-one, 2-Oxovaleric acid and/or oxo-



methylbutanoic acid, Acetoacetic acid, Hexadecenoyl-eicosatetraenoyl-sn-glycerol, Di-(octadecadienoyl)-sn-glycerol, Sphingosine 1-phosphate, Sphinganine 1-phosphate and Vitamin D3 derivatives.

Myers *et al* conducted a proteomics pipeline study which reported 8 validated models that predicted 50-56% cases of pre-eclampsia, centred on insulin like growth factor acid labile subunit<sup>15</sup>. A recent study suggested an association between predisposition to pre-eclampsia and cardiovascular disease, suggesting that this field may contribute to risk stratification of pregnant women for long term morbidities<sup>62</sup>. Pregnancy specific glycoproteins have also been investigated as potential predictors of early-onset pre-eclampsia.<sup>63</sup>

### **Placental-derived and other proteins**

Other biomarkers proposed for pre-eclampsia include disintegrin and metalloprotease 12 (ADAM-12), inhibin-A and activin-A, pregnancy associated plasma protein A (PAPP-A) and placental protein 13 (PP-13).<sup>64</sup> PP-13 is a galectin that binds to proteins on the extracellular matrix between the placenta and the endometrium, and through its action as a lysophospholipase-A assists in placental implantation and maternal artery remodelling. PP-13 appears to be down-regulated in women with pre-eclampsia requiring early delivery.<sup>65</sup> A nested case control study of 446 controls and 44 cases who subsequently developed preterm pre-eclampsia (delivery prior to 35 weeks) yielded areas under the receiver operating characteristics curve for first trimester PP-13, PAPP-A and second trimester uterine artery pulsatility index respectively of 0.71 (95% CI,

0.63–0.79;  $P < 0.001$ ), 0.59 (95% CI, 0.51–0.68;  $P = 0.076$ ) and 0.86 (95% CI, 0.77–0.94;  $P < 0.001$ ). Combining PP-13 and pulsatility index using logistic regression analysis yielded an area under the curve of 0.90 (95% CI, 0.84–0.96;  $P < 0.001$ ) and a sensitivity of 0.79.<sup>66</sup> In one prospective study of early pregnancy activin A and inhibin A were significantly higher in preeclamptic patients than in the other groups (activin A: normotension:  $p < 0.005$ ; gestational hypertension:  $p < 0.001$ ; chronic hypertension:  $p < 0.005$ ) (inhibin A: normotension:  $p < 0.005$ ; gestational hypertension:  $p < 0.001$ ; chronic hypertension:  $p < 0.01$ )<sup>67</sup> but these changes have not translated into a clinically applicable predictive or diagnostic test.

### **Misfolded proteins**

Endoplasmic reticulum stress in the placenta, as in other cell types, leads to upregulation of the unfolded protein response pathway.<sup>68-70</sup> The unfolded protein response is a common cellular defence mechanism that promotes removal of unfolded or misfolded proteins to prevent potentially toxic accumulation. Buhimschi and colleagues recently proposed that the presence of elevated urine congophilia (which is thought to indicate the presence of amyloid protein, an aggregate of misfolded proteins) assessed using the Congo red 'dot' test was useful in the diagnosis of pre-eclampsia and the prediction of medically indicated delivery. In a prospective cohort study ( $n = 563$ ) they demonstrated that Congo red retention (CRR) using a cut-off of  $\geq 15\%$  had a sensitivity of 85.9% (95% CI, 81.1 to 89.9), specificity of 85.0% (95% CI, 80.4 to 88.8), positive likelihood ratio of 5.7 (95% CI, 4.4 to 7.5), and negative likelihood ratio of 0.17

(95% CI, 0.1 to 0.2) in predicting pre-eclampsia requiring medically indicated delivery.

The association of Congo red retention with medically indicated delivery remained significant after controlling for gestational age and maternal demographic characteristics in multiple logistic regression. Women with pre-eclampsia were sampled at a median of 34-36 weeks' gestation in this cohort.<sup>71,72</sup> However, a more recent study demonstrated that in addition to women with pre-eclampsia, women with chronic kidney disease without pre-eclampsia have elevated urine Congo Red retention compared to healthy pregnant women. Non-pregnant women with lupus nephritis also had elevated urine congophilia compared with healthy controls demonstrating that an elevated Congo Red retention may not be able to differentiate between these conditions.<sup>73</sup>

### **Combined screening**

A recent systematic review and meta-analysis demonstrated the pooled sensitivity of all single biomarkers to be 0.40 at a false positive rate of 10%. The area under the Summary of Receiver Operating Characteristics Curve (SROC) was 0.786 . When a combination model was used, the predictive value improved to an area under the SROC of 0.893 .<sup>19</sup> A combination of biophysical, biochemical and ultrasound markers may provide a more useful screening test than a test of either component alone.

Screening by maternal characteristics and medical history alone identifies approximately 35% of all cases of pre-eclampsia and about 40% of pre-term pre-eclampsia cases, at a false-positive rate of 10%.<sup>74</sup> First trimester screening by maternal factors with biochemical (PAPP-A and B-hCG) and biophysical markers (blood pressure

and uterine artery Doppler) yields detection rates of all pre-eclampsia and pre-eclampsia requiring delivery before 37 and 34 weeks' gestation of 54%, 75% and 88%, respectively, at a false-positive rate of 10%.<sup>75,76</sup>

The Screening for Pregnancy Endpoints (SCOPE) study developed a predictive model that utilised clinical risk factors and putative biomarkers to predict pre-eclampsia in low risk nulliparous women<sup>30</sup>. A cohort of 5628 women was used to develop and validate a model to predict early onset pre-eclampsia, which included placental growth factor, mean arterial pressure, BMI at 14 to 16 weeks' gestation, the consumption of  $\geq 3$  pieces of fruit per day and mean uterine artery resistance index. The area under the ROC for this model in training and validation cohorts was 0.73 and 0.68 respectively, a modest prediction at best. This study concluded the model did not perform at a level acceptable for introduction to clinical practice, and that due to the low prevalence of pre-eclampsia, the test resulted in an unacceptable number of false-positive rates.

More recently, an analysis of a prospective cohort study of 35,948 women attending for their routine first hospital visit at 11-13 weeks gestation at two maternity hospitals in England was undertaken. . This cohort included 1058 pregnancies with pre-eclampsia (2.9%) demonstrated that prospective combined screening by maternal factors, uterine artery pulsatility index, mean arterial pressure, and placental growth factor predicted 75% (95% confidence interval, 70-80%) of preterm-pre-eclampsia and 47% (95% confidence interval, 44-51%) of term-pre-eclampsia, at a false-positive rate of 10%.<sup>77</sup>

**Conclusion**

Substantial progress in research has been made in recent years, which has considerably improved our understanding of the aetiology of pre-eclampsia. Findings from initial case-control studies require validation in prospective cohort studies and robust evaluation through randomised controlled trials (of a predictive or diagnostic test) before routine introduction into clinical practice. Increasingly, robust clinical studies are being performed assessing a variety of tools for the first and second trimester prediction of pre-eclampsia and point of care tests for assessing need for delivery. Measurement of angiogenic and anti-angiogenic factors remains the leading commercially available test to aid in the short-term diagnosis of pre-eclampsia requiring need for delivery within two weeks. Ongoing clinical trials will provide robust data on the clinical potential for first and second trimester prediction of pre-eclampsia.

## References

1. Tranquilli AL, Dekker G, Magee L, et al. The classification, diagnosis and management of the hypertensive disorders of pregnancy: A revised statement from the ISSHP. *Pregnancy hypertension*. 2014;4(2):97-104.
2. Tranquilli A, Brown M, Zeeman G, Dekker G, Sibai B. The definition of severe and early-onset preeclampsia. Statements from the International Society for the Study of Hypertension in Pregnancy (ISSHP). *Pregnancy Hypertension: An International Journal of Women's Cardiovascular Health*. 2013;3(1):44-47.
3. Redman C. Pre-eclampsia: a complex and variable disease. *Pregnancy Hypertension*. 2014;4(3):241-242.
4. Wildman K, Bouvier-Colle MH. Maternal mortality as an indicator of obstetric care in Europe. *Bjog*. 2004;111(2):164-169.
5. Souza JP, Gulmezoglu AM, Vogel J, et al. Moving beyond essential interventions for reduction of maternal mortality (the WHO Multicountry Survey on Maternal and Newborn Health): a cross-sectional study. *Lancet*. 2013;381(9879):1747-1755.
6. Burton GJ, Jauniaux E. Placental oxidative stress: from miscarriage to preeclampsia. *Journal of the Society for Gynecologic Investigation*. 2004;11(6):342-352.
7. Yung HW, Atkinson D, Campion-Smith T, Olovsson M, Charnock-Jones DS, Burton GJ. Differential activation of placental unfolded protein response pathways implies heterogeneity in causation of early- and late-onset pre-eclampsia. *The Journal of pathology*. 2014;234(2):262-276.
8. Redman CW, Sargent IL. Placental stress and pre-eclampsia: a revised view. *Placenta*. 2009;30 Suppl A:S38-42.
9. Pijnenborg R, Vercruyse L, Hanssens M. The uterine spiral arteries in human pregnancy: facts and controversies. *Placenta*. 2006;27(9-10):939-958.
10. Roberts JM, Hubel CA. Is oxidative stress the link in the two-stage model of pre-eclampsia? *Lancet*. 1999;354(9181):788-789.

11. Chaiworapongsa T, Chaemsaitong P, Yeo L, Romero R. Pre-eclampsia part 1: current understanding of its pathophysiology. *Nat Rev Nephrol.* 2014;10(8):466-480.
12. Craici IM, Wagner SJ, Weissgerber TL, Grande JP, Garovic VD. Advances in the pathophysiology of pre-eclampsia and related podocyte injury. *Kidney international.* 2014;86(2):275-285.
13. Redman C, Sargent L, Staff A. IFPA Senior Award Lecture: Making sense of pre-eclampsia –  
Two placental causes of preeclampsia? *Placenta.* 2014;24(Supplement):S20-S25.
14. Cuffe JSM HO, Salomon C , Rice GE , Perkins AV. Review: Placental derived biomarkers of pregnancy disorders. *Placenta.* 2017(54):104-110.
15. Myers JE TR, Thomas G, Laroy W, Kas K, Vanpoucke G, Roberst CT, Kenny LC, Simpson NA, Baker PN, North RA. Integrated proteomics pipeline yields novel biomarkers for predicting preeclampsia. *Hypertension.* 2013;61(6):1281-1288.
16. Navaratnam K, Alfirevic Z, Baker PN, et al. A multi-centre phase IIa clinical study of predictive testing for preeclampsia: improved pregnancy outcomes via early detection (IMPROVED). *BMC Pregnancy Childbirth.* 2013;13:226.
17. Health NCCfWsaCs. NICE guideline: Hypertension in Pregnancy. . In. London: RCOG Press; 2011.
18. Duley L, Henderson-Smart DJ, Meher S, King JF. Antiplatelet agents for preventing pre-eclampsia and its complications. *The Cochrane database of systematic reviews.* 2007(2):CD004659.
19. Wu P, van den Berg C, Alfirevic Z, et al. Early Pregnancy Biomarkers in Pre-Eclampsia: A Systematic Review and Meta-Analysis. *International journal of molecular sciences.* 2015;16(9):23035-23056.
20. Maynard SE, Min JY, Merchan J, et al. Excess placental soluble fms-like tyrosine kinase 1 (sFlt1) may contribute to endothelial dysfunction, hypertension, and proteinuria in preeclampsia. *J Clin Invest.* 2003;111(5):649-658.

21. Tsatsaris V, Goffin F, Munaut C, et al. Overexpression of the soluble vascular endothelial growth factor receptor in preeclamptic patients: pathophysiological consequences. *J Clin Endocrinol Metab.* 2003;88(11):5555-5563.
22. Lu F, Longo M, Tamayo E, et al. The effect of over-expression of sFlt-1 on blood pressure and the occurrence of other manifestations of preeclampsia in unrestrained conscious pregnant mice. *Am J Obstet Gynecol.* 2007;196(4):396 e391-397; discussion 396 e397.
23. Maynard SE, Venkatesha S, Thadhani R, Karumanchi SA. Soluble Fms-like tyrosine kinase 1 and endothelial dysfunction in the pathogenesis of preeclampsia. *Pediatr Res.* 2005;57(5 Pt 2):1R-7R.
24. Charnock-Jones DS, Sharkey AM, Boocock CA, et al. Vascular endothelial growth factor receptor localization and activation in human trophoblast and choriocarcinoma cells. *Biol Reprod.* 1994;51(3):524-530.
25. Crocker IP, Strachan BK, Lash GE, Cooper S, Warren AY, Baker PN. Vascular endothelial growth factor but not placental growth factor promotes trophoblast syncytialization in vitro. *J Soc Gynecol Investig.* 2001;8(6):341-346.
26. Athanassiades A, Hamilton GS, Lala PK. Vascular endothelial growth factor stimulates proliferation but not migration or invasiveness in human extravillous trophoblast. *Biol Reprod.* 1998;59(3):643-654.
27. Shaarawy M, Al-Sokkary F, Sheba M, Wahba O, Kandil HO, Abdel-Mohsen I. Angiogenin and vascular endothelial growth factor in pregnancies complicated by preeclampsia. *International journal of gynaecology and obstetrics: the official organ of the International Federation of Gynaecology and Obstetrics.* 2005;88(2):112-117.
28. Andraweera PH, Dekker GA, Roberts CT. The vascular endothelial growth factor family in adverse pregnancy outcomes. *Human reproduction update.* 2012;18(4):436-457.
29. Myatt L, Clifton RG, Roberts JM, et al. Can changes in angiogenic biomarkers between the first and second trimesters of pregnancy predict development of pre-eclampsia in a low-risk nulliparous patient population? *BJOG : an international journal of obstetrics and gynaecology.* 2013;120(10):1183-1191.



30. Kenny LC, Black MA, Poston L, et al. Early pregnancy prediction of preeclampsia in nulliparous women, combining clinical risk and biomarkers: the Screening for Pregnancy Endpoints (SCOPE) international cohort study. *Hypertension*. 2014;64(3):644-652.
31. Skrastad RB, Hov GG, Blaas HG, Romundstad PR, Salvesen KA. Risk assessment for preeclampsia in nulliparous women at 11-13 weeks gestational age: prospective evaluation of two algorithms. *BJOG : an international journal of obstetrics and gynaecology*. 2015;122(13):1781-1788.
32. Zeisler H, Llurba E, Chantraine F, et al. Predictive Value of the sFlt-1:PIGF Ratio in Women with Suspected Preeclampsia. *The New England journal of medicine*. 2016;374(1):13-22.
33. Chappell LC, Duckworth S, Seed PT, et al. Diagnostic accuracy of placental growth factor in women with suspected preeclampsia: a prospective multicenter study. *Circulation*. 2013;128(19):2121-2131.
34. Duckworth S, Griffin M, Seed PT, et al. Diagnostic Biomarkers in Women With Suspected Preeclampsia in a Prospective Multicenter Study. *Obstet Gynecol*. 2016;128(2):245-252.
35. Widmer M, Cuesta C, Khan KS, et al. Accuracy of angiogenic biomarkers at 20weeks' gestation in predicting the risk of pre-eclampsia: A WHO multicentre study. *Pregnancy hypertension*. 2015;5(4):330-338.
36. Lei J, Li Y, Zhang S, Wu Y, Wang P, Liu H. The Prognostic Role of Angiotensin II Type 1 Receptor Autoantibody in Non-Gravid Hypertension and Pre-eclampsia: A Meta-analysis and Our Studies. *Medicine*. 2016;95(17):e3494.
37. Aggarwal S, Sunderland N, Thornton C, Xu B, Hennessy A, Makris A. A longitudinal analysis of Angiotensin II type 1 receptor antibody and angiogenic markers in pregnancy. *American journal of obstetrics and gynecology*. 2016.
38. Zhong Y, Zhu F, Ding Y. Serum screening in first trimester to predict pre-eclampsia, small for gestational age and preterm delivery: systematic review and meta-analysis. *BMC pregnancy and childbirth*. 2015;15:191.

39. Caniggia I, Mostachfi H, Winter J, et al. Hypoxia-inducible factor-1 mediates the biological effects of oxygen on human trophoblast differentiation through TGFbeta(3). *J Clin Invest*. 2000;105(5):577-587.
40. Caniggia I, Winter J, Lye SJ, Post M. Oxygen and placental development during the first trimester: implications for the pathophysiology of pre-eclampsia. *Placenta*. 2000;21 Suppl A:S25-30.
41. Caniggia I, Grisaru-Gravnosky S, Kuliszewsky M, Post M, Lye SJ. Inhibition of TGF-beta 3 restores the invasive capability of extravillous trophoblasts in preeclamptic pregnancies. *J Clin Invest*. 1999;103(12):1641-1650.
42. Daniel Y, Kupferminc MJ, Baram A, et al. Plasma soluble endothelial selectin is elevated in women with pre-eclampsia. *Hum Reprod*. 1998;13(12):3537-3541.
43. Henriksen T. The role of lipid oxidation and oxidative lipid derivatives in the development of preeclampsia. *Semin Perinatol*. 2000;24(1):29-32.
44. Clausen T, Djurovic S, Brosstad FR, Berg K, Henriksen T. Altered circulating levels of adhesion molecules at 18 weeks' gestation among women with eventual preeclampsia: indicators of disturbed placentation in absence of evidence of endothelial dysfunction? *Am J Obstet Gynecol*. 2000;182(2):321-325.
45. Martin A, Krishna I, Badell M, Samuel A. Can the quantity of cell-free fetal DNA predict preeclampsia: a systematic review. *Prenatal diagnosis*. 2014;34(7):685-691.
46. Rolnik DL, O'Gorman N, Fiolna M, van den Boom D, Nicolaides KH, Poon LC. Maternal plasma cell-free DNA in the prediction of pre-eclampsia. *Ultrasound in obstetrics & gynecology : the official journal of the International Society of Ultrasound in Obstetrics and Gynecology*. 2015;45(1):106-111.
47. Kim SY, Kim HJ, Park SY, Han YJ, Choi JS, Ryu HM. Early Prediction of Hypertensive Disorders of Pregnancy Using Cell-Free Fetal DNA, Cell-Free Total DNA, and Biochemical Markers. *Fetal diagnosis and therapy*. 2016.
48. Hahn S, Rusterholz C, Hosli I, Lapaire O. Cell-free nucleic acids as potential markers for preeclampsia. *Placenta*. 2011;32(Supplemental 1):S17-S20.

49. Levine RJ, Qian C, Leshane ES, et al. Two-stage elevation of cell-free fetal DNA in maternal sera before onset of preeclampsia. *American journal of obstetrics and gynecology*. 2004;190(3):707-713.
50. Cotter AM, Martin CM, O'Leary J J, Daly SF. Increased fetal DNA in the maternal circulation in early pregnancy is associated with an increased risk of preeclampsia. *American journal of obstetrics and gynecology*. 2004;191(2):515-520.
51. Purwosunu Y, Sekizawa A, Okazaki S, et al. Prediction of preeclampsia by analysis of cell-free messenger RNA in maternal plasma. *American journal of obstetrics and gynecology*. 2009;200(4):386 e381-387.
52. Redman CW, Sacks GP, Sargent IL. Preeclampsia: an excessive maternal inflammatory response to pregnancy. *Am J Obstet Gynecol*. 1999;180(2 Pt 1):499-506.
53. Redman CW, Sargent IL. Pre-eclampsia, the placenta and the maternal systemic inflammatory response--a review. *Placenta*. 2003;24 Suppl A:S21-27.
54. Martinez MC, Tesse A, Zobairi F, Andriantsitohaina R. Shed membrane microparticles from circulating and vascular cells in regulating vascular function. *Am J Physiol Heart Circ Physiol*. 2005;288(3):H1004-1009.
55. Meziani F, Tesse A, David E, et al. Shed membrane particles from preeclamptic women generate vascular wall inflammation and blunt vascular contractility. *Am J Pathol*. 2006;169(4):1473-1483.
56. King A, Burrows T, YW L. Human uterine natural killer cells. *Nature Immunology*. 1996-1997;15:41-52.
57. Lockwood CJ, Huang SJ, Chen CP, et al. Decidual cell regulation of natural killer cell-recruiting chemokines: implications for the pathogenesis and prediction of preeclampsia. *The American journal of pathology*. 2013;183(3):841-856.
58. Myatt L, Cui X. Oxidative stress in the placenta. *Histochem Cell Biol*. 2004;122(4):369-382.
59. Roberts JM, Taylor RN, Goldfien A. Clinical and biochemical evidence of endothelial cell dysfunction in the pregnancy syndrome preeclampsia. *Am J Hypertens*. 1991;4(8):700-708.

60. Dey M, Arora D, Narayan N, Kumar R. Serum Cholesterol and Ceruloplasmin Levels in Second Trimester can Predict Development of Pre-eclampsia. *North American journal of medical sciences*. 2013;5(1):41-46.
61. Kenny LC, Broadhurst DI, Dunn W, et al. Robust early pregnancy prediction of later preeclampsia using metabolomic biomarkers. *Hypertension*. 2010;56(4):741-749.
62. Blumenstein M MM, Black MA, Wu S, Prakash R, Cooney J, McCowan LM, Cooper GJ, North RA. A proteomic approach identifies early pregnancy biomarkers for preeclampsia: novel linkages between a predisposition to preeclampsia and cardiovascular disease. *Proteomics*. 2009;9(11):2929-2945.
63. Blankley RT FC, Westwood M, North R, Baker PN, Walker MJ, Williamson A, Whetton AD, Lin W, McCowan L, Roberts CT, Cooper GJ, Unwin RD, Myers JE. A label-free SRM workflow identifies a subset of pregnancy specific glycoproteins as potential predictive markers of early-onset pre-eclampsia. *Molecular and cellular proteomics*. 2013;12(11):3148-3159.
64. Grill S, Rusterholz C, Zanetti-Dallenbach R, et al. Potential markers of preeclampsia--a review. *Reproductive biology and endocrinology : RB&E*. 2009;7:70.
65. Burger O, Pick E, Zwickel J, et al. Placental protein 13 (PP-13): effects on cultured trophoblasts, and its detection in human body fluids in normal and pathological pregnancies. *Placenta*. 2004;25(7):608-622.
66. Spencer K, Cowans NJ, Chefetz I, Tal J, Meiri H. First-trimester maternal serum PP-13, PAPP-A and second-trimester uterine artery Doppler pulsatility index as markers of pre-eclampsia. *Ultrasound in obstetrics & gynecology : the official journal of the International Society of Ultrasound in Obstetrics and Gynecology*. 2007;29(2):128-134.
67. Hanisch CG, Pfeiffer KA, Schlebusch H, Schmolling J. Adhesion molecules, activin and inhibin--candidates for the biochemical prediction of hypertensive diseases in pregnancy? *Archives of gynecology and obstetrics*. 2004;270(2):110-115.

68. Wu J, Kaufman RJ. From acute ER stress to physiological roles of the Unfolded Protein Response. *Cell death and differentiation*. 2006;13(3):374-384.
69. Eknoyan G, Levin N, Nissenson A, Owen W, Jr., Levey AS, Bolton K. NKF and RPA collaborating on clinical practice guidelines for chronic kidney disease. *Nephrology news & issues*. 2001;15(8):13, 54.
70. Nissenson AR. Darbepoetin Alfa: A Viewpoint by Allen R. Nissenson. *Drugs*. 2001;61(14):2105-2106.
71. Nissenson AR. Daily hemodialysis: challenges and opportunities in the delivery and financing of end-stage renal disease patient care. *Advances in renal replacement therapy*. 2001;8(4):286-292.
72. Buhimschi IA, Nayeri UA, Zhao G, et al. Protein misfolding, congophilia, oligomerization, and defective amyloid processing in preeclampsia. *Sci Transl Med*. 2014;6(245):245ra292.
73. McCarthy FP, Adetoba A, Gill C, et al. Urinary congophilia in women with hypertensive disorders of pregnancy and preexisting proteinuria or hypertension. *American journal of obstetrics and gynecology*. 2016;215(4):464 e461-467.
74. Wright D, Syngelaki A, Akolekar R, Poon LC, Nicolaides KH. Competing risks model in screening for preeclampsia by maternal characteristics and medical history. *American journal of obstetrics and gynecology*. 2015;213(1):62 e61-10.
75. Akolekar R, Syngelaki A, Sarquis R, Zvanca M, Nicolaides KH. Prediction of early, intermediate and late pre-eclampsia from maternal factors, biophysical and biochemical markers at 11-13 weeks. *Prenatal diagnosis*. 2011;31(1):66-74.
76. Scaccocchio E, Figueras F, Crispi F, et al. Performance of a first-trimester screening of preeclampsia in a routine care low-risk setting. *American journal of obstetrics and gynecology*. 2013;208(3):203 e201-203 e210.
77. O'Gorman N, Wright D, Syngelaki A, et al. Competing risks model in screening for preeclampsia by maternal factors and biomarkers at 11-13 weeks gestation. *American journal of obstetrics and gynecology*. 2016;214(1):103 e101-103 e112.

**Table 1. Selected biomarkers and their clinical efficacy**

	Clinical cutoff	Primary outcome	Sensitivity (95% CI)	Specificity (95% CI)	OR (95% CI)	ROC (95% CI)	+ LR (95% CI)	- LR (95% CI)	P value
<b>Angiogenic markers</b>									
	PIGF <sup>32</sup>	<5th centile	Any PE diagnosis at 20-35/40 gestation	0.96 (0.89-0.99)	0.55(0.48-0.61)			0.98 (0.93-1.00)	
	sFlt-1 : PIGF ratio <sup>31</sup>	Ratio>38	Prediction of PE in next 7 days	0.80 (0.52-0.96)	0.78 (0.75-0.82)			0.99 (0.98-1.00)	
	sFlt-1, endoglin <sup>33</sup>		Prediction of preterm PE requiring delivery<14 days			0.83 (0.78-0.88)		0.83 (0.79-0.88)	
	PIGF, sFlt, sFLT-1:PIGF, endoglin <sup>34</sup>	Combined predictor	Prediction of PE at <20/40 and <34/40 gestation			(0.52-0.59)		(0.50-0.63)	
<b>Endothelial dysfunction</b>									

VCAM-1 <sup>44</sup>	Mean concentration	Comparison of women with hypertensive diseases to normotensive women, >32/40				<0.001
<b>Cell free fetal DNA</b> <sup>48</sup>	Mean concentration	Nested case control of women with PE, cffDNA concentrations at 29-41/40				<0.001
<b>Cell free RNA</b> <sup>50</sup>	Panel of mRNA markers, FPR 10%	Case control of women who developed early or late onset PE		0.93 DR 84 (71.8-91.5)		<0.001
<b>Metabolomics</b>						
14 metabolites* <sup>60</sup>	Predictive model	Discovery study, 15/40 gestation	36 (12-108)	0.94		
14 metabolites* <sup>60</sup>	Predictive model	Validation study, 15/40 gestation, case control	23 (7-73)	0.92		
<b>Placental derived proteins</b>						
PAPP-A <sup>65</sup>	Mean concentration	Nested case control, cases delivered <35/40 due to PE		0.59 (0.51-0.68)		0.076
PP-13 + UA pulsatility index <sup>65</sup>	Combined predictor	Nested case control, cases delivered <35/40 due to PE	0.79	0.90 (0.84-0.96)		<0.001
<b>Misfolded proteins</b>						
Urine congophilia <sup>71</sup>	Congo red retention >15%	Prediction of PE requiring medically indicated delivery, sampled at 34-36/40	85.9 (81.1 - 89.9)	5.7 (4.4-7.5)	0.17 (0.1-0.2)	
<b>Combined screening</b>						
Combined screening* <sup>77</sup>	FPR of 10%	Preterm PE; Term PE,	75.0 (70.0-80.0); 47.0 (44.0-51.0)			
Combined screening <sup>§29</sup>		Prediction of PE in low risk nulliparous women		0.68 (0.63-0.74)		

OR odds ratio; + LR positive likelihood ratio; - LR negative likelihood ratio; ROC area under the receiver operating characteristic curve; DR detection rate; PE: pre-eclampsia; \*5-Hydroxytryptophan, Monosaccharide(s), Decanoylcarnitine, Methylglutaric acid and/or adipic acid, Oleic acid, Docosahexaenoic acid and/or docosatrienoic acid,  $\gamma$ -Butyrolactone and/or oxolan-3-one, 2-Oxovaleric acid and/or oxo-methylbutanoic acid, Acetoacetic acid, Hexadecenoyl-eicosatetraenoyl-sn-glycerol, Di-(octadecadienoyl)-sn-glycerol, Sphingosine 1-phosphate, Sphinganine 1-phosphate and Vitamin D3 derivatives; † maternal factors, uterine artery pulsatility index, mean arterial pressure, and placental growth factor; § placental growth factor, mean arterial pressure, BMI at 14 to 16 weeks' gestation, the consumption of  $\geq 3$  pieces of fruit per day and mean uterine artery resistance index

## Highlights

- Preterm preeclampsia remains an important condition associated with substantial maternal and fetal morbidity and mortality, particularly in the developing world.
- Preeclampsia is thought to occur as a result of a two stage process: impaired trophoblast invasion resulting in placental ischemia, coupled and interacting with maternal constitutional factors including genetic, behavioural and environmental influences.
- Conflicting evidence exists regarding aetiology and potential biomarkers, which likely reflects the heterogeneity of the condition and the historical grouping of term and preterm preeclampsia in studies.
- The development of tests to date has focused on first or second trimester screening, and diagnostic tests to aid in the prediction of time to delivery.
- Placental growth factor and soluble fms-like tyrosine kinase-1 have provided the most promising results as a test to assist in determining need to delivery within a short time-frame.
- Placental growth factor in combination with clinical parameters is also useful in the first trimester prediction of preeclampsia; however, larger robust clinical trials are required.
- A phase IIa clinical study is currently being performed using a metabolomic platform. Robust clinical studies similar to this are required to examine other biomarkers.



- There is a need to address the translational gap between biomarkers which have shown promising results and the introduction using clinical trials into clinical practice.

ACCEPTED MANUSCRIPT