

Title	Physical activity, sedentary behavior and the risk of overweight and obesity in school-aged children
Author(s)	Keane, Eimear; Li, Xia; Harrington, Janas M.; Fitzgerald, Anthony P.; Perry, Ivan J.; Kearney, Patricia M.
Publication date	2017-08
Original citation	Keane, E., Li, X., Harrington, J. M., Fitzgerald, A. P., Perry, I. J. and Kearney, P. M. (2017) 'Physical activity, sedentary behavior and the risk of overweight and obesity in school-aged children', <i>Pediatric Exercise Science</i> , 29(3), pp. 408-418. doi:10.1123/pes.2016-0234
Type of publication	Article (peer-reviewed)
Link to publisher's version	http://dx.doi.org/10.1123/pes.2016-0234 Access to the full text of the published version may require a subscription.
Rights	© 2017, Human Kinetics, Inc. Accepted author manuscript version reprinted, by permission, from <i>Pediatric Exercise Science</i> , 29(3), pp. 408-418. https://doi.org/10.1123/pes.2016-0234
Item downloaded from	http://hdl.handle.net/10468/5727

Downloaded on 2019-03-21T05:43:55Z

1 **Title Page**

2

3 **Title:** Physical activity, sedentary behaviour and the risk of overweight and obesity in school aged

4 children

5 **Preferred running head:** Objective physical activity and childhood obesity

6

7

8

9

10 **Abstract**

11 **Purpose:** Globally, public health policies are targeting modifiable lifestyle behaviours. We explore the
12 independent association of moderate-to-vigorous physical activity (MVPA) and sedentary behaviour
13 on the risk of childhood overweight/obesity.

14

15 **Method:** A cross-sectional survey of children aged 8-11 years (N=826). Objective body mass index was
16 used to classify children as normal weight or overweight/obese. Children wore wrist-worn Geneactiv
17 accelerometers for 7-days and thresholds were applied to categorise MVPA and sedentary time.
18 Screen time (ST) was parent reported. Poisson regression examined the independent association of
19 (1) MVPA, (2) objective sedentary time and (3) ST on the risk of overweight/obesity.

20

21 **Results:** Overall, 23.7% (95% CI, 20.8-26.6%) of children were overweight/obese. On average, children
22 spent 10.8% of waking time at MVPA and 61.3% sedentary. One-fifth (22.1%, 95% CI, 19.3-25.0%) of
23 children achieved MVPA recommendations (≥ 60 minutes each day) and 17.5% (95% CI, 14.9-20.1%)
24 met ST recommendations (< 2 hours per day). Time spent at MVPA was inversely associated with the
25 risk of overweight/obese independent of total sedentary time. Total time spent sedentary was not
26 associated with overweight/obese independent of MVPA. ST was associated with an increased risk of
27 overweight/obese independent of physical activity.

28

29 **Conclusion:** Few schoolchildren met physical activity and screen time recommendations suggesting
30 population based measures are needed.

31 **Background**

32 The World Health Organisation (WHO) has recommended that children achieve at least 60 minutes of
33 moderate-to-vigorous physical activity (MVPA) daily (World Health Organization, 2010). However,
34 large proportions of children fail to meet these guidelines. Recent estimates from a European study
35 from five countries reported that only 4.6% of girls and 16.8% of boys aged 10-12 years met MVPA
36 guidelines (Verloigne et al., 2012). Regular physical activity is beneficial for cardiovascular health, bone
37 health, mental health and weight status (Poitras et al., 2016). Studies assessing the association
38 between MVPA and childhood obesity have been relatively consistent with less active children at an
39 increased risk of obesity when compared to more active children (Jiménez-Pavón, Kelly, & Reilly, 2010;
40 Wittmeier, Mollard, & Kriellaars, 2008).

41

42 Sedentary behaviour can be defined whereby very little energy is being expended (≤ 1.5 metabolic
43 equivalent units [METs]) during waking time, including sitting and lying down (Mark S Tremblay et al.,
44 2011). Children have become increasingly sedentary in recent decades (Bucksch et al., 2016) and the
45 association between sedentary behaviour and unfavourable health outcomes including obesity have
46 received increased attention (Carson et al., 2016). Screen time (ST) is a distinct sub-domain of
47 sedentary behaviour (Sigman, 2012). Current recommendations suggest that children should limit
48 recreational ST to no more than two hours per day (Mark S Tremblay et al., 2016). However, more
49 than half of all children exceed ST recommendations (Fakhouri, Hughes, Brody, Kit, & Ogden, 2013;
50 LeBlanc et al., 2015).

51

52 An emerging evidence base suggests that physical activity and sedentary behaviour are separate
53 constructs independently impacting on obesity risk, rather than sedentary behaviour simply
54 representing low MVPA (Carson & Janssen, 2011; Pearson, Braithwaite, Biddle, Sluijs, & Atkin, 2014;

55 Salmon, Tremblay, Marshall, & Hume, 2011; Mark Stephen Tremblay, Colley, Saunders, Healy, &
56 Owen, 2010). Children can meet MVPA guidelines but can also spend a high proportion of time
57 sedentary and visa-versa (Eisenmann, Barteel, Smith, Welk, & Fu, 2008; Herman, Sabiston, Mathieu,
58 Tremblay, & Paradis, 2014; Laurson et al., 2008). A recent systematic review suggested that further
59 research is needed to clarify the interplay of physical activity and sedentary behaviour due to
60 conflicting findings in the current literature (Pearson et al., 2014). Understanding the relationship
61 between physical activity and sedentary behaviour may help inform the design of policies and
62 interventions to tackle poor lifestyle behaviours and obesity in children.

63

64 Few studies have assessed the independent association of physical activity and sedentary time on
65 adiposity in schoolchildren using objective measurements (Ekelund, Hildebrand, & Collings, 2014),
66 with the majority of evidence from the UK and USA (Ekelund et al., 2014). To date, studies have
67 consistently found that physical activity is associated with obesity indicators independent of sedentary
68 time (Ekelund et al., 2014; Katzmarzyk et al., 2015). However, the association between objective
69 sedentary time and obesity markers independent of physical activity is less clear (Cliff et al., 2016;
70 Ekelund et al., 2014). Generally, the association between objective sedentary time and obesity is
71 attenuated after adjustment for MVPA (Chaput et al., 2012; Ekelund et al., 2012; J. Mitchell, Pate,
72 Beets, & Nader, 2013; J. A. Mitchell et al., 2009; Steele, van Sluijs, Cassidy, Griffin, & Ekelund, 2009).
73 Other studies have assessed associations between subjectively measured sedentary behaviour and
74 obesity independent of objective physical activity. Findings from these studies are mixed with some
75 studies reporting positive associations (Coombs & Stamatakis, 2015; Ekelund et al., 2006) and others
76 reporting no association (Colley et al., 2012; Ortega, Ruiz, & Sjörström, 2007).

77

78 The Cork Children's Lifestyle (CCLaS) Study is the first population based sample of schoolchildren in
79 Ireland to collect objective physical activity data. We hypothesised that schoolchildren with
80 overweight/obesity would be less active and more sedentary than those who are normal weight. This
81 study describes objectively measured physical activity and sedentary time throughout the week, by
82 gender and weight status. Second, we describe the characteristics of children who meet MVPA and ST
83 recommendations (neither, one or both). Third, we explore the independent association of MVPA and
84 sedentary behaviour (objective and questionnaire based data) on the risk of childhood
85 overweight/obesity whilst considering demographic characteristics and other lifestyle factors (diet
86 and parent reported sleep time).

87

88 **Methods**

89 **Study design**

90 Details of the CCLaS Study have been described elsewhere (Keane, Kearney, Perry, Browne, &
91 Harrington, 2014). Briefly, the study aimed to recruit 1,000 schoolchildren to estimate the prevalence
92 of overweight/obesity in Irish children with a precision of $\pm 2.7\%$ assuming a 26% prevalence rate
93 within the study sample. Information on primary schools in Cork City (an urban area) and
94 Mitchelstown (a rural area) was obtained from the Department of Education and Skills website.
95 Children in 3rd and 4th classes (years 5 and 6 of enrollment into primary school) were the target
96 population. Schools were recruited from Cork City using probability proportionate-to-size and
97 purposive sampling. All schools in one rural area in Cork County were invited to participate. Data were
98 collected in schools during school term times between April 2012 and June 2013. Overall, 1,075 8-11
99 year old children living in Cork, Ireland took part in the survey. At the school level, a response rate of
100 65.5% (N=27) was achieved, while 59% (N=1,075) of invited children participated in the study. Data
101 were collected using physical measurements, child and parent questionnaires, 3 day food diaries and
102 accelerometers. Ethical clearance was granted from the Clinical Research Ethics committee of the Cork

103 Teaching Hospitals, Cork, Ireland. Only children whose parents/guardians provided informed consent
104 participated in the study (Keane et al., 2014).

105

106 **Anthropometric measurements and obesity definition**

107 Trained researchers measured child height and weight using standard methods (Keane et al., 2014).

108 Height was measured to the nearest millimetre using a Leicester portable height stick and weight was

109 measured to the nearest 0.1 kilogram using a Tanita WB100MA mechanical scale. Measurements were

110 taken without shoes and in light clothing. Age and gender specific International Obesity Taskforce

111 definitions for body mass index (BMI) were used to categorise children as (1) normal weight or (2)

112 overweight/obese (Cole, Bellizzi, Flegal, & Dietz, 2000).

113

114 **Physical activity and sedentary variables**

115 Accelerometers

116 Free living physical activity was measured over seven consecutive days using a wrist worn validated

117 tri-axial Geneactiv accelerometer (Esliger et al., 2011; Phillips, Parfitt, & Rowlands, 2012). The

118 accelerometers were set to record at 100Hz for seven days using the 'on button press' setting of

119 Geneactiv software version 2.2. The accelerometers were placed on the non-dominant hand of each

120 participating child. Children were asked to wear the accelerometers 24 hours a day each day. The

121 children were advised that the accelerometers were waterproof and were only to be removed if

122 necessary.

123

124 The accelerometer data were downloaded using Geneactiv software version 2.2 and saved on hard

125 drives. The day and month the accelerometer was worn was extracted from the accelerometers during

126 processing. Days of the week were categorised as total week (Sunday-Saturday), week days (Monday-

127 Friday) or weekend days (Saturday-Sunday). Accelerometer wear month was categorised to reflect
128 the school year as January-March, April-June or October-December.

129

130 Parent questionnaire data were used to identify the most frequent waking time and bed time on week
131 and weekend days. These data were then used to estimate the number of waking hours each day. The
132 mean waking time was 14 hours per day. To be included in the analysis, children needed to have
133 recorded ≥ 600 minutes of waking time data for each of the seven days. For children included in the
134 analysis who had missing data, data were scaled to full waking time. Non-wear time was determined
135 using an algorithm by van Hees et al, 2011 (van Hees et al., 2011).

136

137 The raw data files (*.bin) were first read into minute by minute data files (*.csv) using R software and
138 then summarised using SAS 9.3 (Statistical Analysis System Inc., 2002). Minutes per day spent
139 sedentary and at light, moderate and vigorous physical activity were categorised using classification
140 thresholds defined by Phillips et al., 2012 (Phillips et al., 2012). As our data were collected at 100Hz,
141 the classification thresholds were adjusted from 80Hz to 100Hz (Phillips et al., 2012). Sedentary time
142 minutes were divided by 60 and presented as hours per day. Children who achieved ≥ 60 minutes of
143 MVPA on all seven days were categorised as meeting MVPA guidelines (World Health Organization,
144 2010).

145

146 Parent reported screen time

147 Two parent reported responses to questions were used to define ST. The first question asked how
148 many hours per weekday the study child spends watching television, videos or DVDs and the second
149 asked how many hours per weekday the study child spends using computer and non-active game
150 consoles. Both questions had 6 level responses; none, less than an hour, 1 hour to less than 3 hours,

151 3 hours to less than 5 hours, 5 hours to less than 7 hours, or 7 hours or more. The responses were
152 combined and those who engaged in <1 hour in both behaviours were categorised as <2 hours per day
153 and the remainder of children were categorised as >2 hours per day (Mark S Tremblay et al., 2016).

154

155 Combined index for meeting public health recommendations

156 Children were categorised into four groups based on MVPA and ST recommendations as (1) met both,
157 (2) met MVPA only, (3) met ST only or (4) met neither recommendation. This variable was
158 dichotomised as met both recommendations versus met neither/one recommendation for the
159 regression analysis (Table 4) due to the small numbers of children with overweight/obesity who met
160 both recommendations.

161

162 **Covariates**

163 Gender was recorded as girl or boy by trained researchers during physical measurements. Child age
164 was calculated using date of physical measurement and date of birth (parent reported). Parent
165 reported highest level of maternal education is used as a proxy measure of socio-economic status. The
166 variable was coded as lower secondary education or less, higher secondary education, post-secondary
167 education, third level education or missing. A missing maternal education category (N=77) was created
168 to retain the maximum number of children in the analysis.

169

170 Consecutive three day estimated food diaries were completed by the children and assessed for
171 completeness with the children by trained research assistants (Keane et al., 2014). The food diary
172 data were entered into WISP version 4 (Tinuviel Software, Anglesey, UK). Average daily energy (kcal)
173 intake, average daily fruit and vegetable (g/day) intake, and average daily sugar sweetened beverage

174 (SSB; ml/day) intake were estimated. An energy intake to estimated basal metabolic rate (BMR) ratio
175 was calculated, to classify under reporters (energy intake). BMR was calculated using equations by
176 Schofield, 1985 (Schofield, 1984). Cut-off values for energy intake to BMR were defined using an
177 equation proposed by Goldberg as either (1) under reporters or (2) plausible reporters (Black, 2000;
178 Goldberg et al., 1991).

179

180 Sleep time was parent reported. Parents were asked to report when the child usually wakes up and
181 goes to bed on week and weekend days separately. Average sleep time per night was calculated and
182 sleep time was categorised as either adequate (>10 hours for 5-10 year olds, >9 hours for 10+ year
183 olds) or not adequate (\leq 10 hours for 5-10 year olds, \leq 9 hours for 10+ year olds) (Chen, Beydoun, &
184 Wang, 2008).

185

186 **Statistical analysis**

187 Prevalence estimates with 95% confidence intervals were calculated. Continuous variables were
188 assessed for Normality using histograms. Normally distributed data are presented as means and
189 standard deviations and non-Normally distributed data are presented as medians and quartiles.
190 Differences in time spent sedentary and at each physical activity intensity between children who were
191 normal weight and those affected by overweight/obesity were tested using gender and age adjusted
192 ANCOVA while gender differences were tested using age adjusted ANCOVA.

193

194 Poisson regression was used to examine the risk of overweight/obesity by (1) average daily quartile of
195 sedentary time, (2) average daily quartile of MVPA time, (3) meeting MVPA recommendations, (4)
196 meeting ST recommendations and (5) meeting both recommendations. Models 1 were adjusted for
197 gender, child age, maternal education and accelerometer wear month. Models 2 were further

198 adjusted for average daily kcal intake, average daily fruit and vegetable intake, average daily SSB
199 intake and parent reported sleep time. To assess the independent association of sedentary time and
200 MVPA on childhood overweight/obesity, the quartiles of MVPA time and quartiles of sedentary time
201 variables were adjusted for one another whilst including all other covariates. The MVPA and ST
202 recommendation variables were also adjusted for one another. Tests for trend were obtained by
203 including sedentary and MVPA quartiles values as continuous variables. Possible multicollinearity
204 between MVPA and sedentary time was assessed using correlation coefficients and variance inflation
205 factor (VIF). All regression analysis accounted for the clustering of children within schools. Regression
206 analysis excluding children who under reported their average daily kcals intake was also undertaken.
207 Statistical analysis was conducted in Stata 12 IC (StataCorp LP, USA).

208

209

210 **Results**

211 Of the 1,075 children who participated in the study, 1,029 had accelerometer data available. The case
212 base for the current paper is the 826 children who provided valid physical activity (≥ 600 minutes of
213 waking time data recorded for each of the seven days) and BMI data. The median accelerometer wear
214 time for the 826 children was 99.2% (IQR 98.5-99.7%) and missing wear time data were scaled to full
215 wear time for each child. Table 1 describes the characteristics of the participating children by gender.
216 Overall, 56.3% of the participating children were boys and mean age was 9.9 years (SD 0.7). Median
217 daily fruit and vegetable intake in girls and boys was 112.0g and 95.3g, respectively. Median daily SSB
218 intake was 166.7ml in girls and 194.7ml in boys. Overall, 94.6% of girls and 89.6% of boys had adequate
219 parent reported sleep time. One hundred and ninety six children (23.7%, 95% CI, 20.8-26.6%) were
220 affected by overweight/obesity.

221

222 Figure 1 presents the proportion of children achieving WHO recommended MVPA levels, by gender
223 and weight status. In total, 22.1% (95% CI, 19.2-24.9%) of children achieved the recommended
224 physical activity levels on each of the seven days. A higher proportion of boys met the
225 recommendations than girls (26.6% vs. 16.3%, $p<0.001$). A higher proportion of normal weight
226 children met the MVPA recommendations than children with overweight/obesity (26.0% vs. 9.7%,
227 $p<0.001$).

228

229 Table 2 presents mean time spent engaging in each physical activity intensity and mean time spent
230 sedentary over the total week, on weekdays and on weekend days in the total sample and by weight
231 status. Over the total week, children spent on average 27.9% (234.6 minutes) of their time at light
232 physical activity, 10.8% (90.6 minutes) of their time at MVPA and 61.3% (8.6 hours) of their time
233 sedentary (not including sleeping time). Boys spent more time engaged in MVPA than girls on
234 weekdays ($p<0.001$), weekend days ($p=0.002$) and over the total week ($p<0.001$) and also spent slightly
235 more time sedentary than girls over the total week (8.6 v 8.5 hours, $p=0.06$).

236

237 Time spent at light physical activity did not differ between the normal weight children and children
238 with overweight/obesity. Normal weight children spent an average of 20 minutes longer per day
239 engaging in MVPA than children with overweight/obesity over the total week (95.4 v 75.1 minutes,
240 $p<0.001$). Normal weight children spent less time sedentary than children with overweight/obesity on
241 weekdays (8.6 v 8.9 hours, $p=0.008$), weekend days (8.2 v 8.6 hours, $p=0.001$) and over the total week
242 (8.5 v 8.8 hours, $p=0.001$) (Table 2).

243

244 Table 3 presents the proportion and characteristics of children who met neither, one or both
245 recommendations (physical activity and ST) over a total week. Overall, 5.2% of children met both

246 recommendations, 17.0% met MVPA recommendations only, 12.3% met ST recommendations only,
247 and 65.5% met neither recommendation. More children with overweight/obesity met neither
248 recommendation than normal weight children (81.4% v 60.6%, $p < 0.001$). Children who achieved
249 neither recommendation spent less time at MVPA, more time sedentary, had a lower median fruit and
250 vegetable intake, a higher median SSB intake and fewer achieved sleep time guidelines than those
251 who met both recommendations.

252

253 Table 4 presents the independent association of MVPA and total sedentary time on the risk of
254 childhood overweight/obesity over the total week. Children in the lowest MVPA quartile compared to
255 the highest quartile were at a 4.29 (95% CI, 2.34-7.88, $p\text{-trend} < 0.001$) times increased risk of
256 overweight/obesity compared to children who were normal weight, independent of sedentary time.

257 There was a graded association between total time spent sedentary and obesity risk whilst controlling
258 for demographic and lifestyle factors (diet and sleep) though this association did not persist following
259 adjustment for MVPA. Independent of MVPA, the risk of overweight/obesity in children in the most
260 sedentary quartile was 0.74 (95% CI, 0.41-1.34, $p\text{-trend} = 0.5$) when compared to the least sedentary
261 quartile. Children who did not achieve MVPA recommendations were at a 2.45 (95% CI, 1.59-3.77)
262 times increased risk of overweight/obesity compared to those who met MVPA recommendations.

263 Children who had greater than two hours of ST per day were at a 1.90 (95% CI, 1.21-3.02) times
264 increased risk of overweight/obesity compared with those who had less than 2 hours of ST per day.

265 Similar findings were observed when the analysis was conducted on the 560 children considered as
266 plausible energy intake reporters (data not shown).

267

268 Discussion

269 Main findings

270 This study examined the independent association of MVPA and sedentary behaviour (objective and
271 questionnaire based data) on the risk of childhood overweight/obesity. There are three key findings.
272 First, children spent 10.8% of their waking time at MVPA and 61.3% sedentary, with those who were
273 overweight/obese less active than children who were normal weight. Second, few children met
274 physical activity (22.1%) or ST recommendations (17.5%), with only 5.2% of children meeting both
275 recommendations. Furthermore, a higher proportion of children with overweight/obesity met neither
276 recommendation than those with a normal weight (81.4% v 60.6%). Third, MVPA was associated with
277 an increased risk of childhood overweight/obesity independent of sedentary time. In contrast, total
278 sedentary time was not associated with childhood overweight/obesity independent of MVPA.
279 However, children who did not meet ST recommendations were at an increased risk of
280 overweight/obesity, independent of physical activity.

281

282 On individual days of the week, up to three quarters of children met MVPA guidelines though only
283 22% of children met WHO recommendations of 60 minutes of MVPA each day suggesting that children
284 are not consistently active over a full week. Children with a normal weight were also more active and
285 spent less time sedentary than those with overweight/obesity on week and weekend days. Over the
286 full week, normal weight children spent approximately 20 additional minutes per day engaging in
287 MVPA when compared to children with overweight/obesity. Though the findings of this study may not
288 be generalisable to children of all ages, increasing MVPA levels in children with overweight/obesity to
289 reach levels in normal weight children may be an achievable goal for policy makers.

290

291 Having used a wrist worn accelerometer, our study adds to the current evidence base as many cross
292 sectional associations to date are from studies using hip worn monitors (Ekelund et al., 2014). Our
293 findings suggest that lack of physical activity and high ST are both predominant risk factors for

294 childhood overweight/obesity, independent of one another and other factors including dietary intake
295 and parent reported sleep time. Furthermore, a dose response relationship was evident for increasing
296 levels of MVPA and the risk of overweight/obesity. The association between total sedentary time and
297 overweight/obesity was attenuated once MVPA was accounted for in the model. However, there is
298 lack of consensus on the appropriateness of including two accelerometer derived activity behaviours
299 in one model. Some studies have reported similar findings where sedentary time was either weakly or
300 not associated with overweight/obesity independent of MVPA (Ekelund et al., 2014). However, we
301 found that meeting ST recommendations was associated with overweight/obesity independent of
302 physical activity, which contributes to the current evidence base. Our finding suggests that how
303 sedentary time is spent (e.g. ST) may be more important than total time spent sedentary. However,
304 as this study only collected data on weekday ST, this may have influenced the prevalence estimates
305 and the subsequent associations with weight status. Some research suggests that diet may help
306 explain the association between time at screen based activities and obesity (Prentice-Dunn & Prentice-
307 Dunn, 2012). Further research is warranted to assess the potential mediating role of diet on the
308 association between ST and obesity.

309

310 Due to the low number of children with overweight/obesity meeting both recommendations in this
311 study, we were unable to assess any additive effect of meeting both recommendations on the risk of
312 childhood overweight/obesity. However, children who achieved neither recommendation (physical
313 activity or ST) were less active, more sedentary, had a poorer dietary intake and fewer achieved
314 adequate sleep compared to those who met both recommendations. This suggests that targeting
315 multiple lifestyle behaviours with a focus on increasing physical activity and reducing ST behaviour
316 may be important to tackle childhood overweight/obesity. However, this area of research would
317 benefit from further study, particularly with a longitudinal approach.

318

319 **Strengths and limitations**

320 To our knowledge, this is the first population based study to report levels of objectively measured
321 physical activity in Irish children. This is one of few population based studies which included objectively
322 measured height, weight, sedentary time and physical activity along with 3 day food diaries. Parent
323 reported data were also collected which allowed a variable to reflect ST recommendations to be
324 constructed.

325

326 This study also has a number of limitations. As the study is cross-sectional, the direction of the
327 relationships between activity, ST and BMI cannot be proven, and may be reciprocal throughout
328 childhood. Decisions made during data processing of accelerometers can have a methodological
329 influence on whether children are found to comply with recommendations (Cain, Sallis, Conway, Van
330 Dyck, & Calhoun, 2013). Data processing decisions can also influence whether time is categorised as
331 either sedentary or non-wear time. Further, the classification thresholds for the non-dominant wrist
332 may overestimate sedentary time (van Loo et al., 2016). The parental questionnaire did not collect
333 data on all electronic devices that can be used for ST purposes. Furthermore, information on sedentary
334 multi-tasking was not collected as part of the study. Though the ST variables in the study were based
335 on parent reported weekday ST patterns, it is arguably appropriate for ranking individuals. However,
336 children typically have more ST on weekends than weekdays (Bucksch et al., 2016). Finally, the validity
337 and reliability of the food diary needs to be tested though we assessed for measurement error in
338 energy intake reporting in this study.

339

340 **Conclusions**

341 Less than one quarter of schoolchildren met international physical activity (22.1%) recommendations
342 and less than one-fifth (17.5%) met ST recommendations. Overall, two-thirds (65.5%) of children met
343 neither recommendation. Low MVPA was associated with an increased risk of childhood

344 overweight/obesity independent of sedentary time. Further, meeting ST recommendations (but not
345 objective sedentary time) was associated with childhood overweight/obesity independent of physical
346 activity. As physical inactivity and sedentary behaviour are common modifiable behaviours in children,
347 effective population based strategies are urgently needed. Improving compliance to health
348 recommendations and targeting the mean difference of 20 minutes per day spent at MVPA between
349 children who are normal weight and those with overweight/obesity are achievable goals for policy
350 makers.

351 **References**

- 352 Black, A. E. (2000). Critical evaluation of energy intake using the Goldberg cut-off for energy intake:
 353 basal metabolic rate. A practical guide to its calculation, use and limitations. *International*
 354 *journal of obesity and related metabolic disorders: journal of the International Association for*
 355 *the Study of Obesity*, 24(9), 1119-1130.
- 356 Bucksch, J., Sigmundova, D., Hamrik, Z., Troped, P. J., Melkevik, O., Ahluwalia, N., . . . Inchley, J. (2016).
 357 International Trends in Adolescent Screen-Time Behaviors From 2002 to 2010. *Journal of*
 358 *Adolescent Health*, 58(4), 417-425. doi:http://dx.doi.org/10.1016/j.jadohealth.2015.11.014
- 359 Cain, K. L., Sallis, J. F., Conway, T. L., Van Dyck, D., & Calhoun, L. (2013). Using accelerometers in youth
 360 physical activity studies: a review of methods. *J Phys Act Health*, 10(3), 437-450.
- 361 Carson, V., Hunter, S., Kuzik, N., Gray, C. E., Poitras, V. J., Chaput, J.-P., . . . Connor Gorber, S. (2016).
 362 Systematic review of sedentary behaviour and health indicators in school-aged children and
 363 youth: an update 1. *Applied Physiology, Nutrition, and Metabolism*, 41(6), S240-S265.
- 364 Carson, V., & Janssen, I. (2011). Volume, patterns, and types of sedentary behavior and cardio-
 365 metabolic health in children and adolescents: a cross-sectional study. *BMC public health*,
 366 11(1), 274.
- 367 Chaput, J. P., Lambert, M., Mathieu, M. E., Tremblay, M. S., O' Loughlin, J., & Tremblay, A. (2012).
 368 Physical activity vs. sedentary time: independent associations with adiposity in children.
 369 *Pediatric Obesity*, 7(3), 251-258. doi:10.1111/j.2047-6310.2011.00028.x
- 370 Chen, X., Beydoun, M. A., & Wang, Y. (2008). Is Sleep Duration Associated With Childhood Obesity? A
 371 Systematic Review and Meta-analysis. *Obesity*, 16(2), 265-274. doi:10.1038/oby.2007.63
- 372 Cliff, D. P., Hesketh, K., Vella, S. A., Hinkley, T., Tsiros, M. D., Ridgers, N. D., . . . Hardy, L. L. (2016).
 373 Objectively measured sedentary behaviour and health and development in children and
 374 adolescents: systematic review and meta - analysis. *Obesity Reviews*, 17(4), 330-344.
- 375 Cole, T. J., Bellizzi, M. C., Flegal, K. M., & Dietz, W. H. (2000). Establishing a standard definition for child
 376 overweight and obesity worldwide: international survey. *BMJ*, 320(7244), 1240.
- 377 Colley, R. C., Wong, S. L., Garriguet, D., Janssen, I., Gorber, S. C., & Tremblay, M. S. (2012). Physical
 378 activity, sedentary behaviour and sleep in Canadian children: parent-report versus direct
 379 measures and relative associations with health risk. *Health reports*, 23(2), A1.
- 380 Coombs, N. A., & Stamatakis, E. (2015). Associations between objectively assessed and questionnaire-
 381 based sedentary behaviour with BMI-defined obesity among general population children and
 382 adolescents living in England. *BMJ open*, 5(6), e007172.
- 383 Eisenmann, J. C., Bartee, R. T., Smith, D., Welk, G., & Fu, Q. (2008). Combined influence of physical
 384 activity and television viewing on the risk of overweight in US youth. *International journal of*
 385 *obesity*, 32(4), 613-618.
- 386 Ekelund, U., Brage, S., Froberg, K., Harro, M., Anderssen, S. A., Sardinha, L. B., . . . Andersen, L. B.
 387 (2006). TV viewing and physical activity are independently associated with metabolic risk in
 388 children: the European Youth Heart Study. *PLoS Med*, 3(12), e488.
- 389 Ekelund, U., Hildebrand, M., & Collings, P. J. (2014). Physical activity, sedentary time and adiposity
 390 during the first two decades of life. *Proc Nutr Soc*, 73(2), 319-329.
- 391 Ekelund, U., Luan, J. a., Sherar, L. B., Esliger, D. W., Griew, P., Cooper, A., & Collaborators, I. C. s. A. D.
 392 (2012). Moderate to vigorous physical activity and sedentary time and cardiometabolic risk
 393 factors in children and adolescents. *Jama*, 307(7), 704-712.
- 394 Esliger, D. W., Rowlands, A. V., Hurst, T. L., Catt, M., Murray, P., & Eston, R. G. (2011). Validation of the
 395 GENE Accelerometer. *Medicine & Science in Sports & Exercise*, 43(6), 1085.
- 396 Fakhouri, T. H., Hughes, J. P., Brody, D. J., Kit, B. K., & Ogden, C. L. (2013). Physical activity and screen-
 397 time viewing among elementary school-aged children in the United States from 2009 to 2010.
 398 *Jama Pediatrics*, 167(3), 223-229.
- 399 Goldberg, G., Black, A., Jebb, S., Cole, T., Murgatroyd, P., Coward, W., & Prentice, A. (1991). Critical
 400 evaluation of energy intake data using fundamental principles of energy physiology: 1.

- 401 Derivation of cut-off limits to identify under-recording. *European Journal of Clinical Nutrition*,
 402 45(12), 569-581.
- 403 Herman, K. M., Sabiston, C. M., Mathieu, M.-E., Tremblay, A., & Paradis, G. (2014). Sedentary behavior
 404 in a cohort of 8-to 10-year-old children at elevated risk of obesity. *Preventive medicine*, 60,
 405 115-120.
- 406 Jiménez-Pavón, D., Kelly, J., & Reilly, J. J. (2010). Associations between objectively measured habitual
 407 physical activity and adiposity in children and adolescents: Systematic review. *International*
 408 *Journal of Pediatric Obesity*, 5(1), 3-18. doi:10.3109/17477160903067601
- 409 Katzmarzyk, P. T., Barreira, T. V., Broyles, S. T., Champagne, C. M., Chaput, J.-P., Fogelholm, M., . . .
 410 Kurpad, A. (2015). Physical activity, sedentary time, and obesity in an international sample of
 411 children. *Medicine and science in sports and exercise*, 48(10), 2062-2069.
- 412 Keane, E., Kearney, P. M., Perry, I. J., Browne, G. M., & Harrington, J. M. (2014). Diet, Physical Activity,
 413 Lifestyle Behaviors, and Prevalence of Childhood Obesity in Irish Children: The Cork Children's
 414 Lifestyle Study Protocol. *JMIR Res Protoc*, 3(3), e44. doi:10.2196/resprot.3140
- 415 Laurson, K. R., Eisenmann, J. C., Welk, G. J., Wickel, E. E., Gentile, D. A., & Walsh, D. A. (2008).
 416 Combined influence of physical activity and screen time recommendations on childhood
 417 overweight. *The Journal of pediatrics*, 153(2), 209-214.
- 418 LeBlanc, A. G., Katzmarzyk, P. T., Barreira, T. V., Broyles, S. T., Chaput, J.-P., Church, T. S., . . . Kuriyan,
 419 R. (2015). Correlates of total sedentary time and screen time in 9–11 year-old children around
 420 the world: The International Study of Childhood Obesity, Lifestyle and the Environment. *PLoS*
 421 *one*, 10(6), e0129622.
- 422 Mitchell, J., Pate, R., Beets, M., & Nader, P. (2013). Time spent in sedentary behavior and changes in
 423 childhood BMI: a longitudinal study from ages 9 to 15 years. *International journal of obesity*,
 424 37(1), 54-60.
- 425 Mitchell, J. A., Mattocks, C., Ness, A. R., Leary, S. D., Pate, R. R., Dowda, M., . . . Riddoch, C. (2009).
 426 Sedentary behavior and obesity in a large cohort of children. *Obesity*, 17(8), 1596-1602.
- 427 Ortega, F. B., Ruiz, J. R., & Sjöström, M. (2007). Physical activity, overweight and central adiposity in
 428 Swedish children and adolescents: the European Youth Heart Study. *International Journal of*
 429 *Behavioral Nutrition and Physical Activity*, 4(1), 1.
- 430 Pearson, N., Braithwaite, R., Biddle, S., Sluijs, E., & Atkin, A. (2014). Associations between sedentary
 431 behaviour and physical activity in children and adolescents: a meta - analysis. *Obesity*
 432 *Reviews*.
- 433 Phillips, L. R. S., Parfitt, G., & Rowlands, A. V. (2012). Calibration of the GENE accelerometer for
 434 assessment of physical activity intensity in children. *Journal of Science and Medicine in Sport*,
 435 16(2), 124-128.
- 436 Poitras, V. J., Gray, C. E., Borghese, M. M., Carson, V., Chaput, J.-P., Janssen, I., . . . Kho, M. E. (2016).
 437 Systematic review of the relationships between objectively measured physical activity and
 438 health indicators in school-aged children and youth 1. *Applied Physiology, Nutrition, and*
 439 *Metabolism*, 41(6), S197-S239.
- 440 Prentice-Dunn, H., & Prentice-Dunn, S. (2012). Physical activity, sedentary behavior, and childhood
 441 obesity: a review of cross-sectional studies. *Psychology, health & medicine*, 17(3), 255-273.
- 442 Salmon, J., Tremblay, M. S., Marshall, S. J., & Hume, C. (2011). Health risks, correlates, and
 443 interventions to reduce sedentary behavior in young people. *American Journal of Preventive*
 444 *Medicine*, 41(2), 197-206.
- 445 Schofield, W. (1984). Predicting basal metabolic rate, new standards and review of previous work.
 446 *Human nutrition. Clinical nutrition*, 39, 5-41.
- 447 Sigman, A. (2012). Time for a view on screen time. *Archives of disease in childhood*, 97(1), 935-942.
- 448 Steele, R. M., van Sluijs, E. M., Cassidy, A., Griffin, S. J., & Ekelund, U. (2009). Targeting sedentary time
 449 or moderate- and vigorous-intensity activity: independent relations with adiposity in a
 450 population-based sample of 10-y-old British children. *The American journal of clinical*
 451 *nutrition*, 90(5), 1185-1192. doi:10.3945/ajcn.2009.28153

- 452 Tremblay, M. S., Carson, V., Chaput, J.-P., Connor Gorber, S., Dinh, T., Duggan, M., . . . Janson, K. (2016).
453 Canadian 24-Hour Movement Guidelines for Children and Youth: An Integration of Physical
454 Activity, Sedentary Behaviour, and Sleep 1. *Applied Physiology, Nutrition, and Metabolism*,
455 *41*(6), S311-S327.
- 456 Tremblay, M. S., Colley, R. C., Saunders, T. J., Healy, G. N., & Owen, N. (2010). Physiological and health
457 implications of a sedentary lifestyle. *Applied Physiology, Nutrition, and Metabolism*, *35*(6),
458 725-740.
- 459 Tremblay, M. S., LeBlanc, A. G., Janssen, I., Kho, M. E., Hicks, A., Murumets, K., . . . Duggan, M. (2011).
460 Canadian sedentary behaviour guidelines for children and youth. *Applied Physiology,*
461 *Nutrition, and Metabolism*, *36*(1), 59-64.
- 462 van Hees, V. T., Renström, F., Wright, A., Gradmark, A., Catt, M., Chen, K. Y., . . . Wareham, N. J. (2011).
463 Estimation of daily energy expenditure in pregnant and non-pregnant women using a wrist-
464 worn tri-axial accelerometer. *PLoS one*, *6*(7), e22922.
- 465 van Loo, C., Okely, A. D., Batterham, M. J., Hinkley, T., Ekelund, U., Brage, S., . . . Janssen, X. (2016).
466 Wrist Accelerometer Cut-points for Classifying Sedentary Behavior in Children. *Medicine and*
467 *science in sports and exercise*.
- 468 Verloigne, M., Van Lippevelde, W., Maes, L., Yildirim, M., Chinapaw, M., Manios, Y., . . . Brug, J. (2012).
469 Levels of physical activity and sedentary time among 10-to 12-year-old boys and girls across 5
470 European countries using accelerometers: an observational study within the ENERGY-project.
471 *International Journal of Behavioral Nutrition and Physical Activity*, *9*(1), 34.
- 472 Wittmeier, K. D. M., Mollard, R. C., & Kriellaars, D. J. (2008). Physical Activity Intensity and Risk of
473 Overweight and Adiposity in Children. *Obesity*, *16*(2), 415-420. doi:10.1038/oby.2007.73
- 474 World Health Organization. (2010). *Global recommendations of physical activity for health*. Retrieved
475 from Geneva:
- 476

477 **Table 1. Characteristics of the girls and boys participating in the Cork Children's Lifestyle (CCLaS)**
 478 **Study**

	Girls	Boys
	N=362	N=464
	Mean \pm SD/ Median	Mean \pm SD/ Median
	(IQR)	(IQR)
Age (continuous)	9.9 \pm 0.7	9.9 \pm 0.6
Height (cm)	139.7 \pm 7.3	139.9 \pm 6.9
Weight (kg)	34.4 (29.9, 40.8)	34.8 (30.0, 39.5)
BMI (kg/m²)	17.7 (16.1, 20.0)	17.4 (16.1, 19.5)
Average daily energy intake (kcal)	1434.0 (1151.3, 1735.0)	1588.3 (1301.7, 1924.0)
Average daily sugar sweetened beverage intake (ml)	166.7 (50.0, 329.3)	194.7 (55.7, 403.3)
Average daily fruit and vegetable intake (g)	112.0 (48.3, 209.3)	95.3 (37.0, 181.5)
Time spent sedentary**		
Total week (mins/day)	509.8 \pm 65.7	518.8 \pm 67.1
(hours/day)	8.5 \pm 1.1	8.6 \pm 1.1
Week days (mins/day)	517.8 \pm 66.1	524.1 \pm 68.4
(hours/day)	8.6 \pm 1.1	8.7 \pm 1.1
Weekend days (mins/day)	489.8 \pm 87.2	505.7 \pm 93.4
(hours/day)	8.2 \pm 1.5	8.4 \pm 1.6
Time spent at MVPA**		
Total week (mins/day)	78.7 \pm 31.3	99.9 \pm 36.7
Week days (mins/day)	80.4 \pm 30.1	101.4 \pm 37.1
Weekend days (mins/day)	74.3 \pm 45.2	96.1 \pm 54.3

ST recommendations (<2 hours per day)	N (%)	N (%)
Met recommendations	72 (20.6)	68 (15.1)
Didn't meet recommendations	277(79.4)	383 (84.9)
MVPA recommendations (≥60 mins each day)		
Met recommendations	59 (16.3)	124 (26.7)
Didn't meet recommendations	303 (83.7)	340 (73.3)
Average daily MVPA ≥60 mins		
Yes	246 (68.0)	398 (85.8)
No	116 (32.0)	66 (14.2)
Parent reported sleep time		
Adequate	334 (94.6)	407 (89.6)
Not adequate	19 (5.4)	47 (10.4)
Weight status*		
Normal weight	270 (74.6)	360 (77.6)
Overweight	68 (18.8)	85 (18.3)
Obese	24 (6.6)	19 (4.1)

479 **Footnotes:** MVPA= moderate to vigorous physical activity, ST= screen time. Values are either
480 presented as means and SD, medians and IQR or N and %. Data on ST was missing for 26 children.
481 *Weight status calculated from BMI and categorised using IOTF definitions, **Based on an average of
482 14 hours of waking time over total day, an average of 14.2 hours of waking time on week days and an
483 average of 13.4 hours of waking time on weekend days.

484

Table 2. Mean time spent sedentary and at each physical activity intensity in school aged children, by weight status

	Total sample (mean, 95% CI)	Normal weight (mean, 95% CI)	Overweight/obese (mean, 95%CI)	P-value
	N=826	630 (76.3%)	196 (23.7%)	
Total week**				
Sedentary (mins/day)	514.9 (510.3-519.4)	510.7 (505.5-515.8)	528.4 (519.0-537.8)	<0.001
(hours/day)	8.6 (8.5-8.7)	8.5 (8.4-8.6)	8.8 (8.7-9.0)	
Light PA (mins/day)	234.6 (231.2-237.9)	234.0 (230.2-237.8)	236.5 (229.5-243.6)	0.92
Moderate PA (mins/day)	79.8 (77.8-81.8)	83.4 (81.1-85.8)	68.1 (64.4-71.8)	<0.001
Vigorous PA (mins/day)*	9.5 (5.1-16.3)	11.1 (6.0-17.7)	6.0 (3.9-10.7)	<0.001
MVPA (mins/day)	90.6 (88.2-93.1)	95.4 (92.6-98.3)	75.1 (70.8-79.4)	<0.001
Total weekdays**				
Sedentary (mins/day)	521.3 (516.7-525.9)	517.9 (512.6-523.1)	532.4 (522.9-542.0)	0.003
(hours/day)	8.7 (8.6-8.8)	8.6 (8.5-8.7)	8.9 (8.7-9.0)	
Light PA (mins/day)	238.5 (235.1-241.9)	237.4 (233.6-241.3)	241.9 (234.5-249.3)	0.53
Moderate PA (mins/day)	80.9 (78.9-82.9)	84.2 (81.9-86.5)	70.2 (66.5-73.8)	<0.001
Vigorous PA (mins/day)*	9.8 (5.0-17.0)	11.0 (6.0-18.4)	5.8 (3.4-11.0)	<0.001

MVPA (mins/day)	92.2 (89.8-94.7)	96.8 (93.9-99.6)	77.6 (73.4-81.9)	<0.001
Total weekend days**				
Sedentary (mins/day)	498.7 (492.5-504.9)	492.7 (485.6-499.7)	518.2 (505.4-531.1)	<0.001
(hours/day)	8.3 (8.2-8.4)	8.2 (8.1-8.3)	8.6 (8.4-8.9)	
Light PA (mins/day)	224.8 (220.5-229.1)	225.3 (220.4-230.2)	223.0 (214.3-231.8)	0.34
Moderate PA (mins/day)	77.0 (74.1-80.0)	81.5 (78.0-84.9)	62.8 (57.2-68.3)	<0.001
Vigorous PA (mins/day)*	6.4 (2.0-14.6)	7.1 (2.5-16.5)	4.5 (1.0-8.6)	<0.001
MVPA (mins/day)	86.5 (83.0-90.0)	92.0 (87.9-96.1)	68.8 (62.5-75.1)	<0.001

Footnotes: PA = physical activity, MVPA= moderate to vigorous physical activity. All values except vigorous PA are means and 95% CI. Weight status comparisons for means were made using age and gender adjusted ANCOVA. *Medians and IQR. Weight status comparisons for medians were made using Wilcoxon's rank sum test. **Based on an average of 14 hours of waking time over total day, an average of 14.2 hours of waking time on week days and an average of 13.4 hours of waking time on weekend days.

Table 3. Proportion and lifestyle characteristics of children who meet moderate to vigorous physical activity and screen time recommendations over a total week

	N (%)	Sedentary hours/day Mean (SD)	MVPA minutes/day Mean (SD)	Average daily fruit and vegetable intake (g) Median (IQR)	Average daily sugar sweetened beverage intake (ml) Median (IQR)	Sleep time not adequate %
Met physical activity and screen time recommendations						
Met both	42 (5.3)	7.5 (0.7)	130.7 (26.1)	192.0 (91.7, 301.7)	83.3 (0, 202.7)	4.8
Met MVPA only	136 (17.0)	7.7 (0.9)	128.9 (25.2)	112.8 (50.5, 234.5)	182.6 (53.7, 370.3)	11.0
Met ST only	98 (12.3)	8.6 (1.1)	81.1 (33.4)	126.0 (70.7, 252.3)	150.0 (17.7, 353.3)	5.1
Met neither	524 (65.5)	8.9 (1.0)	79.0 (30.1)	91.7 (33.0, 166.7)	200.0 (59.3, 378.0)	8.2

Footnote: MVPA= moderate to vigorous physical activity, ST=screen time.

1 **Table 4. Independent associations between sedentary time, moderate to vigorous physical activity**
 2 **and the risk of childhood overweight/obesity over the total week**

		Model 1¹	Model 2²	Model 3
		Adjusted for demographic factors	Adjusted for demographics and lifestyle factors	Fully adjusted
		RR (95% CI)	RR (95% CI)	
Objective sedentary time	Quartile 1 (least sedentary)	1	1	1 ³
	Quartile 2	0.89 (0.61-1.30)	0.89 (0.59-1.33)	0.69 (0.45-1.05)
	Quartile 3	1.54 (1.07-2.21)	1.47 (0.98-2.18)	0.86 (0.51-1.45)
	Quartile 4 (most sedentary)	1.72 (1.18-2.50)	1.63 (1.09-2.45)	0.74 (0.41-1.34)
		p-trend<0.001	p-trend=0.002	p-trend=0.5
Objective MVPA time	Quartile 4 (most MVPA)	1	1	1 ⁴
	Quartile 3	1.96 (1.25-3.07)	1.92 (1.25-2.95)	2.06 (1.25-3.38)

	Quartile 2	2.73 (1.92-3.87)	2.47 (1.81-3.36)	2.73 (1.71-4.35)
	Quartile 1 (least MVPA)	3.99 (2.53-6.29)	3.71 (2.43-5.67)	4.29 (2.34-7.88)
		p-trend<0.001	p-trend<0.001	p-trend<0.001
Met MVPA recommendations	Yes (≥ 60 mins each day)	1	1	1 ⁵
	No (<60 mins each day)	2.63 (1.72-4.00)	2.42 (1.61-3.64)	2.45 (1.59-3.77)
Met ST recommendations	Yes (< 2 per hours)	1	1	1 ⁶
	No (> 2 per hours)	1.96 (1.22-3.16)	1.90 (1.21-2.96)	1.90 (1.21-3.02)
Met both recommendations	Met both	1	1	
	Met one/ neither	2.13 (1.47-3.07)	2.12 (1.47-3.07)	-

3 **Footnotes:** ¹Model 1 adjusted for gender, child age, highest level of maternal education, and
4 accelerometer wear month. ²Model 2 adjusted for gender, child age, highest level of maternal
5 education, accelerometer wear month, average daily kcal intake, average daily fruit and vegetable
6 intake, average daily SSB intake and parent reported sleep time. ³Partially adjusted model plus quartile
7 of MVPA time. ⁴ Partially adjusted model plus quartile of sedentary time. ⁵ Partially adjusted model

- 8 plus whether participants met ST recommendations. ⁶ Partially adjusted model plus whether
- 9 participants met MVPA recommendations.

Figure 1. Proportion of children achieving recommended level of moderate to vigorous physical activity by (1) gender and (2) weight status

