

**UCC Library and UCC researchers have made this item openly available.
Please [let us know](#) how this has helped you. Thanks!**

Title	Association between psychological distress and cancer type in patients referred to a psycho-oncology service
Author(s)	Lavelle, C.; Ismail, Muhammad Fahmi; Doherty, Kieran; Bowler, A.; Mohamad, M. M.; Cassidy, Eugene M.
Publication date	2017-06
Original citation	Lavelle, C., Ismail, M., Doherty, K., Bowler, A., Mohamad, M. and Cassidy, E. (2017) 'Association between psychological distress and cancer type in patients referred to a psycho-oncology service', Irish Medical Journal, 110(6), 579 (8pp). doi:10147/621506
Type of publication	Article (peer-reviewed)
Link to publisher's version	http://dx.doi.org/10147/621506 Access to the full text of the published version may require a subscription.
Rights	© 2017, Irish Medical Journal. All rights reserved.
Item downloaded from	http://hdl.handle.net/10468/5885

Downloaded on 2021-03-07T08:31:23Z



UCC

University College Cork, Ireland
Coláiste na hOllscoile Corcaigh



This Month

Archive

Subscribe

For Authors

About Us

My CPD

Hosted Content

Home

Association between psychological distress and cancer type in patients referred to a psycho-oncology service

C Lavelle¹, MF Ismail², K Doherty¹, A Bowler¹, MM Mohamad³, EM Cassidy².

¹South Lee Mental Health Service, Cork University Hospital, Wilton, Cork, Ireland

²Department of Psychiatry and Neurobehavioural Science, University College Cork, Cork University Hospital, Wilton, Cork, Ireland

³Cognitive Impairment Research Group, Graduate Entry Medical School, University of Limerick, Limerick, Ireland

Abstract

Psychological distress is common in patients with cancer and psychological well-being is increasingly seen as an important component of cancer care. The aim of this study was to examine the relationship between cancer type and subjective distress. The following data were collected from a database of consecutive psycho-oncology referrals to the Liaison Psychiatry service in Cork University Hospital from 2006 to 2015: demographics, cancer diagnosis, Distress Thermometer (DT) score. 2102 out of 2384 referrals were assessed. Of those assessed, the most common cancer diagnoses were breast (23%, n=486) followed by haematological (21%, n=445). There were significant difference in DT score between the different cancer types, ($\chi^2(13)=33.685$, $p=0.001$, Kruskal–Wallis test). When adjusted for age, gender and whether or not the cancer was recently diagnosed, there was no

significant association between cancer type and psychological distress. In conclusion, cancer type is not associated with level of distress in cancer.

Introduction

Psychological distress is common among cancer patients. Receiving a cancer diagnosis is often life-altering and can be life-limiting. Cancer treatment including medication regimens and necessary lifestyle modifications can be challenging. As a consequence, cancer care has increasingly included psychological well-being as one of its key targets. Studies suggest that mental health disorders can independently affect survival in patients with advanced cancer¹. Subjective distress and mental health disorders are more likely to be identified if screened for routinely and then treated appropriately². It has been suggested that certain cancers may be associated with higher levels of distress. Linden et al⁴ have demonstrated links between different cancer types and psychological distress in particular with lung, haematological and gynaecological cancers. The aim of this study was to examine whether particular cancer type may be associated with higher levels of subjective distress in a large Irish sample of cancer patients at a regional cancer treatment centre.

Methods

Data were collected from an electronic database from January 2006 to December 2015 that was populated prospectively from consecutive psycho-oncology referrals to the Liaison Psychiatry service in Cork University Hospital over that time period. Cork University Hospital (CUH) is the main acute general hospital in the southern region and is one of eight regional cancer centres in Ireland. A multidisciplinary Liaison Psychiatry service includes a Psycho-oncology Liaison Clinical Nurse Specialist who provides assessment and management of adults who are attending for cancer treatment at CUH Regional Cancer centre. Patients referred to the psycho-oncology service who were 18-years-old and above with a cancer diagnosis were included in the study. For this study the following data was collected from inpatient and outpatient assessments: demographics details, cancer diagnosis, and distress thermometer (DT) score.

The Distress Thermometer (DT) was endorsed by the National Comprehensive Cancer Network (NCCN) as a screening tool for measuring subjective distress⁵. It consists of a single statement: "Please circle the number below that best describes how much distress you have been experiencing in the past week including today". Patients indicate their level of distress on a visual analogue scale which is presented in the style of a thermometer with a mark between 0 ('no distress') and 10 ('extreme distress'). The DT is completed by the patient voluntarily and is part of the brief distress assessment by Oncology front line clinical staff at CUH. When considering their patients needs

and specifically the need for Psycho-oncology referral, the Oncology clinical staff use the DT to complement the identified clinical need rather than being a screening tool per se. Statistical analysis was performed using PC SPSS Statistics version 21.0 (IBM). The Mann-Whitney U test was used to compare differences between gender, new diagnosis and DT score. A simple linear regression was calculated to predict DT score based on age. T-tests for independent samples were used to compare age between groups. The DT score was compared across cancer types using Kruskal–Wallis test. A one-way ANCOVA was performed to determine whether there was a statistically significant difference between cancer type and DT score controlling for age, gender and new cancer diagnosis. The likelihood of a ‘Zero’ distress DT score was compared between men and women using the Chi-Square test.

Results

Over a ten-year period, the Psycho-oncology liaison service received a total of 2,384 patient referrals, of whom 2,102 (88%) were assessed. The most common reason for not completing an assessment was being too physically unwell or terminally ill (82%, n=232). Of those assessed, 60% (n=1,257) were female and 40% (n=845) were male. The mean age was 56 years old (SD=13.5, range 18, 88). 60% (n=1,264) had been recently diagnosed with cancer (i.e. during the previous three months). The most common cancer diagnoses were breast and haematological (Table 1).

Table 1. Basic demographic details

Demographics	N (%)
Gender	
Male	845 (40)
Female	1257 (60)
Cancer diagnosis < 3 months	
Yes	1264 (60)
No	838 (40)
Source of referral	
Inpatient	1272 (61)
Outpatient	830 (39)
Cancer Diagnosis	
Breast	486 (23)
Haematology	445 (21)
Lung & Thoracic	280 (13)
Gastrointestinal	234 (11)
Genital	226 (11)
Others	431 (21)
Completed Distress Score	
Yes	1419 (68)
No	683 (32)

Sixty-eight% (n=1419) of those assessed had also completed the DT questionnaire. Patients who did not complete the DT Questionnaire were older in age when compared with those who did complete the DT (57.8 SD 13.6 v 55.5 SD 13.4 years, independent samples t-test, $p < 0.001$). Males were also less likely to complete the DT than females (Pearson Chi-Square 21.9, $p < 0.001$ Fishers Exact test). Of those who did complete the DT Questionnaire, median DT score of the entire sample was 6 (range 0, 10). Females had significantly higher DT scores than males (Mann-Whitney U test, $U = 183943$, $p < 0.001$). There was no significant difference in DT score between those with recent (ie < 3 months) and non-recent cancer diagnosis (Mann-Whitney U test, $U = 226130.5$, $p = 0.911$).

Table 2. Demographics and distress score according to cancer diagnosis

Cancer Diagnosis	N (%)	Mean age (S.D)	Female (%)	Cancer diagnosis <3 months (%)	Median Distress Score
Breast	486 (23.1)	55 (11.5)	483 (99)	276 (57)	7.0 (1-10)
Haematological	445 (21.2)	53 (15.4)	191 (43)	241 (54)	6.0 (0-10)
Lung & thoracic	280 (13.3)	62 (9.1)	124 (44)	221 (79)	6.0 (0-10)
Gastrointestinal	234 (11.1)	60 (11.8)	100 (43)	146 (62)	6.0 (0-10)
Genital	226 (10.8)	56 (13.8)	170 (75)	123 (54)	6.0 (1-10)
Central nervous system	96 (4.6)	53 (17.0)	35 (36)	65 (68)	5.0 (0-10)
Head and neck	88 (4.2)	57 (12.7)	31 (35)	72 (89)	6.0 (2-10)
Skin	77 (3.7)	54 (14.9)	47 (61)	31 (40)	7.0 (1-10)
Urinary tract	53 (2.5)	64 (8.5)	13 (25)	18 (34)	7.0 (3-10)
Mesothelial and soft tissue	31 (1.5)	58 (17.1)	11 (35)	18 (58)	5.0 (0-10)
Endocrine	24 (1.1)	52 (17.9)	17 (71)	19 (79)	7.5 (1-10)
Bone and cartilage	23 (1.1)	57 (15.3)	12 (52)	9 (39)	6.0 (3-10)
Unknown primary	24 (1.1)	56 (12.3)	15 (63)	20 (83)	6.5 (3-10)
Multiple primary	15 (0.7)	59 (13.3)	8 (53)	5 (33)	6.0 (5-8)

A significant regression equation to predict DT score based on age (years) was found ($F(1,1417) = 9.905$, $p = 0.02$), with an R^2 of 0.007. Predicted DT score was equal to $6.982 - 0.016$ (age). DT score reduced by 0.016 for each year of age. There was a significant difference in DT score between the different cancer types, ($\chi^2(13) = 33.685$, $p = 0.001$, Kruskal–Wallis test). Skin, breast, endocrine, urinary tract and unknown primary were associated with a higher median distress score (Table 2). A small number (n= 43) of patients who had been referred for psycho-oncology assessment and completed the DT, rated themselves as not being distressed at all (i.e. DT score of zero). This group were older (59.3 SD 10.7 v 55.4 SD 13.5 years, independent samples t-test, $p < 0.05$) and males were over-represented compared to females (Pearson Chi-Square 34.2, $p < 0.001$ Fishers Exact test).

The relationship between DT score and cancer type was compared using an ANCOVA model, controlling for gender, age and new cancer diagnosis. There was no significant effect of cancer type on DT score after controlling

for age, gender and new cancer diagnosis, $F(13,1405)=1.303$, $p=0.204$.

Discussion

The purpose of this study was to investigate whether or not an association exists between different cancer type/site and self-rated psychological distress in patients attending cancer treatment services. Previous work in this area⁷ suggests that cancer patients do not form a homogenous group when it comes to levels of psychological distress; subjective distress has been shown to differ according to the specific type of cancer, such as prostate and breast cancer, amongst others^{8,9}. In order to carry out our own enquiry, we sourced data from a large electronic database of consecutive psycho-oncology referrals at a Regional Cancer centre over a ten-year period. The majority of these patients referred also completed a validated measure of distress, the Distress Thermometer (DT)⁵. Breast cancer patients were the largest group referred to our service, consistent with the overall prevalence of the disease. Breast cancer patients were also among the cancer types with the highest subjective distress scores although this effect was no longer apparent when gender and age were controlled for.

Our study suggests that age and gender are associated with subjective distress in cancer in line with previous studies^{9,11}. Our gender subgroup analysis suggests that women are 50% more likely to report higher distress levels than their male counterparts. With regard to age, we found that an inverse relationship exists between age and distress, with older patients reporting less distress whereas younger patients are more likely to report higher distress levels^{7,11,12}. Our findings are consistent with previous work⁹ which has suggested that the combination of gender and age is what actually matters: the emotional distress of younger patients tends to be greater than that of their older counterparts for a variety of reasons, including the negative impact of a cancer diagnosis on family life and career, as well as the attendant physical consequences of cancer treatments for young women in particular.

A zero score on the distress thermometer is a curious finding in patients referred because of concerns about their psychological well-being. While a majority of those with cancer are clearly resilient, at least some distress should be expected and therefore its absence could be viewed as maladaptive¹⁴. The finding that men were over-represented in this group is consistent with the literature on male denial of distress¹⁵. It follows that optimal distress assessment especially of older men should include where possible taking a collateral history from a significant other.

One of the limitations of this study is that it did not include illness-related variables such as cancer staging and phase of treatment, which have been found to be key predictors of psychological distress at various points in the course of the disease¹⁰. It also did not include the types of treatment for cancer that have been shown to impact on levels of distress according to the severity of the treatment modality⁹. Other psychosocial variables such as marital status, children, education and employment status were also not examined. Knowledge of such characteristics might have highlighted particular risk factors for developing psychological distress which might otherwise be overlooked. The strengths of the study are the large sample size and the applicability of the findings to Regional Cancer centres all of

whom should offer distress assessment and management to their patients in line with National policy¹⁶ and international best practice⁵.

In conclusion, in our large Irish sample of adults with active cancer, the main finding was that subjective distress was related to gender and age but not to cancer type. A secondary finding was that older males in particular are less likely to complete a distress assessment tool and when they do are less likely to rate themselves as being distressed suggesting the need to adapt the method of distress assessment beyond the distress assessment tool in this cohort.

Conflict of Interest:

The authors declare that they have no conflict of interests.

Correspondence:

Dr Eugene Cassidy, Department of Psychiatry and Neurobehavioural Science, University College Cork, Cork University Hospital, Wilton, Cork, Ireland

Email: eugene.cassidy@hse.ie

References

1. Lloyd- Williams M, Sheils C, Taylor F, Dennis M. Depression – an independent predictor of early death in patients with advanced carcinoma. *J Affect Disord* 1999; 113 (1) 127–132. Doi10.1016/j.jad.2008.04.002
2. Meijer A, Roseman M, Delisle VC, Milete K, Levis B, Syamchandra A, Stefanek ME, Stewart DE, De Jonge P, Coyne JC, Thombs BD. Effects of screening for psychological distress on patient outcomes in cancer A systematic review. *J Psychosom Res* 2013; 75 1-17. Doi10.1016/j.jpsychores.2013.01.012
3. Curtis R, Groarke A, Sullivan F. Stress and self-efficacy predict psychological adjustment at diagnosis of prostate cancer. *Scientific Reports* 4, Article number 5569(2014) doi10.1038/srep05569
4. Linden W, Vodermaier A, Mackenzie R, Greig D. (2012) Anxiety and depression after cancer diagnosis Prevalence rates by cancer type, gender, and age. *J Affect Disord* 2012. 141 343-351
5. Holland JC, Andersen B, Breitbart WS, Compas B, Dudley MM, Fleishman S, Fulcher CD, Greenberg DB, Greiner CB, Handzo GF, Hoofring L, Jacobsen PB, Knight SJ, Learson K, Levy MH, Loscalzo MJ, Manne S, McAllister-Black R, Riba MB, Roper K, Valentine AD, Wagner LI, Zevon MA. NCCN Distress Management Panel. *J Natl Compr Canc Netw*. 2010;8(4)448–485
6. Ryan DA, Gallagher P, Wright S, Cassidy EM. Sensitivity and specificity of the Distress Thermometer and a 2-item depression screen (Patient Health Questionnaire-2) with a “help” question for psychological distress and psychiatric morbidity in patients with advanced cancer. *Psycho-Oncology* 2012; 211275–1284. doi 10.1002/pon.2042
7. Zabora J, BrintzenhofeSzoc K, Curbow B, Hooker C, Piantadosi S. The Prevalence of Psychological Distress by

Cancer Site. *Psycho-Oncology* 2001; 10 19-28.

8. Brintzenhofe-Szoc KM, Levin TT, Li Y, Kissane DW, Zabora JR. Mixed Anxiety/Depression Symptoms in a Large Cancer Cohort: Prevalence by Cancer Type. *Psychosomatics* 2009; 50 383-91. doi 10.1176/app I.psy.50.4.383.

9. Admiraal JM, Reyners AK, Hoekstra-Weebers JE. Do cancer and treatment type affect distress? *Psycho-Oncology* 2012; 22 1766-1773. doi 10.1002/pin.3211. Epub 2012 Oct 29.

10. Vodermaier A, Linden W, MacKenzie R, Greig D, Marshall C. Disease Stage Predicts Post-Diagnosis Anxiety and Depression only in Some Types of Cancer. *Br J Cancer* 2011; 105 1814-7.

11. Thomas BC, NandaMohan V, Nair MK, Pandey M. Gender, Age and Surgery as a Treatment Modality leads to Higher Distress in Patients with Cancer. *Support Care Cancer* 2010; 19 239-250. Doi:10.1007/s00520-009-0810-4. Epub 2010 Feb 23.

12. Carlson LE, Angen M, Cullum J, Goodey E, Koopmans J, Lamont L, MacRae JH, Martin M, Pelletier G, Robinson J, Simpson JS, Speca M, Tillotson L, Bultz BD. High Levels of Untreated Distress and Fatigue in Cancer Patients. *Br J Cancer* 2004; 90 2297-2304.

13. Groarke A, Curtis R, Kerin M. Global Stress Predicts both Positive and Negative Emotional Adjustment at Diagnosis and Post-Surgery in Women with Breast Cancer. *Psycho-Oncology* 2013; 22 177-85. Doi:10.1002/pon.2071. Epub 2011 Oct 17.

14. Molina Y, Yi JC, Martinez-Gutierrez J, Reding KW, Yi-Frazier JP, Rosenberg AR. Resilience Among Patients Across the Cancer Continuum Diverse Perspectives. *Clinical journal of oncology nursing*. 2014;18(1)93-101. doi10.1188/14.CJON.93-101.

15. Ketterer MW, Denollet J, Chapp J, Thayer B, Keteyian S, Clark V, John S, Farha AJ, Deveshwar S. Men deny and women cry, but who dies? Do the wages of "denial" include early ischemic coronary heart disease? *J Psychosom Res*. 2004 Jan;56(1)119-23.

16. Department of Health and Children (DOHC). A strategy for cancer control in Ireland / National Cancer Forum. 2006.

P579



