

**UCC Library and UCC researchers have made this item openly available.
Please [let us know](#) how this has helped you. Thanks!**

Title	Multi-wavelength emission from a single InGaN/GaN nanorod analyzed by cathodoluminescence hyperspectral imaging
Author(s)	Kusch, Gunnar; Conroy, Michele; Li, Haoning; Edwards, Paul R.; Zhao, Chao; Ooi, Boon S.; Pugh, Jon; Cryan, Martin J.; Parbrook, Peter J.; Martin, Robert W.
Publication date	2018
Original citation	Kusch, G., Conroy, M., Li, H., Edwards, P. R., Zhao, C., Ooi, B. S., Pugh, J., Cryan, M. J., Parbrook, P. J. and Martin, R. W. (2018) 'Multi-wavelength emission from a single InGaN/GaN nanorod analyzed by cathodoluminescence hyperspectral imaging', Scientific Reports, 8(1), 1742 (8pp). doi: 10.1038/s41598-018-20142-5
Type of publication	Article (peer-reviewed)
Link to publisher's version	https://www.nature.com/articles/s41598-018-20142-5 http://dx.doi.org/10.1038/s41598-018-20142-5 Access to the full text of the published version may require a subscription.
Rights	© 2018, the Authors. Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. Te images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit http://creativecommons.org/licenses/by/4.0/ . http://creativecommons.org/licenses/by/4.0/
Item downloaded from	http://hdl.handle.net/10468/6348



Downloaded on 2024-01-25T18:19:54Z

Multi-wavelength emission from a single InGaN/GaN nanorod analyzed by cathodoluminescence hyperspectral imaging

Gunnar Kusch^{1,*}, Michele Conroy^{2,3,4}, Haoning Li^{2,3}, Paul R. Edwards¹, Chao Zhao⁵, Boon S. Ooi⁵, Jon Pugh⁶, Martin J. Cryan⁶, Peter J. Parbrook^{2,3}, and Robert W. Martin¹

¹Department of Physics, SUPA, University of Strathclyde, Glasgow G4 0NG, United Kingdom

²Tyndall National Institute, University College Cork, Lee Maltings, Dyke Parade, Cork, Ireland

³School of Engineering, University College Cork, College Road, Cork, Ireland

⁴Pacific Northwest National Laboratory, Richland, WA, United States of America

⁵Photonics Laboratory, King Abdullah University of Science and Technology (KAUST), Thuwal 23955-6900, Saudi Arabia

⁶Department of Electrical and Electronic Engineering, University of Bristol, Bristol BS8 1UB, United Kingdom

*gunnar.kusch@strath.ac.uk

ABSTRACT




Figure 1. Mode positions as calculated by FDTD for varying nanorod length and constant nanorod diameter. Weak modes are shown as open symbols.




Figure 2. Mode positions as calculated by the plane wave model for different rod sidelength, TE modes are shown as dotted, TM modes as full lines.




Figure 3. Nanorod geometry and input spectra used for the FDTD simulation of the nanorod. The source in the shown case is located in the apex of the nanorod representing the *c*-plane QW emission, for the modelling of the behaviour of the semipolar QWs the source was moved into the pyramidal section of the nanorod.