

<b>Title</b>	Multi-wavelength emission from a single InGaN/GaN nanorod analyzed by cathodoluminescence hyperspectral imaging
<b>Author(s)</b>	Kusch, Gunnar; Conroy, Michele; Li, Haoning; Edwards, Paul R.; Zhao, Chao; Ooi, Boon S.; Pugh, Jon; Cryan, Martin J.; Parbrook, Peter J.; Martin, Robert W.
<b>Publication date</b>	2018
<b>Original citation</b>	Kusch, G., Conroy, M., Li, H., Edwards, P. R., Zhao, C., Ooi, B. S., Pugh, J., Cryan, M. J., Parbrook, P. J. and Martin, R. W. (2018) 'Multi-wavelength emission from a single InGaN/GaN nanorod analyzed by cathodoluminescence hyperspectral imaging', Scientific Reports, 8(1), 1742 (8pp). doi: 10.1038/s41598-018-20142-5
<b>Type of publication</b>	Article (peer-reviewed)
<b>Link to publisher's version</b>	<a href="https://www.nature.com/articles/s41598-018-20142-5">https://www.nature.com/articles/s41598-018-20142-5</a> <a href="http://dx.doi.org/10.1038/s41598-018-20142-5">http://dx.doi.org/10.1038/s41598-018-20142-5</a> Access to the full text of the published version may require a subscription.
<b>Rights</b>	© 2018, the Authors. Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. Te images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <a href="http://creativecommons.org/licenses/by/4.0/">http://creativecommons.org/licenses/by/4.0/</a> . <a href="http://creativecommons.org/licenses/by/4.0/">http://creativecommons.org/licenses/by/4.0/</a>
<b>Item downloaded from</b>	<a href="http://hdl.handle.net/10468/6348">http://hdl.handle.net/10468/6348</a>




Downloaded on 2019-02-19T17:46:25Z

**UCC**


University College Cork, Ireland  
Coláiste na hOllscoile Corcaigh







**Figure 2.** Mode positions as calculated by the plane wave model for different rod sidelength, TE modes are shown as dotted, TM modes as full lines.



**Figure 3.** Nanorod geometry and input spectra used for the FDTD simulation of the nanorod. The source in the shown case is located in the apex of the nanorod representing the *c*-plane QW emission, for the modelling of the behaviour of the semipolar QWs the source was moved into the pyramidal section of the nanorod.