

**UCC Library and UCC researchers have made this item openly available.
Please [let us know](#) how this has helped you. Thanks!**

Title	Appropriate use of elective coronary angiography in patients with suspected stable coronary artery disease
Author(s)	Hanrahan, Michael T.; O'Flynn, Anne Marie; McHugh, Sheena M.; Kearney, Peter; Kearney, Patricia M.
Publication date	2019-01-05
Original citation	Hanrahan, M. T., O'Flynn, A. M., McHugh, S. M., Kearney, P., Kearney, P. M. (2019) 'Appropriate use of elective coronary angiography in patients with suspected stable coronary artery disease', Irish Journal of Medical Science, pp. 1-7. doi:10.1007/s11845-018-1953-4
Type of publication	Article (peer-reviewed)
Link to publisher's version	http://dx.doi.org/10.1007/s11845-018-1953-4 Access to the full text of the published version may require a subscription.
Rights	© 2019, Royal Academy of Medicine in Ireland. Published by Springer Nature Switzerland AG. This is a post-peer-review, pre-copyedit version of an article published in the Irish Journal of Medical Science. The final authenticated version is available online at: http://dx.doi.org/10.1007/s11845-018-1953-4
Embargo information	Access to this article is restricted until 12 months after publication by request of the publisher.
Embargo lift date	2020-01-05
Item downloaded from	http://hdl.handle.net/10468/7432

Downloaded on 2022-10-02T06:09:22Z

Table 2. Appropriateness Categories

Indication #	Indication	Total N = 259	Inappropriate N (%) 54 (21%)	Uncertain N (%) 97 (37%)	Appropriate N (%) 108 (42%)
No non-invasive testing: Asymptomatic					
5	Asymptomatic: Low SCORE risk	3	3	-	-
6	Asymptomatic: Intermediate SCORE risk	2	2	-	-
7	Asymptomatic: High SCORE risk	1	-	1	-
No non-invasive testing: Symptomatic					
9	Symptomatic: Intermediate pretest probability	9	-	9	-
10	Symptomatic: High pretest probability	29	-	-	29
Non-invasive testing					
11	Low-risk findings (e.g., DTS \geq 5)	54	49	5	-
12	Intermediate-risk findings (e.g., DTS 4 to -10)	78	-	78	-
13	High-risk findings (e.g., DTS \leq -11)	10	-	-	10
14	Other high-risk findings (ST-segment elevation, hypotension with exercise, ventricular tachycardia, prolonged ST-segment depression)	67	-	-	67
24	Newly recognized LV systolic dysfunction (i.e., LVEF \leq 40%) with an unknown etiology	2	-	1	1
25	Newly recognized LV systolic dysfunction (i.e., LVEF 41% to 49%) with an unknown etiology	2	-	2	0
26	New regional wall motion abnormality with an unknown etiology and normal LV systolic function	1	-	1	0
33	Coronary CTA: Lesion \geq 50% non-left main	1	-	0	1