

**UCC Library and UCC researchers have made this item openly available.  
 Please [let us know](#) how this has helped you. Thanks!**

<b>Title</b>	Pregnancy outcomes in women with severe fear of childbirth
<b>Author(s)</b>	O'Connell, Maeve A.; Leahy-Warren, Patricia; Kenny, Louise C.; Khashan, Ali S.
<b>Publication date</b>	2019-03-14
<b>Original citation</b>	O'Connell, M. A., Leahy-Warren, P., Kenny, L. C. and Khashan, A. S. (2019) 'Pregnancy outcomes in women with severe fear of childbirth', Journal of Psychosomatic Research, 120, pp. 105-109. doi: 10.1016/j.jpsychores.2019.03.013
<b>Type of publication</b>	Article (peer-reviewed)
<b>Link to publisher's version</b>	<a href="http://www.sciencedirect.com/science/article/pii/S0022399918310365">http://www.sciencedirect.com/science/article/pii/S0022399918310365</a> <a href="http://dx.doi.org/10.1016/j.jpsychores.2019.03.013">http://dx.doi.org/10.1016/j.jpsychores.2019.03.013</a> Access to the full text of the published version may require a subscription.
<b>Rights</b>	© 2019, Elsevier Inc. All rights reserved. This manuscript version is made available under the CC BY-NC-ND 4.0 license. <a href="https://creativecommons.org/licenses/by-nc-nd/4.0/">https://creativecommons.org/licenses/by-nc-nd/4.0/</a>
<b>Embargo information</b>	Access to this article is restricted until 12 months after publication by request of the publisher.
<b>Embargo lift date</b>	2020-03-14
<b>Item downloaded from</b>	<a href="http://hdl.handle.net/10468/7760">http://hdl.handle.net/10468/7760</a>

Downloaded on 2021-01-24T00:56:44Z

1 Abstract

2 Objective

3 To compare pregnancy outcomes for women with and without severe fear of childbirth (FOC)  
4 reported in the second trimester of pregnancy.

5 Methods

6 In a prospective cohort study, 389 singleton pregnancies were followed up using medical  
7 records of participants in a study investigating FOC in Cork, Republic of Ireland. FOC was  
8 measured using the Wijma Delivery Experience Questionnaire Part A (W-DEQ A). Severe  
9 FOC was defined as W-DEQ A  $\geq 85$ , moderate FOC, W-DEQ-A 66-84 and low FOC, W-DEQ  
10 A 0-65. Outcome measures were birthweight, birthweight centile, gestational age, and Apgar  
11 scores at 1 minute and Apgar at 5 minutes. Linear regression was used to assess the association  
12 between FOC and each outcome measure with adjustment for maternal age, smoking, parity  
13 and marital status.

14 Results

15 There was no statistically significant difference in mean birthweight (mean difference = -0.03;  
16 [95% CI: -444.69, 315.82]), mean birthweight centile (mean difference= 0.03; [95%CI: -15.97,  
17 23.53]), or mean gestational age (mean difference= -0.06; [95%CI: -11.69, 4.82]) in women  
18 with severe FOC (n=18) compared with women with low FOC (n=371). In the adjusted models,  
19 there was only a slight correlation between severe FOC and Apgar scores at 1 minute (mean  
20 difference= -0.09 [95%CI: -1.28, 0.32]) and Apgar scores at 5 minutes (mean difference= -0.18  
21 [95%CI: -1.16, 1.08]).

22

23 Conclusion While a slight association was noted between severe FOC and Apgar scores, overall  
24 findings are reassuring and could inform educational interventions which may alleviate FOC.  
25 Awareness of FOC for health care professionals is vital to consider women's mental well-being.

26 Keywords

27 Pregnancy, fear of childbirth, tocophobia, outcomes, epidemiology

28 Introduction

29 Fear is a primal and basic emotion experienced universally [1]. Fear exists on a spectrum,  
30 ranging from worries and minor fears, to high fear, and severe phobia[2]. Pregnant women  
31 often experience worries and fear, including fear of childbirth (FOC). Severe FOC impacts  
32 women's experience of pregnancy, manifesting in sleep disturbance and physical complaints  
33 [3-5]. A Swedish study reported that 80% of pregnant women express some level of FOC, thus  
34 it could be considered normal [6], but a recent meta-analysis suggested that up to 14% of  
35 pregnant women could experience severe FOC worldwide [7].

36 FOC is categorised under the general umbrella of anxiety disorders in pregnancy [8] but is  
37 considered a psychological domain in its own right [2]. A meta-analysis [9] examining the  
38 difference between trait fear and trait anxiety concluded that fear has a distinct neurological  
39 mechanism, separate from anxiety and is, therefore, a separate emotion. Thus, various tools  
40 exist specifically to measure FOC [7]. The Wijma Delivery Experience Questionnaire Part A  
41 (W-DEQ A) with a cut-off greater than 85 defining severe FOC is considered the gold standard  
42 [10]. Psychometric analysis of the W-DEQ A [11] indicated the optimal cut-off value of 85 to  
43 detect fear of childbirth which is clinically relevant according to the psychiatric DSM-5  
44 diagnosis of fear of childbirth with 100% sensitivity and 93.8% specificity in an Italian  
45 longitudinal study of nulliparous women (n=106).

46 Only one study to our knowledge previously examined the relationship between FOC and  
47 pregnancy outcomes [12]. Rather than using the validated tool (the W-DEQ A) to assess  
48 women's FOC levels, the previous study [12] was conducted by defining FOC using the  
49 International Classification of Diseases code O99.80, a code allocated to women who attended  
50 dedicated clinics for FOC using data from the Finnish Medical Birth Register to look at all  
51 singleton births during the period 1997 to 2010 (n=788, 317). Findings of this study concluded

52 that both nulliparous and multiparous women with FOC had an association with lower  
53 incidence of low birthweight, small for gestational age babies, preterm birth and low Apgar  
54 score at one minute [12]. While this study was large, the definition of FOC used in the study is  
55 a limitation, since it restricts the results to those who were diagnosed or who requested a  
56 Caesarean and were thus referred to phobia clinics and excluded those who attended primary  
57 care. It is possible that a true association was not captured due to an underestimation of the  
58 incidence of FOC using the ICD-10, thus using the W-DEQ A  $\geq 85$  is a more robust definition.

59 We hypothesise that severe FOC may have an adverse impact on pregnancy outcomes. Various  
60 factors may contribute to the possibility of adverse pregnancy outcomes in women with FOC.  
61 FOC may be associated with increased risk of Caesarean Section [13], unintended pregnancy,  
62 intimate partner violence [14] and a history of sexual abuse (adult or childhood) [15, 16]. Some  
63 evidence proposes there is a relationship between a history of childhood sexual abuse and  
64 preterm birth [17], and intimate partner violence has been correlated with low birthweight and  
65 preterm birth [18]. Moreover, unintended pregnancy could mean that women are less likely to  
66 have modified lifestyle behaviours such as smoking and alcohol consumption in early  
67 pregnancy, which are well-established as deleterious [19]. Therefore, the aim of this study was  
68 to compare the risk of adverse pregnancy outcomes for women with severe FOC as measured  
69 using W-DEQ A  $\geq 85$  during pregnancy compared to women with lower levels of FOC.

## 70 Materials and Methods

71 This was a prospective cohort study of 389 women recruited in a maternity unit in the Republic  
72 of Ireland. The study primary aims were to establish the prevalence and risk factors of FOC in  
73 an Irish context [20]. A convenience sample of women attending routine antenatal care were  
74 recruited by a research midwife undertaking doctoral studies, and by undergraduate students,  
75 who were trained by the midwife to recruit participants, in 2015 and 2016. Findings and full

76 recruitment details are published elsewhere [20]. Full ethics approval was obtained from the  
77 Cork Research Ethics Committee for the Teaching and Learning Hospitals [ECM 4 (06/01/15)  
78 and ECM 3 (03/03/15)].

79 Inclusion criteria were; pregnant women  $\geq 18$  years, 12-24 weeks' pregnant and booked to give  
80 birth in a large university-based tertiary maternity hospital (approximately 8,000 births  
81 annually). Exclusion criteria were; women who self-determined they had insufficient English  
82 to independently carry out the Questionnaire. Questionnaires were completed in clinics, after  
83 research assistants gained written informed consent. Women were invited to provide their  
84 medical records number to allow follow-up. Each woman completed a questionnaire including  
85 socio-demographic and obstetric questions and the W-DEQ A. The W-DEQ A [10] consists of  
86 33 questions using a Likert scale. A total score was calculated; with scores between 0 and 165  
87 possible, scores 0-65, low fear,  $\geq 66$ , moderate fear, and a score  $\geq 85$  defining severe FOC [4,  
88 10]. In Ireland at the time of the study, there were no phobia clinics available to women with  
89 FOC and a formal diagnosis of FOC would be unusual due to a lack of awareness of perinatal  
90 mental health [21].

91 Of 690 women invited to participate, 451 gave consent to postnatal data collection (65%).  
92 Women who had incomplete W-DEQ A scores (n=29), stillbirths (n=2) and miscarriages (n=1)  
93 were excluded due to incomplete datasets, and 21 women were lost to follow-up. For the final  
94 analysis we excluded twin pregnancies (n=9), limiting to singleton pregnancies, in order to  
95 increase homogeneity of the sample. Stillbirth was defined per the World Health Organisation  
96 (WHO) definition [22] as the birth at, or after 28 weeks gestation of a baby with no signs of  
97 life. Although there are various definitions of miscarriage, in this study, miscarriage was  
98 defined as spontaneous fetal loss, from conception to 24 completed weeks gestation [23]. The  
99 final study population consisted of 389 women.

100 Pregnancy outcome data were extracted from medical records by hand, directly from medical  
101 records where possible, or from delivery logbooks and e-health record (Maternal and New-  
102 born-Clinical Management System) as necessary in July 2017. Birthweight centiles were  
103 calculated using a customised centile calculator for Irish mothers [24]. Outcome data were  
104 entered into a secure encrypted SPSS file by the first author.

105 The following pregnancy outcomes were investigated for their association with severe FOC;  
106 birthweight in grams, birthweight centile, gestational age in days, and Apgar scores at 1 minute  
107 and 5 minutes.

108 Statistical analysis was performed using SPSS Version 22.0 Software programme (Chicago,  
109 USA). Continuous variables were tested for normality using histograms and box plots, and  
110 described using means and standard deviation (SD) if normally distributed, and median and  
111 interquartile range (IQR) if not normally distributed. Due to non-normal distribution of the  
112 data, a non-parametric technique (Kruskall-Wallis test) was used to test the hypothesis in  
113 relation to Apgar scores. Analyses were conducted separately for nulliparous and multiparous  
114 women to investigate outcomes in each group. A linear regression model was performed to  
115 investigate the relationship between antenatal experience of FOC and neonatal outcome  
116 (birthweight, birthweight centile, gestational age, and Apgar scores). Models were adjusted for  
117 potential confounding factors: maternal age (<35 years vs ≥35 years), marital status (partner  
118 vs no partner), smoking (smoker vs non-smoker) and parity (nulliparous vs multiparous).  
119 Results were reported using the mean difference and 95% confidence intervals (CIs). For the  
120 comparison of normally distributed continuous variables, the independent t-test was used and  
121 Mann-Whitney U Test was performed for non-normally distributed data. An overall  
122 significance level  $p \leq 0.05$  was considered to be statistically significant and  $p \leq 0.05$  also  
123 considered significant for individuals mean difference of each analysis.

124 Results

125 In the final cohort, eighteen women (4.6%) had W-DEQ A  $\geq 85$ , 103 (26.5%) women had W-  
126 DEQ A  $\geq 66$ , and 268 (68.9%) women had W-DEQ A  $\leq 65$ . Mean W-DEQ A score for the whole  
127 sample was 55.42 (SD= 18.43). Women under 25 years had the highest mean W-DEQ A score  
128 (60.53, SD=17.72). Married women had a lower mean W-DEQ A score (54.87, SD=18.37)  
129 when compared with single women (60.52, SD=18.49). Nulliparous women had a higher mean  
130 W-DEQ A score (59.17, SD=16.64) when compared with multiparous women (52.93,  
131 SD=19.73). There was no difference in mean W-DEQ A score in women with no pregnancy  
132 loss (55.67, SD= 17.96) versus those with one pregnancy loss (55.71, SD= 17.79). Women  
133 with two or more pregnancy loss had a slightly lower W-DEQ A score (53.24, SD=22.49).

134 The mean birthweight in the total sample was 3521g (SD=542.41), mean birthweight centile  
135 was 44.86 (SD=29.04), median gestational age was 279 days (IQR=12), median Apgar score  
136 at 1 minute were 9.00 (IQR=1) and Apgar score at 5 minutes were 10.00 (IQR= 1) (Table 1).  
137 In the exposure group (W-DEQ A $\geq 85$ ), birthweight, mean gestational age, Apgar score at 1  
138 minute and Apgar score at 5 minutes were similar overall (Table 1). There was an increase in  
139 the mean birthweight and birthweight centile for nulliparous women with severe FOC (n=7),  
140 3786g (SD=415.19), 45.59 (SD=24.39), in comparison with nulliparous women with low  
141 exposure 3386g (SD=562.08), 36.17, (SD=25.97), but the number of women in this group is  
142 too small to be reliable. Apgar score at 1 minute and Apgar score at 5 minutes were similar in  
143 all groups except the severe FOC group, which had a mean Apgar score at 1 minute of 8.11 and  
144 mean Apgar score at 5 minutes of 9.11. The results of the linear regression showed a significant  
145 correlation between the exposure (severe FOC) and Apgar scores at 1 minute (mean  
146 difference= -0.09 [95%CI -1.28, 0.32]) and Apgar scores at 5 minutes (mean difference= -0.18  
147 [95%CI: -1.16, 1.08]) when adjusted for possible confounders (Table 2).



148 When labour and delivery outcomes were compared for women with W-DEQ A $\geq$ 85 versus  
149 those with W-DEQ A 0-84, there was no statistical difference in use of epidural analgesia,  
150 induction of labour or Caesarean Section (Table 3).

## 151 Discussion

152 Overall, there was no evidence of an association between FOC and birthweight, birthweight  
153 centile, or gestational age. There was a statistically significant difference in relation to severe  
154 FOC and Apgar scores however, this association is not clinically relevant. This study rejects  
155 our hypothesis that there is an association between antenatal experience of severe FOC and  
156 adverse pregnancy outcomes.

157 One possible explanation of this finding that FOC may not be associated with negative  
158 outcomes is that women have increased opportunities during the second trimester to ask doctors  
159 and midwives questions, which may alleviate FOC and provide reassurance, rather than earlier  
160 on in pregnancy, when typically women have few antenatal appointments.

161 Only one previous study [12], to our knowledge investigated a relationship between FOC and  
162 pregnancy outcomes. Our study confirms the findings of this large population-based  
163 epidemiological study [12] conducted using the Finnish Medical Birth Register which found  
164 no relationship between severe FOC and pregnancy outcomes.

## 165 Strengths and Limitations

166 The main strength of the present study is that, to our knowledge, it is the first to investigate  
167 FOC and pregnancy outcomes using the W-DEQ A.

168 Data were complete for the majority of variables. Study limitations must be acknowledged.  
169 The W-DEQ A was measured once, in the second trimester, but FOC may be triggered at any  
170 point during pregnancy, thus a study which measured FOC in the first and/ or third trimester

171 may find different results. The study used a convenience sample which limits the  
172 generalizability of the findings. The sample consisted of mainly Caucasian women, therefore  
173 a study including a more heterogeneous sample or women with a different ethnicity may result  
174 in different findings. The analysis was not adjusted for potential confounding factors related to  
175 pregnancy complications or high risk pregnancy. It must be acknowledged that the number of  
176 women with severe FOC in the sample were small (n=18), therefore the study was not  
177 adequately powered which led to wide confidence intervals. However, the prevalence of  
178 women with FOC (4.3%) in this study is similar to the findings of previous studies in other  
179 countries which also found a prevalence of approximately 5% [7]. Finally, the Finnish study  
180 [12] reported other pregnancy outcomes which we did not, such as incidence of low birthweight  
181 (<2500g), and small for gestational age babies.

## 182 Conclusions

183 This study suggests maternal exposure to severe FOC in the second trimester of pregnancy has  
184 no adverse impact on birth weight, birth weight centile, and gestational age or Apgar scores.  
185 Findings of this study are reassuring and may be useful to inform women and clinicians, adding  
186 to our limited understanding of severe FOC in an Irish context, highlighting similarities  
187 between Finnish and Irish populations. Awareness of FOC in health care professionals is vital  
188 to integrate management of FOC in antenatal care and enhance emotional support for women,  
189 which may result in a reduction in medical interventions and Caesarean Section rates. Further  
190 research should focus on investigating pregnancy outcomes in other countries and in different  
191 ethnic groups. In addition, future studies should evaluate the pregnancy outcomes of women  
192 with FOC in the first or third trimester.

## 193 Conflicts of interests

194 The authors have no potential conflicts of interest to disclose. The authors have no financial  
195 relationships relevant to this article to disclose.

196 Acknowledgements

197 We are grateful to all the participants who made the study possible. The author would like to  
198 acknowledge the following students from the Dept. of Epidemiology and Public Health at  
199 University College Cork who worked on the project in 2015/2016; Kristina Mendelis, Nicola  
200 Kelleher, Eimear Carr, Stefanie Boehm, from Johannes Gutenberg University Mainz, as well  
201 as Dr Michelle McCarthy from the Dept. of Obstetrics and Gynaecology at University College  
202 Cork for their support.

203 Maeve O'Connell, Prof Louise Kenny, Dr Ali Khashan and Dr Patricia Leahy-Warren  
204 conceived and designed the study. Maeve O'Connell recruited participants, collected the data  
205 and performed the analysis with advice from Dr Ali Khashan. All authors read and contributed  
206 to the manuscript. This study was carried out as part of doctoral studies at The Irish Centre for  
207 Fetal and Neonatal Translational Research (INFANT Centre), which is supported by Science  
208 Foundation Ireland (grant no. 12/ RC/ 2272)

Table 1. Gestational age, birthweight, birthweight centile and Apgar scores and antenatal experience of fear of childbirth

Variable	Gestational Age, days median, IQR (n)			Birthweight, g mean, SD (n)			Birthweight centile mean, SD (n)			Apgar at 1 minute median, IQR (n)			Apgar at 5 minutes median, IQR (n)		
	Total Sample	Nulliparous women	Multiparous women	Total Sample	Nulliparous women	Multiparous women	Total Sample	Nulliparous women	Multiparous women	Total Sample	Nulliparous women	Multiparous women	Total Sample	Nulliparous women	Multiparous women
<b>Overall Sample W-DEQ 0-165</b>	279, 12 (389)	281, 14 (120*)	278, 10 (266)	3521± 542 (389)	3422±553(122)	3568± 532 (266)	45±29 (389)	36±6 (120*)	49±29 (265*)	9., 0 (389)	9, 0 (122)	9, 0 (264*)	10, 1 (389)	10, 1 (122)	10, 1 (264*)
<b>Low Exposure W-DEQ A 0-65</b>	278, 11 (268)	280, 15 (76*)	277, 10 (189)	3529± 562 (268)	3387±600 (78)	3590± 537(189)	46±29 (268)	36±26 (76*)	50±29 (189)	9, 0 (268)	9, 0 (78)	9, 0 (187*)	10, 1 (268)	10, 0 (78)	10, 1 (187*)
<b>Moderate Exposure W-DEQ A 66-84</b>	280, 11 (103)	284, 12 (37)	278, 11 (65*)	3492± 478 (103)	3427±448 (37)	3529± 494 (66)	40±29 (103)	33±26 (37)	44±30 (65*)	9, 0 (103)	9, 0 (37)	9, 0 (66)	10, 1 (103)	9, 1 (37)	10, 1 (66)
<b>Exposure W-DEQ A 85-165</b>	281, 16 (18)	285, 8 (7)	274, 26 (11)	3566± 609 (18)	3788±415 (7)	3425± 686 (11)	54±28 (18)	46±24 (7)	60±30 (11)	9, 1 (18)	9, 0 (7)	9, 1 (11)	9, 1 (18)	9, 1 (7)	9, 1 (11)

SD=Standard Deviation, W-DEQ A= Wijma Delivery Experience Questionnaire Part A, \*=missing data

Table 2. Results of linear regression predicting gestational age, birthweight, birthweight centile and Apgar score

Variable N	Gestational Age, days		Birthweight, g		Birthweight Centile		Apgar at 1 minute		Apgar at 5 minutes	
	Co-efficient (95%CI)		Co-efficient (95%CI)		Co-efficient (95%CI)		Co-efficient (95%CI)		Co-efficient (95%CI)	
	Adjusted	Unadjusted	Adjusted	Unadjusted	Adjusted	Unadjusted	Adjusted	Unadjusted	Adjusted	Unadjusted
Overall Sample (continuous) W-DEQA 0-165 N=389	0.04 (-0.09, 0.14)	0.06 (-0.03, 0.10)	0.12 (-1.94, 9.69)	0.02 (-2.39, 3.48)	0.11 (-0.11, 0.45)	-0.11 (-0.18, 0.14)	-0.04 (-0.01, 0.01)	-0.06 (-0.01, 0.01)	0.36 (-0.01, 0.01)	-0.09 (-0.01, 0.01)
Low Exposure W-DEQ A 0-65 n=268	-	-	-	-	-	-	-	-	-	-
Moderate Exposure W-DEQ A 66-84 n=103	0.08 (-2.23, 6.43)	0.11 (0.27, 5.54)	-0.09 (-305.61, 92.91)	-0.03 (-161.42, 83.87)	-0.14 (-19.68, 1.01)	-0.10 (-13.29, - 0.18)	0.06 (-0.27, 0.56)	0.04 (-0.15, 0.36)	-0.07 (-0.34, 0.14)	-0.02 (-0.18, 0.11)
Exposure W-DEQ A 85-165 n=18	-0.06 (-11.69, 4.82)	-0.06 (-8.75, 2.37)	-0.03 (-444.69, 315.82)	0.02 (-210.22, 305.13)	0.03 (-15.97, 23.53)	0.07 (-4.14, 23.51)	-0.09 (-1.28, 0.32)	-0.11 (-1.16, - 0.08)	-0.18 (-1.03, - 0.12)	-0.16 (-0.81, - 0.19)

**Table Legend:** W-DEQ A= Wijma Delivery Experience Questionnaire Part A  
Adjusted for Age, Marital Status, Parity, Smoking

Table 3. Comparison of labour and delivery outcomes of women with and without a severe fear of childbirth

<b>Labour and delivery outcome</b>	<b>W-DEQ A <math>\geq</math>85, n (%)</b>	<b>W-DEQ A <math>\leq</math>84, n (%)</b>	<b>p</b>
<b>Epidural analgesia</b>	7 ( 1.8)	140 ( 35.9)	0.39
<b>Induction of labour</b>	5 ( 1.3)	130 (33.4 )	0.57
<b>Pre-labour Caesarean</b>	5 ( 1.3)	44 ( 11.3)	0.06
<b>Caesarean in labour</b>	4 ( 1.0)	53 (13.6 )	0.31

**Table Legend:** W-DEQ A= Wijma Delivery Experience Questionnaire Part A  
p<0.05= significant

## 1 References

- 2 [1] I. Milosevic, R.E. McCabe, *Phobias: The Psychology of Irrational Fear*, Greenwood, California,  
3 2015.
- 4 [2] E. Rondung, J. Thomtén, Ö. Sundin, Psychological perspectives on fear of childbirth, *Journal of*  
5 *Anxiety Disorders* 44 (2016) 80-91.
- 6 [3] Y. Richens, D.T. Lavender, D.M. Smith, *Fear of Birth in Clinical Practice: A Structured Review*  
7 *of Current Measurement Tools*, Sexual & Reproductive Healthcare (2018).
- 8 [4] E.L. Ryding, B. Wijma, K. Wijma, H. Rydhstrom, Fear of childbirth during pregnancy may  
9 increase the risk of emergency cesarean section, *Acta Obstetrica et Gynecologica Scandinavica* 77  
10 (1998) 542-547.
- 11 [5] M. Lukasse, B. Schei, E.L. Ryding, G. Bidens Study, Prevalence and associated factors of fear of  
12 childbirth in six European countries, *Sexual and Reproductive Healthcare* 5(3) (2014) 99-106.
- 13 [6] H.L.R. Melender, Experiences of fears associated with pregnancy and childbirth: a study of 329  
14 pregnant women, *Birth* 29(2) (2002) 101-111.
- 15 [7] M.A. O'Connell, P. Leahy-Warren, A.S. Khashan, L.C. Kenny, S.M. O'Neill, Worldwide  
16 prevalence of tocophobia in pregnant women: systematic review and meta-analysis, *Acta Obstetrica*  
17 *et Gynecologica Scandinavica* 96(8) (2017) 907-920.
- 18 [8] S. Nath, E.G. Ryan, K. Trevillion, D. Bick, J. Demilew, J. Milgrom, A. Pickles, L.M. Howard,  
19 Prevalence and identification of anxiety disorders in pregnancy: the diagnostic accuracy of the two-  
20 item Generalised Anxiety Disorder scale (GAD-2), *BMJ open* 8(9) (2018) e023766.
- 21 [9] P. Sylvers, S.O. Lilienfeld, J.L. LaPrairie, Differences between trait fear and trait anxiety:  
22 Implications for psychopathology, *Clinical psychology review* 31(1) (2011) 122-137.
- 23 [10] K. Wijma, B. Wijma, M. Zar, Psychometric aspects of the W-DEQ; a new questionnaire for the  
24 measurement of fear of childbirth, *Journal of Psychosomatic Obstetrics & Gynecology* 19 (1998) 84-  
25 97.
- 26 [11] E. Calderani, L. Giardinelli, S. Scannerini, S. Arcabasso, E. Compagno, F. Petraglia, V.J.J.o.p.r.  
27 Ricca, Tocophobia in the DSM-5 era: Outcomes of a new cut-off analysis of the Wijma delivery  
28 expectancy/experience questionnaire based on clinical presentation, 116 (2019) 37-43.
- 29 [12] S. Raisanen, S. Lehto, H. Nielsen, M. Gissler, M. Kramer, S. Heinonen, Fear of childbirth in  
30 nulliparous and multiparous women: a population-based analysis of all singleton births in Finland in  
31 1997-2010, *BJOG : an international journal of obstetrics and gynaecology* (2014).
- 32 [13] E.L. Ryding, M. Lukasse, A. Van Parys, A.M. Wangel, H. Karro, H. Kristjansdottir, A.M.  
33 Schroll, B. Schei, *Fear of Childbirth and Risk of Cesarean Delivery: A Cohort Study in Six European*  
34 *Countries*, *Birth* 42(1) (2015).
- 35 [14] E. Miller, M.R. Decker, H.L. McCauley, D.J. Tancredi, R.R. Levenson, J. Waldman, P.  
36 Schoenwald, J.G.J.C. Silverman, Pregnancy coercion, intimate partner violence and unintended  
37 pregnancy, *Contraception* 81(4) (2010) 316-322.
- 38 [15] R. Heimstad, R. Dahloe, I. Laache, E. Skogvoll, B. Schei, Fear of childbirth and history of abuse:  
39 implications for pregnancy and delivery, *Acta Obstetrica et Gynecologica Scandinavica* 85(4) (2006)  
40 435-40.
- 41 [16] B. Leeners, G. Görres, E. Block, M.P. Hengartner, Birth experiences in adult women with a  
42 history of childhood sexual abuse, *Journal of Psychosomatic Research* 83 (2016) 27-32.
- 43 [17] B. Leeners, R. Stiller, E. Block, G. Görres, W. Rath, Pregnancy complications in women with  
44 childhood sexual abuse experiences, *Journal of Psychosomatic Research* 69(5) (2010) 503-510.
- 45 [18] Y. Neggers, R. Goldenberg, S. Cliver, J. Hauth, Effects of domestic violence on preterm birth  
46 and low birth weight, *Acta Obstetrica et Gynecologica Scandinavica* 83(5) (2004) 455-460.
- 47 [19] L.M. O'Keeffe, D.L. Dahly, M. Murphy, R.A. Greene, J.M. Harrington, P. Corcoran, P.M.  
48 Kearney, Positive lifestyle changes around the time of pregnancy: a cross-sectional study, *BMJ Open*  
49 6(5) (2016) e010233.
- 50 [20] M.A. O'Connell, Leahy-Warren, P., Kenny, L., O'Neill, S.M., Khashan, A.S., The prevalence and  
51 risk factors of fear of childbirth among pregnant women: a cross sectional study in Ireland, *Acta*  
52 *Obstetrica et Gynecologica Scandinavica* (2019).

- 53 [21] H.S. Executive, Specialist Perinatal Mental Health Services: Model Of Care for Ireland, in:  
54 H.M.H. Services (Ed.) HSE, Dublin, 2017.
- 55 [22] W.H. Organisation, Stillbirth, Maternal, newborn, child and adolescent health Geneva, 2016.
- 56 [23] R.C.o.O.G. (RCOG), The investigation and treatment of couples with recurrent first-trimester  
57 and second-trimester miscarriage, Green-top Guideline No 17, 2011.
- 58 [24] GROW, Customised Centile Calculator, 2018. [www.gestation.net](http://www.gestation.net).

59