

Title	Lung cancer in young patients: tumour characteristics and treatment in an Irish population
Authors	Cullivan, Sarah;Ní Mhaolcatha, Sarah;Henry, Michael T.;Muhammad, Nazri Kaderi;Mullally, William;Bambury, Richard M.;Burke, Louise;Kennedy, Marcus P.
Publication date	2019-08
Original Citation	Cullivan, S., Ní Mhaolcatha, S., Henry, M. T., Muhammad, N. K., Mullally, W., Bambury, R. M., Burke, L. and Kennedy, M. P. [2019] 'Lung cancer in young patients: tumour characteristics and treatment in an Irish population', <i>Translational Cancer Research</i> , 8(4), pp. 1109-1115. doi: 10.21037/tcr.2019.06.22
Type of publication	Article (peer-reviewed)
Link to publisher's version	<a href="http://tcr.amegroups.com/article/view/29925">http://tcr.amegroups.com/article/view/29925</a> - 10.21037/tcr.2019.06.22
Rights	© Translational Cancer Research. All rights reserved.
Download date	2024-04-24 08:04:26
Item downloaded from	<a href="https://hdl.handle.net/10468/8490">https://hdl.handle.net/10468/8490</a>

# Lung cancer in young patients: tumour characteristics and treatment in an Irish population

Sarah Cullivan<sup>1</sup>, Sarah Ni Mhaolcatha<sup>2</sup>, Michael T. Henry<sup>1</sup>, Nazri Kaderi Muhammad<sup>1</sup>, William Mullally<sup>3</sup>, Richard M. Bambury<sup>3</sup>, Louise Burke<sup>2</sup>, Marcus P. Kennedy<sup>1</sup>

<sup>1</sup>Department of Respiratory Medicine, <sup>2</sup>Department of Histopathology, <sup>3</sup>Department of Medical Oncology, Cork University Hospital, Cork, Ireland

**Contributions:** (I) Conception and design: S Cullivan, NK Muhammad, MP Kennedy; (II) Administrative support: All authors; (III) Provision of study materials or patients: All authors; (IV) Collection and assembly of data: S Cullivan, NK Muhammad; (V) Data analysis and interpretation: All authors; (VI) Manuscript writing: All authors; (VII) Final approval of manuscript: All authors.

**Correspondence to:** Dr. Sarah Cullivan, MB, BAO, BCh, MRCPI. Cork University Hospital, Wilton, Cork, Ireland. Email: sarah.cullivan@ucdconnect.ie.

**Background:** Lung cancer is the leading cause of cancer death in both sexes in Ireland. Studies suggest that lung cancer in younger patients has distinct characteristics. The aim of this study is to define the characteristics of lung cancer in patients 55-year-old or younger in an Irish population.

**Methods:** Data was collected retrospectively from local medical records and the hospital electronic database regarding all patients diagnosed with lung cancer aged 55-year-old and younger, from 2010–2016. Information regarding patient demographics, smoking status, tumour histology, molecular analysis, stage and location, diagnostic modality and initial treatment choice was collected. In all cases the diagnosis of lung cancer was confirmed at the regional lung cancer multidisciplinary team (MDT) meeting.

**Results:** In total, 8% (n=130) of all cases of lung cancer diagnosed from 2010 to 2016 in our center occurred in patients aged 55 years old or younger; 83% (n=108) were 45 to 55-year-old, 15% (n=19) were 35 to 44-year-old and 2% (n=3) were younger than 35-year-old; 88% (n=115) of patients reported a smoking history. There was a female preponderance (58%, n=76), higher rates of NSCLC non-squamous subtype (53%, n=69) and an upper lobe predominance (42%, n=54); 53% (n=68) of patients had IV or extensive disease at presentation. Epidermal growth factor receptor (EGFR), anaplastic lymphoma kinase (ALK) and Kirsten rat sarcoma oncogene (KRAS) mutation rates were 9% (n=4) and 3% (n=1) and 80% (n=4) respectively.

**Conclusions:** Lung cancer in younger patients has distinct characteristics. This study suggests a female preponderance, high smoking rates and a predilection for the upper lobes. Further large-scale multicenter studies are required to verify these results and to clarify the responsible mechanisms.

**Keywords:** Lung cancer; younger patients; upper lobe

Submitted Dec 12, 2018. Accepted for publication May 28, 2019.

doi: 10.21037/tcr.2019.06.22

View this article at: <http://dx.doi.org/10.21037/tcr.2019.06.22>

## Introduction

Lung cancer is one of the leading causes of cancer mortality worldwide. While most cases of lung cancer occur in patients of advanced age, there is an important subgroup of younger patients who are affected by the condition. Studies suggest that lung cancer in younger patients has distinguishable characteristics, particularly regarding patient gender, smoking status and tumour morphology (1-4).

National and international data demonstrate increasing rates of lung cancer in women, while the incidence in men continues to fall. There is evidence to suggest that this trend is replicated in younger patients (5,6). The exact mechanism for this is unknown, however this highlights important questions regarding gender specific lung cancer risk factors, which could have important implications for risk stratification and potential screening. Furthermore

**Table 1** Baseline characteristics

Baseline characteristics	N (%)
Patients, n (%)	130 (100%)
Age (year)	
45–55	108 (83%)
35–44	19 (15%)
<35	3 (2%)
Male, n (%)	54 (42%)
Smoking history (current or former)	115 (88%)

higher rates of adenocarcinoma have been demonstrated in younger patients, irrespective of gender, which has important treatment consequences (1–4). More patients are receiving lung cancer directed treatment, and therapeutic options are expanding in light of recent advances in lung cancer molecular diagnostics and treatments. These provide exciting opportunities to explore such therapies in a younger cohort. This retrospective single center study was designed to define the characteristics of lung cancer in patients 55-year-old or younger in an Irish population.

## Methods

This is a retrospective, single center study of patients aged 55 years or younger with a confirmed diagnosis of lung cancer. With institutional approval, data was collected from local medical records and the hospital electronic database regarding patient age at lung cancer diagnosis, gender, smoking history, diagnostic modality, stage, tumour histological subtype and location and initial treatment choice, from January 2010 until December 2016. Patients with a diagnosis other than lung cancer or age greater than 55 years old were excluded from the study. Selected patients were subdivided into three categories according to age; <35 years, 35–44 years and 45–55 years old. The diagnosis of lung cancer was based on clinical, radiological and histological data and multidisciplinary team (MDT) discussion at the regional lung cancer MDT meeting in Cork University Hospital, Ireland.

Tumour histology was divided into subgroups of non-small cell lung cancer (NSCLC) non squamous, NSCLC squamous, NSCLC not otherwise specified (NOS), small cell lung cancer and carcinoid (typical and atypical). The international TNM (tumour-node-metastasis) system, seventh edition, was used to define tumour stage and

molecular test results were noted when assessed.

Statistical analyses were performed on combined age groups using 2×2 contingency tables in GraphPad software to compare clinicopathological features. In all analysis a P value less than 0.05 was considered statistically significant.

## Results

### Demographics

A total of 1,635 patients were diagnosed with lung cancer from January 2010 to December 2016 in Cork University Hospital. A total of 130 patients met the inclusion criteria and were selected for further analysis. Patient characteristics are depicted in *Table 1*. There was no significant difference between gender related smoking rates, as 87% (n=47) of males and 89% (n=68) of females were current or former smokers (P=0.7289).

### Diagnostic modality

All patients (n=130) had a CT thorax as part of the diagnostic workup. Cytopathology and histopathology specimens were obtained using many modalities, including bronchoscopic endobronchial and transbronchial biopsy (27%), endobronchial ultrasound guided transbronchial needle aspiration (20%), CT guided lung biopsy (15%), surgical biopsy (15%) and biopsy of distant metastases (16%) An alternative method was used in 7% (n=9) (*Table 2*).

### Tumour location

Regarding tumour location, the majority of tumours were located in the upper lobes, as 23% (n=30) were located in the right upper lobe and 19% (n=24) in the left upper lobe. The remaining tumours were located in hilar region (22%), lower lobes (19%) and other pulmonary locations in 17% (*Table 2*).

### Tumour histology and stage

NSCLC non squamous was diagnosed in 53% (n=69) of cancers, NSCLC squamous in 15% (n=19) and NSCLC NOS in a further 7% (n=9). Small cell lung cancer was diagnosed in 14% (n=18) and carcinoid in 11% (n=15) (*Table 2*). Regarding tumour stage, 41% (n=53) of patients had stage IV disease and 12% (n=15) had extensive small cell carcinoma at diagnosis (*Table 2*). All histological

**Table 2** Describing diagnostic modality, tumour location, tumour histology and stage

Results	N (%)
<b>Diagnostic modality</b>	
Bronchoscopy	35 (27%)
Endobronchial ultrasound	26 (20%)
CT guided biopsy	20 (15%)
Surgery	19 (15%)
Biopsy of distant metastases	21 (16%)
Other	9 (7%)
<b>Tumour location</b>	
Upper lobes	54 (42%)
Middle lobe & lingual	12 (9%)
Lower lobes	25 (19%)
Hilar region	29 (22%)
Mediastinal	3 (2%)
Endobronchial	5 (4%)
Unknown	2 (2%)
<b>Histology</b>	
NSCLC non squamous	69 (53%)
Adenocarcinoma	64 (49%)
Large-cell carcinoma	4 (3%)
Pleomorphic carcinoma	1 (1%)
NSCLC squamous	19 (15%)
Small cell lung cancer	18 (14%)
Carcinoid	15 (11%)
NSCLC NOS	9 (7%)
<b>Tumour stage</b>	
NSCLC & carcinoid	
Stage I	32 (25%)
Stage II	7 (5%)
Stage III	20 (15%)
Stage IV	53 (41%)
Small cell carcinoma	
Limited	3 (2%)
Extensive	15 (12%)

NSCLC, non small cell lung cancer.

subtypes increased with increasing patient age ( $P=0.0536$ ) (*Figure 1A*). *Figure 1B* depicts tumour histology according to patient gender and demonstrates higher numbers of NSCLC and small cell lung cancer in young female patients relative to their male counterparts, which did not meet statistical significance ( $P=0.6077$ ).


Furthermore, epidermal growth factor receptor (EGFR), anaplastic lymphoma kinase (ALK), Kirsten rat sarcoma oncogene (KRAS) and c-ros oncogene 1 (ROS1) mutations were assessed in individual cases, guided by lung cancer MDT recommendations. Molecular analysis was performed in 47 cases (36%) and all positive cases were mutually exclusive. EGFR mutations were identified in 4 (9%) of 47 cases and an ALK translocation was identified in 1 (3%) of 30 samples. In Ireland assessment of KRAS mutations and PDL-1 expression are only recommended in cases of advanced NSCLC, with no evidence of EGFR or ALK mutations KRAS mutations in 4 (80%) of 5 analysed specimens (*Table 3*). Expression of programmed death-ligand 1 (PDL-1) was assessed in 6 cases. High PDL-1 (>50%) expression was present in 1 case (16.7%) and low expression (1–50%) in a further case (16.7%). The remaining 4 cases tested for PDL-1 expression were negative (66.7%) (*Table 4*).

### Treatment

The most common initial treatment for stage I–II disease was surgery ( $n=33$ , 25%) and for stage III ( $n=7$ , 5%) and IV ( $n=31$ , 24%) were chemotherapy. Patients with small cell lung cancer were typically treated initially with combined chemotherapy and radiation therapy or chemotherapy alone ( $n=17$ , 13%) (*Table 5*).

### Discussion

Lung cancer is one of the leading causes of cancer mortality worldwide and lung cancer in the young can be a devastating, life-limiting disease. Research in this cohort is currently limited by certain factors, including ambiguity regarding the precise definition of ‘young’ and the small and frequently retrospective nature of these studies. Nonetheless, this multicenter retrospective study highlights some interesting characteristics in this population and to our knowledge, is the first study in an Irish population describing lung cancer in patients aged 55-year-old or younger.



**Figure 1** Illustrates lung cancer characteristics in young patients with regarding to patient age and gender. (A) demonstrates tumour histology according to patient age. All histological subtypes increased with increasing patient age. This did not meet statistical significance ( $P=0.0536$ ). (B) shows tumour histology according to patient gender. Higher numbers of NSCLC and small cell lung cancer were demonstrated in young female patients relative to their male counterparts, which did not meet statistical significance ( $P=0.6077$ ). NSCLC, non small cell lung cancer.

**Table 3** Results of molecular testing

Molecular testing	Mutation detected, N (%)	Mutation not detected, N (%)	Inadequate sample, N (%)
EGFR (n=47)	4 (9%)	40 (85%)	3 (6%)
ALK (n=30)	1 (3%)	27 (90%)	2 (7%)
KRAS (n=5)	4 (80%)	1 (20%)	0 (0%)
ROS-1 (n=3)	0 (0%)	2 (67%)	1 (33%)

EGFR, epidermal growth factor receptor; ALK, anaplastic lymphoma kinase; KRAS, Kirsten rat sarcoma oncogene; ROS1, c-ros oncogene 1.

**Table 4** Results of PDL-1 assessment

PDL-1 expression	Low <50%	High >50%	None
PDL-1 (n=6)	1 (16.7%)	1 (16.7%)	4 (66.7%)

PDL-1, programmed death-ligand 1.

At present marked heterogeneity exists regarding the definition of ‘young’ lung cancer patients in the literature; for example Rocha *et al.* [1994] selected a threshold of 40-year-old, Kreuzer *et al.* [1998] chose 45 years and Franceschini *et al.* [2017] adopted a threshold of 55 years to define ‘young’ (1-3). The rationale behind these choices are frequently unclear. Arnold *et al.* [2016] selected an age threshold in their study by calculating the age two standard deviations below the mean age of lung cancer diagnosis in their population (4), which may be an appropriate, population centered approach. Clarification of this definition is essential

to guide the design of future studies and to facilitate subsequent comparisons. The National Cancer Registry Ireland (NCRI) 2011 report noted that 9% of patients diagnosed with lung cancer between 1994 and 2008 were less than 55-year-old and therefore a threshold of 55-year-old was chosen in our study in an Irish population (7).

This study describes interesting characteristics regarding gender, tumour location and histology. Firstly this study suggests a female preponderance (59%, n=76). This is concordant with Irish and international data, demonstrating rising incident rates of lung cancer in women. Specifically, the National Cancer Registry Ireland (NCRI) reported declining incident rates of lung cancer in men (-0.8% per year) and rising rates in women (+2.0% per year) from 1994 to 2013 (5). A similar trend was noted by Jemal *et al.* in American population, which reported higher incidence rates of lung cancer among women (6). Increased smoking rates in women partly explain this trend and our study

**Table 5** Initial treatment according to patient stage

Treatment	Stage I	Stage II	Stage III	Stage IV
NSCLC and carcinoid (n=112, 86%)	32 (25%)	7 (5%)	20 (15%)	53 (41%)
Surgery	30 (23%)	3 (2%)	4 (3%)	2 (1.5%)
Chemotherapy	1 (1%)	4 (3%)	7 (5%)	31 (24%)
Radiotherapy	0 (0%)	0 (0%)	3 (2%)	5 (4%)
Chemoradiotherapy	0 (0%)	0 (0%)	5 (4%)	13 (10%)
Supportive care	0 (0%)	0 (0%)	1 (1%)	2 (1.5%)
Other	1 (1%)	0 (0%)	0 (0%)	0 (0%)
Small cell lung cancer (n=18, 14%)	Limited (n=3)		Extensive (n=15)	
Chemotherapy	1 (1%)		12 (9%)	
Chemoradiotherapy	2 (1.5%)		2 (1.5%)	
Supportive care	0 (0%)		1 (1%)	

NSCLC, non small cell lung cancer.

emphasizes the persistent relevance of this risk factor, as 88% (n=115) of young lung cancer patients were current or former smokers. Authors such as Campling *et al.* [2019] have demonstrated significant age differences between current and former smokers (8), suggesting that ongoing tobacco exposure may accelerate the propensity to develop lung cancer in continuing smokers. This finding was not replicated in our study, as the median age at lung cancer diagnosis was 52 and 53-year-old in current and ex-smokers respectively (P=0.5617). While these findings reinforce the value of sustained anti-tobacco measures, this difference in lung cancer incident rates is not fully explained by sex difference in smoking behavior. Further research into sex specific risk factors for lung cancer, including susceptibility to tobacco carcinogens, should be pursued to establish precise mechanisms (6,9).

Additionally tumour location is an important factor to consider in the assessment of suspected lung cancer cases. The 42% (n=54) of lung cancers in this study were identified in the upper lobes. The American College of Chest Physicians and The British Thoracic Society have identified upper lobe location as an independent predictor of malignancy for solitary pulmonary nodules (10,11). Some studies suggest that this association is potentially more marked in younger patients (12,13), however this finding is inconsistent (14).

Tumour histology and stage have important treatment and prognostic implications. Unfortunately the diagnosis

of lung cancer is occasionally delayed in younger patients, as the clinical suspicion for lung cancer is often lower. This was highlighted by Bryant *et al.* [2008] and Dell'Amore *et al.* [2015] as younger patients were frequently referred to thoracic surgeons later than older patients (15,16). In our study the diagnosis of lung cancer was based on pathological specimens in all cases. This presumably reflects a favorable performance status in younger patients due to fewer comorbidities and a possible tendency to investigate these patients more aggressively due to their age. This contrasts practice regarding the entire lung cancer population in our center, as in 2016, 15% (n=53) of all lung cancer patients did not undergo invasive tests due to a poor performance status or patient preference and therefore received a clinical diagnoses of lung cancer only (17).

Multiple studies have demonstrated a predominance of adenocarcinoma in young lung cancer patients and a tendency to obtain a diagnosis at an advanced stage (18-21). These findings are replicated in our study, as adenocarcinoma was the dominant subtype (46%, n=64) and 53% (n=68) were diagnosed with stage IV NSCLC or extensive small cell lung cancer at presentation. This poses interesting questions regarding lung cancer screening in younger patients. Current evidence, namely the National Lung Cancer Screening Trial and the NELSON trial suggest a role for lung cancer screening in high-risk groups, which are frequently identified as 'older', with an associated high pack year history (22,23). This model has the potential

to neglect 'high risk' younger patients.

Molecular analysis was performed in 36% (n=47) of cases in this study. These rates are comparable to recent data published by Bigay-Gamé *et al.* [2018] in a French cohort of young lung cancer patients, as only one third of their population underwent molecular profiling (18), reflecting the non-mandatory nature of biological profiling in both countries during this period and restricted therapeutic options. For example, in Ireland assessment of KRAS mutations and PDL-1 expression are only recommended in cases of advanced NSCLC, with no evidence of EGFR or ALK mutations. Interestingly, a recent multicenter study exploring the incidence of molecular mutations in an Irish population noted a KRAS gene mutation incidence of 41.6% in their entire lung cancer cohort (24). Therefore further data regarding molecular diagnostics, PD-L1 expression and tumour mutational burden in young lung cancer patients is not only desirable but essential and could have meaningful clinical implications (25).

Given the retrospective nature of this study, documentation of specific patient risk factors such as family history and illicit drug use was incomplete and therefore not included. This is a possible limitation, as prior studies have highlighted the potential importance of these risk factors in this cohort (18,26). A further limitation is the lack of a control group of older patients to facilitate direct comparison within our populations. Finally, molecular analysis was performed in approximately one third of cases and further information regarding biological profiling should be the focus of future studies in this growing cohort of younger patients

### Summary

This study reinforces existing data, which demonstrates distinct characteristics in young lung cancer patients, including a female preponderance, high rates of adenocarcinoma and frequent advanced stage at diagnosis. Further large scale multicenter studies are required to verify these results and to delineate the responsible mechanisms. It is clear from this data that we should continue to aggressively address smoking behavior, while simultaneously exploring these findings.

### Acknowledgments

None.

### Footnote

*Conflicts of Interest:* The authors have no conflicts of interest to declare.

*Ethical Statement:* The study was approved by Cork Research Ethics Committee [No. ECM 6 (q) 01/03/16]. The authors are accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

### References

1. Rocha MP, Fraire AE, Guntupalli KK, et al. Lung cancer in the young. *Cancer Detect Prev* 1994;18:349-55.
2. Kreuzer M, Kreienbrock L, Gerken M, et al. Risk factors for lung cancer in young adults. *Am J Epidemiol* 1998;147:1028-37.
3. Franceschini JP, Jamnik S, Lopes Santoro I. Survival in a cohort of patients with lung cancer: the role of age and gender in prognosis. *J Bras Pneumol* 2017;43:431-6.
4. Arnold BN, Thomas DC, Rosen JE, et al. Lung Cancer in the Very Young : Treatment and Survival in the National Cancer Data Base. *J Thorac Oncol* 2016;11:1121-31.
5. National Cancer Registry. 2015. Cancer in Ireland 1994-2013: Annual Report of the National Cancer Registry. NCR, Cork, Ireland.
6. Jemal A, Miller KD, Ma J, et al. Higher Lung Cancer Incidence in Young Women Than Young Men in the United States. *N Engl J Med* 2018;378:1999-2009.
7. National Cancer Registry. 2011. Cancer in Ireland 1994-2008: Annual Report of the National Cancer Registry. NCR, Cork, Ireland.
8. Campling BG, Ye Z, Lai Y, et al. Disparity in age at lung cancer diagnosis between current and former smokers. *J Cancer Res Clin Oncol* 2019;145:1243-51.
9. Tobacco and Inequities: Guidance for addressing inequities in tobacco related harm. World Health Organization 2014. Available online: [http://www.euro.who.int/\\_\\_data/assets/pdf\\_file/0005/247640/tobacco-090514.pdf](http://www.euro.who.int/__data/assets/pdf_file/0005/247640/tobacco-090514.pdf). Sept 2018.
10. Callister ME, Baldwin DR, Akram AR, et al. British Thoracic Society guidelines for the investigation and management of pulmonary nodules. *Thorax* 2015;70:ii1-ii54.
11. Gould MK, Donington J, Lynch WR, et al. Evaluation of individuals with pulmonary nodules: when is it lung

- cancer? Diagnosis and management of lung cancer, 3rd ed: American College of Chest Physicians evidencebased clinical practice guidelines. *Chest* 2013;143:e93S-e120S.
12. Byers TE, Vena JE, Rzepka TF. Predilection of Lung Cancer for the Upper Lobes: An Epidemiologic Inquiry, *J Natl Cancer Inst* 1984;72:1271-5.
  13. Kaneda H, Uemura Y, Nakano T, et al. Lesions in Patients with Multifocal Adenocarcinoma Are More Frequently in the Right Upper Lobes. *Interact Cardiovasc Thorac Surg* 2012;15:627-32.
  14. Thomas A, Chen Y, Yu T, et al. Trends and Characteristics of Young Non-Small Cell Lung Cancer Patients in the United States. *Front Oncol* 2015;5:113.
  15. Bryant AS, Cerfolio RJ. Differences in outcomes between younger and older patients with non-small cell lung cancer. *Ann Thorac Surg* 2008;85:1735-9.
  16. Dell'Amore A, Monteverde M, Martucci N, et al. Surgery for non-small cell lung cancer in younger patients: what are the differences? *Heart Lung Circ* 2015;24:62-8.
  17. Lung cancer service key performance indicators annual report 2016, regional cancer center south, Cork University. 2016.
  18. Bigay-Gamé L, Bota S, Greillier L, et al. Characteristics of Lung Cancer in Patients Younger than 40 Years: A Prospective Multicenter Analysis in France. *Oncology* 2018;95:337-43.
  19. Maruyama R, Yoshino I, Yohena T, et al. Lung cancer in patients younger than 40 years of age. *J Surg Oncol* 2001;77:208-12.
  20. Skarin AT, Herbst RS, Leong TL, et al. Lung cancer in patients under age 40. *Lung Cancer* 2001;32:255-64.
  21. Ramalingam S, Pawlish K, Gadgeel S, et al. Lung cancer in young patients: analysis of a Surveillance, Epidemiology, and End Results database. *J Clin Oncol* 1998;16:651-7.
  22. National Lung Screening Trial Research Team, Aberle DR, Adams AM, et al. Reduced lung-cancer mortality with low-dose computed tomographic screening. *N Engl J Med* 2011;365:395-409.
  23. Wille MM, Dirksen A, Ashraf H, et al. Results of the randomized Danish lung cancer screening trial with focus on high-risk profiling. *Am J Respir Crit Care Med* 2016;193:542-51.
  24. Mullally W, Kee W, Dennehy C, et al. Real-world practice for non-small cell lung cancer with targetable genetic alterations in an Irish setting 2018. Abstract no 128. EORTC/AACR/NCI Conference PB-079.
  25. Román M, Baraibar I, Lopez I, et al. KRAS oncogene in non-small cell lung cancer: clinical perspectives on the treatment of an old target. *Mol Cancer* 2018;17:33.
  26. Radzikowska E, Roszkowski K, Glaz P. Lung cancer in patients under 50 years old. *Lung Cancer* 2001;33:203-11.

**Cite this article as:** Cullivan S, Ni Mhaolcatha S, Henry MT, Muhammad NK, Mullally W, Bambury RM, Burke L, Kennedy MP. Lung cancer in young patients: tumour characteristics and treatment in an Irish population. *Transl Cancer Res* 2019. doi: 10.21037/tcr.2019.06.22