

**UCC Library and UCC researchers have made this item openly available.
Please [let us know](#) how this has helped you. Thanks!**

Title	Social, biological, behavioural and psychological factors related to physical activity during early pregnancy in the Screening for Pregnancy Endpoints (Cork, Ireland) cohort study
Author(s)	Flannery, Caragh; Dahly, Darren L.; Byrne, Molly; Khashan, Ali S.; McHugh, Sheena; Kenny, Louise C.; McAuliffe, Fionnuala; Kearney, Patricia M.
Publication date	2019-06-20
Original citation	Flannery, C., Dahly, D., Byrne, M., Khashan, A., McHugh, S., Kenny, L. C., McAuliffe, F. and Kearney, P. M. (2019) 'Social, biological, behavioural and psychological factors related to physical activity during early pregnancy in the Screening for Pregnancy Endpoints (Cork, Ireland) cohort study', <i>BMJ Open</i> , 9(6), e025003. (9pp.) DOI: 10.1136/bmjopen-2018-025003
Type of publication	Article (peer-reviewed)
Link to publisher's version	https://bmjopen.bmj.com/content/9/6/e025003 http://dx.doi.org/10.1136/bmjopen-2018-025003 Access to the full text of the published version may require a subscription.
Rights	©Author(s) (or their employer(s)) 2019. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ. This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: http://creativecommons.org/licenses/by-nc/4.0/ . https://creativecommons.org/licenses/by-nc/4.0/
Item downloaded from	http://hdl.handle.net/10468/8501




Downloaded on 2021-09-18T10:58:11Z

UCC


Supplementary file 4: Model fit statistics and three class model definitions

Latent class models were estimated using Mplus version 8.0. The model allowed the two thresholds for each of the 3-level response variables be freely estimated within each specified latent class. Maximum likelihood estimated was employed, using 10k random starts and 1k final stage optimizations. The best observed final stage log likelihood was consistently repeated across random starts. The Mplus code used to estimate the models, as well as additional information on model fit and information criteria and a graphical summary of the three latent classes we focused on, are given below.

a. Model fit statistics



b. Response probability for the three classes



Class 3 High Physical Activity Level: Active group (majority high level moderate exercises and walkers with some mid/high level vigorous exercisers)

Class 2 Moderate Physical Activity Level: Moderately active exercisers (majority mid-level moderate exercisers and walkers)

Class 1 Low Physical Activity Level: Predominantly low exercisers (with some mid/high level walkers)

c. Mplus code

```
Names =
regid vigorous moderate rec_walk;
Missing are . ;
Usevariables =
regid vigorous moderate rec_walk;
Categorical =
    vigorous moderate rec_walk;
Idvariable = regid;
Classes = c(3);
ANALYSIS: Type    = Mixture;
           Starts = 10000 1000;
           processors = 3;

MODEL:
%OVERALL%
%c#1%
[vigorous$1* vigorous$2*];
[moderate$1* moderate$2*];
[rec_walk$1* rec_walk$2*];
%c#2%
[vigorous$1* vigorous$2*];
[moderate$1* moderate$2*];
[rec_walk$1* rec_walk$2*];
%c#3%
[vigorous$1* vigorous$2*];
[moderate$1* moderate$2*];
[rec_walk$1* rec_walk$2*];
```
