

**UCC Library and UCC researchers have made this item openly available.
Please [let us know](#) how this has helped you. Thanks!**

Title	Role of the Photorhabdus Dam methyltransferase during interactions with its invertebrate hosts
Author(s)	Payelleville, Amaury; Blackburn, Dana; Lanois, Anne; Pagès, Sylvie; Cambon, Marine C.; Ginibre, Nadege; Clarke, David J.; Givaudan, Alain; Brillard, Julien
Publication date	2019-10-09
Original citation	Payelleville, A., Blackburn, D., Lanois, A., Pagès, S., Cambon, M. C., Ginibre, N., Clarke, D. J., Givaudan, A. and Brillard, J. (2019) 'Role of the Photorhabdus Dam methyltransferase during interactions with its invertebrate hosts', PLOS ONE, 14(10), e0212655. (14pp.) DOI: 10.1371/journal.pone.0212655
Type of publication	Article (peer-reviewed)
Link to publisher's version	https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0212655 http://dx.doi.org/10.1371/journal.pone.0212655 Access to the full text of the published version may require a subscription.
Rights	©2019 Payelleville et al. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited http://creativecommons.org/licenses/by/4.0/
Item downloaded from	http://hdl.handle.net/10468/8872

Downloaded on 2020-10-30T17:55:13Z

Fig. S2




Figure S2. Motility and pathogenicity of *Chr_dam* strain.

(A) Violin-plot of motility halo size for *Chr_dam* and *Chr_gfp* strain after 36 hours of growth on motility medium. The difference between the two strains was significant (Wilcoxon test, p -value <0.001). (B) Survival of *S. littoralis* larvae after injection of 10^4 CFU of *Chr_gfp* (green) or *Chr_dam* (blue). *Chr_dam* strain was significantly delayed (2hours) in the time needed to kill 50 % of the larvae (Wilcoxon test, p -value <0.001).