

UCC Library and UCC researchers have made this item openly available. Please [let us know](#) how this has helped you. Thanks!

Title	Role of the Photorhabdus Dam methyltransferase during interactions with its invertebrate hosts
Author(s)	Payelleville, Amaury; Blackburn, Dana; Lanois, Anne; Pagès, Sylvie; Cambon, Marine C.; Ginibre, Nadege; Clarke, David J.; Givaudan, Alain; Brillard, Julien
Publication date	2019-10-09
Original citation	Payelleville, A., Blackburn, D., Lanois, A., Pagès, S., Cambon, M. C., Ginibre, N., Clarke, D. J., Givaudan, A. and Brillard, J. (2019) 'Role of the Photorhabdus Dam methyltransferase during interactions with its invertebrate hosts', PLOS ONE, 14(10), e0212655. (14pp.) DOI: 10.1371/journal.pone.0212655
Type of publication	Article (peer-reviewed)
Link to publisher's version	https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0212655 http://dx.doi.org/10.1371/journal.pone.0212655 Access to the full text of the published version may require a subscription.
Rights	©2019 Payelleville et al. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited http://creativecommons.org/licenses/by/4.0/
Item downloaded from	http://hdl.handle.net/10468/8872

Downloaded on 2020-10-30T15:56:22Z

Fig. S3




Figure S3. Emerging IJs from in vitro symbiosis association.

For each *P. luminescens* strain, three independent cultures were used to establish symbiosis with *H. bacteriophora*. Emerging IJs were then counted and the mean of three biological replicates is represented for Chr_dam strain (dark grey) and Chr_gfp (light grey) (see Materials & Methods section for details). The level of emergence between the 2 strains was not significantly different (Wilcoxon, $p=0.63$).