

Title	Palaeogeographical reconstruction and hydrology of glacial Lake Purcell during MIS 2 and its potential impact on the Channeled Scabland, USA
Authors	Peters, Jared L.;Brennand, Tracy A.
Publication date	2020-01-31
Original Citation	Peters, J. L. and Brennand, T. A. (2020) 'Palaeogeographical reconstruction and hydrology of glacial Lake Purcell during MIS 2 and its potential impact on the Channeled Scabland, USA', Boreas. doi: 10.1111/bor.12434
Type of publication	Article (peer-reviewed)
Link to publisher's version	https://onlinelibrary.wiley.com/doi/abs/10.1111/bor.12434 - 10.1111/bor.12434
Rights	© 2020, John Wiley & Sons, Inc. This is the peer reviewed version of the following article: Peters, J. L. and Brennand, T. A. (2020) 'Palaeogeographical reconstruction and hydrology of glacial Lake Purcell during MIS 2 and its potential impact on the Channeled Scabland, USA', Boreas, doi: 10.1111/bor.12434, which has been published in final form at https://doi.org/10.1111/bor.12434 . This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Use of Self-Archived Versions.
Download date	2024-05-14 06:09:17
Item downloaded from	https://hdl.handle.net/10468/9651



UCC

University College Cork, Ireland
Coláiste na hOllscoile Corcaigh

1 **Palaeogeographical reconstruction and hydrology of glacial Lake Purcell**
2 **during MIS 2 and its potential impact on the Channeled Scabland, USA**

3 JARED L. PETERS AND TRACY A. BRENNAND

4 Peters, J. L. & Brennand, T. A.: Palaeogeographical reconstruction and hydrology of glacial Lake
5 Purcell during MIS 2 and its potential impact on the Channeled Scablands.

6 Large, ice-marginal lakes that were impounded by the maximally-extended Cordilleran Ice Sheet
7 (CIS) provided source waters for the extraordinarily large floods that formed the Channeled
8 Scabland of Washington and Idaho, USA. However, flood flows that drained CIS meltwater and
9 contributed to landscape evolution during later stages of deglaciation have hitherto been poorly
10 investigated. This paper provides the first evidence for such a late deglacial floodwater source:
11 glacial Lake Purcell (gLP). Sedimentary evidence records the northward extension of gLP from
12 Idaho, USA into British Columbia, Canada and establishes its minimum palaeogeographical
13 extent. Sedimentary evidence suggests that the deglacial Purcell Lobe was a capable ice dam that
14 impounded large volumes of gLP water. A review of glacioisostatically affected lakes during CIS
15 deglaciation suggests that gLP could have been subjected to tilts ranging from $0 - >1.25 \text{ m km}^{-1}$.
16 Sedimentary evidence suggests high lake plane tilts ($\approx 1.25 \text{ m km}^{-1}$) are the most likely to have
17 affected gLP. Using this, the palaeogeography and volume of gLP are modelled, revealing that
18 $\sim 116 \text{ km}^3$ of water was susceptible to sudden drainage into the Channeled Scabland via the
19 Columbia River system. This calculation is supported by sedimentary and geomorphic evidence
20 compatible with energetic flood flows along the gLP drainage route and suggests gLP drained
21 suddenly, causing significant landscape change.

22 *Jared L. Peters (jared.peters@ucc.ie), Department of Geography, Simon Fraser University, 8888*
23 *University Dr, Burnaby, BC V5A 1S6, Canada and School of Biological, Earth and Environmental*
24 *Sciences, University College Cork, Cork, Ireland; Tracy A. Brennand, Department of Geography,*
25 *Simon Fraser University, 8888 University Dr, Burnaby, BC V5A 1S6, Canada.*

26 Growing concerns over the stability of future hydrosphere-cryosphere interactions and our ability
27 to accurately predict the behaviour of modern glaciers and ice sheets (e.g. Bamber *et al.* 2009;
28 Gardner *et al.* 2013) highlight the importance of a complete understanding of Cordilleran Ice Sheet
29 (CIS) decay. Ice-marginal lakes play an important role in this improved understanding because
30 they affect ice dynamics (Carrivick & Tweed 2013), are effective sediment traps that record
31 detailed glacial histories (e.g. Larsen *et al.* 2011; Liermann *et al.* 2012), and are prone to
32 catastrophic drainage that can influence regional sediment transport and drainage systems (Korup
33 2012). Considering the important effects that glacial lakes can impose on ice sheet decay and
34 landscape evolution, and their increasing abundance and size along modern, deglaciating ice
35 margins (Carrivick & Tweed 2013), the importance of developing a thorough understanding of
36 their role during the deglaciation of the CIS is evident.

37 At the Local Last Glacial Maximum (LLGM) large glacial lakes, like the $\sim 2\,600\text{ km}^3$ glacial Lake
38 Missoula (O'Connor & Baker 1992; Miyamoto *et al.* 2006, 2007), formed when the southern
39 margin of the CIS disrupted regional drainage patterns (Baker 2009). Some of these lakes drained
40 catastrophically and contributed to the formation of the Channeled Scabland (Fig. 1A), a
41 megaflood landscape that geomorphically and sedimentologically records flood flows of nearly
42 unprecedented Earthly scale with maximum discharges of 10-20 Sverdrups (Benito & O'Connor
43 2003; Denlinger & O'Connor 2010). Whereas geological and sedimentological signatures of
44 enormous jökulhlaups (glacial lake outburst floods) entering the Channeled Scabland are abundant
45 and well documented (e.g. Bretz 1925, 1969; Baker 2009; Benito & O'Connor 2003), the potential
46 for post-LLGM flood flows from the drainage of glacial lakes in British Columbia has been
47 proposed (Shaw *et al.* 1999; Lesemann & Brennand 2009; Waitt *et al.* 2009; Waitt 2016) but
48 remains relatively poorly understood.

49 The role of ice-marginal lake formation within the Purcell Trench during CIS deglaciation has
50 received inconstant speculation. Alden (1953) first contemplated a glacial lake in the Purcell
51 Trench and its possible drainage into the Columbia River system via the Kootenay River valley
52 (Fig. 1B). Most researchers (e.g. Alden 1953; Johns 1970; R. Fulton, pers. comm. 2010)
53 speculated that the glacial lake in the Purcell Trench was shallow and primarily ice marginal or
54 supraglacial, owing to stagnant ice occupying the Purcell Trench. These authors also suggest that
55 glacial lake water in the Purcell Trench likely drained gradually past a spillway in the south (the
56 Elmira spillway) and the downwasting ice in the north (Fig. 1B). However, Waitt *et al.* (2009)
57 and Waitt (2016) propose that more energetic drainage of a proglacial lake in the Purcell Trench
58 may have supplied post-Missoula flood flows to the Columbia River.

59 This study provides the first comprehensive investigation of glacial lake evolution in the Purcell
60 Trench. We use geological evidence and previous records of CIS glacioisostatic tilt to inform a
61 palaeogeographic reconstruction of a large lake, named here glacial Lake Purcell (gLK). We
62 explore evidence for ice damming of the lake and its drainage through the Kootenay River valley.
63 These analyses are used to assess the potential for energetic flood flows from the Purcell Trench
64 into the Channeled Scabland after the final drainage of glacial lakes Missoula and Columbia.

65 **Previous work on Purcell Lobe ice-marginal lakes**

66 Previous studies near the Purcell Trench have reconstructed glacial Lake Kootenai (gLK) from
67 thick deposits of lake bed sediments (sand and silt) in valley systems in northern Idaho and
68 northwestern Montana (Alden 1953; Johns 1970; Smith 2006; Fig. 1A). This lake formed when
69 river systems were impounded by the retreating Purcell Lobe (Alden 1953; Johns 1970; Smith
70 2006; Fig. 2). The sediments recording glacial Lake Kootenai are over 90 m thick in some areas
71 and record rapid deposition proximal to inflows (Alden 1953; Smith 2006). Valley-side benches

72 composed of lake bed sediments attributed to gLK range in elevation from 700-740 m a.s.l. in
73 Idaho and from 730-762 m a.s.l. in Montana due to different spillway heights (Alden 1953). The
74 Bull River spillway (Fig. 1B) in Montana was the first flow to be activated and would have
75 commenced following a lowering of the final stage of gLM in the Clark Fork River valley to the
76 south of gLK (Alden 1953). After sufficient northward retreat of the Purcell Lobe, gLK decanted
77 into the southern Purcell Trench, forming a large flood-related fan on the valley floor and an
78 unnamed proglacial lake. Lake levels in the Purcell Trench were dictated by the Elmira spillway
79 (ibid). The geomorphology of the Elmira spillway suggests that its original height was ~710 m
80 a.s.l. and that incision from lake drainage is responsible for its current elevation of 655 m a.s.l.
81 (ibid).

82 The naming conventions used by Alden (1953) and adopted by Johns (1970) and Smith (2006) are
83 abandoned in this study because they ambiguously describe distinct water bodies with a single
84 name (gLK). Furthermore, the name ‘glacial Lake Kootenay’ employed by Waitt *et al.* (2009) is
85 not used, as its closeness to Alden’s lake name is a potential source of confusion. Instead a naming
86 system is employed that distinguishes the discrete and possibly contemporaneous lakes that
87 occupied separate basins (Fig. 2). This new naming scheme retains Alden’s glacial Lake Kootenai
88 moniker in Montana, USA (where most of his research was conducted) but designates the unnamed
89 lake and its northern expansion in the Purcell Trench “glacial Lake Purcell” (Fig. 2).

90 The volumes of these glacial lakes have also been speculated upon and several researchers have
91 pointed out that volume was contingent on the style of CIS retreat through the Purcell Trench. If
92 Purcell Lobe retreat was dominated by stagnation and downwasting, the ice would have likely
93 displaced much of the volume available to any glacial lake. Fulton (1967, 1991) proposes a CIS
94 deglacial model dominated by stagnant, residual ice occupying valley systems resulting from a

95 rapid rise of the equilibrium line due to rapid climate amelioration. Sedimentary evidence for this
96 stagnation, and resultant downwasting, has been reported in the interior of British Columbia (Eyles
97 & Clague 1991; Ryder *et al.* 1991). During ice stagnation, glacial lake volume would have been
98 minimized by valley occupying ice. However marginal areas of the CIS may have experienced a
99 more complex pattern of decay (Fulton 1967) and these complications may have been further
100 exacerbated in mountainous terrain by late deglacial alpine ice advances (Lakeman *et al.* 2008).
101 Such complexities, along with potential inconsistencies in regional glacioisostatic response from
102 crustal heterogeneities (cf. Thorson 1989), may have enabled the formation of a deep, high-volume
103 gLP and highlight the need for investigations in the Purcell Trench.

104 Initial evidence for a high-volume, late-deglacial gLP has been supplied by Waitt *et al.* (2009) and
105 Waitt (2016), who suggest that a glacial lake in the Purcell Trench was a potential water source
106 for flood flow(s) in the Columbia River valley. Putative geomorphic evidence for post-Missoula,
107 late-Wisconsin Glacial Lake Outburst Floods (GLOFs) in the Columbia River valley includes two
108 megaflood bars marked by dune-scale bedforms (“giant current dunes”) near Chelan Falls,
109 Washington (Waitt *et al.* 2009; Fig. 1A). These dune-scale bedforms are tephrostratigraphically
110 dated to <13.5 cal. ka BP (Kuehn *et al.* 2009), after the final drainage of glacial lakes Missoula
111 and Columbia and Lake Bonneville (Waitt *et al.* 1994, 2009). Age constraints on the deglacial
112 CIS are compatible with the tephrostratigraphic age of the dune-scale bedforms and place the
113 Purcell Lobe ice margin near the Kootenay River valley by ~13.5 cal. ka BP (Dyke *et al.* 2003).

114 **Study area**

115 Data were gathered for this study within the Purcell Trench, its high-relief tributary valleys, and
116 along the Kootenay River valley (KRv; Fig. 1B). Much of the floor of the Purcell Trench in
117 Canada is occupied by Kootenay Lake, which is a ribbon-shaped lake >100 km long with an

118 average width of ~6.5 km (Fig. 1B). Kootenay Lake's water surface elevation is controlled by the
119 Corra Linn Dam in the Kootenay River valley to an elevation of ~532 m a.s.l. (Davis 1920; Kyle
120 1938; Fig. 1B). Kootenay Lake marks a change in spelling from the Kootenai River to the
121 Kootenay River (Fig. 1B) and is essentially a stagnation point in the flow of the Kootenai/y River
122 along its circuitous westward route from the Rocky Mountain Trench, British Columbia through
123 the Columbia Mountains. Kootenay Lake drains out of its West Arm via the Kootenay River,
124 which is the first major tributary of the Columbia River. In this study, the West Arm of Kootenay
125 Lake and the Kootenay River are jointly referred to as the Kootenay River valley (KRv; Fig. 1B).

126 **Methods**

127 **Geomorphology and sedimentology**

128 Geomorphic analyses and preliminary investigations to identify potential field sites were carried
129 out using publicly available digital elevation models from Geobase (from Natural Resources
130 Canada) and the National Elevation Database (NED, from the United States Geological Survey).
131 The two datasets were compiled and re-gridded into a single, 25-m resolution Digital Elevation
132 Model (DEM).

133 Sedimentary investigations entailed lithofacies identification, gravel fabric and ripple palaeoflow
134 measurements, and centimetre-scale logging of exposures. Sedimentary data are presented as
135 stereograms, rose diagrams, and exposure photographs. For each fabric analysis ≥ 30 clast a-b
136 plane attitude measurements (maximum dip and down-dip direction of the a-b plane) were taken
137 in $< 0.5 \text{ m}^2$ areas of exposures. The data were then plotted on lower hemisphere, equal area
138 (Schmidt) diagrams as contoured stereonet using the cosine sums method (Stereo32 software)
139 with a cosine exponent of 20 (cf. Roeller 2008). Fabrics from kame deposits are designated K1,
140 K2, K3, K4, K5 and palaeoflow data recording drainage from the Purcell Trench are designated

141 Dr1, D1, D2 and D3 (Fig. 1B). Clast a-axis position relative to the direction of a-b plane maximum
142 dip in gravel fabrics was used to determine the likely mode of clast mobilization. A dominance of
143 clast a-axes transverse to dip direction (a(t)) suggests clasts rolled along the bed, whereas a
144 dominance of clasts parallel to dip direction (a(p)) implies clast sliding across the bed or deposition
145 from suspension in a hyperconcentrated flow (cf. Brennand 1994).

146 **Palaeogeographical modelling of gLP**

147 Typically, glacial lake extent is reconstructed from the distribution of lake bed sediments and by
148 correlating water-plane indicators (e.g., deltas, shorelines; cf. Johnsen & Brennand 2004).
149 However, a dearth of gLP water-plane indicators were identified on the steep bedrock valley walls
150 of the Purcell Trench. There is also a poor potential for lacustrine sediment preservation within
151 the floodplain of the modern Kootenai River and Kootenay Lake may cover significant areas of
152 gLP sediment (because they share a common basin). Thus, after confirming minimum extents
153 with sedimentary data, gLP extent and volume are estimated by assessing modelled lake surface
154 planes against limited evidence and in comparison to contemporaneous and geographically close
155 palaeolakes.

156 Two critical palaeogeographical elements enabled this modelling of gLP: (i) the reconstruction of
157 the pre-incision lake bed and (ii) the application of an appropriate glacioisostatic adjustment
158 (GIA) to the water plane. These reconstructions were used to produce a combination of rasters
159 that were used in conjunction with modern topography to define gLP palaeogeography and
160 calculate its volume (cf. Leverington *et al.* 2002; DeVogel *et al.* 2004). The pre-incision palaeo-
161 lake bed was reconstructed using an inverse distance weighting function to interpolate a series of
162 pre-incision elevations measured using the composite DEM. In the model, the boundaries of this
163 pre-incision lake bed surface were defined by its intersection with topographic highs on the

164 composite DEM of the Purcell Trench. A plausible range of GIA was derived from a survey of
165 previously reported glacioisostatic tilts for CIS glacial lake planes. The published ages of these
166 lakes are also reported, which are derived using disparate methods with varying accuracy. These
167 ages were recalibrated for this study with Calib software (Stuiver & Reimer 1993) to 2σ confidence
168 using the IntCal13 radiocarbon curve (Reimer *et al.* 2013) to improve comparability of the ages
169 (cf. Peters *et al.* 2016).

170 The resultant array of plausible lake plane tilts was projected along the Purcell Trench from the
171 Elmira Spillway, which controlled gLP lake levels prior to drainage into the Kootenay River valley
172 (Alden 1953; Johns 1970; Smith 2006). A DEM of modern Kootenay Lake bathymetry
173 (bathymetric DEM) was constructed from interpolated individual soundings using a GIS and was
174 used to estimate the total volume of gLP (i.e. the volume of modern Kootenay Lake was added to
175 the calculated drainable volume of gLP based on topographic DEMs).

176 **Results and interpretations**

177 **The northward extension of gLP**

178 Evidence for the northward extension of gLP is mainly preserved in large (>60 km long and up to
179 ~20 km wide) sediment benches that occupy the floor of the Purcell Trench adjacent to the
180 Kootenai River (Fig. 1B). These benches extend from the Elmira spillway (~10 km south of
181 Bonners Ferry, Idaho) to ~5 km north of Creston, British Columbia (Fig. 1B), and reach an
182 elevation of up to 706 m a.s.l. (Fig. 3), >176 m above the modern water level of Kootenay Lake.
183 Bench tread elevations are incrementally lower towards the Kootenai River floodplain (Fig. 3).
184 Some treads exhibit channels on their surfaces (channels A and B, Fig. 3).

185 The benches are composed of massive or laminated silt and clay rhythmites (sites 3, 4, 5, 15; Fig.
186 4A) containing occasional pebble- to cobble-sized clasts (lonestones) that display striated facets
187 and plucked ends (sites 5, 15). Lonestones occur in relatively high abundance within massive silt
188 (Fig. 4B). Rare deposits of massive, silty coarse sand, interbedded with silt and clay laminae that
189 drape lonestones are also present (Fig. 4C). Silt and clay rhythmites are exposed north of the
190 contiguous benches at an elevation of 675 m a.s.l. (site 2, Fig. 1B). Gravel deposits composed of
191 dipping (apparent 30° downwards dip towards 221°), normally-graded, tabular beds that reach
192 thicknesses of ≥ 10 m are also exposed in the Idaho sediment benches (Fig. 4G). These gravel
193 deposits are typically capped by laminated or massive silt.

194 The highest bench surfaces are composed of flat-topped silt deposits that share similar elevations,
195 recording contiguous lake bed deposits (cf. Ryder *et al.* 1991; Johnsen & Brennand 2004). The
196 flat-topped, occasionally channelised bench segments (e.g. Fig. 3) record remnant lake bed
197 sediments following fluvial incision (cf. Clague 1986). This interpretation of terrace formation is
198 supported by the close proximity of the Kootenai River.

199 Rhythmites record lake bed sedimentation and suggest a record of varying sediment influx from
200 suspension settling and underflows (Smith & Ashley 1985). Clasts with striated facets and plucked
201 ends are interpreted as glacially modified (Sharp 1982) and suggest an ice-proximal sediment
202 source. Correspondingly, lonestones within laminated and massive silt and sand deposits (Fig.
203 4C) are interpreted as dropstones (Lønne 1995) and their occurrence indicates that lake bed
204 sedimentation took place in an ice-marginal environment with water deep enough to induce calving
205 (Pelto & Warren 1991; Boyce *et al.* 2007; Tsutaki *et al.* 2011). Abundant dropstones within
206 massive silt (Fig. 4B) record iceberg rollover events (cf. Winsemann *et al.* 2004).

207 The inclined gravel deposits (Fig. 4G) are consistent with alluvial fan progradation (Blair &
208 McPherson 1994) and their increased occurrence in the south of the study area (site 10, Figs 1B,
209 3) implies that they record deposition during the decanting of gLK into gLP (unnamed lake, Alden
210 1953). These inclined gravels may also record deltaic deposition, however no topsets have been
211 identified, so an interpretation of alluvium, or potentially expansion bar deposits, is preferred.
212 Thus, the lake bed benches and bench segments confirm the northward extension of gLP through
213 the Purcell Trench and suggest that it was an ice-contact, proglacial lake.

214 **The Purcell Lobe as an ice dam**

215 A series of elevated gravel benches occupy interfluvies along the valley walls of the Purcell Trench
216 (kame sediment, Fig. 1B). The gravel benches are ≥ 6 -16 m thick and reach elevations of 600-725
217 m a.s.l. They are typically composed of normally-graded beds of planar-stratified and trough
218 cross-stratified sand and gravel lithofacies (Fig. 4E) that occasionally exhibit faulting. Small
219 pebbles to large cobbles are typically well rounded and imbricated with gravel fabrics revealing
220 predominantly a(t) pebble orientations and valley-parallel southerly and northerly palaeoflow
221 directions (stereograms K1-K5, Fig. 1B). Unconsolidated, poorly sorted sand clasts are
222 occasionally preserved in the gravel with little evidence of rounding (Fig. 4D).

223 The normally-graded, planar- and trough cross-stratified gravel beds (Fig. 4E) are consistent with
224 fluvial deposition in a gravel-bed stream (Miall 1977). Gravel fabrics record valley-parallel stream
225 flows and traction transport (rolling; Brennand 1994) that is anomalous to modern topography.
226 Furthermore, the northward flows are also irreconcilable with an interpretation of remnant advance
227 outwash deposits and the elevation of these deposits (up to 725 m a.s.l.) place them ~200 m above
228 the surface of the modern Kootenay River valley where post-gLP damming is unlikely (Peters
229 2012). Thus, the gravels are interpreted as kame terraces deposited against the valley walls by ice-

230 marginal meltwater streams (cf. Terpiłowski 2007; Pisarska-Jamroży *et al.* 2010). This
231 interpretation is further supported by the incorporation of unconsolidated, unrounded sand-clasts
232 (Fig. 4D), which may record rapid deposition by turbidites in a deltaic ice-marginal environment
233 (cf. Winsemann *et al.* 2018); alternatively, the angularity of these deposits and lack of cohesive
234 material (e.g. silt) suggest that they could have been preserved during mobilisation and
235 incorporation into the gravel because they were frozen (Menzies 1990). The distribution of the
236 kame deposits (Fig. 1B) suggests the Purcell Lobe dammed northern flow of gLP at its maximum
237 extent.

238 **Glacial Lake Purcell reconstruction**

239 Glacioisostatic adjustments (GIA) derived from CIS palaeolake plane data range from horizontal
240 to $\sim 2.1 \text{ m km}^{-1}$ (Table 1). This dataset was assessed for outliers using 1.5x the inter-quartile range
241 (low cut-off -0.5, high cut-off 3.1), which revealed that all the tilts assessed were mathematically
242 relevant (all values fall between the limits defining outliers). GIA data were plotted against time
243 (Fig. 5) to elucidate possible patterns in glacioisostatic behaviour during MIS 2 deglaciation.
244 Glacial lakes Arrow and Invermere formed closest to gLP (within ~ 50 and 70 km of the Purcell
245 Trench, respectively; Fulton *et al.* 1989; Sawicki & Smith 1992) and thus are more likely to have
246 had rates of glacioisostatic rebound governed by similar lithospheric properties (Clague & James
247 2002) and the Clayhurst stage of glacial Lake Peace was chronologically nearest to the proposed
248 dates for gLP (Mathews 1978; Table 1; Fig. 5). Together, these records suggest that potential GIA
249 of gLP water planes ranged from 0 m km^{-1} to $\sim 1 \text{ m km}^{-1}$ but could have been as high as 2 m km^{-1}
250 (Fig. 5). A projected array of tilted water planes ($0.0, 0.5, 0.75, 1.0$ and 1.25 m km^{-1}) produces a
251 range of modelled gLP extents (Fig. 6) that encompass all but three of the previously reported
252 GIAs with age constraints (Fig. 5). The calculated volume of gLP for the modelled array of tilts

253 (assuming a steep Purcell Lobe ice margin and a position consistent with estimates by Dyke *et al.*
254 2003) ranges from 40-142 km³ (Table 2).

255 The intersection of the modelled lake extent and the geomorphology of the Kootenai River flood-
256 related fan suggests that with low GIAs, gLP could have drained past the Elmira Spillway via a
257 south-flowing stream that drained along the west side of the Purcell Trench (Fig. 6). However,
258 sedimentary evidence suggests a steeper GIA may be more accurate. Exposures of the Kootenai
259 River flood-related fan in the southern Purcell Trench (site 10, Figs 1B, 3) are topped by silt
260 deposits (e.g. Fig. 4G) that suggest a lacustrine environment. Furthermore, the contiguous lake
261 sediment benches recording the minimum extent of gLP (Fig. 1B) cover a larger area than
262 modelled extents with low GIAs (Fig. 6). Thus, a steep GIA is deemed most likely to have
263 influenced gLP's lake plane (i.e. at least 1.25 m km⁻¹; Fig. 6). A GIA of 1.25 m km⁻¹, like the tilt
264 that affected glacial Lake Bretz (Table 1), would have resulted in a total gLP volume of 142 km³
265 (Table 2). Its surface elevation against a northern ice dam would have been >800 m a.s.l. and it
266 would have reached depths of >400 m (Table 2, Fig. 7).

267 **GLOF evidence in the Kootenay River valley**

268 The Kootenay River valley contains elevated (>100 m above the modern Kootenay River
269 floodplain) sediment benches or terraces (GLOF sediment, Fig. 1B), and alluvial fans, which are
270 remnants of a thicker valley fill (Figs 1B, 8). The highest truncated and bisected alluvial fan
271 remnants reach elevations of up to ~675 m a.s.l. (Fig. 8B). The highest terrace occupies both sides
272 of the Kootenay River valley with tread elevations from 642 m a.s.l. near the Purcell Trench to
273 ~600 m a.s.l. near the Kootenay River valley confluence with the Columbia River valley (Fig. 8A,
274 B) and a down-valley slope of ~1.5 m km⁻¹. The terraces typically contain normally-graded planar-
275 stratified and trough cross-stratified sand, imbricated gravel with occasional massive gravel beds,

276 and, occasionally, diffusely graded, sinusoidally stratified sand beds (Fig. 4I). Trough-cross
277 stratified gravel is most common near the top of the valley fill, where smoothed and potholed
278 bedrock also exists at elevations up to 624 m a.s.l. (Figs 1B, 8A). Type-a ripples overlain by type-
279 s ripples (Ashley *et al.* 1982; Fig. 4F) are exposed in inset terraces lower than 560 m a.s.l. in the
280 Kootenay River valley valley fill and towards the valley centreline (Fig. 1B). Gravel fabrics from
281 planar-stratified gravels and type-a ripple measurements record westward palaeoflows through the
282 Kootenay River valley (Fig. 1B, Stereograms D1, D2 and Dr1).

283 The normally-graded, planar and cross-stratified beds of sand and imbricated gravel with westward
284 palaeoflows in the terraces are compatible with observations of sand and/or gravel dune or bar
285 formation (e.g. Carrivick *et al.* 2004; Rushmer 2006; Russell 2009) during a confined GLOF from
286 gLP. Diffusely graded, sinusoidally stratified sand records stationary and breaking antidunes,
287 deposited rapidly from supercritical hyperconcentrated GLOF flows (cf. Lang & Winsemann
288 2013; Lang *et al.* 2017). Occasional massive gravel beds are associated with deposition of traction
289 load during GLOFs (Carrivick *et al.* 2004; Russell 2009) or rapid deposition (Rushmer 2006).
290 Ripples in fine sediments have been interpreted to record deposition during waning GLOF flows
291 (Russell 2009) or hydraulic damming during the latter stages of floods (cf. Touchet Beds in the
292 Channeled Scabland, Waitt 1980). Together, this sedimentary sequence is interpreted to record a
293 period of sand and gravel aggradation during energetic GLOF flows through the Kootenay River
294 valley, followed by fine sediment deposition during waning flood flows or in areas of local
295 hydraulic damming (cf. Winsemann *et al.* 2016).

296 The Kootenay River valley highest terrace, just below the alluvial fan remnants, is interpreted to
297 record the pre-GLOF valley bottom because the relative prevalence of alluvial fans that have
298 aggraded onto it (Fig 8B) suggests this surface was exposed for a significant amount of time. This

299 terrace's elevation suggests a pre-incision down-valley slope of $\sim 1.5 \text{ m km}^{-1}$ and a pre-flood valley
300 bottom elevation of $\sim 640 \text{ m a.s.l.}$ near the Purcell Trench (i.e., Kootenay Lake, Fig. 8A). Terrace
301 geomorphology indicates that the valley fill (pre-GLOF and GLOF sediments) has experienced
302 $\sim 110 \text{ m}$ of incision near the Purcell Trench and $\sim 150 \text{ m}$ near the Columbia River valley (Fig. 8).

303 Although it is likely that the highest terraces were formed during erosive stages of a gLP GLOF,
304 it is unknown exactly which terraces formed by channelising GLOF flows, and which by
305 postglacial fluvial incision. However, recent geomorphic examinations suggest the bedrock around
306 Cora Linn Dam ($\sim 530 \text{ m a.s.l.}$, Figs 1B, 8) was sculpted by energetic flows (Waite 2016). The
307 potholed bedrock near the top of the valley fill sequence (624 m a.s.l. , Fig 8A) also suggests
308 bedrock erosion during a period of energetic flow that likely removed pre-GLOF valley fill
309 sediment (Zen & Prestegard 1994; Fig. 8A). Taken together, these observations suggest that the
310 rising stage of a gLP GLOF was responsible for alluvial fan truncation, the creation of elevated
311 potholes (624 m a.s.l.), removal of pre-GLOF sediment to bedrock (at least 90 m thickness at Corra
312 Linn Dam), and bedrock sculpting around Corra Linn Dam (Waite 2016). In this scenario, the
313 terraces formed after GLOF aggradation during incision by waning-stage channelising GLOF
314 flows or post-glacial river flows. More conservatively, the rising stage of the GLOF may only have
315 been responsible for alluvial fan truncation, elevated pothole formation and partial incision through
316 valley fill. In this scenario, waning-stage channelizing GLOF flows and post-glacial river flows
317 may have been responsible for the bulk of sediment removal, terracing and the bedrock sculpting
318 around Corra Linn Dam.

319 The large, fan-shaped landform situated at the confluence of the Kootenay River valley and the
320 Columbia River valley (Playmor Junction, Fig. 1B) exhibits multiple terraces (Fig. 8C) and reaches
321 an elevation of approximately 490 m a.s.l. , which is $\sim 110 \text{ m}$ lower than the highest terrace treads

322 recording the pre-GLOF valley bottom in the western Kootenay River valley (Fig. 8A). It is largely
323 composed of very coarse, poorly-sorted, clast-supported massive gravel with abundant well-
324 rounded boulders (up to 0.5 m b-axes) (Fig. 4H). Clast a-axes are randomly oriented and gravel
325 fabric measurements have a polymodal distribution (D3, Fig. 1B). The coarse, fan-building
326 gravels are capped by ~10 m of trough cross-stratified sand and cobble gravel (Peters 2012).

327 The fan shape and composition of the gravel deposit at Playmor Junction (Fig. 1B) suggests
328 deposition as a large alluvial fan (Blair & McPherson 1994) or GLOF expansion bar (e.g. Baker
329 1984; Benito 1997). The well-rounded gravel that comprises the fan indicates fluvial mobilisation
330 and its massive, poorly-sorted structure with boulder-sized clasts is consistent with deposition
331 during high-energy flows associated with GLOFs (Cenderelli & Wohl 2003; Harrison *et al.* 2006;
332 Russell 2009). The polymodal fabric (D3, Fig. 1B; cf. Morison & Hein 1987; Meetei *et al.* 2007)
333 with randomly-oriented clast a-axes, poorly sorted texture, and massive structure of the gravel
334 suggests rapid deposition in a turbulent, possibly hyperconcentrated flow (cf. Brennand & Shaw
335 1996). This boulder gravel is coarser than any gravels observed in the Kootenay River valley,
336 suggesting deposition during the most energetic GLOF flows responsible for pothole erosion,
337 alluvial fan truncation, and pre-GLOF valley-fill incision in the Kootenay River valley. The
338 capping, trough cross-stratified sand and gravel indicates a subsequent period of braided stream or
339 alluvial fan deposition (cf. Allen 1983; Blair 1987; Kjær *et al.* 2004) or possibly deposition during
340 waning flood flows (Winsemann *et al.* 2016) and the multiple terraces preserved on the fan suggest
341 incremental incision by fluvial erosion over time (Fig. 8C). Thus, we interpret the Playmor
342 Junction deposit to most likely have been deposited as a GLOF expansion bar that was capped by
343 alluvium and subsequently incised by the postglacial Kootenay River. The ~110 m height

344 difference between its highest surface (~490 m a.s.l.) and the local pre-GLOF valley bottom (~600
345 m a.s.l.; Fig. 8A) indicates deposition after an initial period of erosive GLOF flows.

346 **Discussion**

347 **Summary of gLP evolution**

348 Sedimentary and geomorphic evidence records gLP evolution from a small ice-contact proglacial
349 lake (described by Alden 1953) to a large, valley-filling ice-contact proglacial lake that drowned
350 >100 km length of the Purcell Trench (Fig. 2). At one stage in its evolution, gLK (Alden 1953)
351 decanted much of its volume into gLP (Fig. 2B). This drainage is recorded by the flood-related
352 fan (Alden 1953) that has since been incised by the Kootenai River (Figs. 1B, 6). This fan formed
353 the southern shore of gLP and forced southward drainage towards the Elmira spillway against the
354 western valley wall (Fig. 6). The northern extent of gLP was dictated by the northward-retreating
355 Purcell Lobe of the CIS (Figs 2, 6).

356 No direct geochronological ages are available for gLP or gLK. However, tephrostratigraphic ages
357 that constrain the geomorphic evidence for Columbia River valley flood flows provided by Waitt
358 *et al.* (2009) can provide some tentative, preliminary gLP chronology. The dune-scale bedforms
359 described by Waitt *et al.* (2009) and Waitt (2016) were deposited <13.5 cal. ka BP, which is
360 younger than the final drainage of other regional palaeolakes (Missoula, Columbia, and
361 Bonneville), but compatible with our reconstructions of gLP and reconstructions of the deglacial
362 Purcell Lobe's ice margin, which was near the Kootenay River valley by ~12 cal. ka BP (Dyke *et*
363 *al.* 2003; Fig. 6). Thus, in the absence of independent dating, gLP sedimentation is assumed to
364 have occurred during MIS 2 with a maximum possible age between 14 750±375 and 15 200±400
365 cal. a (LLGM; Atwater 1987). Drainage most likely occurred after ~13.5 cal. ka BP (Waitt *et al.*
366 1994, 2009; Dyke *et al.* 2003).

367 **GLP volume and local glaciological impacts**

368 Most of the lake bed sediments that comprise the contiguous benches in the southern Purcell
369 Trench are not significantly deformed, suggesting deposition in an ice-distal environment without
370 remobilisation from removal of buried ice. This suggests previous hypotheses of a low-volume,
371 ice-marginal, or supraglacial lake that formed during CIS downwasting in the Purcell Trench are
372 incorrect (e.g. Alden 1953; Eyles & Clague 1991; Ryder *et al.* 1991). Furthermore, iceberg rain
373 out sediment interpreted in this study suggests that gLP's lake depth (>400 m, Table 2) was enough
374 to force ice-marginal flotation and induce calving retreat through the Purcell Trench (Carrivick &
375 Tweed 2013). Thus, the Purcell Lobe would have likely formed a steep terminus (Fig. 7) and been
376 unable to displace significant amounts of gLP volume. This interpretation is compatible with the
377 lack of ice-marginal landforms (kame terraces and moraines) in the southern Purcell Trench.
378 Furthermore, kame terrace deposits (Figs 1B, 6D, E) confirm that the Purcell Lobe was sufficiently
379 sealed to the Purcell Trench valley-wall for a period that allowed at least 16 m of glaciofluvial
380 deposition. The relatively low position of the kame terraces within the valley (600-725 m a.s.l.)
381 indicates that the seal existed late in the deglaciation of the Purcell Lobe. This seal, although not
382 likely to be concurrent with the lacustrine deposits of gLP (based on elevation, Table 2), provides
383 evidence that the Purcell Lobe could have dammed large volumes of water long after the CIS
384 margin retreated northward into British Columbia. This evidence of a high-volume gLP elucidates
385 important potential for large flood flows late in CIS deglaciation.

386 A deep, high-volume gLP (>400 m, almost 150 km³, respectively, Table 2) would have held a
387 similar amount of water as modern Lake Tahoe, or ~30% more than the Dead Sea. Such a lake
388 would have mechanically exacerbated local CIS mass loss through calving, thereby steepening the
389 ice margin causing increased ice flow velocities (Carrivick & Tweed 2013). Mass loss would also

390 have been accelerated in the Purcell Trench by thermal erosion, because ice-marginal lakes deliver
391 heat to glacier termini. Such thermal erosion can undercut the ice margin at the water line (e.g.
392 Kirkbride & Warren 1999; Röhl 2006) further steepening the terminus and intensifying calving
393 retreat. These feedbacks suggest that the deglacial Purcell Lobe would have had a steep ice-front
394 prior to gLP drainage (Fig. 7).

395 **GLP drainage and impacts on landscape evolution**

396 GLP was confined to the Purcell Trench until the Purcell Lobe's calving margin retreated
397 sufficiently northward to allow drainage into the Kootenay River valley (Figs 1B, 2). At this time,
398 in order to drown gLP lake bed sediments and the flood-related fan in the southern Purcell Trench
399 (Alden 1953; Fig. 2B), the gLP water surface was most likely tilted $\sim 1.25 \text{ m km}^{-1}$, relative to the
400 modern landscape (Fig. 6). Applying this tilt to a modelled gLP lake plane results in a surface
401 elevation against the ice dam near the Kootenay River valley of 817 m a.s.l. and a drainable volume
402 of $\sim 116 \text{ km}^3$ (Table 2). This elevated lake surface is $\sim 180 \text{ m}$ above the pre-GLOF valley bottom
403 in the Kootenay River valley (Table 2, Fig. 8A), suggesting that gLP water likely drained suddenly
404 into the Kootenay River valley following catastrophic ice-dam failure.

405 The $\sim 180 \text{ m}$ elevation difference between the gLP water surface and the pre-GLOF valley bottom
406 in the Kootenay River valley (Table 2) suggests that this sudden drainage would have generated
407 extremely high specific and total stream powers, capable of eroding large amounts of boulder-
408 sized sediment (cf. Cenderelli & Wohl 2003). This erosive GLOF is recorded by truncated alluvial
409 fans and potholed bedrock $\sim 140 \text{ m}$ and $\sim 90 \text{ m}$, respectively, above the modern Kootenay River
410 valley floor (cf. Winsemann *et al.* 2016). If the fluvially-eroded bedrock described by Waitt (2016)
411 at the Corra Linn Dam is attributed to a gLP GLOF, $>90 \text{ m}$ depth of sediment would have been
412 removed by the GLOF at this location, which is comparable to previous models of GLOF erosion

413 (e.g. Winsemann *et al.* 2016; Lang *et al.* 2019; Fig. 8A). After incising the Kootenay River valley
414 fill, the flood flows debouched into the larger Columbia River valley at Playmor Junction, where
415 a large, fan-shaped expansion bar was formed from cobble- and boulder-sized bedload (cf. Baker
416 1984; Benito 1997; Figs 1B, 4H, 8C).

417 The flood flows generated by gLP drainage would have entered the Columbia River valley via the
418 Kootenay River valley (Figs 1B, 8C). Whether or not these flows would have been capable of
419 enough geomorphic work to have formed dune-scale bedforms in the Channeled Scabland near
420 Chelan Falls, Washington (Waite *et al.* 1994, 2009) depends on flow attenuation along the ~500-
421 km long flood route (defined by the lengths of the modern Kootenay River and Columbia River
422 from the Kootenay confluence). However, because gLP likely drained after ~13.5 cal. ka BP
423 (Kuehn *et al.* 2009; Waite *et al.* 2009), drained a large volume of water (~116 km³, Table 2), and
424 likely drained suddenly (based on the modelled ~180 m elevation difference between the gLP
425 water surface and the top of the highest terrace in the Kootenay River valley), it is possible that
426 gLP flood flows induced late-Pleistocene geomorphic changes in the Channeled Scabland (Fig.
427 1A; cf. Waite *et al.* 2009); however, hydraulic modelling should be performed to assess this (e.g.
428 Winsemann *et al.* 2016).

429 Regardless of the role the gLP GLOF may have played in the Channeled Scabland, its regional
430 effects on postglacial fluvial systems are evidenced geomorphically and sedimentologically. The
431 low (<560 m a.s.l.) terraces located along the Kootenay River valley (Fig. 8A, B) likely record
432 postglacial (late-Pleistocene and Holocene) fluvial incision by the Kootenay River, which
433 remobilised the waning flood-flow deposits towards the Kootenay River valley confluence with
434 the Columbia River valley (Figs 1B, 8C). As the sediment-laden Kootenay River exited the narrow
435 Kootenay River valley, it deposited its bedload as ~10 m of trough cross-stratified sand and gravel

436 alluvium (Peters 2012) over the surface of the boulder-gravel expansion bar (cf. Kehew *et al.*
437 2010). Finally, when the Kootenay and Columbia rivers neared their modern elevations and the
438 Kootenay River reached the Kootenay River valley's bedrock and/or its specific sediment yield
439 relaxed following postglacial incision (Church & Slaymaker 1989), alluvial deposition over the
440 expansion bar was replaced with incision, forming extensive fluvial terraces (Fig. 8C).

441 **Conclusions**

- 442 • GLP was a large (~1 152 km², 142 km³) ice-contact proglacial lake that most likely reached
443 water depths of >400 m. This deep water induced calving retreat along the Purcell Lobe
444 terminus, evidenced by iceberg rain-out deposits and dropstones within the gLP lakebed
445 sediments. This evidence contradicts previous hypotheses that propose stagnant ice filled
446 the valley limiting lake volume.
- 447 • Kame terraces were formed by ice-marginal stream deposition along the flanks of the
448 deglacial Purcell Lobe north of the Kootenay River valley, indicating that an ice-valley
449 wall seal was maintained throughout much of CIS deglaciation in the Purcell Trench. This
450 suggests that the Purcell Lobe could have effectively dammed gLP within the Purcell
451 Trench without allowing significant gradual drainage into the Kootenay River valley.
- 452 • The Purcell Lobe's terminus was altered mechanically by its calving margin and thermally
453 by heat exchange with gLP. These processes likely exacerbated the northward rate of
454 Purcell Lobe retreat and formed a steep ice front in the Purcell Trench. This steep ice
455 margin dammed the northern extent of gLP prior to its drainage into the Kootenay River
456 valley.
- 457 • The gLP lake surface was likely >800 m a.s.l. against its dam prior to its final drainage into
458 the Kootenay River valley after 13.5 cal. ka BP, which is ~180 m above the top of the
459 Kootenay River valley's pre-GLOF valley fill. This height discrepancy suggests gLP could
460 have drained 116 km³ of water into the Columbia River via the Kootenay River valley.
461 This large volume of water likely drained suddenly following catastrophic ice dam failure.
- 462 • The initial flood flows caused by the gLP GLOF may have eroded up to ~150 m of pre-
463 existing sediment from the Kootenay River valley, scouring to bedrock in places and
464 producing an expansion bar at its junction with the Columbia River, before depositing
465 GLOF sand and gravel in the Kootenay River valley.

- 466 • The timing of gLP drainage into the Kootenay River valley (based on CIS reconstructions)
467 is compatible with tephrostratigraphic age constraints from dune-scale bedforms along the
468 Columbia River at Chelan in the Channeled Scabland, Washington. Considering that gLP
469 drainage likely supplied a $>100 \text{ km}^3$ pulse of water into the Columbia River system, it is
470 conceivable that this GLOF formed these Channeled Scabland dune-scale bedforms;
471 however hydraulic modelling of flow attenuation should be performed to verify this
472 hypothesis.
- 473 • Following the catastrophic drainage of gLP, the Kootenay River incised into the GLOF
474 sediments, leaving a series of terraces formed by the GLOF and later, postglacial fluvial
475 incision at elevations from $\sim 600 - 550 \text{ m a.s.l.}$ ($\sim 20 - 60 \text{ m}$ above the modern river). A
476 $\sim 10\text{-m}$ thick deposit of alluvium was deposited over the expansion bar at the confluence of
477 the Kootenay River valley with the Columbia River valley, which was also incised as the
478 Kootenay River approached its modern elevation, leaving a series of fluvial terraces.
- 479 • Overall, these findings suggest that previous hypotheses favouring stagnant ice during CIS
480 deglaciation may underestimate the potential hydrological impacts of transient, late-
481 deglacial lakes. Furthermore, it seems likely that CIS GLOFs may have effected changes
482 in the Channeled Scabland after glacial lakes Missoula and Columbia had drained.

483 Acknowledgements – This research was funded by a NSERC discovery grant (194107) to TAB
484 and a GSA graduate research grant to JLP. Andrew Perkins, Mathew Burke, and Aaron Dixon
485 provided manual labour and helpful insights in the field. We appreciate insightful reviews from
486 Jutta Winsemann and an anonymous reviewer.

487 Data availability – We agree to make data available upon request.

488 Author contributions – JLP collected the data used for this research with occasional supervision
489 from TAB. JLP digitised and processed the data. Both authors interpreted the results, discussed
490 their overarching scientific relevance, and contributed to the writing of this manuscript.

491

492 **References**

- 493 Alden, W. C. 1953: Physiography and glacial geology of western Montana and adjacent areas. *U.S.*
494 *Geological Survey Professional Paper 231*, 142-153.
- 495 Allen, J. R. L. 1983: Studies in fluvial sedimentation; bars, bar-complexes and sandstone sheets (low-
496 sinuosity braided streams) in the brownstones (L. Devonian), Welsh borders. *Sedimentary Geology*
497 *33*, 237-293.
- 498 Ashley, G. M., Southard, J. B. & Boothroyd, J. C. 1982: Deposition of climbing-ripple beds: a flume
499 simulation. *Sedimentology* *29*, 67-79.
- 500 Atwater, B. F. 1987: Status of glacial Lake Columbia during the last floods from glacial Lake Missoula.
501 *Quaternary Research* *27*, 182-201.
- 502 Baker, V. R. 1984: Flood sedimentation in bedrock fluvial systems. *Sedimentology of Gravels and*
503 *Conglomerates – Memoir 10*, 87-98.
- 504 Baker, V. R. 2009: The Channeled Scabland: a retrospective. *Annual Review of Earth and Planetary*
505 *Sciences* *37*, 393-411.
- 506 Bamber, J. L., Riva, R. E., Vermeersen, B. L. & LeBrocq, A. M. 2009: Reassessment of the potential sea-
507 level rise from a collapse of the West Antarctic Ice Sheet. *Science* *324*, 901-903.
- 508 Benito, G. 1997: Energy expenditure and geomorphic work of the cataclysmic Missoula flooding in the
509 Columbia River Gorge, USA. *Earth Surface Processes and Landforms* *22*, 457-472.
- 510 Benito, G. & O'Connor, J. E. 2003: Number and size of last-glacial Missoula floods in the Columbia River
511 valley between the Pasco Basin, Washington, and Portland, Oregon. *Geological Society of America*
512 *Bulletin* *115*, 624-638.

- 513 Blair, T. C. 1987: Sedimentary processes, vertical stratification sequences, and geomorphology of the
514 Roaring River alluvial fan, Rocky Mountain National Park, Colorado. *Journal of Sedimentary*
515 *Research* 57, 1-18.
- 516 Blair, T. C. & McPherson, J. G. 1994: Alluvial fans and their natural distinction from rivers based on
517 morphology, hydraulic processes, sedimentary processes, and facies assemblages. *Journal of*
518 *Sedimentary Research* 64, 450-489.
- 519 Boyce, E. S., Motyka, R. J. & Truffer, M. 2007: Flotation and retreat of a lake-calving terminus, Mendenhall
520 Glacier, southeast Alaska, USA. *Journal of Glaciology* 53, 211-224.
- 521 Brennand, T. A. 1994: Macroforms, large bedforms and rhythmic sedimentary sequences in subglacial
522 eskers, south-central Ontario: implications for esker genesis and meltwater regime. *Sedimentary*
523 *Geology* 91, 9-55.
- 524 Brennand, T. A. & Shaw, J. 1996: The Harricana glaciofluvial complex, Abitibi region, Quebec: its genesis
525 and implications for meltwater regime and ice-sheet dynamics. *Sedimentary Geology* 102, 221-
526 262.
- 527 Bretz, J. H. 1925: The Spokane flood beyond the Channeled Scablands. *Journal of Geology* 33, 97-115.
- 528 Bretz, J. H. 1969: The lake Missoula floods and the Channeled Scabland. *Journal of Geology* 77, 505-543.
- 529 Carrivick, J. L. & Tweed, F. S. 2013: Proglacial lakes: character, behaviour and geological importance.
530 *Quaternary Science Reviews* 78, 34-52.
- 531 Cenderelli, D. A. & Wohl, E. E. 2003: Flow hydraulics and geomorphic effects of glacial-lake outburst
532 floods in the Mount Everest region, Nepal. *Earth Surface Processes and Landforms* 28, 385-407.
- 533 Church, M. & Slaymaker, O. 1989: Disequilibrium of Holocene sediment yield in glaciated British
534 Columbia. *Nature* 337, 452-454.

- 535 Clague, J. J. 1986: The Quaternary stratigraphic record of British Columbia-evidence for episodic
536 sedimentation and erosion controlled by glaciation. *Canadian Journal of Earth Sciences* 23, 885-
537 894.
- 538 Clague, J. J. & James, T. S. 2002: History and isostatic effects of the last ice sheet in southern British
539 Columbia. *Quaternary Science Reviews* 21, 71-87.
- 540 Davis, W. M. 1920: Features of glacial origin in Montana and Idaho: A Shaler memorial study. *Annals of*
541 *the Association of American Geographers* 10, 75-147.
- 542 Denlinger, R. P. & O'Connell, D. 2010: Simulations of cataclysmic outburst floods from Pleistocene Glacial
543 Lake Missoula. *Geological Society of America Bulletin* 122, 678-689.
- 544 DeVogel, S. B., Magee, J. W., Manley, W. F. & Miller, G. H. 2004: A GIS-based reconstruction of late
545 Quaternary palaeohydrology: Lake Eyre, arid central Australia. *Palaeogeography,*
546 *Palaeoclimatology, Palaeoecology* 204, 1-13.
- 547 Dyke, A. S., Moore, A. & Robertson, L. 2003: Deglaciation of North America. *Geological Survey of*
548 *Canada. Open File*, 1574, <https://doi.org/10.4095/214399>.
- 549 Eyles, N. & Clague, J. J. 1991: Glaciolacustrine Sedimentation During Advance and Retreat of the
550 Cordilleran Ice Sheet in Central British Columbia. *Géographie physique et Quaternaire* 45, 317-
551 331.
- 552 Fulton, R. J. 1967: Deglaciation studies in Kamloops region, an area of moderate relief, British Columbia.
553 *Geological Survey of Canada Bulletin* 154, 1-36.
- 554 Fulton, R. J. 1969: Surficial Geology, Shuswap Lake, British Columbia. *Geological Survey of Canada, "A"*
555 *Series Map* 1244A.

- 556 Fulton, R. J. 1991: A conceptual model for growth and decay of the Cordilleran Ice Sheet. *Géographie*
557 *physique et Quaternaire* 45, 281-286.
- 558 Fulton, R. J. & Walcott, R. I. 1975: Lithospheric flexure as shown by deformation of glacial lake shorelines
559 in southern British Columbia. *Geological Society of America, Memoir 142*, 163-173.
- 560 Fulton, R. J., Warner, B. G., Kubiw, H. J. & Achard, R. A. 1989: Geology and palaeoecology of early
561 Holocene lacustrine deposits in the Columbia River valley near Fauquier, southern British
562 Columbia. *Canadian Journal of Earth Sciences* 26, 257-265.
- 563 Fulton, R. J., Ryder, J. M. & Tsang, S. 2004: The Quaternary glacial record of British Columbia, Canada.
564 In Ehlers, J. & Gibbard P. L. (eds.): *Quaternary Glaciations - Extent and Chronology. Part II:*
565 *North America*, 39-50. Elsevier, Amsterdam.
- 566 Gardner, A. S., Moholdt, G., Cogley, J. G., Wouters, B., Arendt, A. A., Wahr, J., Berthier, E., Hock, R.,
567 Pfeffer, W. T., Kaser, G., Ligtenberg, S. R. M., Bolch, T., Sharp, M. J., Hagen, J. O., van den
568 Broeke, M. R. & Paul, F. 2013: A reconciled estimate of glacier contributions to sea level rise:
569 2003 to 2009. *Science* 340, 852-857.
- 570 Government of Canada. 2019: Open Government Portal. Available at:
571 <https://open.canada.ca/data/en/dataset?keywords=GeoBase> (accessed 07.10.2019).
- 572 Harrison, S., Glasser, N., Winchester, V., Haresign, E., Warren, C. & Jansson, K. 2006: A glacial lake
573 outburst flood associated with recent mountain glacier retreat, Patagonian Andes. *The Holocene*
574 16, 611-620.
- 575 Hickin, A. S., Lian, O. B., Levson, V. M. & Cui, Y. 2015: Pattern and chronology of glacial Lake Peace
576 shorelines and implications for isostasy and ice-sheet configuration in northeastern British
577 Columbia, Canada. *Boreas* 44, 288-304.

- 578 Johns, W. M. 1970: Geology and mineral deposits of Lincoln and Flathead counties, Montana. *Montana*
579 *Bureau of Mines and Geology, Bulletin 79*, 10-13.
- 580 Johnsen, T. F. & Brennand, T. A. 2004: Late-glacial lakes in the Thompson basin, British Columbia:
581 Paleogeography and evolution. *Canadian Journal of Earth Sciences 41*, 1367-1383.
- 582 Kehew, A. E., Milewski, A. & Soliman, F. 2010: Reconstructing an extreme flood from boulder transport
583 and rainfall–runoff modelling: Wadi Isla, South Sinai, Egypt. *Global and Planetary Change 70*,
584 64-75.
- 585 Kirkbride, M. P. & Warren, C. R. 1999: Tasman Glacier, New Zealand: 20th-century thinning and predicted
586 calving retreat. *Global and Planetary Change 22*, 11-28.
- 587 Korup, O. 2012: Earth's portfolio of extreme sediment transport events. *Earth-Science Reviews 112*, 115-
588 125.
- 589 Kjær, K. H., Sultan, L., Krüger, J. & Schomacker, A. 2004: Architecture and sedimentation of outwash fans
590 in front of the Mýrdalsjökull ice cap, Iceland. *Sedimentary Geology 172*, 139-163.
- 591 Kuehn, S. C., Froese, D. G., Carrara, P. E., Foit, F. F., Pearce, N. J. & Rotheisler, P. 2009: Major-and trace-
592 element characterization, expanded distribution, and a new chronology for the latest Pleistocene
593 Glacier Peak tephras in western North America. *Quaternary Research 71*, 201-216.
- 594 Kyle, G. W. 1938: *1938 Kootenay Lake order*. 1-7 pp. International Joint Commission, New York.
- 595 Lakeman, T. R., Clague, J. J. & Menounos, B. 2008: Advance of alpine glaciers during final retreat of the
596 Cordilleran ice sheet in the Finlay River area, northern British Columbia, Canada. *Quaternary*
597 *Research 69*, 188-200.

- 598 Lang, J. & Winsemann, J. 2013: Lateral and vertical facies relationships of bedforms deposited by
599 aggrading supercritical flows: from cyclic steps to humpback dunes. *Sedimentary Geology* 296, 36-
600 54.
- 601 Lang, J., Brandes, C. & Winsemann, J. 2017: Erosion and deposition by supercritical density flows during
602 channel avulsion and backfilling: Field examples from coarse-grained deepwater channel-levée
603 complexes (Sandino Forearc Basin, southern Central America). *Sedimentary Geology* 349, 79-102.
- 604 Larsen, D. J., Miller, G. H., Geirsdóttir, Á. & Thordarson, T. 2011: A 3000-year varved record of glacier
605 activity and climate change from the proglacial lake Hvítárvatn, Iceland. *Quaternary Science*
606 *Reviews* 30, 2715-2731.
- 607 Lesemann, J. & Brennand, T. A. 2009: Regional reconstruction of subglacial hydrology and glaciodynamic
608 behaviour along the southern margin of the Cordilleran Ice Sheet in British Columbia, Canada and
609 northern Washington State, USA. *Quaternary Science Reviews* 28, 2420-2444.
- 610 Leverington, D. W., Teller, J. T. & Mann, J. D. 2002: A GIS method for reconstruction of late Quaternary
611 landscapes from isobase data and modern topography. *Computers & Geosciences* 28, 631-639.
- 612 Liermann, S., Beylich, A. A. & van Welden, A. 2012: Contemporary suspended sediment transfer and
613 accumulation processes in the small proglacial Sætrevatnet sub-catchment, Bødalen, western
614 Norway. *Geomorphology* 167, 91-101.
- 615 Lønne, I. 1995: Sedimentary facies and depositional architecture of ice-contact glaciomarine systems.
616 *Sedimentary Geology* 98, 13-43.
- 617 Mathews, W. H. 1978: *Quaternary Stratigraphy and Geomorphology of Charlie Lake (94A) Map-area,*
618 *British Columbia.* 1-32 pp. Energy, Mines and Resources Canada.

- 619 Meetei, L. I., Pattanayak, S. K., Bhaskar, A., Pandit, M. K. & Tandon, S. K. 2007: Climatic imprints in
620 Quaternary valley fill deposits of the middle Teesta valley, Sikkim Himalaya. *Quaternary*
621 *International* 159, 32-46.
- 622 Menzies, J. 1990: Sand intraclasts within a diamicton mélange, southern Niagara Peninsula, Ontario,
623 Canada. *Journal of Quaternary Science* 5, 189-206.
- 624 Miall A. D. 1977: A review of the braided-river depositional environment. *Earth Science Reviews* 13, 1-
625 62.
- 626 Miyamoto, H., Itoh, K., Komatsu, G., Baker, V. R., Dohm, J. M., Tosaka, H. & Sasaki, S. 2006: Numerical
627 simulations of large-scale cataclysmic floodwater: A simple depth-averaged model and an
628 illustrative application. *Geomorphology* 76, 179-192.
- 629 Miyamoto, H., Komatsu, G., Baker, V. R., Dohm, J. M., Ito, K. & Tosaka, H. 2007: Cataclysmic Scabland
630 flooding: Insights from a simple depth-averaged numerical model. *Environmental Modelling &*
631 *Software* 22, 1400-1408.
- 632 Morison, S. R. & Hein, F. J. 1987: Sedimentology of the White Channel gravels, Klondike area, Yukon
633 Territory: fluvial deposits of a confined valley. In Ethridge, F. G., Flores, R. M., Harvey, M. D. &
634 Weaver, J. N. (eds.): *Recent Developments in Fluvial Sedimentology*, 206-216. *Society of Economic*
635 *Paleontologists and Mineralogists, Special Publication* 39.
- 636 Nasmith, H. (1962). Late glacial history and surficial deposits of the Okanagan Valley, British Columbia.
637 *British Columbia Department of Mines and Petroleum Resources, Bulletin* 46.
- 638 O'Connor, J. E. & Baker, V. R. 1992: Magnitudes and implications of peak discharges from glacial Lake
639 Missoula. *Geological Society of America Bulletin* 104, 267-279.

- 640 Pelto, M. S. & Warren, C. R. 1991: Relationship between tidewater glacier calving velocity and water depth
641 at the calving front. *Annals of Glaciology* 15, 115-118.
- 642 Peters, J. L. 2012: *Late Pleistocene evolution of glacial Lake Purcell: a potential floodwater source to the*
643 *Channeled Scabland*. M.Sc. thesis, Simon Fraser University, 1-224 pp.
- 644 Peters, J. L., Benetti, S., Dunlop, P., Ó Cofaigh, C., Moreton, S. G., Wheeler, A. J. & Clark, C. D. 2016:
645 Sedimentology and chronology of the advance and retreat of the last British-Irish Ice Sheet on the
646 continental shelf west of Ireland. *Quaternary Science Reviews* 140, 101-124.
- 647 Pisarska-Jamroży, M., Machowiak, K. & Krzyszkowski, D. 2010: Sedimentation style of a Pleistocene
648 kame terrace from the Western Sudety Mountains, S Poland. *Geologos* 16, 101-110.
- 649 Porter, S. C. & Swanson, T. W. 1998: Radiocarbon age constraints on rates of advance and retreat of the
650 Puget lobe of the Cordilleran ice sheet during the last glaciation. *Quaternary Research* 50, 205-
651 213.
- 652 Reimer, P. J., Bard, E., Bayliss, A., Beck, J. W., Blackwell, P. G., Ramsey, C. B., Buck, C. E., Hai, C.,
653 Edwards, R. L., Friedrich, M., Grootes, P. M., Guilderson, T. P., Haflidason, H., Hajdas, I., Hatte,
654 C., Heaton, T. J., Hoffmann, D. L., Hogg, A. G., Hughen, K. A., Kaiser, K. F., Kromer, B.,
655 Manning, S. W., Niu, M., Reimer, R. W., Richards, D. A., Scott, E. M., Southon, J. R., Staff, R.
656 A., Turney, C. S. M. & van der Plicht, J. 2013: IntCal13 and Marine13 radiocarbon age calibration
657 curves 0-50,000 years cal BP. *Radiocarbon* 55, 1869-1887.
- 658 Roeller, K. 2008: *Software for Geoscientists: Stereo32 and StereoNett*. Ruhr University Bochum, Bochum.
- 659 Röhl, K. 2006: Thermo-erosional notch development at fresh-water-calving Tasman Glacier, New Zealand.
660 *Journal of Glaciology* 52, 203-213.

- 661 Rushmer, E. L. 2006: Sedimentological and geomorphological impacts of the jökulhlaup (glacial outburst
662 flood) in January 2002 at Kverkfjöll, northern Iceland. *Geografiska Annaler: Series A, Physical*
663 *Geography* 88, 43-53.
- 664 Russell, A. J. 2009: Jökulhlaup (ice-dammed lake outburst flood) impact within a valley-confined sandur
665 subject to backwater conditions, Kangerlussuaq, West Greenland. *Sedimentary Geology* 215, 33-
666 49.
- 667 Ryder, J. M., Fulton, R. J. & Clague, J. J. 1991: The Cordilleran Ice Sheet and the glacial geomorphology
668 of southern and central British Columbia. *Géographie physique et Quaternaire* 45, 365-377.
- 669 Sawicki, O. & Smith, D. G. 1992: Glacial Lake Invermere, upper Columbia River valley, British Columbia:
670 a paleogeographic reconstruction. *Canadian Journal of Earth Sciences* 29, 687-692.
- 671 Sharp, M. 1982: Modification of clasts in logement tills by glacial erosion. *Journal of Glaciology* 28, 475-
672 480.
- 673 Shaw, J., Munro-Stasiuk, M., Sawyer, B., Beaney, C., Lesemann, J., Musacchio, A. & Young, R. R. 1999:
674 The Channeled Scabland: Back to Bretz? *Geology* 27, 605-608.
- 675 Smith, L. N. 2006: Pleistocene glacial deposits in the Libby and Lake River valley areas, Lincoln County,
676 Montana. *Northwest Geology* 35, 87-90.
- 677 Smith, N. D. & Ashley, G. M. 1985: Proglacial lacustrine environment. *Society of Economic Petrologists*
678 *and Mineralogists, Short Course* 16, 135-215.
- 679 Stuiver, M. & Reimer, P. J. 1993: Extended ¹⁴C database and revised CALIB radiocarbon calibration
680 program. *Radiocarbon* 35, 215-230.
- 681 Terpiłowski, S. 2007: Genesis of transverse kame trains in eastern Poland. *Sedimentary Geology* 193, 59-
682 69.

- 683 Thorson, R. M. 1989: Glacio-isostatic response of the Puget Sound area, Washington. *Geological Society*
684 *of America Bulletin* 101, 1163-1174.
- 685 Tsutaki, S., Nishimura, D., Yoshizawa, T. & Sugiyama, S. 2011: Changes in glacier dynamics under the
686 influence of proglacial lake formation in Rhonegletscher, Switzerland. *Annals of Glaciology* 52,
687 31-36.
- 688 USGS. (2019). National Geospatial Program. Available at: [https://www.usgs.gov/core-science-](https://www.usgs.gov/core-science-systems/national-geospatial-program/national-map)
689 [systems/national-geospatial-program/national-map](https://www.usgs.gov/core-science-systems/national-geospatial-program/national-map) (accessed 07.10.2019).
- 690 Waitt Jr, R. B. 1980: About forty last-glacial Lake Missoula jökulhlaups through southern Washington. *The*
691 *Journal of Geology* 88, 653-679.
- 692 Waitt, R. B. 2016: Megafloods and Clovis cache at Wenatchee, Washington. *Quaternary Research* 85, 430-
693 444.
- 694 Waitt, R. B., O'Connor, J. E. & Benito, G. 1994: *Scores of gigantic, successively smaller Lake Missoula*
695 *floods through Channeled Scabland and Columbia valley*. Paper presented at the Geologic field
696 trips in the Pacific Northwest (1994 Geological Society of America Annual Meeting), Seattle,
697 University of Washington.
- 698 Waitt, R. B., Denlinger, R. P. & O'Connor, J. E. 2009: Many monstrous Missoula floods down Channeled
699 Scabland and Columbia valley. *Field Guides* 15, 775-844.
- 700 Winsemann, J., Aspiron, U. & Meyer, T. 2004: Sequence analysis of early Saalian glacial lake deposits
701 (NW Germany): evidence of local ice margin retreat and associated calving processes. *Sedimentary*
702 *Geology* 165, 223-251.

703 Winsemann, J., Alho, P., Laamanen, L., Goseberg, N., Lang, J. & Klostermann, J. 2016: Flow dynamics,
704 sedimentation and erosion of glacial lake outburst floods along the Middle Pleistocene
705 Scandinavian Ice Sheet (northern central Europe). *Boreas* 45, 260-283.

706 Winsemann, J., Lang, J., Polom, U., Loewer, M., Igel, J., Pollok, L. & Brandes, C. 2018: Ice-marginal
707 forced regressive deltas in glacial lake basins: geomorphology, facies variability and large-scale
708 depositional architecture. *Boreas* 47, 973-1002.

709 Zen, E. & Prestegard, K. L. 1994: Possible hydraulic significance of two kinds of potholes: Examples from
710 the paleo-potomac river. *Geology* 22, 47-50.

711

712 **List of captions**

713 **Tables**

714 Table 1: Previously reported differential glacioisostatic adjustments associated with the CIS during MIS 2 assessed for this study.

Tilt ¹ (m km ⁻¹)	Location (glacial lake name)	Distance from LLGM limit ² (km)	Upslope direction	Reported age	Calibrated age range (median) cal. ka BP ³	Reference
0	Rocky Mountain Trench (Invermere)	120	NA	10±0.14 (ka BP)	11.2 – 12.0 (11.6)	Sawicki & Smith (1992)
0.2	SW Yukon Territory (Champagne)	<100	South	Unknown	NA	Gilbert & Desloges (2006)
~0.4	Peace Basin (Peace, Clayhurst stage)	100	West	11.6 (ka BP)	13.1 – 13.7 (13.4) ⁴	Mathews (1978)
0.46	Peace Basin (Peace Phase II)	100	230.9°±5°	<16.0±2.5 (optical)	NA	Hickin <i>et al.</i> (2015)
0.6	Shuswap Basin (Shuswap, Rocky Point stage)	200	East	Unknown	NA	Fulton (1969)
0.7	Okanogan Valley (Penticton, B.X. stage)	140	North	8.41±0.1 (ka BP)	9.1 – 9.5 (9.4)	Nasmith (1962)
0.85	Puget Lowland (Russell-Hood)	20	North	13.7 (ka BP)	16.1 – 17.0 (16.6) ⁴	Thorson (1989)
0.9	Columbia River valley (“glacial Lake Arrow”)	100	North	10.1±0.15 (ka BP)	11.2 – 12.2 (11.7)	Fulton <i>et al.</i> (1989)
1.15	Puget Lowland (Bretz, western)	10	North	13.5 (ka BP)	15.8 – 16.8 (16.3) ⁴	Thorson (1989)
1.6 ± 0.7	Nicola Basin (Hamilton, lower stage)	200	North (347°±7°)	Unknown	NA	Fulton & Walcott (1975)
1.7 ± 0.4	Thompson Basin (Deadman, lowest stage)	200	Northwest (321°±6.1°)	~10.9 – 11.1 (cal. ka BP)	10.9 – 11.1 (11.0)	Johnsen & Brennand (2004)
1.8 ± 0.6	Nicola Basin (Merritt)	200	Northwest (341°±18°)	Unknown	NA	Fulton & Walcott (1975)
1.8 ± 0.6	Nicola Basin (Hamilton, upper stage)	200	North (354°±11°)	Unknown	NA	Fulton & Walcott (1975)
1.8 ± 0.7	Thompson Basin (Thompson, high stage)	200	Northwest (332°±9.9°)	>10.9 – 11.1 (cal. ka BP)	10.9 – 11.1 (11.0)	Johnsen & Brennand (2004)
~2.1	Nicola Basin (Quilchena)	200	Northwest	≥8 900	9.5 – 10.3 (10.0)	Fulton & Walcott (1975)

715 ¹ Land surface tilt caused by differential glacioisostatic adjustment, as recorded by palaeo-lake-level indicators.

716 ² Approximate distances between the glacial lake’s nearest margin of the CIS during the LLGM from Fulton *et al.*
717 (2004). GLP was ~40 km from the LLGM limit, perhaps <13.5 cal. ka BP (Waite *et al.* 2009).

718 ³ Radiocarbon ages calibrated for this study with Calib software version 7.10 (Stuiver & Reimer 1993) using the
719 IntCal13 radiocarbon curve (Reimer *et al.* 2013). Reported as the 2σ median probability (e.g. Peters *et al.* 2016).

720 ⁴ Calibration was performed using an assumed standard error of ±160 (the highest reported in this review, to avoid
721 spurious precision) because insufficient information was reported.

722

723 Table 2: Dimensions (area and volume) of gLP and its ice dam for the tested range of glacioisostatic tilts.

Water-plane tilt (m km ⁻¹)	Lake area ¹ (km ²)	Lake volume ¹ (km ³)	Maximum drainable volume ² (km ³)	Water surface elevation ³ (m a.s.l.)	Water depth ⁴ (m)	Minimum ice dam thickness ⁵ (m)	Height of water surface above pre-GLOF valley bottom ⁶ (m)
0	600	40	14	655	257	286	15
0.50	663	69	43	723	325	361	83
0.75	667	73	47	756	358	398	116
1.00	671	93	67	788	390	433	148
1.25	1152	142	116	817	419	466	177

724 ¹ Location of ice dam placed at 49°36'55.7"N, 116°52'21.9"W (~132 km north of the Elmira spillway); a calving
725 margin is assumed (see text).

726 ² Lake volume minus the volume of Kootenay Lake within the palaeolake extent.

727 ³ Maximum elevations, as measured against the ice dam.


728 ⁴ Maximum depths, measured against the ice dam.

729 ⁵ Clean ice thickness required to resist flotation at the dam following the 9/10th ratio of ice to water densities
730 (Thorarinsson 1939; Fowler 1999).

731 ⁶ Height of water surface above the height of the highest terrace in the Kootenay River valley (640 m a.s.l.).


732

733 **Figures**



735 Fig. 1: A. Study area (red box) at the western USA-Canada border showing the Channeled
736 Scabland and CIS extent at the Last Glacial Maximum (LGM; Fulton *et al.* 2004). The Purcell
737 Lobe's southern terminus at ~13.5 ka BP (dashed black line) is mapped after Dyke *et al.* (2003).
738 CF = Chelan Falls. B. Schematic map of the study area depicting the locations of sedimentary
739 deposits (kame; glacial Lake Purcell (gLP) lake bed sediments; gLP drainage sediments; gLP
740 Glacial Lake Outburst Flood (GLOF) sediments; glacial Lake Kootenai flood-related fan),
741 palaeoflow measurements (lower hemisphere, equal area projection and rose diagram) and other
742 locations discussed in the text. Note that GLOF sediments are present throughout the Kootenay
743 River valley, but only major exposures are mapped. Also, the West Arm of Kootenay Lake (east
744 of the Corra Linn Dam) and the Kootenay River are jointly referred to in the text as the Kootenay
745 River valley (KRv) and changes in river width between these water bodies are not represented to
746 enhance illustrative clarity. Red box delineates the area shown in Fig. 3. BF = Bonners Ferry; PJ
747 = Playmor Junction.


748



749
 750 Fig. 2: Schematic representation of glacial lake evolution showing. A. Discrete stages of glacial
 751 Lakes Purcell (gLP) and Kootenai (gLK) (dark blue) impounded behind the Purcell Lobe (light
 752 blue, PL) sometime after the Last Late Glacial Maximum (LLGM) (likely <math>< 17.4 \text{ cal. ka BP}</math>);

753 Atwater 1987; Porter & Swanson 1998; Clague & James 2002). Both the Elmira spillway (ES) at
 754 710 m a.s.l. and Bull River spillway (BRS) at 732 m a.s.l. are active (shown as blue arrows).
 755 During this stage Alden (1953) considered gLP an unnamed proglacial lake. B. Continued retreat
 756 of the Purcell Lobe causes the growth of gLP and its northward expansion. Eventually the portion
 757 of gLK above 710 m a.s.l. (the water above the Elmira spillway) decanted into gLP (shown as red
 758 arrow), rapidly reducing gLK's volume. C. Glacial Lake Kootenai has largely drained into gLP
 759 and may not exist at all. The late deglacial Purcell Lobe retreats northward towards the Kootenay
 760 River valley (KRv) until it fails to dam gLP. GLP debouches into the Kootenay River valley (red
 761 arrow) and eventually its floodwaters reach the Channeled Scabland via the Columbia River (Fig.
 762 1). Note that the naming scheme of Alden (1953) is abandoned and replaced with the one depicted
 763 by this schematic because it ignores gLP's nascent formation and considers later stages of gLP to
 764 be the same lake as gLK. BC = British Columbia; WA = Washington; ID = Idaho; MT = Montana.

765




766

767 Fig. 3: Hillshaded DEM from a composite of Geobase (Government of Canada 2019) and National
 768 Elevation Dataset (United States Geological Survey) data (USGS 2019) revealing benches A and

769 B incised into the gLP lake bed deposits. Dashed lines delineate the contact between the lake bed
770 sediment and the valley walls. Point elevations (white dots with Xs) that typify elevation data
771 used to reconstruct the pre-incision gLP lake bed are provided and highlight the relative flatness
772 of the deposit's surface across the Purcell Trench. Channels 'A' and 'B' correspond in elevation
773 with benches 'A' and 'B', respectively. Site 10 is shown as a labelled white dot within Channel
774 B.

775




776

777 Fig. 4: Examples of sediment deposits. Sites are located in Fig. 1B. A. Massive and laminated
 778 silt deposits are common throughout the lake bed bench (photograph is from site 5). B. Massive
 779 silt with abundant, outsized clasts (lonestones) at site 15. Knife blade is ~9 cm long. C. Massive,
 780 coarse sand laminae, interlaminated with silt and clay laminae that conformably overlie a well-

781 rounded, granitic lonestone at site 5. D. Angular, unconsolidated sand clasts (outlined by white
 782 dashed lines) in an interfluvial-occupying, valley-wall deposit (kame) near gravel fabric K1. Knife
 783 handle is ~9 cm long. E. Planar-stratified sand and gravel in a kame terrace. Arrow marks metre
 784 stick for scale. Location of gravel fabric K2 is shown as a labelled white dot. F. Climbing ripples
 785 (after Ashley *et al.* 1982) measured for palaeoflow Dr1 from a ~560 m a.s.l. gLP GLOF terrace in
 786 the Kootenay River valley. G. Inclined gravel beds (white dashed lines highlight two lower
 787 contacts) overlain by massive silt at site 10 (Fig. 3). Arrow points to a person for scale. H. Poorly
 788 sorted cobble and boulder gravel at ~490 m a.s.l. in a gLP GLOF expansion bar measured for
 789 gravel fabric D3. Ruler is 36 cm long. I. Planar-stratified sand and gravel and diffusely graded,
 790 sinusoidally stratified sand composes a ~640 m a.s.l. gLP GLOF terrace in the Kootenay River
 791 valley. Location of gravel fabric D1 shown as a labelled white dot. Arrow marks metre stick for
 792 scale. p.flow is palaeoflow.

793




794

795 Fig. 5: Plot of the ten glacioisostatic tilts derived from CIS palaeolake planes with reported ages
 796 (black and grey dots, Table 1). Glacial lakes Arrow (Ar) and Invermere (Iv) are highlighted grey
 797 because they are geographically near the the Purcell Trench and Glacial Lake Peace, Clayhurst

798 stage (PC), because it's chronologically closest to putative ages for gLP (delineated by the labelled
 799 grey bar; Dyke *et al.* 2003; Waitt *et al.* 2009; Table 1). The four undated GIAs reviewed are
 800 marked as chevrons. The estimated age of gLP drainage is derived from previous CIS
 801 reconstructions (Dyke *et al.* 2003) and tephrochronologic ages (Waitt *et al.* 2009). The distribution
 802 of the previously reported lake tilts reveals the propriety of the modelled GIAs.


803



804

805 Fig. 6: Extent of gLP based on the array of tested glacioisostatic tilts (Table 2). Note that only the
 806 steepest tilt (1.25 m km^{-1}) covers the gLP lake bed sediment bench (red dashed line; Fig. 1A) and
 807 allows gLP to overtop the flood-related fan formed by gLK's drainage (Alden 1953) which is
 808 capped by silty lake bed sediments at site 10 (red dot). This suggests that the fan was partially
 809 inundated by gLP. The locations of modern Kootenay Lake (black), the Kootenay River valley
 810 (KRv), and the Purcell Lobe <13.5 cal. ka BP (Dyke *et al.* 2003) are also shown. Only the
 811 northernmost extent of the gLP lakebed is outlined for visual clarity with other aspects of this
 812 figure; see Fig. 1A for the complete extent.


813



814

815 Fig. 7: Schematic representation of three modelled gLP water surface tilts (Table 2) projected
 816 northwards from the Elmira spillway, past the alluvial fan deposited by the drainage of gLK (Alden
 817 1953) and over the reconstructed gLP lake bed. The elevations of the pre-incision Kootenay River
 818 valley sediment fill and modern Kootenay Lake are also depicted (note the depth of Kootenay
 819 Lake during glaciation is schematically represented).

820



821
 822 Fig. 8: Kootenay River valley (KRv; Fig. 1B) geomorphology. A. Kootenay River valley long
 823 profile cartoon depicting the relationship between the most likely gLP elevation prior to drainage

824 (817 m a.s.l. at a 1.25 m km^{-1} tilt), the highest truncated alluvial fan deposits ($\sim 675 \text{ m a.s.l.}$), the
825 highest valley-flanking terrace (depicting the pre-GLOF valley bottom), and the modern Kootenay
826 River (bold blue line). Low-elevation, untruncated alluvial fans aggrading into the Kootenay River
827 valley are shown as brown polygons. Locations of GLOF palaeoflow measurements and potholes
828 are provided (see Fig. 1B for map view). B. Eight Kootenay River valley, cross-sectional profiles
829 (from Geobase, Natural Resources Canada DEM; Government of Canada 2019), grouped to
830 represent three reaches of the Kootenay River valley. The groups are shown with stacked profiles
831 that reveal trends in bench and remnant fan elevations and allow estimations of pre-incision valley-
832 fill elevations. Estimated extents of truncated alluvial fan remnants are shown as dotted lines. C.
833 Terrace treads incised into an expansion bar at the Kootenay River valley (KRv)-Columbia River
834 valley (Rv) confluence (see Fig. 1B for regional map) (DEM (Geobase, Natural Resources Canada)
835 overlain by a georeferenced aerial photograph (National Aerial Photograph Library, Environment
836 Canada)). Contacts between the valley-fill sediment and the bedrock valley walls are shown as
837 dashed lines. Solid black lines outline individual terraces identified using the DEM. The location
838 of gravel fabric D3 (Figs 1B, 6H) is shown as a labelled red dot.

839