

Title	Exploiting the continuous in situ generation of mesyl azide for use in a telescoped process
Authors	O'Mahony, Rosella M.;Lynch, Denis;Hayes, Hannah L. D.;Ní Thuama, Eilís;Donnellan, Philip;Jones, Roderick C.;Glennon, Brian;Collins, Stuart G.;Maguire, Anita R.
Publication date	2017-08-21
Original Citation	O'Mahony, R. M., Lynch, D., Hayes, H. L. D., Ní Thuama, E., Donnellan, P., Jones, R. C., Glennon, B., Collins, S. G. and Maguire, A. R. [2017] 'Exploiting the Continuous in situ Generation of Mesyl Azide for Use in a Telescoped Process', European Journal of Organic Chemistry, 2017(44), pp. 6533-6539. doi: 10.1002/ejoc.201700871
Type of publication	Article (peer-reviewed)
Link to publisher's version	https://chemistry-europe.onlinelibrary.wiley.com/doi/full/10.1002/ejoc.201700871 - 10.1002/ejoc.201700871
Rights	© 2017 Wiley-VCH Verlag GmbH & Co. KGaA, Weinheim. This is the peer reviewed version of the following article: O'Mahony, R.M. et al. (2017), Exploiting the Continuous in situ Generation of Mesyl Azide for Use in a Telescoped Process. Eur. J. Org. Chem., 2017, which has been published in final form at https://doi.org/10.1002/ejoc.201700871 This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Self-Archiving.
Download date	2024-04-20 01:03:35
Item downloaded from	https://hdl.handle.net/10468/10769



University College Cork, Ireland
Coláiste na hOllscoile Corcaigh

SUPPORTING INFORMATION

Title: Exploiting the Continuous in situ Generation of Mesyl Azide for Use in a Telescoped Process

Author(s): Rosella M. O'Mahony, Denis Lynch, Hannah L. D. Hayes, Eilís Ní Thuama, Philip Donnellan, Roderick C. Jones, Brian Glennon, Stuart G. Collins,* Anita R. Maguire*

Contents

1. Experimental Continuous-flow Setup.....	3
2. Copies of ^1H and ^{13}C NMR Spectra for Compounds 4g , 7a-g and 8	5

1. Experimental Continuous-flow Setup

All continuous processes were performed using either a Vapourtec R-Series flow reactor or a Vapourtec E-Series flow reactor.

The R-Series flow reactor consists of four piston pumps and up to four temperature controlled tubular reactors. To prepare the reactor for operation pumps were purged with the solvent to be used in the reaction prior to use. All reaction tubing, coils, inlets and connections were also purged thoroughly in a similar manner.

Table S1: General specifications for R-Series continuous-flow reactor

General Specifications	
Material of tubing	PFA
Diameter of tubing	1 mm
Working flow rates	0.05 mL/min – 9.99 mL/min
Tubular reactor working volume	10 mL
Temperature range	-70 °C to 250 °C

The Vapourtec E-Series flow reactor was used for the final telescoped reactions alongside the R-series. The E-Series flow reactor consists of three peristaltic pumps and up to two temperature controlled tubular reactors. To prepare the reactor for operation pumps were, again, purged with the solvent to be used in the reaction prior to use. All reaction tubing, coils, inlets and connections were also purged thoroughly in a similar manner.

Table S2: General specifications for E-Series continuous-flow reactor

General Specifications	
Material of tubing	PFA
Diameter of tubing	1 mm
Working flow rates	0.02 mL/min – 10.0 mL/min
Tubular reactor working volume	10 mL
Temperature range	-70 °C to 250 °C




Figure S1: Experimental Continuous Flow Setup.

2. Copies of ^1H and ^{13}C NMR Spectra for Compounds 4g and 7a–g




Figure S2.1: ^1H NMR (400 MHz, CDCl_3) spectrum of **4g**




Figure S2.2: ^{13}C NMR (100.6 MHz, CDCl_3) spectrum of **4g**.




Figure S2.3: ¹H NMR (400 MHz, CDCl₃) spectrum of **7a**.




Figure S2.4: ¹³C NMR (100.6 MHz, CDCl₃) spectrum of **7a**.




Figure S2.5: ¹H NMR (400 MHz, CDCl₃) spectrum of **7b**.





Figure S2.6: ¹³C NMR (100.6 MHz, CDCl₃) spectrum of **7b**.






Figure S2.9: ¹H NMR (400 MHz, CDCl₃) spectrum of **7d**.







Figure S2.10: ¹³C NMR (100.6 MHz, CDCl₃) spectrum of **7d**.









Figure S2.15: ^1H NMR (400 MHz, CDCl_3) spectrum of **7g**.






Figure S2.17: ¹H NMR (400 MHz, CDCl₃) spectrum of **8**.




Figure S2.18: ¹³C NMR (100.6 MHz, CDCl₃) spectrum of **8**.