CORA Cork Open Research Archive
Cartlann Taighde Oscailte Chorcaí

Title	Lactobacillus gasseri APC 678 reduces shedding of the pathogen Clostridium difficile in a murine model
Authors	Quigley, Lisa;Coakley, Mairead;Alemayehu, Debebe;Rea, Mary C.;Casey, Patrick G.;O'Sullivan, Órla;Murphy, Eileen;Kiely, Barry;Cotter, Paul D.;Hill, Colin;Ross, R. Paul
Publication date	2019-02-20
Original Citation	Quigley, L., Coakley, M., Alemayehu, D., Rea, M.C., Casey, P.G., O'Sullivan, O., Murphy, E., Kiely, B., Cotter, P.D., Hill, C. and Ross, P., 2019. Lactobacillus gasseri APC 678 Reduces Shedding of the Pathogen Clostridium difficile in a Murine Model. Frontiers in microbiology, 10, (273). DOI:10.3389/fmicb.2019.00273
Type of publication	Article (peer-reviewed)
Link to publisher's version	https://www.frontiersin.org/articles/10.3389/fmicb.2019.00273 - 10.3389/fmicb.2019.00273
Rights	© 2019 Quigley, Coakley, Alemayehu, Rea, Casey, O'Sullivan, Murphy, Kiely, Cotter, Hill and Ross - https://creativecommons.org/licenses/by/4.0/
Download date	2026-11-07 10:27:13
Item downloaded from	https://hdl.handle.net/10468/9152

UCC
University College Cork, Ireland
Coláiste na hOllscoile Corcaigh

Supplementary Material

Lactobacillus gasseri APC 678 Reduces Shedding of the Pathogen *Clostridium difficile* in a Murine Model


Lisa Quigley[#], Mairéad Coakley[#], Debebe Alemayehu, Mary C. Rea, Patrick G. Casey, Órla O'Sullivan, Eileen Murphy, Barry Kiely, Paul D. Cotter, Colin Hill and R. Paul Ross*

***Correspondence:**
Prof. R. Paul Ross
p.ross@ucc.ie


[#] LQ and MC are joint first authors

1.1 Supplementary Figures


Supplementary Figure 1: Relative abundance (%) of bacterial phyla in the cecum at Day 7 of the control and test mice (*Lactobacillus gasseri* APC 678, *Lactobacillus rhamnosus* DPC 6111 and *Lactobacillus gasseri* ATCC 33323).



Supplementary Figure 2: Relative abundance (%) of bacterial families in the cecum at Day 7 of the control mice and test mice (*Lactobacillus gasseri* APC 678, *Lactobacillus rhamnosus* DPC 6111 and *Lactobacillus gasseri* ATCC 33323).



Supplementary Figure 3: Relative abundance (%) of bacterial genera in the cecum at Day 7 of the control mice and test mice (*Lactobacillus gasseri* APC 678, *Lactobacillus rhamnosus* DPC 6111 and *Lactobacillus gasseri* ATCC 33323).



1.2 Supplementary Tables

Supplementary Table 1: Source and origin of bacterial strains used during this study.

Bacterial strain	Source	Origin
Clostridium difficile EM304 (ribotype 027)	ELDERMET Culture Collection[1]	elderly adult feces
Clostridium difficile VPI 10463 (ATCC® 43255-FZ™)	American Type Culture Collection[2]	human abdominal wound
L. gasseri APC 678	APC Culture Collection[1]	human feces
L. paracasei APC 1483	APC Culture Collection	human feces
L. rhamnosus DPC 6111	DPC Culture Collection[1]	human feces
L. gasseri DPC 6112	DPC Culture Collection	human feces
L. gasseri ATCC 33323	American Type Culture Collection	human gastrointestinal tract

[1]housed at Teagasc, Moorepark, Fermoy, Co. Cork, Ireland; [2]Manassas, VA 20110, USA.

Supplementary Table 2: Target strains, growth medium and incubation conditions for well diffusion assays to detect bacteriocin production by *L. gasseri* APC 678 *in vitro*.

Target strain	**Culture Collection code**	**Incubation conditions**	**Growth medium**
Enterococcus faecium	LMG[1] 11423	37°C, anaerobic	MRS
Enterococcus mundtii	LMG 10758	37°C, anaerobic	MRS
Enterococcus saccharolyticus	LMG 11427	37°C, anaerobic	MRS
Lactobacillus acidophilus	LMG 9433	37°C, anaerobic	MRS
Lactobacillus agilis	LMG 9186	37°C, anaerobic	MRS
Lactobacillus amylovorus	LMG 9496	37°C, anaerobic	MRS
Lactobacillus casei	LMG 6904	37°C, anaerobic	MRS
Lactobacillus crispatus	LMG 9479	37°C, anaerobic	MRS
Lactobacillus delbrueckii subsp. *bulgaricus*	LMG 6901	37°C, anaerobic	MRS
Lactobacillus delbrueckii subsp. *lactis*	LMG 7942	37°C, anaerobic	MRS
Lactobacillus fermentum	LMG 6902	37°C, anaerobic	MRS
Lactobacillus plantarum	LMG 6907	37°C, anaerobic	MRS
Lactobacillus rhamnosus	ATCC[2] 53103	37°C, anaerobic	MRS
Listeria innocua	DPC[3] 3572	37°C, aerobic	BHI*
Micrococcus luteus	DPC 6275	30°C, aerobic	BHI
Salmonella typhimurium	DPC 6048	37°C, aerobic	BHI
Staphylococcus aureus	DPC 5246	37°C, aerobic	BHI

[1]LMG Culture Collection, B-9000 Gent, Belgium; [2]American Type Culture Collection, Manassas, VA 20110, USA and [3]DPC Culture Collection, Teagasc, Moorepark, Fermoy, Co. Cork, Ireland. *BHI: Brain heart infusion medium, Merck, Darmstadt, Germany.

Supplementary Table 3: Relative abundance (%) of bacterial phyla in the cecum at Day 7 of the control and test mice (*Lactobacillus gasseri* APC 678, *Lactobacillus rhamnosus* DPC 6111 and *Lactobacillus gasseri* ATCC 33323). Average and standard error of the mean (SEM) of results for 10 mice per treatment are represented.

Group	Control	APC 678	DPC 6111	ATCC 33323
	relative abundance (%)			
Firmicutes	54.51±7.32	37.22±4.18	28.44±2.32*	31.29±3.63*
Bacteroidetes	43.17±7.19	62.24±4.19	70.62±2.31*	66.25±3.88*
Verrucomicrobia	1.53±1.42	0.08±0.03	0.35±0.12	0.93±0.76
Proteobacteria	0.47±0.08	0.19±0.06*	0.09±0.03*	1.29±0.44
Actinobacteria	0.26±0.05	0.18±0.03	0.19±0.04	0.13±0.03*
Tenericutes	0.01±0.01	0.03±0.02	0.26±0.14	0.10±0.07
Cyanobacteria	0.00±0.00	0.00±0.00	0.00±0.00	0.00±0.00
Other	0.04±0.01	0.05±0.01	0.05±0.01	0.02±0.01

*significantly different compared to the control; significance was determined by $p<0.05$.

Supplementary Table 4: Relative abundance (%) of bacterial families in the cecum at Day 7 of the control and test mice (*Lactobacillus gasseri* APC 678, *Lactobacillus rhamnosus* DPC 6111 and *Lactobacillus gasseri* ATCC 33323). Average and standard error of the mean (SEM) of results for 10 mice per treatment are represented.

Group	**Control**	**APC 678**	**DPC 6111**	**ATCC 33323**
	relative abundance (%)			
Lachnospiraceae	40.346±5.285	27.461±3.459	20.141±2.049*	23.600±3.273*
S24-7	24.104±5.359	31.182±2.779	40.720±3.787*	23.214±4.942
Bacteroidaceae	9.289±3.278	21.603±4.812	18.614±3.699	28.214±5.646*
Porphyromonadaceae	7.918±2.407	1.611±0.396	4.213±1.364	6.710±1.656
Ruminococcaceae	7.743±1.458	6.576±1.054	5.874±0.974	4.957±0.678
uncultured *Clostridiales*	2.008±0.655	0.639±0.130	0.213±0.082*	0.582±0.284*
Rikenellaceae	1.859±0.696	7.314±0.900*	6.515±0.753*	8.112±1.439*
Lactobacillaceae	1.835±0.515	1.078±0.480	1.259±0.361	1.036±0.229
Erysipelotrichaceae	1.804±0.370	0.616±0.110*	0.829±0.188*	0.948±0.341*
Verrucomicrobiaceae	1.531±1.419	0.084±0.029	0.354±0.121	0.928±0.755
Peptostreptococcaceae	0.603±0.245	0.078±0.054*	0.019±0.013*	0.050±0.031*
Alcaligenaceae	0.286±0.060	0.085±0.043*	0.069±0.032*	1.235±0.429
Enterobacteriaceae	0.177±0.072	0.010±0.004*	0.017±0.009*	0.055±0.026
Coriobacteriaceae	0.130±0.040	0.140±0.025	0.167±0.036	0.063±0.018
Bifidobacteriaceae	0.126±0.031	0.039±0.016*	0.024±0.011*	0.065±0.041*
Christensenellaceae	0.103±0.046	0.375±0.178	0.000±0.000	0.012±0.006
Enterococcaceae	0.030±0.008	0.023±0.009	0.007±0.003*	0.013±0.007
Clostridiales* Family XIII *Incertae Sedis	0.019±0.006	0.028±0.005	0.029±0.010	0.019±0.005
Clostridiaceae	0.016±0.010	0.000±0.000	0.002±0.002	0.001±0.001
Anaeroplasmataceae	0.013±0.011	0.033±0.023	0.258±0.140	0.097±0.075
Oxalobacteraceae	0.005±0.002	0.000±0.000	0.000±0.000	0.000±0.000
Peptococcaceae	0.004±0.002	0.041±0.009*	0.020±0.007	0.066±0.049
Rhodospirillaceae	0.002±0.001	0.008±0.004	0.000±0.000	0.000±0.000
Staphylococcaceae	0.002±0.001	0.288±0.149	0.045±0.035	0.000±0.000
Streptococcaceae	0.001±0.001	0.001±0.000	0.000±0.000	0.000±0.000
***Rickettsiales* mitochondria**	0.001±0.001	0.000±0.000	0.000±0.000	0.000±0.000
Cyanobacteria 4C0d-2 uncultured	0.001±0.001	0.000±0.000	0.000±0.000	0.000±0.000
Moraxellaceae	0.001±0.001	0.005±0.002	0.000±0.000	0.000±0.000
Planococcaceae	0.000±0.000	0.012±0.005	0.000±0.000	0.000±0.000
Prevotellaceae	0.000±0.000	0.533±0.198	0.561±0.201*	0.001±0.001
***Bacteroidales* ratAN060301C**	0.000±0.000	0.000±0.000	0.000±0.000	0.000±0.000
Bacillaceae	0.000±0.000	0.001±0.001	0.000±0.000	0.000±0.000
Methylobacteriaceae	0.000±0.000	0.000±0.000	0.000±0.000	0.001±0.001
Comamonadaceae	0.000±0.000	0.002±0.001	0.000±0.000	0.000±0.000
Desulfovibrionaceae	0.000±0.000	0.024±0.024	0.000±0.000	0.000±0.000
Xanthomonadaceae	0.000±0.000	0.057±0.048*	0.001±0.001	0.000±0.000
Others	0.045±0.014	0.051±0.005	0.047±0.010	0.018±0.010

*significantly different compared to the control; significance was determined by $p<0.05$.

Supplementary Table 5: Relative abundance (%) of bacterial genera in the cecum at Day 7 of the control and test mice (*Lactobacillus gasseri* APC 678, *Lactobacillus rhamnosus* DPC 6111 and *Lactobacillus gasseri* ATCC 33323). Average and standard error of the mean (SEM) of results for 10 mice per treatment are represented.

Genus	Control	APC 678	DPC 6111	ATCC 33323
	relative abundance (%)			
uncultured *Lachnospiraceae* A	29.985±3.699	20.933±3.149	14.777±2.014*	18.000±3.312*
uncultured S24-7	24.104±5.359	31.182±2.779	40.720±3.787*	23.214±4.942
Lachnospiraceae Incertae Sedis	9.665±2.069	5.321±0.918	4.994±1.722	4.906±0.772
Bacteroides	9.289±3.278	21.603±4.812	18.614±3.699	28.214±5.646*
Parabacteroides	7.917±2.407	1.611±0.396	4.213±1.364	6.710±1.656
Ruminococcaceae Incertae Sedis	5.364±1.393	2.185±0.485*	1.441±0.300*	2.014±0.454*
uncultured *Clostridiales*	2.008±0.655	0.639±0.130	0.213±0.082*	0.582±0.284*
Lactobacillus	1.835±0.515	1.078±0.480	1.259±0.361	1.036±0.229
Akkermansia	1.531±1.419	0.084±0.029	0.354±0.121	0.928±0.755
Alistipes	1.406±0.507	3.395±0.343*	4.949±0.585*	6.034±0.907*
uncultured *Ruminococcaceae*	1.081±0.189	2.031±0.339*	0.771±0.286	1.308±0.441
Erysipelotrichaceae Incertae Sedis	0.953±0.392	0.171±0.062	0.290±0.060	0.268±0.087
Allobaculum	0.691±0.346	0.401±0.133	0.449±0.150	0.294±0.139
Peptostreptococcaceae Incertae Sedis	0.603±0.245	0.078±0.054*	0.019±0.013*	0.050±0.031*
Anaerotruncus	0.601±0.155	1.090±0.252	1.846±0.399*	0.755±0.154
***Rikenellaceae* RC9 gut group**	0.453±0.319	3.895±0.815*	1.566±0.590*	2.078±0.956
Oscillibacter	0.363±0.199	1.151±0.258*	1.805±0.421*	0.787±0.291
Parasutterella	0.286±0.060	0.085±0.043*	0.069±0.032*	1.235±0.429
Blautia	0.267±0.105	0.076±0.072	0.030±0.017	0.132±0.103
Flavonifractor	0.251±0.176	0.012±0.010*	0.007±0.005*	0.087±0.058
Roseburia	0.207±0.083	0.903±0.291*	0.212±0.104	0.288±0.228
Escherichia/Shigella	0.177±0.072	0.010±0.004*	0.017±0.009*	0.055±0.026
uncultured *Erysipelotrichaceae* A	0.138±0.030	0.036±0.008*	0.054±0.018*	0.039±0.009*
Anaerostipes	0.129±0.040	0.064±0.049	0.064±0.023	0.065±0.040
Enterorhabdus	0.128±0.040	0.135±0.023	0.159±0.034	0.059±0.017
Bifidobacterium	0.126±0.031	0.039±0.016*	0.024±0.011*	0.065±0.041*
uncultured *Christensenellaceae*	0.102±0.046	0.375±0.178	0.000±0.000	0.008±0.005
Ruminococcus	0.082±0.045	0.102±0.072	0.001±0.001	0.005±0.004
Coprococcus	0.053±0.030	0.063±0.015	0.017±0.014	0.020±0.009
Marvinbryantia	0.038±0.007	0.099±0.042	0.047±0.031	0.188±0.079
Enterococcus	0.030±0.008	0.023±0.009	0.007±0.003*	0.013±0.007
Clostridiales* Family XIII *Incertae Sedis	0.019±0.006	0.025±0.004	0.029±0.010	0.012±0.004
uncultured *Erysipelotrichaceae* B	0.017±0.007	0.009±0.004	0.026±0.012	0.011±0.008
Clostridium	0.016±0.010	0.000±0.000	0.002±0.002	0.001±0.001
Anaeroplasma	0.013±0.011	0.033±0.023	0.258±0.140	0.097±0.075
Oxalobacter	0.005±0.002	0.000±0.000	0.000±0.000	0.000±0.000
Coprobacillus	0.004±0.004	0.000±0.000	0.010±0.010	0.337±0.275
uncultured *Peptococcaceae*	0.004±0.002	0.025±0.005*	0.006±0.002	0.066±0.049

Acetitomaculum	0.002±0.001	0.000±0.000	0.000±0.000	0.000±0.000
Thalassospira	0.002±0.001	0.008±0.004	0.000±0.000	0.000±0.000
Staphylococcus	0.002±0.001	0.287±0.149	0.045±0.035	0.000±0.000
Streptococcus	0.001±0.001	0.001±0.000	0.000±0.000	0.000±0.000
Christensenella	0.001±0.001	0.000±0.000	0.000±0.000	0.004±0.002
uncultured *Coriobacteriaceae*	0.001±0.001	0.005±0.002	0.007±0.005	0.004±0.003
Idiospermum	0.001±0.001	0.000±0.000	0.000±0.000	0.000±0.000
uncultured Cyanobacteria 4C0d-2	0.001±0.001	0.000±0.000	0.000±0.000	0.000±0.000
Acinetobacter	0.001±0.001	0.005±0.002	0.000±0.000	0.000±0.000
Sporosarcina	0.000±0.000	0.012±0.005	0.000±0.000	0.000±0.000
Barnesiella	0.000±0.000	0.000±0.000	0.000±0.000	0.000±0.000
Prevotella	0.000±0.000	0.533±0.198	0.561±0.201*	0.000±0.000
uncultured *Prevotellaceae*	0.000±0.000	0.000±0.000	0.000±0.000	0.001±0.001
Rikenella	0.000±0.000	0.025±0.021	0.000±0.000	0.000±0.000
uncultured *Bacteroidales* ratAN060301C	0.000±0.000	0.000±0.000	0.000±0.000	0.000±0.000
Bacillus	0.000±0.000	0.001±0.001	0.000±0.000	0.000±0.000
Salinicoccus	0.000±0.000	0.001±0.001	0.000±0.000	0.000±0.000
Anaerovorax	0.000±0.000	0.000±0.000	0.000±0.000	0.001±0.001
uncultured *Clostridiales* Family XIII *Incertae Sedis*	0.000±0.000	0.003±0.002	0.000±0.000	0.007±0.003
uncultured *Lachnospiraceae* B	0.000±0.000	0.002±0.002	0.000±0.000	0.000±0.000
Peptococcus	0.000±0.000	0.016±0.009	0.015±0.006	0.000±0.000
Hydrogenoanaerobacterium	0.000±0.000	0.004±0.002*	0.000±0.000	0.000±0.000
Subdoligranulum	0.000±0.000	0.000±0.000	0.003±0.002	0.000±0.000
Methylobacterium	0.000±0.000	0.000±0.000	0.000±0.000	0.001±0.001
Variovorax	0.000±0.000	0.002±0.001	0.000±0.000	0.000±0.000
Bilophila	0.000±0.000	0.024±0.024	0.000±0.000	0.000±0.000
Stenotrophomonas	0.000±0.000	0.057±0.048*	0.001±0.001	0.000±0.000
Other	0.045±0.014	0.051±0.005	0.047±0.010	0.018±0.010

*significantly different compared to the control; significance was determined by $p<0.05$.