

Title	Short-term consumption of a high-fat diet increases host susceptibility to <i>Listeria monocytogenes</i> infection
Authors	Las Heras, Vanessa;Clooney, Adam G.;Ryan, Feargal J.;Cabrera-Rubio, Raul;Casey, Pat G.;Hueston, Cara M.;Pinheiro, Jorge;Rudkin, Justine K.;Melgar, Silvia;Cotter, Paul D.;Hill, Colin;Gahan, Cormac G.
Publication date	2019
Original Citation	Las Heras, V., Clooney, A. G., Ryan, F. J., Cabrera-Rubio, R., Casey, P. G., Hueston, C. M., Pinheiro, J., Rudkin, J. K., Melgar, S., Cotter, P. D., Hill, C. and Gahan, C. G. M. (2019) 'Short-term consumption of a high-fat diet increases host susceptibility to <i>Listeria monocytogenes</i> infection', <i>Microbiome</i> , 7(1) (12pp). DOI: 10.1186/s40168-019-0621-x
Type of publication	Article (peer-reviewed)
Link to publisher's version	https://microbiomejournal.biomedcentral.com/articles/10.1186/s40168-019-0621-x - 10.1186/s40168-019-0621-x
Rights	© The Author(s). 2019. © The Author(s). 2019Open AccessThis article is distributed under the terms of the Creative Commons Attribution 4.0International License (http://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The Creative Commons Public Domain Dedication waiver(http://creativecommons.org/publicdomain/zero/1.0/) applies to the data made available in this article, unless otherwise stated. - https://creativecommons.org/licenses/by/4.0/
Download date	2025-07-04 16:01:48
Item downloaded from	https://hdl.handle.net/10468/7680



University College Cork, Ireland
Coláiste na hOllscoile Corcaigh

Supplemental data, Las Heras et al Fig S2.




Figure S2. Increased dietary fat from animal source increases host susceptibility to oral infection with *Listeria monocytogenes* EDGe^m pIKM2. IG inoculation of seven-week-old female C57BL/6 mice (ENVIGO, UK) was repeated in a separate duplicate experiment (n=8) using a 200µl inoculum comprising 2.8×10^9 CFU *L. monocytogenes* EGDm::*pIMK2lux*, a bioluminescent murinized strain. After infection (three days), the faecal pellets were collected daily and plated for CFU to determine shedding of *L. monocytogenes*. **A.** Bacterial burden of spleen, liver, lymph nodes and of C57BL/6 mice fed with diets varying in percentage of fat content from the total caloric intake (n=8, standard deviation from the mean, statistical analysis was conducted using Mann Whitney Nonparametric Test). **B.** *Listeria monocytogenes* bacterial shedding per gram of faecal sample in each day after infection. **C.** Comparison between the two trials (EGDe^m and pIKM2). Mann Whitney Nonparametric Test.