


Class-specific histone deacetylase inhibitors promote 11-beta hydroxysteroid dehydrogenase type 2 expression in JEG-3 cells
Togher, Katie L.;Kenny, Louise C.;O'Keeffe, Gerard W.
2017-02-21
Togher, K. L., Kenny, L. C. and O'Keeffe, G. W. (2017) 'Class-specific histone deacetylase inhibitors promote 11-beta hydroxysteroid dehydrogenase type 2 expression in JEG-3 cells', International Journal of Cell Biology, 6169310 (10 pp). doi:10.1155/2017/6169310
Article (peer-reviewed)
10.1155/2017/6169310
© 2017 Katie L. Togher et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited https://creativecommons.org/licenses/by/4.0/
2025-06-01 01:45:14
https://hdl.handle.net/10468/5342



Supplementary Figures:

Supplementary Figure 1:



Supplementary Figure 1: Expression levels of different classes of HDACs: (A,B) Expression data from the BioGPS database showing the relative expression of (A) Class-I HDACs, HDAC 2, 3 and 8 and (B) Class-I Ia HDACs, HDAC 4, 7 and 9 in the placenta (red) relative to multiple human tissues and fetal brain.