

Title	School-level differences in self-harm and psychosocial problems associated with self-harm in adolescents
Authors	McEvoy, David;Brannigan, Ross;Healy, Colm;Dooley, Niamh;Arensman, E.;Clarke, Mary
Publication date	2026
Original Citation	McEvoy, D, Brannigan, R, Healy, C, Dooley, N, Arensman, E & Clarke, M 2026, 'School-level differences in self-harm and psychosocial problems associated with self-harm in adolescents', Irish Journal of Psychological Medicine, pp. 1-12. https://doi.org/10.1017/ipm.2026.10192
Type of publication	Article (peer-reviewed)
Link to publisher's version	https://www.scopus.com/pages/publications/105037062440 - 10.1017/ipm.2026.10192https://www.scopus.com/pages/publications/105037062440?origin=resultslist
Rights	© 2026, theAuthors. Published by Cambridge University Press on behalf of College of Psychiatrists of Ireland. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (https://creativecommons.org/licenses/by/4.0/), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited. - https://creativecommons.org/licenses/by/4.0/
Download date	2026-10-26 02:16:06
Item downloaded from	https://hdl.handle.net/10468/18820

Original Research

School-level differences in self-harm and psychosocial problems associated with self-harm in adolescents

David McEvoy^{1,2} , Ross Brannigan¹, Colm Healy³ , Niamh Dooley^{2,4} , Ella Arensman^{5,6} and Mary Clarke^{1,2}

¹School of Population Health, Royal College of Surgeons in Ireland, Dublin, Ireland, ²Department of Psychiatry, Royal College of Surgeons in Ireland, Dublin, Ireland, ³Division of Psychiatry, Centre for Clinical Brain Sciences, University of Edinburgh, Edinburgh, UK, ⁴King's College London, UK, ⁵National Suicide Research Foundation, University College Cork, Cork, Ireland and ⁶School of Public Health, University College Cork, Cork, Ireland

Abstract

Background: The school environment plays a key role in adolescents' emotional development and well-being, yet little research has compared self-harm and related psychosocial problems across different secondary school types.

Methods: Using data from the Growing Up in Ireland (GUI) longitudinal cohort, this study examined differences in the prevalence of self-harm and psychosocial risk factors across different school types: single-sex versus coeducational, fee-paying versus non-fee-paying, disadvantaged versus non-disadvantaged, and schools with different religious ethos. Multilevel regression models distinguished school-level from individual-level effects.

Results: Almost all variance in self-harm and most of the variance in psychosocial problems associated with self-harm occurred at the individual level. Higher self-harm prevalence in single-sex girls' schools was accounted for by the greater concentration of girls, who had over twice the odds of self-harm compared with boys (OR 2.1, 95% CI 1.71–2.69). No significant differences in self-harm were found by school socio-economic status or religious ethos. Disadvantaged schools showed higher prevalence in seven of nine psychosocial problems, although only internalising problems and truancy/absenteeism remained significantly associated with disadvantaged schools in the fully adjusted models. Adolescents whose parents reported having a religion were less likely to self-harm (OR 0.62, 95% CI 0.50–0.75).

Discussion: Although schools are important settings for self-harm prevention, findings indicate that interventions should primarily target individuals and high-risk groups. Girls, in particular, may benefit from supports addressing self-harm. Disadvantaged schools, where well-established psychosocial risk factors for self-harm are more common, may benefit from well-being programmes targeting internalising problems and truancy/absenteeism.

Keywords: adolescents; psychosocial problem; schools; self-harm; self-injury

(Received 30 June 2025; revised 12 December 2025; accepted 28 February 2026)

Introduction

Self-harm in adolescents is a major public health concern and a key predictor of suicide (Hawton et al. 2012, O'Connor et al. 2009). Both the economic and social costs of self-harm are considerable, and thus public health approaches to self-harm, and indeed interventions for the psychosocial problems associated with self-harm, must be considered (Moran et al. 2024). The school environment is one of the principal settings that influences the health behaviours, well-being and emotional health of adolescents (Kidger et al. 2015; Pisinger et al. 2019). Moreover, self-harm presentations at emergency departments have been found to be lower during school holidays in adolescents (McEvoy et al. 2024). Furthermore, many of the psychosocial problems associated with self-harm involve, or may occur in, the school setting – such as

bullying, being unpopular with peers, school truancy or dropout, conduct or behavioural issues, and engaging in violence (McEvoy et al. 2023). Given the extensive time spent in school by adolescents, it is both an obvious and convenient setting for implementing universal interventions for self-harm and the psychosocial problems to which self-harm is associated.

Self-harm is defined as intentional self-poisoning or self-injury irrespective of the apparent purpose of the act (National Institute for Health and Care Excellence (NICE) 2022). It typically begins in young people in early adolescence, about the same time adolescents enter secondary or high school (Hawton et al. 2012; Sawyer et al. 2018).

School-based interventions for suicide prevention have been found to be effective in reducing suicidal self-harm (Bailey et al. 2017; Moran et al. 2024; Wasserman et al. 2015). In addition, school-based interventions that address common psychosocial problems associated with self-harm, such as anti-bullying programmes, school-based violence prevention programmes, and emotional learning programmes have been proven to be

Corresponding author: David McEvoy; Email: davidmcevoy@rcsi.com

Cite this article: McEvoy D, Brannigan R, Healy C, Dooley N, Arensman E, and Clarke M. School-level differences in self-harm and psychosocial problems associated with self-harm in adolescents. *Irish Journal of Psychological Medicine* <https://doi.org/10.1017/ipm.2026.10192>

© The Author(s), 2026. Published by Cambridge University Press on behalf of College of Psychiatrists of Ireland. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.

successful in improving the mental health of young people, more generally (Dowling and Barry 2020; Fraguas et al. 2021; Stănicke 2021). Implementing such programmes in schools, like effective anti-bullying measures, could simultaneously improve mental health for young people in school (Fraguas et al. 2021) and possibly reduce self-harm. Hence, school-based interventions targeting psychosocial problems associated with self-harm, as well as interventions for self-harm itself, could help to reduce self-harm in adolescents.

Understanding the extent to which the rates of self-harm and its risk factors (psychosocial problems) vary across different types of schools is important so that, in cases of limited resources, departments or ministries of education could assess which schools have the most pressing need for such public health interventions. Furthermore, identifying which types of schools demonstrated higher levels of self-harm, or psychosocial problems associated with self-harm, could aid in the identification of sub-groups of adolescents at an elevated risk.

School demography, ethos, and policies vary widely and could have an influence on the propensity of students to experience distress and engage in self-harming behaviours. Schools can vary with respect to being coeducational or single-sex, having a religious ethos or being non-denominational, being socio-economically disadvantaged, or being a private fee-paying versus a non-fee-paying school. It is currently unknown to what extent these factors influence the levels of self-harm and the psychosocial problems to which self-harm is associated.

Most of the research concerning self-harm, or the psychosocial problems to which it is associated, has typically been conducted at the individual level rather than the school level (Kim and Chun 2020). Studies that have examined school-level differences for self-harm are rare. One example of such a study by Young et al., found that while holding Catholic religious beliefs were protective for students against suicidal behaviours, paradoxically attending a Catholic school was a risk factor (Young et al. 2011). There were also studies that examined psychosocial problems associated with self-harm across different types of schools. For example, emotional and behavioural health, measured by the Strengths and Difficulties Questionnaire (SDQ), is a key risk factor for self-harm (McEvoy et al. 2023; McEvoy et al. 2024) and was found to vary across different primary schools (Lewer et al. 2024). Furthermore, bullying has been found to vary across different schools in low- and middle-income countries (Kim and Chun 2020).

This study used a nationally representative longitudinal cohort study in Ireland to explore how self-harm, and well-established psychosocial problems associated with self-harm, varied across different types of schools. Psychosocial problems, such as internalising problems or bullying, have been well established as risk factors for self-harm by an umbrella review of systematic reviews (McEvoy et al. 2023). These risk factors are referred to as psychosocial problems in this study since they are also important outcomes in their own right. Types of schools was determined by sex composition, socio-economic status, and religious ethos.

The primary aim of this study was to explore which types of schools had higher proportions of young people with lifetime reported self-harm in adolescents at age 17. The secondary aim of the study was to examine which types of schools had higher proportions of young people exhibiting psychosocial problems (risk factors) associated with self-harm at age 13. The tertiary aim of this study was to examine and separate the school-level and individual-level effects on self-harm and the psychosocial

problems to which it was associated, using multilevel regression models.

Methods

Study population

Growing Up in Ireland (GUI) is a nationally representative longitudinal cohort study in Ireland (Murphy et al. 2019). This study used data from Cohort '98, which recruited $n = 8,568$ nine-year-olds born in 1998 (or the year before or after) and followed them up at ages 13, 17, and 20 years (Murphy et al. 2019). Data collected involved outcomes covering the participants' physical health and development, social and emotional well-being, and educational achievement. This study used data from two waves from the GUI Cohort '98: $n = 7,525$ participants at age 13 and $n = 6,216$ participants followed up at age 17. There were 84 participants that were removed at age 13 for analysis since they were still in primary school, were in a special school or did not provide information on the type of school that they attended. Hence, $n = 7,441$ participants at age 13, clustered within 622 secondary schools, were analysed for this study.

Exposure variables

All of the exposure variables were from the age 13 wave of the data. Typically, adolescents transition from primary school to secondary school in Ireland about the age of 12 and adolescents who self-harm usually first exhibit this behaviour approximately from ages 12 to 14 (Cipriano et al. 2017; Department of Education and Science 2004). At age 13, a number of school-level variables were derived in the GUI dataset using a database of schools from the Department of Education in the Republic of Ireland. The school-level derived variables included whether the school type was coeducational or single-sex; a non-fee-paying school or a fee-paying school; or whether the school was a disadvantaged school. A disadvantaged status of a school was determined if the school was subsidised by the Delivering Equality of Opportunity in Schools (DEIS) programme during the time that data for the age 13 wave of GUI was collected. The DEIS programme was designed to give tailored support to schools that have a high concentration of disadvantage (Department of Education and Skills 2017). Data pertaining to religious ethos of schools were also included. A summary of the exposure and outcome variables is presented in Figure 1.

In order to differentiate between the sex composition at the school-level and individual-level differences with respect to sex, this study used two variables at each level: parent-reported sex of the young person and the sex composition of the school – coeducation, single-sex girls, and single-sex boys, with coeducation school being the reference category.

Two variables were also included to differentiate between school-level disadvantage and individual-level disadvantage. This study also used another derived variable from the GUI dataset – namely, household social class of the young person. This variable was derived using the reported occupations of the primary and secondary care givers, with the following categories in descending order of social status: professional workers, managerial and technical, non-manual, skilled manual, semi-skilled, unskilled, and all others gainfully occupied and unknown occupational status. This six-category ordered variable was collapsed into a three-category ordered variable into 'professional or managerial', 'non-manual or skilled', and 'semi-skilled or unskilled', with the first of




Figure 1. Exposure and outcome variables used in this study.

these being the reference category. The associated school-level variable was the three-outcome school type: disadvantaged school, fee-paying school, and standard school, which was a school that was in neither of the former categories and was the reference group for this variable.

Finally, two variables were also used to differentiate between individual-level religion and the religious ethos of the school. Each school had a religious denomination from the following list: Catholic, inter-denominational, Church of Ireland, multi-denominational, Methodist, and Quaker. This was recoded into three categories: Catholic, multi-/inter-denominational, and Protestant (combining Church of Ireland, Methodist, and Quaker, due to small numbers in each). There was no explicit variable for the young person's religion at age 13 years; rather, the variable that both parents had a religion was used as a proxy variable. This variable was coded as missing if only one parent or neither parent provided a response.

Outcome variables

All of the outcomes for this study were binary. The outcomes were self-harm by age 17 and psychosocial problems (risk factors) associated with self-harm at age 13: internalising problems (depression or anxiety); externalising problems (behavioural or conduct issues); bullying; substance use; school truancy or excessive absence from school; low popularity with peers; being involved in violence; and not getting on with parents.

Lifetime self-harm

Lifetime self-harm data were recorded at age 17. The participants were asked if they had ever hurt themselves on purpose in any way.

Psychosocial outcomes at age 13

Psychosocial risk factors were measured at age 13 to ensure clear temporal ordering between school-type exposure, early psychosocial functioning, and later self-harm. Although we observed cross-sectional associations between school type and psychosocial

problems at age 13, it is unlikely that psychosocial difficulties determine the type of school attended, which is usually established earlier and remains stable. Thus, the more plausible direction is that school environment influences psychosocial problems. Using age 13 measures also avoids reverse causation, providing a valid baseline of psychosocial problems before self-harm by age 17.

The selection of these psychosocial problems was informed by two studies conducted prior to this study (McEvoy et al. 2023, McEvoy et al. 2024). Using an umbrella review examining risk factors for self-harm in young people (McEvoy et al. 2023), a list of psychosocial factors was compiled from the risk factors that had a pooled odds ratio from meta-analyses of at least two and/or were identified as a risk factor in at least four systematic reviews. This list of well-established risk factors for self-harm was then reduced to psychosocial factors for which data was available from the GUI dataset. Then using a latent class analysis of psychosocial factors for self-harm at age 13 using the GUI dataset (McEvoy et al. 2024), it was determined which risk factors from this list should be included in this study (i.e. factors that did not have a low prevalence) – namely, the nine psychosocial factors listed in previously mentioned and listed in Figure 1. Note that being female (a risk factor for self-harm) was included instead as an exposure variable and not an outcome variable.

Internalising problems was measured using the SDQ, which is the emotional problems sub-score added to the peer problems sub-score (Goodman and Goodman 2009). An internalising problems score of 8 or more is considered to be clinically significant and was used as a 'yes' for this variable (Goodman and Goodman 2009).

Externalising problems was also measured using the SDQ, which is the conduct problems sub-score added to the hyperactivity problems sub-score (Goodman and Goodman 2009). An externalising problems score of 11 or more is considered significant and was used as a 'yes' for this variable (Goodman and Goodman 2009). A second measure of school-specific conduct issues (i.e., *school misbehaviour*) was also used as an outcome. For 'school misbehaviour reported by the young person at age 13', the young person had to report either 'often' or 'all the time' to not

following the school rules, unruly behaviour in class, getting detention, being suspended, or receiving other punishments for misbehaviour.

Bullying was defined as a report by the primary care giver that the young person had been bullied in the last three months.

For *school truancy or excessive absence* from school, this variable was recorded as 'yes' if either the young person was absent from school for more than 20 days in the past school year or answered that they occasionally, quite often or always skipped classes. Irish schools are mandated to inform social services if a child has missed 20 days or more in the school year (TUSLA: Education Welfare Service 2025).

For *substance use*, the young person had to answer 'yes' to at least one of: smoked cigarettes at least once a week; consumed alcohol at least once a month; or ever tried using cannabis, sniffing glues or illicit drugs.

Low popularity with peers was determined by a score of ≤ 39 on the Piers-Harris Children's Self-Concept popularity sub-scale (Community-University Partnership for the Study of Children Youth and Families 2011).

Violence or peer conflict was a report by the young person that they had been involved in a serious physical fight where someone got badly hurt or needed to see a doctor. 'Does not get on with a parent' was a report by the young person that they did not get on with their mother, father or both.

Statistical analysis

Cross table proportions were calculated for all of the school factors with lifetime self-harm at age 17 and the psychosocial problems to which self-harm is associated. Sample weights were applied to these proportions to make them representative of the population of the Republic of Ireland. Chi-square tests then were used to determine if there were significant relationships between each of the 3 school variables and the 10 outcome variables.

The Benjamini-Hochberg procedure was used to control the false discovery rate of the $m = 30$ hypothesis tests and is an appropriate method for this type of exploratory analysis (James et al. 2013). This procedure ranks the p -values and compares them to adjusted thresholds, allowing for the identification of statistically significant results while limiting false positives (James et al. 2013).

Next, categories of schools were created using both school sex (girls, boys, or coeducation) and school type (standard, fee-paying, and DEIS) to examine the interactions between these two school variables. Using sample weights, the proportions of lifetime self-harm reported at age 17 was calculated for participants within each of these categories of schools. These proportions were then ranked to determine which categories of schools had the highest proportions of self-harm. For statistical disclosure control reasons, school religion was not included in these interactions due to small numbers in coeducation denominational schools. Proportions for single-sex DEIS schools were also excluded from these analyses due to small raw numbers in these categories. All data wrangling and statistical analyses described thus far were conducted in R.

To disentangle school-level from individual-level effects, multilevel regression models were used with students nested within schools. For each of the ten dependent variables (self-harm and the nine psychosocial problems), a multilevel logistic regression was used, which included both school-level variables and their matching individual-level control variables (see Figure 1). Multilevel logistic regression was conducted using the 'meqrlogit' function using STATA (version 18). We did not apply sample

weights to these models since including them inflated the school-level variance to implausible levels. Within each multilevel regression model, the variance partition coefficient (VPC) or intra-class coefficient (ICC) was used to determine the variation between schools for each outcome variable. Again, we applied the Benjamini-Hochberg procedure to all $m = 100$ p -values to control the false discovery rate. A sensitivity analysis was also conducted by alternately including and excluding school- and individual-level variables for sex, socio-economic status, and religion.

Results

School characteristics and participant distribution

Variation for self-harm and the psychosocial problems across the different school characteristics (unadjusted for individual-level factors) are presented in Table 1. The majority of participants (59%) attended 386 coeducational schools; 21.7% of participants attended 135 single-sex girls' schools; and 19.2% of the participants attended 101 single-sex boys' schools. The majority (72.7%) of participants also attended schools that were neither fee-paying nor DEIS schools (416 'standard' schools); 17.9% of individuals attended 151 DEIS schools; and 9.4% of individuals attended 55 fee-paying schools. With respect to religious ethos of schools, 53.3% of participants attended schools with a Catholic ethos; 41.8% attended multi- or inter-denominational schools; and a small proportion of participants (4.9%) attended schools with a Protestant ethos.

The majority (78.0%) of students attending a Protestant school also attended a fee-paying school, whereas a minority (10.3%) of those attending a Catholic school also attended a fee-paying school. Virtually all multid denominational schools were non-fee-paying. See the supplementary material for further details.

Self-harm variation across different types of schools

A lifetime report of self-harm was reported by 17.7% of all young people. The sex of the school was the only category that had a statistically significant difference in the proportions of young people that reported having ever self-harmed by age 17 (not adjusting for individual-level factors). The highest proportion of young people who reported lifetime self-harm was in girls' schools (24.5%), followed by coeducation schools (17.3%), and then boys' schools (12.4%). There were no statistically significant differences in the proportions of self-harm across type of schools (standard, DEIS, or fee-pay) or across schools of different religious ethos.

We can also see the trend of higher proportions of lifetime self-harm in those who attended girls' schools in Figure 2. Within girls' schools, those who attended fee-paying schools reported lower proportions (albeit non-statistically significant) of lifetime self-harm at age 17 compared to those in standard schools. Conversely, in boys' schools, those in fee-paying schools had higher proportions of lifetime self-harm by age 17 compared to those in standard schools (again non-statistically significant).

School variation of psychosocial problems associated with self-harm at age 13

While single-sex girls' schools had statistically significant higher proportions of lifetime self-harm at age 17, paradoxically, single-sex girls' schools had statistically significant lower proportions in five out of the nine risk factors for self-harm: namely, externalising problems, school misbehaviour, truancy or excessive school absenteeism, substance use, and being involved in violence.

Table 1. School-level characteristics (unadjusted for individual-level factors) and the proportion of self-harm and its associated psychosocial problems within each stratum

	Sex of school				Type of school				Religious ethos of school			
	Coeducation	Boys	Girls	Adjusted <i>p</i> -value*	Standard	Disadvantaged (DEIS)	Fee-pay	Adjusted <i>p</i> -value*	Multi-/inter-denominational	Catholic	Protestant	Adjusted <i>p</i> -value*
Number of schools <i>m</i> (%)	386 (62.1)	101 (16.2)	135 (21.7)	–	416 (66.9)	151 (24.3)	55 (8.8)	–	268 (43.1)	328 (52.7)	26 (4.2)	
Number of participants <i>n</i> (%)	4,170 (59.0)	1,472 (19.2)	1,799 (21.7)	–	5,711 (72.7)	1,009 (17.9)	721 (9.4)	–	2,863 (41.8)	4,251 (53.3)	327 (4.9)	
Female attendance <i>n</i> (%)	2,013 (46.7)	0 (0.0)	1,799 (100.0)	–	3,005 (50.1)	504 (49.9)	303 (41.9)	–	1,402 (47.2)	2,251 (50.3)	159 (54.9)	
Lifetime self-harm at age 17 <i>n</i> (%) = 1,025 (17.7)												
Participants <i>n</i> (%) reporting lifetime self-harm at age 17 of which are female <i>n</i> (%)	549 (17.3)	160 (12.4)	316 (24.5)	<0.001	798 (18.1)	122 (16.5)	105 (18.4)	0.849	374 (16.3)	593 (18.5)	58 (24.1)	0.196
	350 (63.8)	0 (0.0)	316 (100.0)	–	534 (66.9)	74 (60.7)	58 (55.2)	–	234 (62.6)	393 (66.3)	Cannot disclose ‡	
Psychosocial problems (associated with self-harm) at age 13												
<i>Internalising problems</i> <i>n</i> (%) = 538 (8.3)	313 (9.2)	92 (7.1)	133 (7.3)	0.294	369 (7.3)	125 (14.1)	44 (5.3)	<0.001	<240 (<10.0)	285 (7.5)	<30 (<9.2)	0.013
<i>Externalising problems</i> <i>n</i> (%) = 294 (5.2)	184 (5.6)	65 (5.1)	45 (4.1)	0.003	196 (4.5)	<80 (<10.0)	<30 (<4.2)	<0.001	<135 (<7.0)	148 (4.8)	<30 (<9.2)	0.074
<i>Experienced bullying</i> <i>n</i> (%) = 710 (10.3)	420 (10.7)	128 (9.6)	162 (10.0)	0.246	521 (9.6)	132 (13.1)	57 (10.6)	<0.001	<300 (<11.0)	384 (9.8)	<30 (<9.2)	0.200
<i>School misbehaviour</i> <i>n</i> (%) = 1,063 (16.3)	666 (18.1)	228 (16.6)	171 (11.0)	<0.001	756 (14.0)	198 (23.2)	109 (20.9)	<0.001	457 (18.7)	548 (14.2)	58 (18.6)	0.001
<i>Truancy or excessive school absenteeism</i> <i>n</i> (%) = 472 (8.0)	302 (8.8)	91 (7.5)	79 (6.4)	<0.001	323 (7.0)	110 (12.0)	39 (8.6)	<0.001	<210 (<9.0)	240 (7.3)	<30 (<9.2)	0.016
<i>Substance use</i> <i>n</i> (%) = 359 (6.4)	244 (7.0)	72 (5.8)	74 (5.4)	0.032	290 (6.0)	<80 (<9.0)	<30 (<4.2)	0.004	<180 (<8.0)	203 (6.2)	<30 (<9.2)	0.129
<i>Low PH popularity</i> † <i>n</i> (%) = 2,335 (33.0)	1,318 (33.1)	444 (32.0)	573 (33.8)	0.451	1,797 (32.7)	328 (34.8)	210 (32.8)	0.249	944 (33.9)	1,290 (32.1)	101 (35.9)	0.074
<i>Involved in violence</i> <i>n</i> (%) = 236 (4.2)	156 (5.2)	<60 (<4.5)	<30 (<1.7)	<0.001	<180 (<5.0)	54 (6.9)	<30 (<4.2)	<0.001	<120 (<6.0)	118 (3.5)	<30 (<9.2)	0.032
<i>Conflict with parents</i> <i>n</i> (%) = 224 (3.8)	133 (4.0)	34 (2.4)	57 (4.4)	0.200	165 (3.4)	<40 (<5.5)	<30 (<4.2)	0.136	<100 (<4.5)	120 (3.4)	<30 (<9.2)	0.395

Notes: Proportions have been adjusted using sample weights; * chi square test and adjusted *p*-values were computed using the Benjamini–Hochberg False Discovery rate procedure; † Low Piers Harris sub-score for popularity with peers at age 13; ‡ Figure not disclosed since it would be possible to identify other figures less than *n* = 30.




Figure 2. Interactions between sex of school and school type (adjusted using sample weights).

Disadvantaged schools did not have a statistically significant different proportion for lifetime self-harm at age 17 but the participants who attended DEIS schools had statistically higher proportions in seven out of the nine risk factors for self-harm compared to those who attended standard or fee-paying schools (namely, internalising and externalising problems, bullying, school misbehaviour, truancy or excess absenteeism, substance use, and being involved in violence).

Participants attending Catholic schools exhibited a statistically significant lower percentages of school misbehaviour and being involved with violence at age 13.

School-level versus individual-level effects for self-harm

School-level and individual-level effects, separated using multilevel models, are presented in Tables 2 and 3, respectively. In Table 2 (school-level effects), boys in single-sex school were compared to boys in coeducational schools (the reference category). Similarly, girls in single-sex school were compared to girls in coeducational schools. In Table 3 (individual-level effects), boys were compared to girls, regardless of what type of school they attended.

School-level effects (self-harm)

School-level characteristics (adjusted for individual-level effects) for self-harm and the associated psychosocial problems are presented in Table 2. None of the school-level independent variables were significantly associated with self-harm. Specifically, there were no statistically significant associations between the sex, type, or religious ethos of the school and self-harm in young people, when controlling for individual-level associations. Moreover, the majority of the variation for self-harm was between individuals rather than between schools. The ICC for the self-harm model was 0.1% meaning that only 0.1% of the total variance in lifetime reported self-harm in young people was due to differences between schools – that is, virtually all of the variance was at the individual level.

Individual-level effects (self-harm)

Individual-level characteristics (adjusted for school-level effects) for self-harm and the associated psychosocial problems are presented in Table 3. Being female was found to be significantly associated with higher lifetime reported self-harm at age 17: the odds ratio (OR) for females self-harming was 2.14 (95% CI 1.71–2.69) compared to males. Given the significant individual-level effect of being female, the non-significant effect of school sex (coeducational *v.* single-sex), and the fact that virtually all variance

in self-harm lies at the individual level, the higher proportions of self-harm in single-sex girls' schools compared to coeducational schools (and in coeducational relative to single-sex boys' schools) are explained by the larger proportion of girls attending those schools.

Both parents having a religion was associated with lower lifetime reported self-harm in the young person (OR = 0.62, 95% CI 0.50–0.75) compared to young people where one or both of the parents did not have a religion.

A statistically significant association between being in a lower social class and lifetime self-harm was not found.

School-level versus individual-level effects for psychosocial problems associated with self-harm

School-level effects (psychosocial problems)

Like in the self-harm outcome model, the ICC was also less than 1% in the models with internalising problems, bullying, substance use, low popularity, and conflict with parents as dependent variables. The highest ICC values were in the models with externalising problems (11.7%) and truancy/excessive absence (7.2%) as dependent variables.

Being in a disadvantaged (DEIS) school was only associated with two psychosocial problems associated with self-harm: internalising problems (OR = 1.57, 95% CI 1.15–2.14) and truancy or excessive absence from school (OR = 1.66, 95% CI 1.17–2.35). A sensitivity analysis found that if 'both parents having a religion' was excluded, then being in a disadvantaged school was also associated with externalising problems, bullying, and school misbehaviour, suggesting that these relationships were confounded by family background characteristics correlated with parental religiosity.

Individual-level effects (psychosocial problems)

Even though being female had over twice the odds of lifetime self-harm at age 17 compared to males, being female was associated with lower odds of externalising problems, school misbehaviour, and being involved in violence (0.46 (95% CI 0.30–0.71), 0.52 (95% CI 0.42–0.64), and 0.16 (95% CI 0.09–0.28), respectively).

As well as being protective against a lifetime report of self-harm by age 17, having both parents report having a religion was found to be protective in eight out of the nine risk factors for self-harm – only not statistically significant for being involved in violence. The ORs ranged from 0.57 (95% CI 0.43–0.75) for engaging in truancy or having excessive absence from school at age 13 to 0.71 (95% CI 0.56–0.91) for being a bullying victim at age 13.

Table 2. Odds ratios and school-level effects (adjusted for individual-level factors)

	Sex of school (Ref: Coeducational)				Type of school (Ref: Standard)				Religious ethos of school (Ref: Multi-/inter-denominational)				Combined	
	Boys		Girls		Disadvantaged (DEIS)		Fee-pay		Catholic		Protestant		σ_u^2 Between-school variance (SE)	VPC/ICC
	OR (95% CI)	Adjusted <i>p</i> -value *	OR (95% CI)	Adjusted <i>p</i> -value *	OR (95% CI)	Adjusted <i>p</i> -value *	OR (95% CI)	Adjusted <i>p</i> -value *	OR (95% CI)	Adjusted <i>p</i> -value *	OR (95% CI)	Adjusted <i>p</i> -value *		
Lifetime Self-harm at age 17	1.36 (0.99, 1.86)	0.209	1.19 (0.91, 1.55)	0.539	0.93 (0.70, 1.23)	0.791	1.00 (0.73, 1.35)	0.975	0.88 (0.68, 1.13)	0.607	1.13 (0.73, 1.75)	0.791	0.004 (0.048)	0.001
Psychosocial problems (associated with self-harm) at age 13														
Internalising problems	1.11 (0.72, 1.70)	0.791	1.06 (0.72, 1.58)	0.864	1.57 (1.15, 2.14)	0.031	1.23 (0.79, 1.91)	0.626	0.78 (0.55, 1.10)	0.446	0.44 (0.20, 0.95)	0.156	<0.001 (<0.001)	<0.001
Externalising problems	1.02 (0.57, 1.81)	0.975	0.99 (0.51, 1.93)	0.975	1.73 (1.10, 2.73)	0.093	1.20 (0.62, 2.32)	0.791	0.91 (0.54, 1.53)	0.840	1.07 (0.42, 2.70)	0.960	0.437 (0.215)	0.117
Experienced bullying	1.05 (0.75, 1.48)	0.866	0.83 (0.60, 1.15)	0.546	1.36 (1.02, 1.82)	0.156	0.88 (0.60, 1.29)	0.755	1.15 (0.86, 1.53)	0.624	1.14 (0.66, 1.99)	0.791	<0.001 (<0.001)	<0.001
School misbehaviour	0.94 (0.71, 1.25)	0.797	0.86 (0.62, 1.18)	0.615	1.16 (0.90, 1.49)	0.446	1.07 (0.78, 1.48)	0.795	0.94 (0.73, 1.21)	0.791	1.25 (0.80, 1.95)	0.615	0.068 (0.056)	0.020
Truancy or excessive school absenteeism	1.12 (0.72, 1.74)	0.791	0.71 (0.45, 1.13)	0.446	1.66 (1.17, 2.35)	0.031	0.76 (0.44, 1.30)	0.604	1.05 (0.72, 1.54)	0.874	2.00 (1.02, 3.94)	0.177	0.254 (0.127)	0.072
Substance use	1.09 (0.70, 1.71)	0.807	0.89 (0.58, 1.42)	0.791	1.29 (0.90, 1.86)	0.446	0.83 (0.49, 1.41)	0.754	0.84 (0.57, 1.23)	0.647	1.35 (0.68, 2.68)	0.651	0.013 (0.119)	0.004
Low PH popularity †	1.14 (0.92, 1.42)	0.546	1.13 (0.92, 1.40)	0.546	0.95 (0.78, 1.14)	0.791	0.91 (0.72, 1.15)	0.705	0.84 (0.70, 1.00)	0.200	1.01 (0.72, 1.42)	0.975	0.017 (0.026)	0.005
Involved in violence	0.71 (0.42, 1.19)	0.503	0.97 (0.42, 2.23)	0.975	1.65 (1.05, 2.59)	0.150	0.39 (0.16, 0.94)	0.156	1.31 (0.82, 2.11)	0.546	1.37 (0.47, 4.01)	0.791	0.180 (0.213)	0.052
Conflict with parents	0.85 (0.44, 1.67)	0.791	0.97 (0.54, 1.74)	0.972	0.99 (0.57, 1.71)	0.975	1.20 (0.62, 2.34)	0.791	0.89 (0.52, 1.50)	0.792	0.89 (0.34, 2.32)	0.877	<0.001 (<0.001)	<0.001

Notes: * *p*-values were computed using the Benjamini–Hochberg False Discovery rate procedure; † Low Piers Harris sub-score for popularity with peers at age 13.

Table 3. Odds ratios and individual-level effects (adjusted for school-level factors)

	Sex (Ref: Male)		Socio-economic status (Ref: professional or managerial)				Parents religion (Ref: No)	
	Female		Non-manual or skilled		Semi-skilled or unskilled		Yes	
	OR (95% CI)	Adjusted <i>p</i> -value*	OR (95% CI)	Adjusted <i>p</i> -value*	OR (95% CI)	Adjusted <i>p</i> -value*	OR (95% CI)	Adjusted <i>p</i> -value*
Lifetime Self-harm at age 17	2.14 (1.71, 2.69)	<0.001	0.95 (0.79, 1.14)	0.791	0.81 (0.57, 1.15)	0.546	0.62 (0.50, 0.75)	<0.001
Psychosocial problems (associated with self-harm) at age 13								
Internalising problems	1.22 (0.92, 1.63)	0.446	1.07 (0.83, 1.37)	0.791	1.49 (1.03, 2.17)	0.156	0.64 (0.50, 0.84)	0.013
Externalising problems	0.46 (0.30, 0.71)	0.004	1.47 (1.05, 2.07)	0.141	1.88 (1.12, 3.13)	0.092	0.59 (0.41, 0.85)	0.031
Experienced bullying	1.20 (0.93, 1.54)	0.446	1.05 (0.85, 1.29)	0.797	1.05 (0.73, 1.51)	0.874	0.71 (0.56, 0.91)	0.037
School misbehaviour	0.52 (0.42, 0.64)	<0.001	1.13 (0.95, 1.35)	0.446	1.11 (0.82, 1.51)	0.755	0.65 (0.53, 0.78)	<0.001
Truancy or excessive school absenteeism	0.85 (0.63, 1.14)	0.548	1.24 (0.95, 1.61)	0.375	2.19 (1.51, 3.17)	0.001	0.57 (0.43, 0.75)	0.001
Substance use	0.84 (0.61, 1.15)	0.548	1.29 (0.98, 1.70)	0.225	1.29 (0.82, 2.05)	0.548	0.63 (0.47, 0.85)	0.021
Low PH popularity †	1.06 (0.91, 1.23)	0.754	0.94 (0.83, 1.08)	0.651	1.17 (0.94, 1.46)	0.446	0.68 (0.58, 0.79)	<0.001
Involved in violence	0.16 (0.09, 0.28)	<0.001	1.41 (1.00, 2.01)	0.200	1.57 (0.90, 2.74)	0.375	0.79 (0.52, 1.19)	0.546
Conflict with parents	1.30 (0.83, 2.04)	0.546	1.17 (0.79, 1.71)	0.705	1.31 (0.69, 2.47)	0.668	0.58 (0.39, 0.86)	0.045

Notes: **p*-values were computed using the Benjamini–Hochberg False Discovery rate procedure; † Low Piers Harris sub-score for popularity with peers at age 13.

The odds for engaging in truancy or having excessive absence from school at age 13 was 2.19 (95% CI 1.51–3.17) in those from a semi-skilled or unskilled social class compared to those from a professional or managerial social class.

Discussion

While girls' schools showed higher crude proportions of self-harm, this was fully explained by the larger number of girls in those schools. Once individual sex was accounted for, school sex type was no longer associated with self-harm, indicating that the differences seen reflected the sex makeup of the student body rather than school-level factors. Girls report self-harm at roughly twice the rate of boys – as seen in this study (OR = 2.14, 95% CI 1.71–2.69) and in other Irish studies (Doyle et al. 2015; Griffin et al. 2018; McMahon et al. 2014) – so schools with more girls will naturally have higher crude proportions. Virtually all variance in self-harm was at the individual level.

Higher levels of self-harm in adolescent girls compared to boys are not unique to the Irish context; internationally, the girl-to-boy ratio can be as high as five or six to one in early adolescence, narrowing in later years when self-harm increases among boys (Hawton et al. 2012; Trafford et al. 2023). Suggested explanations for this imbalance include girls experiencing higher levels of internalising problems and influences associated with the timing of menarche (Keyes and Platt 2023; Roberts et al. 2020). There is also some evidence to suggest that girls are more susceptible to the social contagion effect of self-harm than boys – that is, being exposed to peers engaging in self-harm may predispose them to engage in this behaviour themselves (Prinstein et al. 2010).

Girls had significantly lower odds of externalising problems, school misbehaviour, and violence. Yet, paradoxically, girls' schools had higher crude proportions of self-harm despite showing significantly lower levels of five of the nine key risk factors – externalising problems, school misbehaviour, truancy or excessive absence, substance use, and involvement in violence. Paradoxes of

this kind are not uncommon in the self-harm and suicide literature: for example, girls consistently report higher levels of self-harm than boys (Doyle et al. 2015; Griffin et al. 2018; Hawton et al. 2012; McMahon et al. 2014; Trafford et al. 2023), whereas boys tend to have higher suicide rates (McMahon et al. 2014) even though self-harm is one of the strongest predictors of suicide (Hawton et al. 2012; O'Connor et al. 2009).

Self-harm is often described as a strategy for affect regulation, helping to reduce overwhelming emotions or regain a sense of control, and is commonly used by adolescent girls (Stănică 2021). The higher prevalence of self-harm among girls, contrasted with higher physical aggression among boys, may reflect social and cultural norms around emotional expression (Paulo et al. 2020). Society may be more accepting of aggressive behaviour in boys and anxious or depressive behaviours in girls, shaping how distress is expressed even when underlying emotions are similar (Paulo et al. 2020). Although school sex type was not significantly associated with self-harm after adjusting for individual sex, the higher crude proportions in girls' schools indicate a clustering of higher-risk individuals (i.e. girls). These schools may therefore be appropriate settings for targeted preventive interventions for girls.

Self-harm proportions in disadvantaged (DEIS) schools were not significantly different from those in fee-paying or standard schools. In another paradox, however, DEIS schools showed significantly higher proportions for seven of the nine risk factors for self-harm, including both internalising and externalising problems, which were more than twice as common as in non-DEIS schools. Although earlier research identified low socio-economic status as a risk factor for self-harm (Aggarwal et al. 2017; Bozzini et al. 2021; Carballo et al. 2020), recent evidence suggests this pattern may be changing, with rising self-harm among girls in less deprived communities (Trafford et al. 2023).

It is unclear why higher proportions of young people with risk factors for self-harm in disadvantaged schools did not translate into higher self-harm rates. One conjecture (Kim and Chun 2020), is that the psychological impact of stressors like bullying may be

reduced in these settings because such behaviours are more common. Another possibility is that other behaviours – such as externalising problems, substance use, violence, truancy or absenteeism – may substitute for self-harm as responses to distress. Because health risk behaviours tend to co-occur and predict one another (Bozzini et al. 2021), elevated levels of these behaviours may diffuse the risk of self-harm. Additional potential explanations include stronger pastoral supports, greater stigma surrounding self-harm, or increased resilience arising from repeated exposure to adversity in disadvantaged schools.

While most self-harm risk factors were more prevalent in disadvantaged schools, only internalising problems and truancy/absenteeism remained significantly associated with school disadvantage in the fully adjusted multilevel model. After adding the variable ‘both parents having a religion’, three previously significant associations – externalising problems, bullying, and school misbehaviour – were substantially attenuated, suggesting that these relationships were confounded by family background characteristics correlated with parental religiosity. Prior research indicates that parental religiosity often serves as a proxy for aspects of the family environment known to influence behavioural risk, including greater parental involvement and monitoring (Bornstein et al. 2017; Kim-Spoon et al. 2014), and higher levels of family cohesion and stability (Spilman et al. 2013). These family-level processes likely account for the higher prevalence of these behaviours in disadvantaged schools observed in the unadjusted analyses.

Overall, although attending a DEIS school was not directly associated with self-harm, the concentration of risk factors suggests these schools may contain clusters of vulnerable students who could benefit from targeted support. Given that internalising problems and truancy/absenteeism disproportionately affect disadvantaged schools, evidence-based well-being programmes should be introduced in accordance with Ireland’s mental health policy (Department of Health 2020–2030), and the School Completion Programme (providing targeted supports for young people at risk of early school leaving) should be continued (TUSLA: Child and Family Service 2025).

Unlike the Young et al., study, which found that attending a Catholic school was a risk factor (Young et al. 2011), the current study found no statistically significant differences in self-harm prevalence across schools with different religious ethos. However, consistent with Young et al., who reported that Catholic religious beliefs were protective against suicidal behaviours, the current study found that both parents having a religious affiliation – used as a proxy for the young person’s own religion – was associated with a 38% lower odds of lifetime self-harm (OR = 0.62, 95% CI 0.50–0.75), compared with young people whose parents did not both identify with a religion.

Parents reporting a religious affiliation may signify aspects of a more positive or cohesive family environment (Bornstein et al. 2017; Kim-Spoon et al. 2014; Spilman et al. 2013), and in an increasingly secular society, it may also reflect cultural belonging or heritage rather than adherence to doctrine (Central Statistics Office 2023; Inglis 2007). Thwarted belongingness is a well-established predictor of suicidal behaviour (O’Connor and Kirtley 2018; Young et al. 2011), suggesting that the protective effect observed from ‘both parents having a religion’ may be explained by greater social connectedness or community identification. Additionally, having a sense of meaning in life is known to protect against mental health problems and suicidal behaviours (Disabato et al. 2017; Lew et al. 2020). Altogether, although parental religious affiliation

appeared strongly protective against both risk factors for self-harm and self-harm itself, there was no evidence that the religious ethos of the school influenced these outcomes.

Overall, almost all of the variance for self-harm, and most of the variance for the risk factors for self-harm, was at the individual level. While schools remain important sites for interventions, the data from this study suggests that interventions should be primarily targeted at individuals in high-risk groups.

Public health implications and further research

Girls may need to be prioritised in both single-sex and coeducation schools for specific interventions for the prevention of self-harm. On the other hand, while there have been well-evaluated school-based programmes for equipping young people to seek help when experiencing mental health difficulties, well-evaluated programmes specific to self-harm are rare (Moran et al. 2024). Whilst the positives of implementing youth self-harm or suicide prevention strategies in schools generally outweigh the negatives, such programmes have had some unanticipated adverse consequences like increasing maladaptive attitudes related to suicide or help-seeking; increasing the endorsement that suicide was a reasonable solution; or increasing maladaptive coping styles like self-harm or suicidal behaviour – sometimes even leading to hospitalisations (Kuiper et al. 2019). More randomised control trials of school-based interventions to reduce self-harm are therefore needed.

With this in mind, this study could inform the implementation of other public health measures, not specific to self-harm, but addressing the psychosocial problems associated with self-harm. This could result in the reduction of self-harm in adolescents and thereby possibly reduce later deaths by suicide, for which self-harm is a key risk predictor (O’Connor et al. 2009). While this study did not find a significant OR for being in a disadvantaged (DEIS) school with self-harm, there were higher proportions in seven out of the nine risk factors examined for self-harm in disadvantaged schools. Therefore, these schools should be prioritised for interventions addressing the psychosocial problems highlighted in this study. Implementing effective anti-bullying measures or interventions aimed at reducing substance use in disadvantaged schools could simultaneously improve the mental health of its students (Fraguas et al. 2021; Van Ryzin et al. 2023). Well-designed evidence-based social and emotional learning programmes, which have been shown to have effective positive outcomes for students (Dowling and Barry 2020), could be implemented in disadvantaged schools to address the higher levels of internalising and externalising problems, or school misbehaviour in their pupils.

Proportions of poor school behaviour, truancy, and violence were highest in disadvantaged schools. School and social policies in response to negative classroom behaviours, such as reduced timetables, suspensions, expulsions, can be ineffective and even counterproductive (Downes et al. 2020; Robison et al. 2017). Indeed, exclusion from school is higher among those from a lower socio-economic status and is associated with higher psychopathology (Ford et al. 2018) – a key risk factor for self-harm (McEvoy et al. 2023). Multidisciplinary teams of counsellors/therapists, speech and language therapists, occupational therapists in schools (particularly disadvantaged schools), and other professionals should have a role in schools to support students with poor classroom behaviours, externalising problems, poor school attendance, mental health problems or students who are bullying victims or perpetrators (Downes et al. 2020; Sourander et al. 2007).

Ultimately, designing and implementing interventions around the risk factors (psychosocial problems) for self-harm may result in the reduction of self-harm itself among adolescents, especially among those in disadvantaged schools.

Much of the literature on risk and protective factors for self-harm has been focused on risk factors (McEvoy et al. 2023). Future studies could explore protective factors at ages 9 and 13 of the GUI '98 Cohort for self-harm at ages 17 and 20.

Limitations

Sample weights in the multilevel regression models were removed. Ignoring sample weights in multilevel models may introduce some bias into the models (Cai 2013), and it also results in lower external validity of the results of the model. In addition, there was also a small number of participants from disadvantaged and protestant schools, and this may have contributed to a lack of statistical power.

It is also possible that the number of participants in the study who self-harmed was underreported for two reasons. Firstly, any participants that reported "prefer not to say" were counted as missing, though it is possible they did self-harm but did not wish to disclose it due to the stigma surrounding self-harm (Moran et al. 2024). Secondly, it is also possible at age 17, some participants may not wish to disclose if they engaged in self-harm in their lifetime, given the stigma of self-harm. Boys, in particular, are less likely to disclose if they had engaged in self-harm (Victor et al. 2018).

Finally, this study controlled for the false discovery rate rather than the family-wise error rate. While this limits the number of type 1 errors, it does not fully eliminate the possibility of these occurring (James et al. 2013). On balance, limiting the number of type 1 errors rather than eliminating them totally was more appropriate since the analyses were exploratory (James et al. 2013).

Conclusion

Virtually all of the variance in self-harm and most of the variance in associated psychosocial problems occur at the individual level. Therefore, interventions should primarily target high-risk individuals rather than school types. Nevertheless, the school context remains important when designing public health strategies to address mental health among young people. Consistent with existing evidence, this study found that the prevalence of self-harm among adolescent girls is more than twice that of boys. Accordingly, girls' schools and coeducational schools should be prioritised for self-harm-specific interventions, although further research is needed to assess the feasibility and effectiveness of such approaches. While the higher prevalence of psychosocial problems associated with self-harm in disadvantaged schools can largely be explained by adolescents' family circumstances, the greater concentration of affected individuals in these schools suggests they should be prioritised for the implementation of well-being programmes and the continued delivery of interventions aimed at improving school attendance.

Supplementary material. The supplementary material for this article can be found at <https://doi.org/10.1017/ipm.2026.10192>.

Data availability statement. The GUI datasets are available upon application to the ISSDA in Ireland.

Acknowledgements. We wish to thank the Central Statistics Office (CSO) in Ireland for providing access to the data. Results are based on analysis of strictly controlled Research Microdata Files provided by the CSO. The CSO does not

take any responsibility for the views expressed or the outputs generated from this research. We also wish to thank all of the participants from the Growing Up in Ireland (GUI) study. We wish to thank Professor Emer Smyth from the Economic and Social Research Institute (ESRI) and Dr Eve Griffin from the National Suicide Research Foundation for their advice and expertise during this project. We also wish to thank Professor George Leckie from the Centre for Multilevel Modelling in the University of Bristol. We also wish to thank Dr Fiona Boland from the Data Science Centre, RCSI.

Funding statement. This was funded by the Health Research Board (HRB) Ireland as part of the SPHeRE Programme (Structured Population and Health-services Research Education): <https://www.sphereprogramme.ie>.

Competing interests. None to declare.

Ethical standards. Ethical approval for GUI was granted by the research ethics committee of the Health Research Board in Ireland. The secondary analysis of this dataset was approved by the Research Ethics Committee for the Royal College of Surgeons in Ireland (Record ID: 212,613,109). The authors assert that all procedures contributing to this work comply with the ethical standards of the relevant national and institutional committee on human experimentation with the Helsinki Declaration of 1975, as revised in 2008.

Consent for publication. Not applicable.

Patient and public involvement. School teachers were consulted prior to the study being conducted on the direction of the study.

References

- Aggarwal S, Patton G, Reavley N, Sreenivasan SA and Berk M (2017) Youth self-harm in low- and middle-income countries: Systematic review of the risk and protective factors. *International Journal of Social Psychiatry* 63, 359–375. <https://doi.org/10.1177/0020764017700175>.
- Bailey E, Spittal MJ, Pirkis J, Gould M and Robinson J (2017) Universal suicide prevention in young people. *Crisis—the Journal of Crisis Intervention and Suicide Prevention* 38, 300–308. <https://doi.org/10.1027/0227-5910/a000465>.
- Bornstein MH, Putnick DL, Lansford JE, Al-Hassan SM, Bacchini D, Bombi AS, Chang L, Deater-Deckard K, Di Giunta L, Dodge KA, Malone PS, Oburu P, Pastorelli C, Skinner AT, Sorbring E, Steinberg L, Tapanya S, Tirado LMU, Zelli A and Alampay LP (2017) 'Mixed blessings': Parental religiousness, parenting, and child adjustment in global perspective. *The Journal of Child Psychology and Psychiatry* 58, 880–892. <https://doi.org/10.1111/jcpp.12705>.
- Bozzini AB, Bauer A, Maruyama J, Simões R and Matijasevich A (2021) Factors associated with risk behaviors in adolescence: A systematic review. *Brazilian Journal of Psychiatry* 43, 210–221. <https://doi.org/10.1590/1516-4446-2019-0835>.
- Cai T (2013). Investigation of ways to handle sampling weights for multilevel model analyses. *Sociological Methodology* 43, 178–219. <https://journals.sagepub.com/doi/abs/10.1177/0081175012460221>.
- Carballo JJ, Llorente C, Kehrmann L, Flamarique I, Zuddas A, Purper-Ouakil D, Hoekstra PJ, Coghill D, Schulze UME, Dittmann RW, Buitelaar JK, Castro-Fornieles J, Lievesley K, Santosh P, Arango C, Curran S, Selema L, Flanagan R, Craig I, Parnell N, Yeboah K, Sala R, Singh J, Fiori F, Pupier F, Vinkenvleugel L, Glennon J, Bakker M, Drent C, Bloem E, Steenhuis M-P, Berg R, Häge A, Dau MB, Mechler K, Rauscher S, Aslan S, Schlanser S, Keller F, Schneider A, Plener P, Fegert Jörg M, Paton J, Macey M, Iessa N, Alfred K, Helen F, Nick P, Baillon C, Peyre H, Cohen D, Bonnot O, Brunelle J, Franc N, Rayssé P, Humbertclaude Véronique, Rodriguez-Quiroga A, Díaz-Caneja CM, Espliego A, Merchán J, Tapia C, Baeza I, Romero S, La Fuente A, Pintor M, Ligas F, Cera FM, Frongia R, Falissard B, Schwalber A, Ditttrich J, Wohner A, Zimmermann K, Schwalber A and Aitchison K (2020) Psychosocial risk factors for suicidality in children and adolescents. *European Child & Adolescent Psychiatry* 29, 759–776. <https://doi.org/10.1007/s00787-018-01270-9>.
- Central Statistics Office. *Census of Population* (2023) Census of population 2022 - diversity, migration, ethnicity, Irish Travellers & religion. Available at <https://www.cso.ie/en/releasesandpublications/ep/p-cpp5/censusofpopulation>

- 2022profile5-diversitymigrationethnicityirishtravellersreligion/religion/ (accessed 29 May 2025).
- Cipriano A, Cella S and Cotrufo P (2017) Nonsuicidal self-injury: A systematic review. *Frontiers in Psychology* 8, 1946. <https://doi.org/10.3389/fpsyg.2017.01946>.
- Community–University Partnership for the Study of Children Youth and Families (2011) *Review of the Piers-Harris Childrens Self-Concept Scale*, 2nd edn. Alberta, Canada, Edmonton. Available at <https://www.ualberta.ca/community-university-partnership/media-library/community-university-partnership/resources/tools—assessment/piers-harris-2may-2012.pdf> (accessed 20 March 2024).
- Department of Education and Science (2004) A brief description of the Irish Education System. Available at <https://assets.gov.ie/24755/dd437da6d2084a49b0ddb316523aa5d2.pdf> (accessed 20 March 2024).
- Department of Education and Skills (2017) DEIS Plan, 2017. Available at <https://www.gov.ie/en/policy-information/4018ea-deis-delivering-equality-of-opportunity-in-schools/> (accessed 20 March 2024).
- Department of Health (2020–2030) Sharing the vision: A mental health policy for everyone. Available at <https://www.hse.ie/eng/about/who/mentalhealth/sharing-the-vision/> (accessed 5 December 2025).
- Disabato DJ, Kashdan TB, Short JL and Jarden A (2017) What predicts positive life events that influence the course of depression? A longitudinal examination of gratitude and meaning in life. *Cognitive Therapy & Research* 41, 444–458. <https://link.springer.com/article/10.1007/s10608-016-9785-x>.
- Dowling K and Barry MM (2020) Evaluating the implementation quality of a social and emotional learning program: A mixed methods approach. *International Journal of Environmental Research & Public Health* 17(9), 3249. <https://doi.org/10.3390/ijerph17093249>.
- Downes P, Pike S and Murphy S (2020) Marginalised students in Primary and Post-Primary DEIS Schools and other settings: System gaps in policy and practice and the priority issues for consideration, with reference to the impact of Covid 19. Available at <https://www.dcu.ie/sites/default/files/inline-files/educationaldisadvantagecentredcuinstituteofeducationjointoireachtascommitteeducationsubmissiondecember2020-1.pdf> (accessed 20 March 2024).
- Doyle L, Treacy MP and Sheridan A (2015) Self-harm in young people: Prevalence, associated factors, and help-seeking in school-going adolescents. *International Journal of Mental Health Nursing* 24, 485–494. <https://doi.org/10.1111/inm.12144>.
- Ford T, Parker C, Salim J, Goodman R, Logan S and Henley W (2018) The relationship between exclusion from school and mental health: A secondary analysis of the British Child and Adolescent Mental Health Surveys 2004 and 2007. *Psychological Medicine* 48, 629–641. <https://doi.org/10.1017/S003329171700215x>.
- Fraguas D, Díaz-Caneja CM, Ayora M, Durán-Cutilla M, Abregú-Crespo R, Ezquiaga-Bravo I, Martín-Babarro J and Arango C (2021) Assessment of school anti-bullying interventions: A meta-analysis of randomized clinical trials. *JAMA Pediatrics* 175, 44–55. <https://doi.org/10.1001/jamapediatrics.2020.3541>.
- Goodman A and Goodman R (2009) Strengths and difficulties questionnaire as a dimensional measure of child mental health. *Journal of the American Academy of Child and Adolescent Psychiatry* 48, 400–403. <https://doi.org/10.1097/chi.0b013e3181985068>.
- Griffin E, McMahon E, McNicholas F, Corcoran P, Perry IJ and Arensman E (2018) Increasing rates of self-harm among children, adolescents and young adults: A 10-year national registry study 2007–2016. *Social Psychiatry & Psychiatric Epidemiology* 53, 663–671. <https://doi.org/10.1007/s00127-018-1522-1>.
- Hawton K, Saunders KE and O'Connor RC (2012) Self-harm and suicide in adolescents. *Lancet* 379, 2373–2382. [https://doi.org/10.1016/S0140-6736\(12\)60322-5](https://doi.org/10.1016/S0140-6736(12)60322-5).
- Inglis T (2007) Catholic identity in contemporary Ireland: Belief and belonging to tradition. *Journal of Contemporary Religion* 22, 205–220. <https://doi.org/10.1080/13537900701331064>.
- James G, Witten D, Hastie T and Tibshirani R (2013) *An Introduction to Statistical Learning: With Applications in R*. 1st Edn. Hastie T, Tibshirani R, Witten D (editors). New York: Springer.
- Keyes KM and Platt JM (2023) Annual research review: Sex, gender, and internalizing conditions among adolescents in the 21st century - trends, causes, consequences. *The Journal of Child Psychology and Psychiatry* 65, 384–407. <https://doi.org/10.1111/jcpp.13864>.
- Kidger J, Heron J, Leon DA, Tilling K, Lewis G and Gunnell D (2015) Self-reported school experience as a predictor of self-harm during adolescence: A prospective cohort study in the South West of England (ALSPAC). *Journal of Affective Disorders* 173, 163–169. <https://doi.org/10.1016/j.jad.2014.11.003>.
- Kim HH-s and Chun J (2020) Bullying victimization, school environment, and suicide ideation and plan: Focusing on youth in low- and middle-income countries. *Journal of Adolescent Health* 66, 115–122. <https://doi.org/10.1016/j.jadohealth.2019.07.006>.
- Kim-Spoon J, Farley JP, Holmes C, Longo GS and McCullough ME (2014) Processes linking parents' and adolescents' religiousness and adolescent substance use: Monitoring and self-control. *Journal of Youth and Adolescence* 43, 745–756. <https://doi.org/10.1007/s10964-013-9998-1>.
- Kuiper N, Goldston D, Godoy Garraza L, Walrath C, Gould M and McKeon R (2019) Examining the unanticipated adverse consequences of youth suicide prevention strategies: A literature review with recommendations for prevention programs. *Suicide and Life-Threatening Behavior* 49, 952–965. <https://doi.org/10.1111/sltb.12492>.
- Lew B, Chistopolskaya K, Osman A, Huen JMY, Abu Talib M and Leung ANM (2020) Meaning in life as a protective factor against suicidal tendencies in Chinese university students. *BMC Psychiatry* 20, 73. <https://bmcpsy psychiatry.biomedcentral.com/articles/10.1186/s12888-020-02485-4>.
- Lewer D, Gilbody S, Lewis G, Pryce J, Santorelli G, Wadman R, Watmuff A and Wright J (2024) How do schools influence the emotional and behavioural health of their pupils? A multi-level analysis of 135 schools in the Born in Bradford inner city multi-ethnic birth cohort. *Social Psychiatry & Psychiatric Epidemiology* 59(8), 1335–1346. <https://doi.org/10.1007/s00127-023-02608-8>.
- McEvoy D, Brannigan R, Cooke L, Butler E, Walsh C, Arensman E and Clarke M (2023) Risk and protective factors for self-harm in adolescents and young adults: An umbrella review of systematic reviews. *Journal of Psychiatric Research* 168, 353–380. <https://doi.org/10.1016/j.jpsychires.2023.01.017>.
- McEvoy D, Brannigan R, Walsh C, Arensman E and Clarke M (2024) Identifying high-risk subgroups for self-harm in adolescents and young adults: A longitudinal latent class analysis of risk factors. *Journal of Affective Disorders* 351, 40–48. <https://doi.org/10.1016/j.jad.2024.01.230>.
- McEvoy D, Joyce M, Mongan D, Clarke M and Codd M (2024) Comparing times of self-harm presentations to hospital emergency departments in children, adolescents, young adults and adults: A national registry study 2007–2019. *BMC Psychiatry* 24, 474. <https://doi.org/10.1186/s12888-024-05921-x>.
- McMahon EM, Keeley H, Cannon M, Arensman E, Perry IJ, Clarke M, Chambers D and Corcoran P (2014) The iceberg of suicide and self-harm in Irish adolescents: A population-based study. *Social Psychiatry & Psychiatric Epidemiology* 49, 1929–1935. <https://doi.org/10.1007/s00127-014-0907-z>.
- Moran P, Chandler A, Dudgeon P, Kirtley OJ, Knipe D, Pirkis J, Sinyor M, Allister R, Ansloos J, Ball MA, Chan LF, Darwin L, Derry KL, Hawton K, Heney V, Hetrick S, Li A, Machado DB, McAllister E, McDaid D, Mehra I, Niederkrotenthaler T, Nock MK, O'Keefe VM, Oquendo MA, Osafo J, Patel V, Pathare S, Peltier S, Roberts T, Robinson J, Shand F, Stirling F, Stoor JP A, Swinger N, Turecki G, Venkatesh S, Waitoki W, Wright M, Yip PS F, Spoelma MJ, Kapur N, O'Connor RC and Christensen H (2024) The Lancet Commission on self-harm. *Lancet* 404, 1445–1492. [https://doi.org/10.1016/S0140-6736\(24\)01121-8](https://doi.org/10.1016/S0140-6736(24)01121-8).
- Murphy D, Williams J, Murray A and Smyth E (2019) *Growing Up in Ireland – National Longitudinal Study of Children: Design, Instrumentation and Procedures for Cohort '98 at 17/18 years of age*. Dublin, Ireland: Department of Children and Youth Affairs. https://www.esri.ie/system/files/publications/BKMNEXT386_0.pdf (accessed 20 March 2024).
- National Institute for Health and Care Excellence (NICE) (2022) Self-harm: Assessment, management and preventing recurrence. Available at <https://www.nice.org.uk/guidance/ng225> (accessed 07 September 2022).

- O'Connor RC and Kirtley OJ (2018) The integrated motivational-volitional model of suicidal behaviour. *Philosophical Transactions of the Royal Society B Biological Sciences* 373 (1754), 20170268. <https://doi.org/10.1098/rstb.2017.0268>.
- O'Connor RC, Rasmussen S, Miles J and Hawton K (2009) Self-harm in adolescents: Self-report survey in schools in Scotland. *British Journal of Psychiatry* 194, 68–72. <https://doi.org/10.1192/bjp.bp.107.047704>.
- Paulo M, Vagos P, Ribeiro Da Silva D and Rijo D (2020) The role of shame and shame coping strategies on internalizing/externalizing symptoms: Differences across gender in adolescents. *European Journal of Developmental Psychology* 17, 578–597. <https://doi.org/10.1080/17405629.2019.1682991>.
- Pisinger VSC, Hawton K and Tolstrup JS (2019) School- and class-level variation in self-harm, suicide ideation and suicide attempts in Danish high schools. *Scandinavian Journal of Public Health* 47, 146–156. <https://doi.org/10.1177/1403494818799873>.
- Prinstein MJ, Heilbron N, Guerry JD, Franklin JC, Rancourt D, Simon V and Spirito A (2010) Peer influence and nonsuicidal self injury: Longitudinal results in community and clinically-referred adolescent samples. *Journal of Abnormal Child Psychology* 38, 669–682. <https://doi.org/10.1007/s10802-010-9423-0>.
- Roberts E, Fraser A, Gunnell D, Joinson C and Mars B (2020) Timing of menarche and self-harm in adolescence and adulthood: A population-based cohort study. *Psychological Medicine* 50, 2010–2018. <https://doi.org/10.1017/S0033291719002095>.
- Robison S, Jagers J, Rhodes J, Blackmon BJ and Church W (2017) Correlates of educational success: Predictors of school dropout and graduation for urban students in the deep south. *Children and Youth Services Review* 73, 37–46. https://aquila.usm.edu/fac_pubs/15328.
- Sawyer SM, Azzopardi PS, Wickremarathne D and Patton GC (2018) The age of adolescence. *Lancet Child and Adolescent Health* 2, 223–228. [https://doi.org/10.1016/s2352-4642\(18\)30022-1](https://doi.org/10.1016/s2352-4642(18)30022-1).
- Sourander A, Jensen P, Rönning JA, Elonheimo H, Niemelä S, Helenius H, Kumpulainen K, Piha J, Tamminen T, Moilanen I, Almqvist F (2007) Childhood bullies and victims and their risk of criminality in late adolescence: The Finnish from a boy to a man study. *Archives of Pediatrics & Adolescent Medicine* 161, 546–552. <https://doi.org/10.1001/archpedi.161.6.546>.
- Spilman SK, Neppl TK, Donnellan MB, Schofield TJ and Conger RD (2013) Incorporating religiosity into a developmental model of positive family functioning across generations. *Developmental Psychology* 49, 762–774. <https://doi.org/10.1037/a0028418>.
- Stänicke LI (2021) The punished self, the unknown self, and the harmed self – toward a more nuanced understanding of self-harm among adolescent girls. *Frontiers in Psychology* 12, 543303. <https://doi.org/10.3389/fpsyg.2021.543303>.
- Trafford AM, Carr MJ, Ashcroft DM, Chew-Graham CA, Cockcroft E, Cybulski L, Garavini E, Garg S, Kabir T, Kapur N, Temple RK, Webb RT and Mok PLH (2023) Temporal trends in eating disorder and self-harm incidence rates among adolescents and young adults in the UK in the 2 years since onset of the COVID-19 pandemic: A population-based study. *The Lancet Child & Adolescent Health* 7, 544–554. [https://doi.org/10.1016/s2352-4642\(23\)00126-8](https://doi.org/10.1016/s2352-4642(23)00126-8).
- TUSLA: Child and Family Service (2025) SCP - school completion programme. Available at <https://www.tusla.ie/services/educational-welfare-services/scp/> (accessed 5 December 2025).
- TUSLA: Education Welfare Service (2025) What will the school do if my child misses a lot of school? Available at <https://www.tusla.ie/tess/information-for-parents-and-guardians-tess/education-welfare-service/what-will-the-school-do-if-my-child-misses-a-lot-of-school/#:~:text=The%20school%20must%20tell%20tess,child%20is%20missing%20at%20school> (accessed 2 December 2025).
- Van Ryzin MJ, Cil G and Roseth CJ (2023) Costs and benefits of cooperative learning as a universal school-based approach to adolescent substance use prevention. *Journal of Community Psychology* 51, 438–452. <https://doi.org/10.1002/jcop.22916>.
- Victor SE, Muehlenkamp JJ, Hayes NA, Lengel GJ, Styer DM and Washburn JJ (2018). Characterizing gender differences in nonsuicidal self-injury: Evidence from a large clinical sample of adolescents and adults. *Comprehensive Psychiatry* 82, 53–60. <https://doi.org/10.1016/j.comppsy.2018.01.009>.
- Wasserman D, Hoven CW, Wasserman C, Wall M, Eisenberg R, Hadlaczky G, Kelleher I, Sarchiapone M, Apter A, Balazs J, Bobes J, Brunner R, Corcoran P, Cosman D, Guillemin F, Haring C, Iosue M, Kaess M, Kahn J-P, Keeley H, Musa GJ, Nemes B, Postuvan V, Saiz P, Reiter-Theil S, Varnik A, Varnik P and Carli V (2015) School-based suicide prevention programmes: The SEYLE cluster-randomised, controlled trial. *Lancet* 385, 1536–1544. [https://doi.org/10.1016/s0140-6736\(14\)61213-7](https://doi.org/10.1016/s0140-6736(14)61213-7).
- Young R, Sweeting H and Ellaway A (2011) Do schools differ in suicide risk? The influence of school and neighbourhood on attempted suicide, suicidal ideation and self-harm among secondary school pupils. *BMC Public Health* 11, 874. <https://doi.org/10.1186/1471-2458-11-874>.