

Title	Diphtheria toxoid dissolving microneedle vaccination: Adjuvant screening and effect of repeated-fractional dose administration
Authors	Leone, M.;Romeijn, S.;Du, G.;Le Dévédec, S. E.;Vrieling, H.;O'Mahony, Conor;Bouwstra, J. A.;Kersten, G.
Publication date	2020-02-28
Original Citation	Leone, M., Romeijn, S., Du, G., Le Dévédec, S. E., Vrieling, H., O'Mahony, C., Bouwstra, J. A. and Kersten, G. (2020) 'Diphtheria toxoid dissolving microneedle vaccination: Adjuvant screening and effect of repeated-fractional dose administration', International Journal of Pharmaceutics, 580, 119182 (11pp). doi: 10.1016/j.ijpharm.2020.119182
Type of publication	Article (peer-reviewed)
Link to publisher's version	10.1016/j.ijpharm.2020.119182
Rights	© 2020, Elsevier B.V. All rights reserved. This manuscript version is made available under the CC BY-NC-ND 4.0 license. - https://creativecommons.org/licenses/by-nc-nd/4.0/
Download date	2024-03-29 05:27:23
Item downloaded from	https://hdl.handle.net/10468/9848



Appendix




Figure S1. DT-specific IgG1 and IgG2a ratio titers on day 63. Bars represent mean \pm SEM, n = 8. No significant difference among the groups was found ($p > 0.05$). hMN: hollow microneedle; DT: diphtheria toxoid; PI: poly(I:C); CT: cholera toxin; Gib: gibbsite; Boe: boehmite; AlPO₄: aluminum phosphate.




Figure S2. DT-specific IgG1 and IgG2a ratio titers measured in BALB/c mice on day 63.

Bars represent mean \pm SEM, n = 8. *p < 0.05, **p < 0.01, ***p < 0.001. SFD: single-full dose; RFrD: repeated-fractional dose; dMN: dissolving microneedles; hMN: hollow microneedle; DT: diphtheria toxoid; PI: poly(I:C); Gib: gibbsite; E: empty dMN; AlPO₄: aluminum phosphate