

Title	Why are vinyl cations sluggish electrophiles?
Authors	Byrne, Peter A.;Kobayashi, Shinjiro;Würthwein, Ernst-Ulrich;Ammer, Johannes;Mayr, Herbert
Publication date	2017-01-01
Original Citation	Byrne, P. A., Kobayashi, S., Würthwein, E.-U., Ammer, J. and Mayr, H. (2017) 'Why Are Vinyl Cations Sluggish Electrophiles?', Journal of the American Chemical Society, 139(4), pp. 1499-1511. doi: 10.1021/jacs.6b10889
Type of publication	Article (peer-reviewed)
Link to publisher's version	<a href="http://pubs.acs.org/doi/abs/10.1021/jacs.6b10889">http://pubs.acs.org/doi/abs/10.1021/jacs.6b10889</a> - <a href="http://pubs.acs.org/doi/pdf/10.1021/jacs.6b10889">10.1021/jacs.6b10889</a>
Rights	© 2017 American Chemical Society. This document is the Accepted Manuscript version of a Published Work that appeared in final form in Journal of the American Chemical Society, copyright © American Chemical Society after peer review and technical editing by the publisher. To access the final edited and published work see <a href="https://pubs.acs.org/doi/pdf/10.1021/jacs.6b10889">https://pubs.acs.org/doi/pdf/10.1021/jacs.6b10889</a>
Download date	2025-04-08 20:55:48
Item downloaded from	<a href="https://hdl.handle.net/10468/7701">https://hdl.handle.net/10468/7701</a>



# UCC

**University College Cork, Ireland**  
 Coláiste na hOllscoile Corcaigh

# Why are Vinyl Cations Sluggish Electrophiles?

Peter A. Byrne,<sup>a</sup> Shinjiro Kobayashi,<sup>a</sup> Ernst-Ulrich Würthwein,<sup>b</sup> Johannes Ammer,<sup>a</sup> Herbert Mayr<sup>a\*</sup>

a) Department Chemie, Ludwig-Maximilians-Universität München,  
Butenandtstr. 5-13, 81377 München, Germany.


b) Organisch-Chemisches Institut, Westfälische Wilhelms-Universität,  
48149 Münster, Germany

**Abstract.** The kinetics of the reactions of the vinyl cations **2** [ $\text{Ph}_2\text{C}=\text{C}^+-(4\text{-MeO-C}_6\text{H}_4)$ ] and **3** [ $\text{Me}_2\text{C}=\text{C}^+-(4\text{-MeO-C}_6\text{H}_4)$ ] (generated by laser flash photolysis) with diverse nucleophiles (e.g. pyrroles, halide ions, alcoholic and aqueous solvents) have been determined photometrically. It was found that the reactivity order of the nucleophiles toward these vinyl cations is the same as that towards diarylcarbenium ions (benzhydrylium ions). However, the reaction rates of vinyl cations are affected only half as much by variation of the nucleophiles as those of the benzhydrylium ions. For that reason, the relative reactivities of vinyl cations and benzhydrylium ions depend strongly on the nature of the nucleophiles. It is shown that vinyl cations **2** and **3** react, respectively, 227 and 14 times more slowly with trifluoroethanol than the parent benzhydrylium ion  $(\text{Ph})_2\text{CH}^+$ , even though in solvolysis reactions (80 % aqueous ethanol at 25 °C) the vinyl bromides leading to **2** and **3** ionize much more slowly (half-lives 1.15 yrs and 33 days) than  $(\text{Ph})_2\text{CH-Br}$  (half-life 23 s). The origin of this counter-intuitive phenomenon was investigated by high-level MO calculations. We report that vinyl cations are not exceptionally high energy intermediates, and that high intrinsic barriers for the  $\text{sp}^2 \rightleftharpoons \text{sp}$  rehybridizations account for the general phenomenon that vinyl cations are formed slowly by solvolytic cleavage of vinyl derivatives, and are also consumed slowly by reactions with nucleophiles.

## Introduction

Over the course of the last century, the properties of carbocations (which frequently are reactive intermediates) have been investigated by diverse techniques in order to understand their role in organic chemical reactions.<sup>1,2,3,4,5,6,7,8,9,10,11</sup> Although there exist several means for quantifying the stabilities of carbocations in solution and in the gas phase,<sup>2-10,12,13</sup> general stability scales for carbocations ( $\text{R}^+$ ) do not exist, i.e., the absolute stability of a given carbocation cannot be uniquely defined.<sup>2,14</sup> Frequently, the equilibrium constants of their reactions with a certain reference Lewis base  $\text{X}^-$  (eq. 1a), i.e. their Lewis acidities with respect to  $\text{X}^-$ , are employed as a measure of their relative stabilities.<sup>2-9,12,15,16,17</sup> Alternatively, carbocations have been ranked


according to their Brønsted acidities with respect to a certain Brønsted base (eq. 1b),<sup>2b,18</sup> and most Brønsted acidity scales refer to the reaction medium (solvent) as the reference base.



Solvolysis reactions often proceed with rate-determining heterolytic cleavage of R-X (reverse of reaction 1a). The initially formed intermediate carbocation  $\text{R}^+$  is then immediately trapped by the solvent and does not recombine with the leaving group.<sup>1,19</sup> As the transition state of this cleavage is generally assumed to be carbocation-like (Hammond's postulate), the rates of the solvolysis reactions of R-X have frequently been considered to be a measure for the relative stabilities of the intermediate carbocations.<sup>1,6c</sup>

In several series of solvolysis reactions, linear relationships between the measured solvolysis rates and the Lewis acidities of the intermediate carbocations (which are derived from equilibrium measurements) have been observed, but deviations from such rate-equilibrium relationships have also been reported.<sup>12,15,20,21,22</sup> In our investigations of the reactivities of benzhydryl derivatives, we have observed, for example, that benzhydrylium ion **1a** is formed 22 times faster than **1b** during solvolysis of the corresponding benzoates (Scheme 1)<sup>20</sup> even though **1a** is the stronger Lewis acid according to equilibrium studies in solution and computational studies for the gas phase (Scheme 1).<sup>12,15</sup>

**Scheme 1.** Rate constants (*k*) and Gibbs energies of reaction ( $\Delta G^0$ ) for heterolysis reactions giving benzhydrylium ions **1a** and **1b**.




<sup>a</sup> Experimental rate constant for reaction at 25 °C in 80:20 MeCN/H<sub>2</sub>O.<sup>20</sup>

<sup>b</sup> Gibbs reaction energy  $\Delta G^0$  at 20 °C calculated using  $\lg K = LA + LB$ ,<sup>12</sup> in which the *LA* values of **1a** and **1b** in CH<sub>2</sub>Cl<sub>2</sub> (−5.39 and −5.72, respectively), and the *LB* value of benzoate in MeCN (17.45) were employed.

<sup>c</sup> Negative of the calculated gas phase methyl anion affinity of the benzhydrylium ion at 20 °C (see reference 15).

What is the origin of this discrepancy? Direct rate measurements in the solvents used for the solvolysis studies showed that most flash photolytically generated benzhydryl cations react

with chloride and bromide ions under diffusion control.<sup>19</sup> As there is obviously no barrier for the ion combination, one can conclude that in the reverse reaction (first step of an S<sub>N</sub>1 reaction) the transition state also corresponds to the ion pair (Figure 1).




**Figure 1.** First step of the S<sub>N</sub>1 reaction of a benzhydryl halide giving benzhydrylium ion and halide anion.

On the other hand, we have measured significant barriers for the combinations of benzhydrylium ions with carboxylate anions (as depicted for the reactions from right to left in Figure 2).<sup>21</sup> From the principle of microscopic reversibility one can now derive that the first step of an S<sub>N</sub>1 reaction of a benzhydryl carboxylate also must proceed via a transition state that is higher in energy than the carbocation (see reactions from left to right in Figure 2).

According to Marcus (eq. 2),<sup>23a-g</sup> the Gibbs activation energy ( $\Delta G^\ddagger$ ) of a chemical reaction can be expressed by a combination of the Gibbs reaction energy ( $\Delta G^\circ$ ) and the intrinsic barrier ( $\Delta G_0^\ddagger$ ), the latter of which corresponds to  $\Delta G^\ddagger$  for an identity reaction. For reaction series where identity reactions cannot be established, e.g. carbocation-anion combinations, the intrinsic barrier  $\Delta G_0^\ddagger$  is commonly obtained by extrapolations to reactions with  $\Delta G^\circ = 0$ .<sup>23h</sup>

$$\Delta G^\ddagger = \Delta G_0^\ddagger + \frac{\Delta G^\circ}{2} + \frac{(\Delta G^\circ)^2}{16\Delta G_0^\ddagger} \quad (2)$$

Consider two heterolysis reactions, involving different species A-X and B-X, that have identical Gibbs energies of reaction ( $\Delta G^\circ$ ) but different Gibbs energies of activation ( $\Delta G^\ddagger(\text{A}) < \Delta G^\ddagger(\text{B})$ ), as shown in Figure 2. Since the thermodynamic contribution ( $\Delta G^\circ$ ) to the Gibbs energy of activation in both cases is identical, the difference between  $\Delta G^\ddagger(\text{A})$  and  $\Delta G^\ddagger(\text{B})$  must arise exclusively from the differences between the intrinsic barriers  $\Delta G_0^\ddagger(\text{A})$  and  $\Delta G_0^\ddagger(\text{B})$  (not shown explicitly in Figure 2). Thus, in the case depicted in Figure 2, the intrinsic barrier  $\Delta G_0^\ddagger(\text{B})$  must be greater than  $\Delta G_0^\ddagger(\text{A})$ . The energy profiles in Figure 2 now illustrate the counterintuitive conclusion that in cases where the difference of the heterolysis rates is mostly due to a difference in intrinsic barriers, the carbocation that is formed more quickly, A<sup>+</sup>, also reacts more quickly than B<sup>+</sup> with X<sup>-</sup>.



**Figure 2.** Comparison of ionizations of the alkyl halides A-X and B-X to give carbocations A<sup>+</sup> and B<sup>+</sup> of equal Lewis acidity with different rates.


Let us now return to the example illustrated in Scheme 1. As the Lewis acidity of carbocation **1a** is slightly higher than that of **1b**, the higher solvolysis rate of **1a-X** compared to **1b-X** must be due to the lower intrinsic barrier for the ionization of **1a-X** than of **1b-X**. With this conclusion, we can explain why carbocation **1a**, which is formed faster in S<sub>N</sub>1 reactions than **1b**, has also been found to generally react faster with nucleophiles than **1b**.<sup>24</sup>

What is the consequence of these considerations for vinyl cation chemistry? The very low S<sub>N</sub>1 reactivities of vinyl halides and vinyl tosylates<sup>25,26,27</sup> have commonly been ascribed to low thermodynamic stabilities of vinyl cations due to the sp-hybridization of the carbenium center.<sup>28,29</sup> However, when one considers that the 1-phenylvinyl cation has a similar hydride affinity to the benzyl cation,<sup>30,31,32</sup> only 2.9 kcal mol<sup>-1</sup> greater than that of the *tert*-butyl cation,<sup>30</sup> the question arises whether the low solvolysis rates of vinyl halides are really mostly due to the low thermodynamic stabilities of dicoordinated carbenium ions. If instead high intrinsic barriers for the sp<sup>2</sup> ⇌ sp rehybridization were responsible for the slow solvolyses of vinyl derivatives, the discussion of Figure 2 implies that also the reverse reaction should be slow, and vinyl cations should not be extraordinarily reactive intermediates but rather sluggish electrophiles. Though rate constants for the reactions of vinyl cations generated by laser-flash photolysis with a variety of nucleophiles have previously been reported,<sup>33,34</sup> a systematic comparison of the electrophilic reactivities of vinyl cations and tricoordinated carbenium ions has not yet been performed. We now report on a systematic analysis of the reactivities of vinyl cations and show that exceptionally high intrinsic barriers account for their extraordinarily slow formation in S<sub>N</sub>1 reactions as well as for their slow reactions with nucleophiles.

## Results

Vinyl cations **2** and **3** (Chart 1), which can be generated by laser flash photolysis from **4-Br** and **5-Br**, were selected as representative vinyl cations to study the rates of the reactions with

nucleophiles **6-22** (Table 1) in MeCN (**13** and **14** in 2,2,2-trifluoroethanol, TFE) and with the solvent mixtures **23-41** listed in Table 2.



**Chart 1.** Structures of vinyl cations **2** and **3** and vinyl derivatives **4-X** and **5-X**.

**Table 1.** Structures and values of the nucleophile-specific parameters  $N$  and  $s_N$  (in MeCN unless otherwise indicated) for nucleophiles **6-22**.

Nucleophile	#	$N$	$s_N$	Ref	Nucleophile	#	$N$	$s_N$	Ref
	<b>6</b>	3.76	0.91	35		<b>15</b>	12.35	0.72	37
	<b>7</b>	4.41	0.96	24		<b>16</b>	13.77	0.70	37
	<b>8</b>	5.21	1.00	24		<b>17</b>	15.10	0.73	37
	<b>9</b>	5.85	1.03	36		<b>18</b>	15.65	0.74	37
	<b>10</b>	8.01	0.96	36		<b>19</b>	17.35	0.68	37
	<b>11</b>	9.11	0.88	24	$n\text{Bu}_4\text{N}^+\text{OAc}^-$	<b>20</b>	16.90	0.75	21
$\text{F}_3\text{C}-\text{CH}_2-\text{NH}_2$	<b>12</b>	10.13	0.75	37	$n\text{Bu}_4\text{N}^+\text{OMs}^-$	<b>21</b>	-	-	-
$n\text{Bu}_4\text{N}^+\text{Cl}^-$	<b>13</b>	10.3 <sup>a</sup>	0.60	19	$n\text{Bu}_4\text{N}^+\text{I}^-$	<b>22</b>	-	-	-
$n\text{Bu}_4\text{N}^+\text{Br}^-$	<b>14</b>	11.7 <sup>a</sup>	0.60	19					

<sup>a</sup> Solvent = 2,2,2-trifluoroethanol (TFE).


**Table 2.** Nucleophile-specific reactivity parameters  $N_1$  and  $s_N$  for solvents and solvent mixtures (v/v) **23-41**.<sup>38</sup>

Nucleophile		$N_1$	$s_N$
TFE <sup>a</sup>	<b>23</b>	1.11 <sup>a</sup>	0.96 <sup>a</sup>
TFE/H <sub>2</sub> O (90:10)	<b>24</b>	2.93	0.88
TFE/H <sub>2</sub> O (60:40)	<b>25</b>	3.42	0.90
MeCN/H <sub>2</sub> O (90:10)	<b>26</b>	4.56	0.94
MeCN/H <sub>2</sub> O (80:20)	<b>27</b>	5.02	0.89
MeCN/H <sub>2</sub> O (67:33)	<b>28</b>	5.02	0.90
MeCN/H <sub>2</sub> O (50:50)	<b>29</b>	5.05	0.89
MeCN/EtOH (90:10)	<b>30</b>	5.19	0.96
MeCN/EtOH (80:20)	<b>31</b>	5.77	0.92
MeCN/EtOH (67:33)	<b>32</b>	6.06	0.90
MeCN/EtOH (50:50)	<b>33</b>	6.37	0.90
MeCN/EtOH (33:67)	<b>34</b>	6.74	0.89
MeCN/EtOH (20:80)	<b>35</b>	6.94	0.90
MeCN/EtOH (10:90)	<b>36</b>	7.10	0.90
EtOH/H <sub>2</sub> O (40:60)	<b>37</b>	5.81	0.90
EtOH/H <sub>2</sub> O (50:50)	<b>38</b>	5.96	0.89
EtOH/H <sub>2</sub> O (60:40)	<b>39</b>	6.28	0.87
EtOH/H <sub>2</sub> O (80:20)	<b>40</b>	6.68	0.85
EtOH/H <sub>2</sub> O (90:10)	<b>41</b>	7.03	0.86

<sup>a</sup> TFE = 2,2,2-trifluoroethanol;  $N_1$  and  $s_N$  are taken from ref 35.

## Product Characterization

**Scheme 2.** Reactions of vinyl cations **2** and **3**, generated by ionization of **4-OMs** and **5-Br**, respectively, in TFE in the presence of different nucleophiles.



Vinyl cations **2** and **3** can be transiently generated in solvents of high ionizing power by heterolysis of precursor vinyl halides or pseudohalides **4-X** and **5-X**.<sup>25,27,39,40a,41</sup> We have found that heterolyses of **4-OMs** and **5-Br** occur at convenient rates at 40 °C in 2,2,2-trifluoroethanol (TFE), a solvent of low nucleophilicity ( $N_1 = 1.11$ ).<sup>35</sup> When the vinyl cations are generated in aqueous TFE, the ketones **42** and **43** are formed quantitatively, as shown in Scheme 2a.

In order to verify that the vinyl cations generated in TFE solution can also be intercepted by other nucleophiles, **4-OMs** and **5-Br** were dissolved in TFE containing  $\geq 1 \text{ mol L}^{-1}$  of **10** or **20** as representative nucleophiles. Following our previously published methodology for carrying out Friedel-Crafts-type chemistry in neutral aqueous or alcoholic solutions,<sup>42</sup> high nucleophile concentrations were employed to avoid trapping by trifluoroethanol or traces of water. In this way, the reactions of solvolytically generated **2** and **3** with pyrrole **10** ( $N = 8.01$  in MeCN) resulted in high conversion to 3-vinylpyrroles **44** and **45**, respectively (Scheme 2b), without formation of hydrolysis products.<sup>43</sup> The reaction of **2** with [<sup>n</sup>Bu<sub>4</sub>N]OAc (**20**;  $N = 16.9$  in MeCN) gave vinyl acetate **46** (Scheme 2c) as the major product, with the formation of a small amount (5%) of hydrolysis product **42**.

*ipso*-Substitution (at C-4 of the anisyl group), which has been observed in reactions of photochemically generated vinyl cations **2** with cyanide or alkoxide,<sup>44,45</sup> but not with alcohols,<sup>45e,f</sup> did not occur under the conditions described in Scheme 2. We therefore conclude,





that all nucleophiles added to the cationic sp-center of **2** and **3** in our kinetic experiments, in line with results from earlier studies carried out under similar conditions.<sup>33,34,39,40,45e,45f,46,47,48,49</sup>

### Kinetic Investigations

Vinyl cations **2** and **3** were generated in MeCN, 2,2,2-trifluoroethanol, or solvent mixtures (Tables 1 and 2) by irradiation of the precursor vinyl bromides **4-Br** and **5-Br** with a 7-ns laser pulse of  $\lambda = 266$  nm. A single signal is observed in the UV-Vis spectrum of each transient cation.<sup>50</sup> In the presence of a large excess of the nucleophiles **6** - **22** (pseudo first-order conditions), the absorbance of the vinyl cation was generally observed to undergo mono-exponential decay, as shown for the reaction of **2** with **9** (Scheme 3) in Figure 3a.<sup>51</sup> Least-squares fitting of the single-exponential function  $A_t = A_0 e^{-k_{\text{obs}} t} + C$  ( $A_0$  and  $A_t$  are the absorbances at time 0 and time  $t$ , respectively, and  $C$  is a constant) to the absorbance decay curve for the reaction of **2** or **3** with a nucleophile yielded  $k_{\text{obs}}$  ( $\text{s}^{-1}$ ) for the particular concentration of nucleophile.

**Scheme 3.** Generation of vinyl cation **2** by laser flash photolysis of **4-Br** in MeCN, and subsequent reaction with pyrrole **9**.






















**Figure 3.** (a) Decay curve ( $\lambda = 350$  nm) for the reaction of vinyl cation **2** with pyrrole **9**; (b) The second order rate constant  $k$  was obtained from the slope of the plot of  $k_{\text{obs}}$  vs.  $[\mathbf{9}]_0$ .

Plots of the pseudo-first-order rate constants  $k_{\text{obs}}$  vs concentrations of the nucleophile were linear (see Figure 3b) and can be expressed by equation 3:

$$k_{\text{obs}} = k[\text{Nu}] + k_{\text{solv}} \quad (3)$$

where  $[\text{Nu}]$  is the molar concentration of the nucleophile,  $k$  (the slope of a plot of  $k_{\text{obs}}$  vs.  $[\text{Nu}]$ ) is the second-order rate constant for the reaction of vinyl cation with nucleophile (values in Table 3), and the intercept is the first-order rate constant for the reaction of the vinyl cation with solvent ( $k_{\text{solv}}$ ). Nine of the 14 correlations in acetonitrile showed intercepts in the range  $(1.4 - 1.7) \times 10^5 \text{ s}^{-1}$ , which we ascribe to the reaction of **2** with acetonitrile. The only strong deviation from this value ( $6 \times 10^6 \text{ s}^{-1}$ , for the reaction of **2** with the strong nucleophile **20**) is likely to be a consequence of the problematic extrapolation of the very large rate constants to the concentration  $[\mathbf{20}] = 0$ . Whereas the intercept for the correlation of  $k_{\text{obs}}$  vs.  $[\text{Bu}_4\text{N}]\text{Cl}$  (**13**) in trifluoroethanol agrees exactly with the previously reported rate constant for the reaction of **2** with trifluoroethanol (nucleophile **23**), for unknown reasons the corresponding plot of  $k_{\text{obs}}$  vs.  $[\text{Bu}_4\text{N}]\text{Br}$  (**14**) gives an intercept which is two times larger.


**Table 3.** Second-order rate constants,  $k$  (20 °C), for the reactions of vinyl cation **2** with nucleophiles **6-22** (solvent MeCN unless otherwise indicated).

Nucleophile			Nucleophile		
		$k$ (L mol <sup>-1</sup> s <sup>-1</sup> )			$k$ (L mol <sup>-1</sup> s <sup>-1</sup> )
	<b>6</b>	1.64 × 10 <sup>5</sup>		<b>15</b>	1.23 × 10 <sup>7</sup>
	<b>7</b>	3.10 × 10 <sup>5</sup>		<b>16</b>	4.64 × 10 <sup>7</sup>
	<b>8</b>	1.30 × 10 <sup>6</sup>		<b>17</b>	1.28 × 10 <sup>8</sup>
	<b>9</b>	1.07 × 10 <sup>8</sup>		<b>18</b>	2.18 × 10 <sup>8</sup>
	<b>10</b>	3.17 × 10 <sup>8</sup>		<b>19</b>	6.88 × 10 <sup>8</sup>
	<b>11</b>	1.50 × 10 <sup>7</sup>		<b>20</b>	4.79 × 10 <sup>9</sup>
	<b>12</b>	6.01 × 10 <sup>6</sup>		<b>21</b>	5.99 × 10 <sup>7</sup>
	<b>13</b>	8.06 × 10 <sup>5</sup> <sup>a</sup>		<b>22</b>	ca. 4.4 × 10 <sup>9</sup> <sup>b</sup>
	<b>14</b>	3.67 × 10 <sup>6</sup> <sup>a</sup>			


<sup>a</sup> Solvent = 2,2,2-trifluoroethanol (TFE).

<sup>b</sup> Approximate rate constant derived from an experiment with a single concentration of [Bu<sub>4</sub>N]I. See details in Supporting Information, p S25.

Monoexponential decays of the absorbances of the vinyl cations **2** and **3** were also observed when they were generated in trifluoroethanol (**23**) and in the solvent mixtures **24-41** (see Figure 4 for an example), and the first-order rate constants  $k_{\text{obs}}$  were obtained from fitting of  $A_t = A_0 e^{-k_{\text{obs}} t} + C$  to the decay curves. However,  $k_{\text{obs}}$  did not increase linearly with [H<sub>2</sub>O] (Figure 5) and remained almost constant when the water content was raised beyond 20% v/v H<sub>2</sub>O, in line with previous reports on the consumption of benzhydrylium<sup>38,52</sup> and tritylium ions<sup>52</sup> in MeCN/H<sub>2</sub>O mixtures. Since a similar situation was also observed for other solvent mixtures, Table 4 lists the first-order rate constants  $k = k_{\text{obs}}$  for reactions of **2** with the solvent nucleophiles **23-41**.



**Figure 4.** Decay curve ( $\lambda = 350$  nm) from the reaction of vinyl cation **2** with 67:33 MeCN/EtOH (**32**).



**Figure 5.** Plot of  $k_{\text{obs}}$  for the consumption of **2** in MeCN/H<sub>2</sub>O solvents vs. concentration of H<sub>2</sub>O.



**Table 4.** Observed first-order rate constants,  $k$  (20 °C), for the reactions of vinyl cation **2** with solvent nucleophiles **23-41**.

Nucleophile		$k$ (s <sup>-1</sup> )
TFE	<b>23</b>	$1.4 \times 10^4$ <sup>a</sup>
TFE/H <sub>2</sub> O (90:10)	<b>24</b>	$2.47 \times 10^4$
TFE/H <sub>2</sub> O (60:40)	<b>25</b>	$6.86 \times 10^4$
MeCN/H <sub>2</sub> O (90:10)	<b>26</b>	$3.22 \times 10^5$
MeCN/H <sub>2</sub> O (80:20)	<b>27</b>	$4.22 \times 10^5$
MeCN/H <sub>2</sub> O (67:33)	<b>28</b>	$4.80 \times 10^5$
MeCN/H <sub>2</sub> O (50:50)	<b>29</b>	$5.24 \times 10^5$
MeCN/EtOH (90:10)	<b>30</b>	$6.06 \times 10^5$
MeCN/EtOH (80:20)	<b>31</b>	$1.05 \times 10^6$
MeCN/EtOH (67:33)	<b>32</b>	$2.01 \times 10^6$
MeCN/EtOH (50:50)	<b>33</b>	$3.02 \times 10^6$
MeCN/EtOH (33:67)	<b>34</b>	$4.13 \times 10^6$
MeCN/EtOH (20:80)	<b>35</b>	$5.14 \times 10^6$
MeCN/EtOH (10:90)	<b>36</b>	$6.93 \times 10^6$
EtOH/H <sub>2</sub> O (40:60)	<b>37</b>	$1.47 \times 10^6$
EtOH/H <sub>2</sub> O (50:50)	<b>38</b>	$1.61 \times 10^6$
EtOH/H <sub>2</sub> O (60:40)	<b>39</b>	$1.77 \times 10^6$
EtOH/H <sub>2</sub> O (80:20)	<b>40</b>	$2.88 \times 10^6$
EtOH/H <sub>2</sub> O (90:10)	<b>41</b>	$4.03 \times 10^6$

<sup>a</sup> Rate constant taken from ref 33.

Second-order rate constants for the reactions of **3** with pyrroles **9** and **10** and first-order rate constants for the reactions of **3** with the solvents **23-27**, **30**, and **31** were derived in a similar manner to that described above for vinyl cation **2**. The rate constants determined in this way are shown in Table 5.

**Table 5.** Rate constants,  $k$  (20 °C), for the reactions of vinyl cation **3** with  $\pi$ -nucleophiles **9** and **10** and with solvent nucleophiles **23-27**, **30**, and **31**.

Nucleophile		$k$
 N-Me	<b>9</b>	$3.19 \times 10^8$ $\text{L mol}^{-1} \text{s}^{-1}{}^a$
 Me	<b>10</b>	$1.21 \times 10^9$ $\text{L mol}^{-1} \text{s}^{-1}{}^a$
TFE	<b>23</b>	$2.3 \times 10^5 \text{s}^{-1}{}^b$
TFE/H <sub>2</sub> O (90:10)	<b>24</b>	$5.85 \times 10^5 \text{s}^{-1}$
TFE/H <sub>2</sub> O (60:40)	<b>25</b>	$1.76 \times 10^6 \text{s}^{-1}$
MeCN/H <sub>2</sub> O (90:10)	<b>26</b>	$1.30 \times 10^7 \text{s}^{-1}$
MeCN/H <sub>2</sub> O (80:20)	<b>27</b>	$1.46 \times 10^7 \text{s}^{-1}$
MeCN/EtOH (90:10)	<b>30</b>	$1.62 \times 10^7 \text{s}^{-1}$
MeCN/EtOH (80:20)	<b>31</b>	$2.70 \times 10^7 \text{s}^{-1}$

<sup>a</sup> Solvent = MeCN

<sup>b</sup> Rate constant taken from ref 33. TFE = 2,2,2-trifluoroethanol.

## Correlations and Discussion


In numerous investigations we have shown that the second-order rate constants  $k$  for the reactions of electrophiles with nucleophiles at 20 °C may be calculated using equation 4,

$$\lg k (20\text{ °C}) = s_N (E + N) \quad (4)$$

where  $E$  characterizes the electrophilicity of the electrophile (treated as being solvent-independent), while  $N$  represents the nucleophilicity of the nucleophile, and  $s_N$  is a nucleophile-specific susceptibility parameter.<sup>24,53</sup> Whereas new  $N$  and  $s_N$  parameters of nucleophiles are derived from linear plots of  $\lg k$  vs. the known  $E$  parameters of the reference electrophiles, new  $E$  parameters of electrophiles have been derived from linear correlations between  $(\lg k)/s_N$  and  $N$  of the reference nucleophiles.


Following this procedure,  $(\lg k)/s_N$  was plotted vs.  $N$  for all investigated nucleophiles with known nucleophilicity parameters (Figure 6). As the published reactivity parameters  $N$  and  $s_N$

for nucleophiles **6** – **20** refer to second-order rate constants in acetonitrile, while those for solvents **23** – **41** (designated  $N_1$ ) refer to first-order rate constants, it is possible to plot the logarithms of the second-order rate constants for the  $\pi$ -systems **6** – **11**, the amines **12** and **15** – **19**, and the anions **13**, **14**, **20** as well as those of the first-order rate constants for the solvents **23** – **41** side by side in Figure 6. The remarkably good correlation for these diverse nucleophiles over a reactivity range of more than 16 orders of magnitude shows that the nucleophilic reactivities toward vinyl cation **2** ( $sp$ -electrophile) follow the same pattern as those toward benzhydrylium ions. However, equation (4) is not fulfilled because the slope of this correlation is 0.53 and not 1.0, as required by equation (4), showing that **2** is substantially less sensitive to changes in the reactivity of the nucleophile than benzhydrylium ions ( $sp^2$ -electrophiles).



**Figure 6.** Plot of  $(\lg k/s_N)$  vs.  $N$  for the reactions of vinyl cation **2** with nucleophiles **6-20** and **23-41**.


A similar plot, shown in Figure 7, was constructed using the first- and second-order rate constants  $k$  (listed in Table 5) for the reactions of vinyl cation **3** with a variety of nucleophiles. Again, a strong linear correlation is observed over a wide range of reactivity, and again a slope of much less than 1 is obtained. It is remarkable that even pyrrole **10** fits this correlation though the rate constant of  $1.21 \times 10^9 \text{ L mol}^{-1} \text{ s}^{-1}$  is already close to the diffusion control limit. Vinyl cation **3** thus shows very similar behavior to **2**. Its higher electrophilic reactivity may be due to reduced steric shielding of the cationic carbon center.



**Figure 7.** Plot of  $(\lg k/s_N)$  vs.  $N$  for vinyl cation **3**, from its reactions with various nucleophiles.

Analogous linear correlations of  $(\lg k/s_N)$  vs.  $N$  with slopes much smaller than 1 were previously found for  $S_N2$  reactions of alkyl halides,<sup>54</sup> indicating that variation of the nucleophiles also had a smaller influence on the rate constants of the  $S_N2$  reactions than on those for the reactions with benzhydrylium ions. Although the correlations shown in Figures 6 and 7 might also be mathematically expressed by adding an additional electrophile-specific susceptibility parameter  $s_E$  to eq. (4),<sup>54</sup> we refrain from deriving electrophilicity parameters  $E$  from the extended correlations. The reason is that  $E$  determined in this way would represent an approximate reactivity ranking toward very weak nucleophiles which react with rate constants close to 1 ( $\lg k = 0$ ), i. e., reactions which have little relevance in practice, because commonly used solvents react much faster.


For that reason, let us directly compare rate constants for the reactions of nucleophiles with vinyl cations **2** and **3** and with benzhydrylium ions **47** – **49** (Chart 2).



**Chart 2.** Benzhydrylium ions **47** - **49**.




In Figure 8, the  $\lg k$  values for the reactions of **2** (values of  $k$  taken from Tables 3 and 4) are plotted against the corresponding  $\lg k$  values for the reactions of benzhydrylium ion **47** ( $k(\mathbf{47})$  from Table S1 on p. 43-44 of the Supporting Information).<sup>55</sup> The plot shows a fair correlation between the two data sets and (similar to Figure 6) that the rate constants for the reactions with the vinyl cation **2** are less affected by variation of the nucleophiles than the corresponding rate constants for **47** (slope = 0.48). If we neglect the two pyrroles (**9** and **10**), which deviate significantly from the correlation, one can see that the vinyl cation **2** reacts faster than the dimethoxybenzhydrylium ion **47** with weak nucleophiles ( $k < 5 \times 10^6 \text{ s}^{-1}$  or  $\text{L mol}^{-1}\text{s}^{-1}$ ), while stronger nucleophiles react faster with benzhydrylium ion **47**. Overall, however, the vinyl cation **2** and the dimethoxybenzhydrylium ion **47** have comparable electrophilic reactivities.



**Figure 8.** Correlation of  $\lg k$  for the reactions of vinyl cation **2** with various nucleophiles (**6-16**, **23-41**) with  $\lg k$  for the analogous reactions of the 4,4'-dimethoxybenzhydrylium ion **47** (from Table S1 in SI). The red line is a plot of  $\lg k(\mathbf{47})$  against itself to highlight the crossing range where nucleophiles react with equal rates with **2** and **47**.

An analogous comparison can be made between vinyl cation **3** and benzhydrylium ion **48**. A plot of  $\lg k$  for the reactions of **3** with various nucleophiles vs.  $\lg k$  for the corresponding reactions of **48** ( $k(\mathbf{48})$  from Table S2 on p. S44 of the Supporting Information) shows a good linear correlation (Figure 9),<sup>56</sup> with a slope significantly less than 1 (slope = 0.56). Vinyl cation **3** thus has an electrophilicity comparable to that of **48**.




**Figure 9.** Correlation of  $\lg k$  for the reactions of vinyl cation **3** with various nucleophiles (**9**, **23-27**, **30**, **31**) with  $\lg k$  for the analogous reactions of the 4,4'-dimethylbenzhydrylium ion **48** (from Table S2 in SI). The red line is a plot of  $\lg k(\mathbf{48})$  against itself to highlight the crossing range where nucleophiles react with equal rates with **3** and **48**.

The observation that vinyl cations **2** and **3** show reactivities toward nucleophiles that are similar to the corresponding reactivities of the donor-stabilized benzhydrylium ions **47** and **48** appears surprising at first glance, as the latter species are formed much faster in  $S_N1$  reactions than the corresponding vinyl derivatives. In fact, the ionizations of the benzhydryl bromides, which yield the highly stabilized benzhydryl cations **47** and **48**, are so fast in aqueous ethanol that it is not possible at present to measure their solvolysis rates.

Therefore, we compare here the solvolyses of the vinyl bromides **4-Br** and **5-Br** with that of  $\text{Ph}_2\text{CHBr}$  (for which experimental data are available). Scheme 4 shows that the benzhydryl bromide solvolyzes  $10^5$  to  $10^6$  times faster in 80% aqueous ethanol at 25 °C than the vinyl bromides **4-Br** and **5-Br**. Despite its much faster rate of formation, the parent benzhydrylium ion reacts one to two orders of magnitude faster with trifluoroethanol at 20 °C than **2** or **3** (compare the second reactions in each of Scheme 4a, 4b and 4c). Similar rate ratios have been found for activation-controlled reactions of these electrophiles with numerous other nucleophiles (i. e. reactions which do not proceed with diffusion controlled rates).

**Scheme 4.** Solvolysis reactions of (a) **4-Br**,<sup>a</sup> (b) **5-Br**,<sup>b</sup> and (c) Ph<sub>2</sub>CHBr<sup>c</sup> in 80:20 EtOH/H<sub>2</sub>O at 25 °C, and reactions of cations **2**, **3**, and **49** with TFE at 20 °C.



<sup>a</sup> For the first step of reaction (a), a value of  $k = 1.92 \times 10^{-4} \text{ s}^{-1}$  was measured at 120 °C;<sup>39d</sup> reported activation parameters allow extrapolation to value at 25 °C given in the Scheme. The rate constant for the second step is taken from reference 33.


<sup>b</sup> For the first step of reaction (b), a value of  $k = 8.78 \times 10^{-4} \text{ s}^{-1}$  was measured at 120 °C;<sup>40c</sup> reported activation parameters allow extrapolation to value at 25 °C given in the Scheme. The rate constant for the second step is taken from reference 33.

<sup>c</sup> The rate constant for the first step of (c) is from reference 57. The rate constant for the second step is from reference 58.

Laser flash experiments have shown that the reactions of the parent benzhydrylium ion **49** with Br<sup>−</sup> are diffusion-controlled in all alcoholic solvents investigated.<sup>19</sup> As there is no barrier for the ion combination, the principle of microscopic reversibility implies that the transition state for the heterolytic cleavage of Ph<sub>2</sub>CHBr corresponds to the ion-pair, as illustrated in Figure 10.

In contrast, the reaction of Br<sup>−</sup> with vinyl cation **2** is activation controlled, proceeding with a rate constant of  $3.7 \times 10^6 \text{ L mol}^{-1} \text{ s}^{-1}$  in TFE (Table 3). For the same reaction in 80 % aqueous ethanol one can calculate  $k = 7.4 \times 10^6 \text{ L mol}^{-1} \text{ s}^{-1}$  from  $N = 14.5$ ,  $s_N = 0.6$  (for Br<sup>−</sup> in 80 % aqueous ethanol)<sup>19</sup> using the correlation equation given in Figure 6. As this reaction is not diffusion controlled, transition state theory can be applied to calculate an activation energy of  $\Delta G^\ddagger = 8.1 \text{ kcal mol}^{-1}$  for the ion combination in 80 % aqueous ethanol (Figure 10, right hand side). The kinetic data implies that the heterolyses of Ph<sub>2</sub>CHBr and **4-Br** have almost identical  $\Delta G^\circ$  values, as shown in Figure 10.

Figure 10 clearly illustrates that the major reason for the different solvolysis rates of  $\text{Ph}_2\text{CHBr}$  and **4-Br** is the difference in the intrinsic barriers (since the Gibbs energies of reaction  $\Delta G^\circ$  are very similar). Hence, the transition state of the first step of the  $\text{S}_{\text{N}}1$  reaction of **4-Br** cannot be carbocation-like i.e. it does not correspond to the ion pair of  $\text{Br}^-$  with vinyl cation **2**. We demonstrate explicitly below using quantum chemical calculations that  $\Delta\Delta G^\circ = \Delta G^\circ_{\text{Ph}_2\text{CHBr}} - \Delta G^\circ_{\text{4-Br}}$  is indeed negligible (*vide infra*), as derived from kinetic data for the construction of Figure 10.




**Figure 10.** Schematic Gibbs energy profiles ( $\text{kcal mol}^{-1}$ ) for the ionization of benzhydryl bromide and vinyl bromide **4-Br** in 80% aqueous ethanol at 25 °C.  $\Delta G^\ddagger$  values were calculated using the Eyring equation (for the solvolysis reactions, rate constants from Scheme 4a and 4c were used; see main text for details of the calculation for addition of  $\text{Br}^-$  to **2**).

### Common Ion Rate Depression

Although  $\text{S}_{\text{N}}1$  reactions are usually accelerated when the polarity of the solvent is increased by salt additives,<sup>1c,d,h,59</sup> in certain cases, the opposite effect is observed. Common ion rate depression is a phenomenon whereby the rate of a solvolysis reaction  $k_{\text{obs}}$  is slowed by addition of a salt containing the anion of the leaving group ( $\text{X}^-$ ).<sup>1h,19,27,60</sup> It is observed when the recombination of the carbocation  $\text{R}^+$  with  $\text{X}^-$  is faster than addition of the solvent to  $\text{R}^+$  (with first order rate constant  $k_{\text{solv}}$ ); i.e., when  $k_{-1}[\text{X}^-] \geq k_{\text{solv}}$  (see Scheme 5, and equation (5), in which the full expression for  $k_{\text{obs}}$  is given, and  $\alpha = k_{-1}/k_{\text{solv}}$ ).

**Scheme 5.** Reversible heterolysis of  $\text{RX}$  to give  $\text{R}^+$ , which may react with solvent (completing the solvolysis) or revert to  $\text{RX}$  by reaction with  $\text{X}^-$ .



$$k_{\text{obs}} = \frac{k_1 k_{\text{solv}}}{k_{\text{solv}} + k_{-1}[\text{X}^-]} = \frac{k_1}{1 + \alpha[\text{X}^-]} \quad (5)$$

As highly reactive carbocations are immediately trapped by the solvent, which is present in high concentration, common ion return is only observed when highly stabilized carbocations are generated in solvents of low nucleophilicity. Since vinyl cations had been considered to be highly reactive because of their slow formation in  $S_N1$  processes, the observation of common ion rate depression in  $S_N1$  processes was highly surprising, and the numerous attempts to rationalize this phenomenon have been summarized.<sup>60b</sup> With the knowledge that vinyl cations **2** and **3** have relatively low electrophilic reactivities, similar to those of the highly stabilized benzhydrylium ions **47** and **48** (see Figures 8 and 9), two systems for which common ion depression has generally been observed,<sup>19</sup> it is no longer surprising that this effect was also found for the solvolyses of **4-X** and **5-X** ( $X = \text{Cl}, \text{Br}$ ).<sup>60b</sup>

In previous work we have demonstrated that the occurrence of common ion rate depression can be derived from the directly measured rate constants of the reactions of the independently generated carbocations with the anionic leaving groups (i. e.  $X^-$ ) and the solvent.<sup>19</sup> From the rate constants in Tables 3 and 4 for the reactions of **2** with  $\text{Cl}^-$  in TFE ( $8.06 \times 10^5 \text{ L mol}^{-1} \text{ s}^{-1}$ ),  $\text{Br}^-$  in TFE ( $3.67 \times 10^6 \text{ L mol}^{-1} \text{ s}^{-1}$ ), and TFE ( $1.40 \times 10^4 \text{ s}^{-1}$ ), respectively, one can calculate  $\alpha$  values of 58 and 262, respectively, in good agreement with common ion rate depressions observed for similar systems under comparable conditions.<sup>61</sup>


## Computational Analysis

Directly measured rate constants as well as the observation of common ion rate depression thus indicate that vinyl cations are sluggish electrophiles, despite their very slow formation in  $S_N1$  reactions. By combining the experimentally determined rate constants for the solvolysis reactions and ion recombinations (see Figure 10 and associated discussion above), we had derived that  $\Delta G^\circ$  for the heterolytic cleavage of the C-Br bond in  $\text{Ph}_2\text{CHBr}$  is almost the same as that for heterolysis of **4-Br**. This conclusion is in line with quantum chemical calculations, as we show below.

Recently, we have calculated<sup>62</sup> gas phase methyl anion affinities of benzhydrylium ions as a measure for their relative Lewis acidities (Scheme 6, table entries 1 – 3).<sup>12</sup> Comparison with entries 4 and 5 shows that the methyl anion affinities of vinyl cations **2** and **3** are closely similar to that of the unsubstituted benzhydrylium ion (**49**), as derived from the rate constants for forward and backward reactions in Figure 10.<sup>63</sup> Kinetic data and quantum chemically calculated Lewis acidities thus agree on the conclusion that the differences in intrinsic barriers account for the fact that vinyl cations **2** and **3**, despite their much higher Lewis acidities, are

not more electrophilic than the highly stabilized benzhydrylium ions **47** and **48**, respectively (Figures 8 – 9 and associated discussion).

**Scheme 6.** Calculated Gibbs energies of reactions<sup>a</sup> of methyl anion with benzhydrylium ions **47-49** and of vinyl cations **2** and **3** (methyl anion affinities) in the gas phase.



Entry		R	$\Delta G^\circ$ (kcal mol <sup>-1</sup> )
1	<b>47</b>	OMe	-178.9 <sup>b</sup>
2	<b>48</b>	Me	-188.6 <sup>b</sup>
3	<b>49</b>	H	-197.7 <sup>b</sup>
4	<b>2</b>	Ph	-198.2 <sup>c</sup>
5	<b>3</b>	Me	-200.2 <sup>c</sup>

<sup>a</sup> Computational method and basis set employed for all values of  $\Delta G^\circ$ : B3-LYP/6-311++G(3df,pd)//B3LYP/6-31G(d,p).<sup>62,64</sup>

<sup>b</sup> Calculated value reported in reference 12.

<sup>c</sup> This work.<sup>62,63,64</sup>

The calculations shown in Table 6 confirm that in the gas phase,  $\text{Ph}_2\text{CH}^+$  and **2** also have almost equal affinities toward  $\text{Br}^-$  ( $\Delta\Delta G^\circ = 0.9$  kcal mol<sup>-1</sup>, entries 1, 2) and  $\text{Br}^-$  solvated by one molecule of water ( $\Delta\Delta G^\circ = 1.1$  kcal mol<sup>-1</sup>, entries 3, 4), in agreement with our observations based on the kinetic data (Figure 10, above). As expected, the ion combinations are calculated to be much less exergonic in solution (entries 5 – 8), but the very small calculated values of  $\Delta G^\circ$  in aqueous solution indicate that the PCM model significantly overestimates the ion solvation energies.<sup>65</sup>


**Table 6.** Calculated Gibbs energies of the reactions<sup>a</sup> of Br<sup>−</sup> (and Br<sup>−</sup> solvated by 1 water molecule) with the parent benzhydrylium ion (**49**) and vinyl cation **2** in the gas phase and in aqueous solution.<sup>b</sup>

$\text{R}^{\oplus} + \text{X}^{\ominus} \xrightarrow{\Delta G^{\circ} \text{ (kcal mol}^{-1}\text{)}} \text{R-X}$				
Entry	R <sup>+</sup>	X <sup>−</sup>	Gas Phase / Solvent Model	ΔG° (kcal mol <sup>−1</sup> )
1	Ph <sub>2</sub> CH <sup>+</sup>	Br <sup>⊖</sup>	Gas	−113.9
2	<b>2</b>	Br <sup>⊖</sup>	Gas	−114.8
3	Ph <sub>2</sub> CH <sup>+</sup>	Br <sup>⊖</sup> (H <sub>2</sub> O) <sub>1</sub>	Gas	−102.1
4	<b>2</b>	Br <sup>⊖</sup> (H <sub>2</sub> O) <sub>1</sub>	Gas	−103.2
5	Ph <sub>2</sub> CH <sup>+</sup>	Br <sup>⊖</sup>	Water	−9.5
6	<b>2</b>	Br <sup>⊖</sup>	Water	−15.3
7	Ph <sub>2</sub> CH <sup>+</sup>	Br <sup>⊖</sup> (H <sub>2</sub> O) <sub>1</sub>	Water	−5.8
8	<b>2</b>	Br <sup>⊖</sup> (H <sub>2</sub> O) <sub>1</sub>	Water	−11.6

<sup>a</sup> Computational method and basis set employed for all values of ΔG°: TPSS/def2TZVP+GD3.<sup>66,67</sup>

<sup>b</sup> PCM; method TPSS/def2TZVP+GD3. See Table S7 (p S83 of Supporting Information) for further quantities determined from these calculations.

When trying to localize the transition states for the heterolytic cleavage reactions of **4-Br** and Ph<sub>2</sub>CHBr in aqueous solution (i.e., the reactions in Figure 10) by the DFT method TPSS/def2TZVP+GD3 using the PCM continuum solvent model,<sup>68</sup> we observed a continuous increase of the potential energy (*E*<sub>tot</sub>) as the C-Br bond length was elongated, and the ion pair was reached without passing through a maximum in *E*<sub>tot</sub> (Figure 11). The same result was found when one molecule of water was explicitly considered (see Figure S4 in SI, p S84). In both cases, the gradient was much larger for the vinyl bromide due to the higher force constant of the C<sub>sp2</sub>-Br bond.




**Figure 11.** Intrinsic reaction coordinate calculations for the cleavage of the C-Br bond in **4-Br** and  $\text{Ph}_2\text{CHBr}$  (Total energies  $E_{\text{tot}}$ ; TPSS/def2TZVP+GD3; PCM (solvent = water)).

The fact that the maximum of total energy in the **4-Br** graph of Figure 7 is slightly lower than  $\Delta E_{\text{tot}}$  calculated for the reaction of **2** with  $\text{Br}^-$  in water ( $-24.9 \text{ kcal mol}^{-1}$ , see Table S7 in SI on p S83) may be due to the PCM-model used for the calculations on the heterolysis reactions and on the separated species and to the fact that for longer C-Br distances the closed-shell restriction does not apply. A computational assessment of the transition state for the vinyl bromide heterolysis (for which the activation barrier was unequivocally deduced from kinetic data – see Figure 10 and associated discussion) might be possible by explicit consideration of more solvent molecules. However, general problems of this type of treatment have recently been pointed out by Singleton,<sup>69</sup> and in any case, this approach is beyond the scope of this investigation.

As mentioned in the Introduction, the intrinsic barrier in the Marcus equation ( $\Delta G_0^\ddagger$ ) corresponds to the Gibbs activation energy of an identity reaction. Since there are no identity reactions for the ion combinations in eq. (1), we analyzed the intrinsic barriers for the identity hydride transfers illustrated in Scheme 7 as models for the transition states of  $\text{sp}^3 \rightleftharpoons \text{sp}^2$  and  $\text{sp}^2 \rightleftharpoons \text{sp}$  rehybridizations (Table 7).<sup>70</sup>



**Scheme 7.** Formation of (a) tricoordinated and (b) dicoordinated hydrido-bridged carbenium ions  $[R\cdots H\cdots R]^+$  from the isolated carbenium ion  $R^+$  and its parent compound, RH (without considering species showing aromatic interactions ( $\pi$ -stacking)).



The left part of Table 7 allows comparison of quantum chemically calculated and experimental gas-phase hydride affinities for several tri- and di-coordinated carbenium ions.<sup>71</sup> While the absolute hydride affinities calculated with different basis sets differ considerably (see Supporting Information, p. S69-70), there is good agreement between relative hydride affinities obtained with different computational methods and experimental data.

**Table 7.** Comparison of Calculated and Experimental Gas Phase Values  $\Delta H_{HA}$  (298 K) and  $\Delta G_{HA}$  (298 K) for Additions of  $H^-$  to Various Carbenium Ions (Hydride Affinity =  $-\Delta H_{HA}$ ), and Calculated Intrinsic Barriers  $\Delta H_{bridge}$  and  $\Delta G_{bridge}$  for the Hydride Transfer Reactions:

$$R-H \xleftarrow[\Delta H_{HA}]{H^-} R^+ \xrightarrow[\Delta H_{bridge}]{R-H} [R\cdots H\cdots R]^+$$

Entry	$R^+$	Experimental $\Delta H_{HA}^a$	Calculated Quantities <sup>b</sup>			
			$\Delta H_{HA}^b$	$\Delta G_{HA}^b$	$\Delta H_{bridge}^b$	$\Delta G_{bridge}^b$
1	$Me-\overset{+}{C}-Me$	-251.8	-259.3	-251.8	-18.5	-8.1
2	$Me-\overset{+}{C}=\equiv$	-264.9	-272.4	-264.2	-14.9	-2.9
3	$Ph-\overset{+}{C}-Me$	-225.1	-234.5	-228.0	-12.2	+0.3
4	$Ph-\overset{+}{C}=\equiv$	-239.7	-247.5	-240.1	-3.8	+8.2
5	$Ph-\overset{+}{C}$	-239.3	-246.8	-240.6	-12.9	-1.9
6	$Ph-\overset{+}{C}-Ph$	-	-220.9	-212.8	-8.0	+3.3

<sup>a</sup> Experimental data (kcal mol<sup>-1</sup>) listed in ref 30.

<sup>b</sup> Calculated values from this work. Method and basis set employed: TPSS/TPSS/def2TZVP+GD3//TPSS/TPSS/def2TZVP+GD3. Units are kcal mol<sup>-1</sup>.

Table 7 shows, for example, that successive replacement of the methyl groups in the isopropyl cation by phenyl (i.e., giving first phenethyl cation and then benzhydryl cation) reduces the hydride affinities by 25 and then 15 kcal mol<sup>-1</sup> in calculated  $\Delta H_{\text{HA}}$  and  $\Delta G_{\text{HA}}$  (entries 1, 3, 6) and in experimentally determined  $\Delta H_{\text{HA}}$ . The hydride affinities of the vinyl cations (entries 2 and 4) are 13 kcal mol<sup>-1</sup> larger than those of the corresponding saturated analogues (entries 1 and 3).

Hydride transfer to tri- or di-coordinated carbenium ions from their parent alkanes or alkenes occurs through hydrido-bridged species (Scheme 7). The enthalpies and Gibbs energies ( $\Delta H_{\text{bridge}}$  and  $\Delta G_{\text{bridge}}$ , respectively) of formation of these entities from the isolated reactants, i. e., the intrinsic barriers for the hydride transfers are shown in the right hand columns of Table 7. Whereas the hydrido-bridged species from isopropyl cation/propane (entry 1) and benzyl cation/toluene (entry 5) are minima on the potential energy surface (gas phase), all other hydrido-bridged species correspond to transition states (Table 7, and Tables S4-S6 in Supporting Information p. S71-S75). All minima and all transition states have a negative  $\Delta H_{\text{bridge}}$  with respect to the isolated reactants. As expected, the tendency to undergo hydrido-bridging decreases with decreasing hydride ion affinity, i.e., for tricoordinated carbenium ions,  $\Delta G_{\text{bridge}}$  increases in the order  $1 < 5 \approx 3 < 6$  and in the order entry 2 < entry 4 for dicoordinated carbenium ions.

In the context of our analysis, the comparisons of entries 2 and 1 and of entries 4 and 3 are of particular importance. Although the vinyl cations in entries 2 and 4 have significantly higher hydride affinities compared to their saturated analogs in entries 1 and 3, respectively, their hydrido-bridging tendencies are much smaller. As the hydrido-bridged complexes represent models for the transition states of  $\text{sp}^3 \rightleftharpoons \text{sp}^2$  and  $\text{sp}^2 \rightleftharpoons \text{sp}$  rehybridizations (Scheme 7), the 8 kcal/mol lower tendency towards hydrido-bridging of the phenyl-substituted vinyl cation in entry 4 compared to the saturated analog in entry 3 reflects the high intrinsic barriers which are responsible for the low rates of vinyl halide heterolyses (low electrofugalities<sup>15</sup> of vinyl cations) and the low rates of nucleophilic addition to vinyl cations (low electrophilicities<sup>15</sup> of vinyl cations).

## Conclusion

Discussions of carbocation reactivities are generally based on the assumption that the transition states of S<sub>N</sub>1 reactions are carbocation-like (Hammond postulate), and that the slower a carbocation is formed in an S<sub>N</sub>1 process, the faster it will react with a nucleophile. Although deviations from this rule have previously been reported, including literature reports on

comparatively low absolute rate constants for reactions of vinyl cations with nucleophiles and common ion rate depression in solvolyses of vinyl derivatives (typical for carbocations of relatively low reactivity), the consequences of these observations for the interpretation of vinyl cation chemistry have rarely been considered.

We have now shown that the transition states of vinyl halide solvolyses are often not carbocation-like. Consequently, the influence of intrinsic barriers on this step cannot be neglected. The approximation to consider only the thermodynamic term (i. e.,  $\Delta G^0$  of the ionization step) for analyzing the kinetic behaviour of carbocations, which works well for the reactions of most weakly stabilized tricoordinated carbenium ions, cannot be applied for vinyl cations. Vinyl cations (i.e., dicoordinated carbenium ions) are weaker electrofuges as well as weaker electrophiles than tricoordinated carbenium ions of similar Lewis acidity because of the high intrinsic barriers for  $sp^2 \rightleftharpoons sp$  rehybridizations.

### Supporting Information

Experimental procedures and characterization of representative products; details of kinetic experiments with vinyl cations **2** and **3**; literature rate constants used for the construction of Figures 8 and 9; copies of NMR spectra; details of quantum chemical calculations. The Supporting Information is available free of charge on the ACS Publications website at DOI: \_

### Author Information

Corresponding Author: \* herbert.mayr@cup.uni-muenchen.de

Notes: The authors declare no competing financial interest.

### Acknowledgements

We thank the Deutsche Forschungsgemeinschaft (SFB 749, Project B1) for financial support, the Humboldt foundation for the provision to PB of a Humboldt Research Fellowship for Postdoctoral Researchers, Giulio Volpin, Dr. David Stephenson, and Claudia Dubler for assistance with NMR spectroscopy, and Dr. A. R. Ofial for helpful comments.

## References and Notes

- 1 General Reviews: (a) *Carbonium Ions*, Vol. 1-5; Olah, G. A.; Schleyer, P. v. R., Eds.; Interscience: New York, 1968-1976; (b) Olah, G. A. *Angew. Chem.* **1973**, *85*, 183-225; *Angew. Chem., Int. Ed. Engl.* **1973**, *12*, 173-212; (c) Streitwieser Jr., A. *Solvolytic Displacement Reactions*; McGraw-Hill: New York, 1962; (d) Ingold, C. K. *Structure and Mechanism in Organic Chemistry*, 2nd ed.; Cornell University Press: Ithaca, NY, 1969; (e) Vogel, P. *Carbocation Chemistry*; Elsevier: Amsterdam, 1985; (f) *Advances in Carbocation Chemistry*, Vol. 1; Creary, X., Ed.; JAI Press: Greenwich, CT, 1989; (g) *Advances in Carbocation Chemistry*, Vol. 2; Coxon, J. M., Ed.; JAI Press: Greenwich, CT, 1995; (h) Raber, D. J.; Harris, J. M.; Schleyer, P. v. R. in *Ions and Ion Pairs in Organic Reactions*; Wiley and Sons: New York; 1974; Volume 2, pp 247-374.
- 2 Electrophilicities of carbocations: (a) Ritchie, C. D. *Can. J. Chem.* **1986**, *64*, 2239-2250, and references therein; (b) Mayr, H. in *Seminars in Organic Synthesis*, 40th "A. Corbella" International Summer School, Gargnano (ISBN 978-88-86208-96-3), Società Chimica Italiana, **2015**, pp 82-107; (c) Mayr, H.; Ofial A. R. in *Carbocations*; Olah, G. A.; Prakash, G. K. S., Eds.; Chapter 13, Wiley: New York, 2004.
- 3  $pK_{R^+}$  values ( $X^- = HO^-$ ): (a) Deno, N. C.; Jaruzelski, J. J.; Schriesheim, A. *J. Am. Chem. Soc.* **1955**, *77*, 3044-3051; (b) Deno, N. C.; Schriesheim, A. *J. Am. Chem. Soc.* **1955**, *77*, 3051-3054; (c) Arnett, E. M.; Bushick, R. D. *J. Am. Chem. Soc.* **1964**, *86*, 1564-1571; (d) Ritchie, C. D.; Fleischhauer, H. *J. Am. Chem. Soc.* **1972**, *94*, 3481-3483; (e) Kitagawa, T.; Takeuchi, K. *J. Phys. Org. Chem.* **1998**, *11*, 157-170, and references therein; (f) Deno, N. C.; Berkheimer, H. E.; Evans, W. L.; Peterson, H. J. *J. Am. Chem. Soc.* **1959**, *81*, 2344-2347; (g) Gandler, J. J. *J. Am. Chem. Soc.* **1985**, *107*, 8218-8223; (h) Amyes, T. L.; Richmond, J. P.; Novale, M. *J. Am. Chem. Soc.* **1992**, *114*, 8032-8041; (i) Ostovic, D.; Lee, I. S. H.; Roberts, R. M. G.; Kreevoy, M. M. *J. Org. Chem.* **1985**, *50*, 4206-4211; (j) Okamoto, K.; Takeuchi, K.; Komatsu, K.; Kubota, Y.; Ohara, R.; Arima, M.; Takahashi, K.; Waki, Y.; Shirai, S. *Tetrahedron* **1983**, *39*, 4011-4024; (k) See also reference 2a.
- 4 Gas phase hydride affinities of carbenium ions ( $X^- = H^-$ ): (a) Wolf, J. F.; Staley, R. H.; Koppel, I.; Taagepera, M.; McIver, R. T.; Beauchamp, J. L.; Taft, R. W. *J. Am. Chem. Soc.* **1977**, *99*, 5417-5429; (b) Lossing, F. P.; Holmes, J. L. *J. Am. Chem. Soc.* **1984**, *106*, 6917-6920; (c) Schultz, J. C.; Houle, F. A.; Beauchamp, J. I. *J. Am. Chem. Soc.* **1984**, *106*, 3917-3927; (d) Karaman, R.; Huang, J.-T. L.; Fry, J. L. *J. Org. Chem.* **1991**, *56*, 188-195; (e) Zhu, X.-Q.; Wang, C.-H. *J. Phys. Chem. A* **2010**, *114*, 13244-13256; (f) Kim, C. K.; Lee, K. A.; Bae, S. Y.; Han, I. S.; Kim, C. K. *Bull. Korean Chem. Soc.* **2004**, *25*, 311-313.
- 5 Solution hydride affinities of carbenium ions ( $X^- = H^-$ ): (a) Cheng, J.-P.; Handoo, K. L.; Parker, V. D. *J. Am. Chem. Soc.* **1993**, *115*, 2655-2660; (b) Arnett, E. M.; Flowers, R. A.; Ludwig, R. T.; Meekhof, A. E.; Walek, S. A. *J. Phys. Org. Chem.* **1997**, *10*, 499-513; (c) See also reference 3i; (d) See also reference 4e.
- 6 Calorimetric determination of enthalpies of ionization of alkyl halides in  $SbF_5/SO_2ClF$  superacid (formally the reverse of the reaction in equation (1b);  $X^- = \text{halide}$ ): (a) Arnett, E. M.; Petro, S. C. *J. Am. Chem. Soc.* **1978**, *100*, 2563-2564; (b) Arnett, E. M.; Petro, S. C. *J. Am. Chem. Soc.* **1978**, *100*, 5402-5407, (c) Arnett, E. M.; Petro, S. C. *J. Am. Chem. Soc.* **1978**, *100*, 5408-5416; (d) Arnett, E. M.; Petro, S. C.; Schleyer, P. v. R. *J. Am. Chem. Soc.* **1979**, *101*, 522-526; (e) Arnett, E. M.; Pienta, N. J. *J. Am. Chem. Soc.* **1980**, *102*, 3329-3334; (f) Arnett, E. M.; Hofelich, T. C. *J. Am. Chem. Soc.* **1983**, *105*, 2889-2895; (g) Arnett, E. M.; Hofelich, T. C. *J. Am. Chem. Soc.* **1982**, *104*, 3522-3524.
- 7 Equilibrium constants for ionization of benzhydryl chlorides ( $X^- = \text{halide}$ ): Schade, C.; Mayr, H.; Arnett, E. M. *J. Am. Chem. Soc.* **1988**, *110*, 567-571.
- 8 Enthalpies and equilibrium constants for reactions of carbocations with carbanions ( $X^- = \text{carbanion}$ ): (a) Troughton, E. B.; Molter, K. E.; Arnett, E. M. *J. Am. Chem. Soc.* **1984**, *105*, 6726-6735; (b) Arnett, E. M.; Chawla, B.; Molter, K.; Amarnath, K.; Healy, M. *J. Am. Chem. Soc.* **1985**, *107*, 5288-5289; (c) Arnett, E. M.; Molter, K. E. *Acc. Chem. Res.* **1985**, *18*, 339-346.
- 9 Gibbs energies of reaction for production of carbocations from neutral precursors in the gas phase, determined by ion cyclotron resonance spectroscopy: (a) Aue, D. H.; Bowers, M. T. in *Gas Phase Ion Chemistry*; Academic Press: New York, 1979; Vol. 2, Chapter 9; (b) Walder, R.; Franklin, J. L. *Int. J. Mass Spectrom.*

- 
- Ion Phys.* **1980**, *36*, 85-112; (c) Kebarle, P. *Annu. Rev. Phys. Chem.* **1977**, *28*, 445-476; (d) Abboud, J.-L. M.; Alkorta, I.; Davalos, J. Z.; Müller, P.; Quintanilla, E.; Rossier, J.-C. *J. Org. Chem.* **2003**, *68*, 3786-3796.
- 10 Destabilized carbocations: Tidwell, T. T. *Angew. Chem.* **1984**, *96*, 16-28; *Angew. Chem., Int. Ed. Engl.* **1984**, *23*, 20-32.
- 11 Gas phase IR-spectroscopy: (a) Crestoni, M. E.; Fornarini, S. *Angew. Chem.* **2012**, *124*, 7487-7489; *Angew. Chem., Int. Ed.* **2012**, *51*, 7373-7375; (b) Chiavolino, B.; Crestoni, M. E.; Dopfer, O.; Maitre, P.; Fornarini, S. *Angew. Chem.* **2012**, *124*, 5031-5033; *Angew. Chem., Int. Ed.* **2012**, *51*, 4947-4949.
- 12 Mayr, H.; Ammer, J.; Baidya, M.; Maji, B.; Nigst, T. A.; Ofial, A. R.; Singer, T. *J. Am. Chem. Soc.* **2015**, *137*, 2580-2599.
- 13 Abboud, J.-L. M.; Alkorta, I.; Dávalos, J. Z.; Müller, P.; Quintanilla, E. *Adv. Phys. Org. Chem.* **2002**, *37* 57-135.
- 14 Hine, J. *Structural Effects on Equilibria in Organic Chemistry*; Robert E. Krieger Publishing Company; Huntington: New York; 1981, pp 215-256.
- 15 Mayr, H.; Ofial, A. R.; *Acc. Chem. Res.* **2016**, *49*, 952-965.
- 16 Equilibrium constants have been reported for reversible C-C bond heterolysis in DMSO to give hydrocarbon salts ( $X^-$  = carbanion) in reference 3e.
- 17 Carbocation reduction potentials ( $E_{red}$ ) have also been used as a measure of their stability: See references 3e, 3i, 3j, and 5b.
- 18 (a) Stewart, R. *The Proton: Applications to Organic Chemistry*; Academic Press: Orlando; 1985, p125; (b) Kresge, A. J.; Chen, H. J.; Hakka, L. E.; Kouba, J. E. *J. Am. Chem. Soc.* **1971**, *93*, 6174-6181; (c) Kresge, A. J.; Mylonakis, S. G.; Sato, Y.; Vitullo, V. P. *J. Am. Chem. Soc.* **1971**, *93*, 6181-6188.
- 19 Minegishi, S.; Loos, R.; Kobayashi, S.; Mayr, H. *J. Am. Chem. Soc.* **2005**, *127*, 2641-2649.
- 20 Streidl, N.; Denegri, B.; Kronja, O.; Mayr, H. *Acc. Chem. Res.* **2010**, *43*, 1537-1549.
- 21 Schaller, H. F.; Tishkov, A. A.; Feng, X.; Mayr, H. *J. Am. Chem. Soc.* **2008**, *130*, 3012-3022.
- 22 (a) Richard, J. P. *Tetrahedron* **1995**, *51*, 1535-1573; (b) Richard, J. P.; Amyes, T. L.; Williams, K. B. *Pure Appl. Chem.* **1998**, *70*, 2007-2014; (c) Amyes, T. L.; Stevens, I. W.; Richard, J. P. *J. Org. Chem.* **1993**, *58*, 6057-6066; (d) Amyes, T. L.; Richard, J. P. *J. Chem. Soc., Chem. Commun.* **1991**, 200-202; (e) Richard, J. P. *J. Am. Chem. Soc.* **1989**, *111*, 1455-1465.
- 23 For details on Marcus theory, see: (a) Marcus, R. A. *Annu. Rev. Phys. Chem.* **1964**, *15*, 155-196; (b) Marcus, R. A. *J. Phys. Chem.* **1968**, *72*, 891-899; (c) Marcus, R. A. *J. Am. Chem. Soc.* **1969**, *91*, 7224-7225; (d) Albery, W. J.; Kreevoy, M. M. *Adv. Phys. Org. Chem.* **1978**, *16*, 87-157; (e) Albery, W. J. *Annu. Rev. Phys. Chem.* **1980**, *31*, 227-263; (f) Marcus, R. A. *Pure Appl. Chem.* **1997**, *69*, 13-29; (g) Marcus, R. A. *Angew. Chem.* **1993**, *105*, 1161-1172; *Angew. Chem., Int. Ed. Engl.* **1993**, *32*, 1111-1121. (h) Peters, K. S.; Gasparrini, S.; Heeb, L. R. *J. Am. Chem. Soc.* **2005**, *127*, 13039-13047.
- 24 Mayr, H.; Bug, T.; Gotta, M. F.; Hering, N.; Irrgang, B.; Janker, B.; Kempf, B.; Loos, R.; Ofial, A. R.; Remennikov, G.; Schimmel, H. *J. Am. Chem. Soc.* **2001**, *123*, 9500-9512.
- 25 *Vinyl Cations*; Stang, P.; Rappoport, Z.; Hanack, M.; Subramanian, L. R., Eds.; Academic Press: New York, 1979.
- 26 (a) *Dicoordinated Cations*; Z. Rappoport, P. J. Stang, Eds.; Wiley: New York, 1997; (b) Grob, C. A. in reference 26a Chapter 1, pp 1-7; (c) Stang, P. J. in *Progress in Physical Organic Chemistry*; Streiwieser Jr., A. J.; Taft, R. W., Eds.; Wiley and Sons: New York; 1973; Vol. 10, pp 205-325; (d) Rappoport, Z. in *Reactive Intermediates*; Abramovitch, R. A., Ed.; Plenum Press: New York; 1983; Vol. 3, pp 427-615; (e) Modena, G.; Tonellato, U. *Adv. Phys. Org. Chem.* **1971**, *9*, 185-280.
- 27 Kitamura, T.; Taniguchi, H. in reference 26a, Chapter 7, pp 321-376.
- 28 Generation of  $\beta$ -silyl-substituted vinyl cations that are stable at low temperature in super-acidic media: (a) Siehl, H.-U.; Kaufmann, F.-P.; Apeloig, Y.; Braude, V.; Danovich, D.; Berndt, A.; Stamatis, N. *Angew. Chem.* **1991**, *103*, 1546-1549; *Angew. Chem., Int. Ed. Engl.* **1991**, *30*, 1479-1482; (b) Siehl, H.-U.; Müller, T.; Gauss,

- J.; Buzek, P.; Schleyer, P. v. R. *J. Am. Chem. Soc.* **1994**, *116*, 6384-6387; (c) F.-P. Kaufmann, H.-U. Siehl, *J. Am. Chem. Soc.* **1992**, *114*, 4937-4939; (d) H.-U. Siehl, F.-P. Kaufmann, K. Hori, *J. Am. Chem. Soc.* **1992**, *114*, 9343-9349.
- 29 Generation of room temperature-stable silyl-substituted vinyl cations: (a) Klaer, A.; Syha, Y.; Nasiri, H. R.; Müller, T. *Chem. Eur. J.* **2009**, *15*, 8414-8423; (b) Müller, T.; Meyer, R.; Lennartz, D.; Siehl, H.-U. *Angew. Chem.* **2000**, *112*, 3203-3206; *Angew. Chem., Int. Ed.* **2000**, *39*, 3074-3077; (c) Müller, T.; Margraf, D.; Syha, Y. *J. Am. Chem. Soc.* **2005**, *127*, 10852-10860; (d) Müller, T.; Juhasz, M.; Reed, C. A. *Angew. Chem.* **2004**, *116*, 1569-1572; *Angew. Chem., Int. Ed.* **2004**, *43*, 1543-1546.
- 30 Aue, D. H. in *Dicoordinated Carbocations*. Rappoport, Z., Stang, P. J., Eds.; Wiley: New York, 1997; pp 105-156. In particular, see pp126-129, Table 2.
- 31 Calculations on stabilities of vinyl cations: (a) Apeloig, Y.; Franke, W.; Rappoport, Z.; Schwarz, H.; Stahl, D. *J. Am. Chem. Soc.* **1981**, *103*, 2770-2780; (b) Apeloig, Y.; Schleyer, P. v. R.; Pople, J. A. *J. Am. Chem. Soc.* **1977**, *99*, 1291-1296; (c) Apeloig, Y.; Schleyer, P. v. R.; Pople, J. A. *J. Org. Chem.* **1977**, *42*, 3004-3011; (d) Apeloig, Y.; Schleyer, P. v. R.; Pople, J. A. *J. Am. Chem. Soc.* **1977**, *99*, 5901-5909; (e) Apeloig, Y.; Stanger, A. *J. Org. Chem.* **1982**, *47*, 1462-1468; (f) van Alem, K.; Lodder, G.; Zuilhof, H. *J. Phys. Chem. A* **2002**, *106*, 10681-10690; (g) Alem, K.; Lodder, G.; Zuilhof, H. *J. Phys. Chem. A* **2000**, *104*, 2780-2787; (h) Mayr, H.; Schneider, R.; Wilhelm, D.; Schleyer, P. v. R. *J. Org. Chem.* **1981**, *46*, 5336-5340.
- 32 Studies of the stabilities of vinyl cations  $[\text{ArC}=\text{CH}_2]^+$  by ion cyclotron resonance techniques: (a) Kobayashi, S.; Matsumoto, T.; Taniguchi, H.; Mishima, M.; Fujio, M.; Tsuno, Y. *Tetrahedron Lett.* **1993**, *34*, 5903-5906; (b) Mishima, M.; Ariga, T.; Fujio, M.; Tsuno, Y.; Kobayashi, S.; Taniguchi, H. *Chem. Lett.* **1992**, 1085-1088.
- 33 Cozens, F. L.; Kanagasabapathy, V. M.; McClelland, R. A.; Steenken, S. *Can. J. Chem.* **1999**, *77*, 2069-2082.
- 34 (a) Kobayashi, S.; Hori, Y.; Hasako, T.; Koga, K.; Yamataka, H. *J. Org. Chem.* **1996**, *61*, 5274-5279. (b) Kobayashi, S.; Schnabel, W.; *Z. Naturforsch. B.: Chem. Sci.* **1992**, *47*, 1319-1323; (c) Kobayashi, S.; Zhu, Q. Q.; Schnabel, W. *Z. Naturforsch. B.: Chem. Sci.* **1988**, *47*, 825-829.
- 35 Ammer, J.; Nolte, C.; Mayr, H. *J. Am. Chem. Soc.* **2012**, *134*, 13902 – 13911.
- 36 Nigst, T.; Westermaier, M.; Ofial, A. R.; Mayr, H. *Eur. J. Org. Chem.* **2008**, 2369-2374.
- 37 Kanzian, T.; Nigst, T. A.; Maier, A.; Pichl, S.; Mayr, H. *Eur. J. Org. Chem.* **2009**, 6379-6385.
- 38 Minegishi, S.; Kobayashi, S.; Mayr, H. *J. Am. Chem. Soc.* **2004**, *126*, 5174-5181.
- 39 (a) Rappoport, Z.; Kaspi, J. *J. Am. Chem. Soc.* **1974**, *96*, 4518-4530; (b) Rappoport, Z.; Kaspi, J. *J. Am. Chem. Soc.* **1970**, *92*, 3220-3221; (c) Rappoport, Z.; Apeloig, Y. *J. Am. Chem. Soc.* **1969**, *91*, 6734-6742; (d) Rappoport, Z.; Gal, A. *J. Am. Chem. Soc.* **1969**, *91*, 5246-5254.
- 40 (a) Rappoport, Z.; Kaspi, J. *J. Chem. Soc., Perkin Trans. 2* **1972**, 1102-1111; (b) Miller, L. L.; Kaufman, D. A. *J. Am. Chem. Soc.* **1968**, *90*, 7282-7287; (c) Rappoport, Z.; Gal, A. *J. Chem. Soc., Perkin Trans. 2* **1973**, 301-310.
- 41 (a) Rappoport, Z.; Kaspi, J., *J. Am. Chem. Soc.* **1975**, *97*, 821-835; (b) Rappoport, Z.; Gal, A., *J. Am. Chem. Soc.* **1969**, *91*, 5246-5254; (c) Rappoport, Z.; Kaspi, J., *J. Am. Chem. Soc.* **1974**, *96*, 586-588.
- 42 Hofmann, M.; Hampel, N.; Kanzian, T.; Mayr, H. *Angew. Chem.* **2004**, *116*, 5518-5521; *Angew. Chem., Int. Ed.* **2004**, *43*, 5402-5405.
- 43 No side products were formed; i.e., other than the 3-vinylpyrrole products, only starting materials remained when the reaction was stopped.
- 44 *ipso*-Substitution with cyanide: Kitamura, T.; Murakami, M.; Kobayashi, S.; Taniguchi, H. *Tetrahedron Lett.* **1986**, *27*, 3885-3888.
- 45 *ipso*-Substitution with alkoxide: (a) Kitamura, T.; Kabashima, T.; Kobayashi, S.; Taniguchi, H. *Tetrahedron Lett.* **1988**, *29*, 6141-6142; (b) Kitamura, T.; Nakamura, I.; Kabashima, T.; Kobayashi, S.; Taniguchi, H. *Chem. Lett.* **1990**, 9-12; (c) Kitamura, T.; Nakamura, I.; Kabashima, T.; Kobayashi, S.; Taniguchi, H. *J. Am. Chem. Soc.*, **1990**, *112*, 6149-6150; (d) Kitamura, T.; Kabashima, T.; Nakamura, I.; Fukada, T.; Taniguchi, H. *J. Am. Chem. Soc.*, **1991**, *113*, 7255-7261; (e) Kitamura, T.; Kabashima, T.; Kobayashi, S.; Taniguchi, H. *Chem. Lett.* **1988**, 1951-1954; (f) Kitamura, T.; Nakamura, I.; Kabashima, T.; Kobayashi, S.; Taniguchi, H. *J. Chem. Soc.*,

- Chem. Commun.* **1989**, 1154-1155; (g) Kitamura, T.; Soda, S.; Nakamura, I.; Fukada, T.; Taniguchi, H. *Chem. Lett.* **1991**, 2195-2198; (h) Hori, K.; Kamada, H.; Kitamura, T.; Kobayashi, S.; Taniguchi, H. *J. Chem. Soc., Perkin Trans. 2* **1992**, 871-877.
- 46 Interception with azide and cyanate nucleophiles: (a) Kitamura, T.; Kobayashi, S.; Taniguchi, H. *Tetrahedron Lett.* **1979**, 20, 1619-1622; (b) Kitamura, T.; Kobayashi, S.; Taniguchi, H. *J. Org. Chem.* **1984**, 49, 4755-4760.
- 47 Interception with thiocyanate: (a) Kitamura, T.; Kobayashi, S.; Taniguchi, H. *Chem. Lett.* **1984**, 1523-1526; (b) Kitamura, T.; Kobayashi, S.; Taniguchi, H. *J. Org. Chem.* **1990**, 55, 1801-1805.
- 48 Interception with acetate or bromide (labelled): van Ginkel, F. I. M.; Cornelisse, J.; Lodder, G. *J. Am. Chem. Soc.* **1991**, 113, 4261-4272.
- 49 Interception with methylphenylsulfide: Kitamura, T.; Kabashima, T.; Taniguchi, H. *J. Org. Chem.* **1991**, 56, 3739-3741.
- 50 The transient UV-Vis spectra of **2** and **3** are included on p. S45 of the Supporting Information.
- 51 The decay curves for the reactions of **2** with nucleophiles **7-9** and **11** ceased to be mono-exponential once a certain conversion had been reached. The use of a particularly large excess of the nucleophile was necessary in each case (involving **7-9** and **11**) to produce decay curves in which the first 50-80% of the decay was mono-exponential (i.e. to overcome the influence of background reactions). In reactions of **3** with **7-9**, the observed rate constants for the decays of the vinyl cations never exceeded the rate of the reaction of the solvent with **3**. We discuss this material in greater detail in the Supporting Information on p. S40-42.
- 52 McClelland, R. A.; Kanagasabapathy, V. M.; Banait, N. S.; Steenken, S. *J. Am. Chem. Soc.* **1989**, 111, 3966-3972.
- 53 (a) Lucius, R.; Loos, R.; Mayr, H. *Angew. Chem.* **2002**, 114, 97-102; *Angew. Chem., Int. Ed.* **2002**, 41, 91-95; (b) Mayr, H. *Tetrahedron* **2015**, 71, 5095-5111.
- 54 Phan, T. B.; Breugst, M.; Mayr, H. *Angew. Chem.* **2006**, 118, 3954-3959; *Angew. Chem., Int. Ed.* **2006**, 45, 3869-3874.
- 55 Experimentally determined rate constants were used where available, otherwise  $\lg k(47)$  was calculated using equation (4), using  $E = 0.00$  (for **47**)<sup>24</sup> and the relevant  $N$  and  $s_N$  values from Table 1. Calculated rate constants for nucleophiles **17-20** exceed the diffusion limit, and hence are not included in the correlation.
- 56 Experimentally determined rate constants were used where available, otherwise  $\lg k(48)$  was calculated using equation (4), using  $E = 3.63$  (for **48**)<sup>35</sup> and the relevant  $N$  and  $s_N$  values from Table 2. Pyrrole **10** is not included in the correlation because the rate constant calculated for its reaction with **48** greatly exceeds the diffusion limit. The absence of this number undoubtedly contributes to the apparent better quality of the correlation in Figure 9 compared to Figure 8.
- 57 Liu, K.-T.; Chin, C.-P.; Lin, Y.-S.; Tsao, M.-L. *Tetrahedron Lett.* **1995**, 36, 6919-6922.
- 58 McClelland, R. A.; Kanagasabapathy, V. M.; Steenken, S. *J. Am. Chem. Soc.* **1988**, 110, 6913-6914.
- 59 (a) Hojo, M.; Ueda, T.; Ueno, E.; Hamasaki, T.; Nakano, T. *Bull. Chem. Soc. Jpn.* **2010**, 83, 401-414; (b) Hojo, M.; Aoki, S. *Bull. Chem. Soc. Jpn.* **2012**, 85, 1023-1030; (c) Manege, L. C.; Ueda, T.; Hojo, M.; Fujio, M. *J. Chem. Soc., Perkin Trans 2* **1998**, 1961-1966.
- 60 (a) See pg. 483 in reference 1d; (b) See reference 25, pp 290-447, and in particular pp 355-391; (c) See reference 26d, pp 583-594.
- 61 See reference 25, p 361, Table 6.25.
- 62 Gaussian 09, Revision D.01, Frisch, M. J.; Trucks, G. W.; Schlegel, H. B.; Scuseria, G. E.; Robb, M. A.; Cheeseman, J. R.; Scalmani, G.; Barone, V.; Mennucci, B.; Petersson, G. A.; Nakatsuji, H.; Caricato, M.; Li, X.; Hratchian, H. P.; Izmaylov, A. F.; Bloino, J.; Zheng, G.; Sonnenberg, J. L.; Hada, M.; Ehara, M.; Toyota, K.; Fukuda, R.; Hasegawa, J.; Ishida, M.; Nakajima, T.; Honda, Y.; Kitao, O.; Nakai, H.; Vreven, T.; Montgomery, J. A., Jr.; Peralta, J. E.; Ogliaro, F.; Bearpark, M.; Heyd, J. J.; Brothers, E.; Kudin, K. N.; Staroverov, V. N.; Kobayashi, R.; Normand, J.; Raghavachari, K.; Rendell, A.; Burant, J. C.; Iyengar, S. S.; Tomasi, J.; Cossi, M.; Rega, N.; Millam, M. J.; Klene, M.; Knox, J. E.; Cross, J. B.; Bakken, V.; Adamo, C.; Jaramillo, J.; Gomperts, R.; Stratmann, R. E.; Yazyev, O.; Austin, A. J.; Cammi, R.; Pomelli, C.; Ochterski,

- 
- J. W.; Martin, R. L.; Morokuma, K.; Zakrzewski, V. G.; Voth, G. A.; Salvador, P.; Dannenberg, J. J.; Dapprich, S.; Daniels, A. D.; Farkas, Ö.; Foresman, J. B.; Ortiz, J. V.; Cioslowski, J.; Fox, D. J. Gaussian, Inc., Wallingford CT, 2009.
- 63 See Supporting Information p. S54-S59 for details of the calculations.
- 64 (a) Becke, A. D. *J. Chem. Phys.* **1993**, *98*, 5648-5652; (b) Lee, C.; Yang, W.; Parr, R. G. *Phys. Rev. B* **1988**, *37*, 785–789.
- 65 As no barrier between the vinyl bromide **4-Br** and the ion pair **{2}**Br<sup>−</sup> is found using the PCM method (see Fig. 11), the  $\Delta G^\circ$  values given in entries 6 and 8 correspond to solvolysis half-lives on the micro- and millisecond time scale, in sharp contrast to the experimental finding that **4-Br** solvolyzes with a half-life of 1 year in 80 % aqueous ethanol at 25 °C (Scheme 4).
- 66 Tao, J. M.; Perdew, J. P.; Staroverov, V. N.; Scuseria, G. E. *Phys. Rev. Lett.* **2003**, *91*, 146401.
- 67 (a) Grimme, S.; Antony, J.; Ehrlich, S.; Krieg, H. *J. Chem. Phys.* **2010**, *132*, 154104; (b) Grimme, S.; Hansen, A.; Brandenburg, J. G.; Bannwarth, C. *Chem. Rev.* **2016**, *116*, 5105-5154.
- 68 Tomasi, J.; Mennucci, B.; Cammi, R. *Chem. Rev.* **2005**, *105*, 2999 – 3093.
- 69 Plata, R. E.; Singleton, D. A. *J. Am. Chem. Soc.* **2015**, *137*, 3811 – 3826.
- 70 See Supporting Information p. S60-75 for details of the calculations of the hydride affinities and analysis of the hydrido-bridged species at different levels of theory.
- 71 For previous experimental and theoretical investigations of hydride affinities of carbocations see refs 4 and 5.