

Title	Trial of feasibility and acceptability of routine low-dose aspirin versus early screening test indicated aspirin for pre-eclampsia prevention (TEST study): a multicentre randomised controlled trial		
Authors	Mone, Fionnuala;Mulcahy, Cecilia;McParland, Peter;Breathnach, Fionnuala M.;Downey, Paul;McCormack, Dorothy;Culliton, Marie;Stanton, Alice;Cody, Fiona;Morrison, John J.;Daly, Sean;Higgins, John R.;Cotter, Amanda;Hunter, Alyson;Tully, Elizabeth C.;Dicker, Patrick;Alfirevic, Zarko;Malone, Fergal D.;McAuliffe, Fionnuala M.		
Publication date	2018		
Original Citation	Mone, F., Mulcahy, C., McParland, P., Breathnach, F., Downey, P., McCormack, D., Culliton, M., Stanton, A., Cody, F., Morrison, J. J., Daly, S., Higgins, J., Cotter, A., Hunter, A., Tully, E. C., Dicker, P., Alfirevic, Z., Malone, F. D. and McAuliffe, F. M. (2018) 'Trial of feasibility and acceptability of routine low-dose aspirin versus Early Screening Test indicated aspirin for pre- eclampsia prevention (TEST study): a multicentre randomised controlled trial', BMJ Open, 8(7), e022056 (9pp). doi: 10.1136/ bmjopen-2018-022056		
Type of publication	Article (peer-reviewed)		
Link to publisher's version	https://bmjopen.bmj.com/content/8/7/e022056 - 10.1136/ bmjopen-2018-022056		
Rights	© 2018, the Author(s) (or their employer(s). This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non- commercial. See: http://creativecommons.org/licenses/by- nc/4.0/ http://creativecommons.org/licenses/by-nc/4.0/		
Download date	2025-09-08 12:04:16		



University College Cork, Ireland Coláiste na hOllscoile Corcaigh

Outcome	Screen positive; Aspirin Group 3A N=13	Screen negative; No Aspirin Group 3B N=171	OR (95% CI)
Preeclampsia No. (%) Pre-eclampsia <34- weeks	2 (15.4) 0 (0)	6 (3.5) 2 (1.2)	5.0 (0.9 – 27.7)
Pre-eclampsia <37- weeks	2 (15.4)	2 (1.2)	15.4 (2.0 – 120)
Birthweight <10 <sup>th</sup> centile No. (%)	4 (30.7)	21 (12.3)	3.2 (0.9 – 11.2)
Pre-term delivery <34 weeks No. (%)	1 (7.7)	1 (0.6)	14.1 (0.8 – 240)
NICU admission No. (%)	0 (0)	9 (5.3)	-
Outcome No. (%) Alive at 6-weeks	13 (100)	169 (98.8)	
Stillbirth Neonatal death	$ \begin{array}{c} 0 (0) \\ 0 (0) \end{array} $	2 (1.2) 0 (0)	

Table S2 - Secondary outcome measures in Group 3 (screen and treat)

(Expressed as average and standard deviation unless otherwise stated)

Note: ORs are not presented when number of events is 0 in the Screen –positive group.