

1    **Evaluation of production of Cheddar cheese from micellar**  
2    **casein concentrate**

3

4    Bozhao Li<sup>1</sup>, David S. Waldron<sup>1</sup>, John T. Tobin<sup>2</sup>, Surabhi Subhir<sup>2</sup>, Alan L. Kelly<sup>1\*</sup>, Paul L. H.  
5    McSweeney<sup>1</sup>

6    *<sup>1</sup>School of Food and Nutritional Sciences, University College Cork, College Road, Cork, T12*  
7    *K8AF, Ireland.*

8    *<sup>2</sup>Department of Food Chemistry and Technology, Teagasc Food Research Centre, Fermoy, Co.*  
9    *Cork, P61 C996, Ireland.*

10

11    \*Corresponding author: Tel: +353-21-4903405

12    Email address: [a.kelly@ucc.ie](mailto:a.kelly@ucc.ie)

13

14    **ABSTRACT**

15    The objective of this study was to evaluate the production of Cheddar cheese using  
16    micellar casein concentrate (MCC), a novel milk ingredient powder with a high casein  
17    content (~92%). Four types of Cheddar cheese were manufactured and ripened for 180  
18    days from the following starting materials: standardized control milk (control), skim  
19    milk with cream (SC), reconstituted MCC with cream (MC) and reconstituted low-heat  
20    skim milk powder with cream (PC). Only minor differences were found in composition  
21    between treatments, but MC cheese showed higher levels of proteolysis compared to  
22    other treatments, linked to significantly higher plasmin and chymosin activities. No  
23    differences were observed in hardness between treatments (60, 120 and 180 days), but  
24    the springiness and cohesiveness of MC and PC cheeses were significantly higher than  
25    that of the control and SC cheeses at 60, 120 and 180 days. In conclusion, the use of  
26    casein-dominant dairy streams has the potential for production of Cheddar cheese with  
27    tailored functionality.

## 1. Introduction

Global milk production was approximately 843 million tonnes in 2018 (FAO, 2019); among dairy products (i.e., cheese, casein and butter), cheese production used the highest proportion of milk. Whey protein powder, production of which is traditionally associated with the manufacture of cheese, represents 57% of the market for global protein supplement for exercising and nutrition, while there is a huge market for whey protein powders in infant formula. Cheese output is increasing at a rate of 2% yearly while the demand for whey protein has been growing at 6-7% yearly (Hoogwegt Group, 2019). Nowadays, whey protein with high quality is required as it can be used to manufacture a range of food ingredients or products with nutrition and functional properties (Boland, 2011). Kelly (2019) reported that the ultrafiltration (UF) properties of sweet and acid whey are influenced by their high or low pH, respectively. However, microfiltration (MF) permeate made directly from skim milk is considered to be an ideal whey source for whey protein ingredients production. Therefore, recovering whey protein from milk rather than cheese whey could improve the whey quality and is an option for whey protein ingredients manufacture.

During membrane filtration of skim milk, native micellar casein is concentrated in the retentate, which could be recovered and concentrated to produce micellar casein concentrate (MCC). This is a novel dairy ingredient powder with a high casein fraction of 85-95% of total protein which may be used in functional and nutritional

applications (Crowley et al., 2018). As the traditional way to manufacture Cheddar cheese is followed by whey protein manufacture, the concept of recovering whey protein before Cheddar cheese manufacture is of interest.

In terms of the protein ingredients being used for Cheddar cheese manufacture, low heat skim milk powder (LHSMP) can be used to enhance cheese yield giving a constant cheese production throughout the year (Freeman et al., 1970). LHSMP is produced from skim milk using low temperatures during manufacture and is mostly used for condensed milk, UHT-treated fluid milk and ice-cream (Augustin & Margetts, 2003). Unlike reconstituted medium- and high-heat skim milk powder that have impaired rennet coagulation ability, the rennet coagulation properties of LHSMP are good as the whey protein is not highly denatured (Ménard et al., 2005). With only physical separation processing, the micellar casein in MCC may have better rennet coagulation properties compared to LHSMP. The rennet coagulation properties and cheese-making potential of MCC may be better than those of LHSMP.

Two main factors that influence cheese yield are lactation and seasonality. The gross composition of bovine milk varies with the stage of lactation. Kuchlík et al. (2008) reported that protein and casein content increase and lactose content decreases through the lactation. After 200 days lactation, cows are in the late lactation stage, which requires the udder tissue to repair and recover for next lactation. During late lactation, milk yield decreases dramatically, which may influence cheese yield, unless milk is standardised. Also, at specific times of the year, the milk volume decreases, and the composition and rennet coagulation properties of milk are

significantly influenced (Freeman et al., 1970; O'Brien et al., 1999). Both cheese yield and manufacturing efficiency are influenced by seasonal variation of milk protein and fat composition (Barbano & Sherbon, 1984).

To solve the problem of low cheese yield, one solution is adding LHSMP to low-protein milk to increase the protein composition (Freeman et al., 1970). Another possibility may be making Cheddar cheese with MCC that is manufactured as a co-product of high-quality whey protein.

This study aimed to evaluate the production of Cheddar cheese from micellar casein concentrate, with LHSMP used for comparison. The consequence of this manufacture was evaluated concerning composition, proteolysis, texture and functionality.

## **2 Materials and Methods**

### *2.1 Preparation of cheese milk*

Micellar casein concentrate (MCC) powder was obtained from Teagasc (Moorepark, Fermoy Co. Cork, Ireland); whereby pasteurized bovine skim milk was microfiltered at 50 °C using 0.14-µm TAMI Isoflux ceramic membranes (TAMI, Lyons, France). MF was performed in a batch mode involving two diafiltration steps to produce a final 3X MF retentate (liquid MCC) which was evaporated at 65 °C using a pilot plant single-effect falling-film evaporator (Anhydro F1 Laboratory;

Copenhagen, Denmark). This was followed by spray-drying of the evaporated liquid MCC in a single-stage spray dryer (Anhydro Laboratory Spray Dryer, SPX Flow Technology, Denmark) equipped with nozzle atomisation at inlet and outlet temperatures of 185 and 85°C, respectively. The protein content (91.6% casein) of MCC was determined by the Kjeldahl method (IDF, 1986). MCC powder and low heat skim milk powder (LHSMP; WPNI=6.0) (Uelzena, Uelzen, Germany) were rehydrated using a Silverson mixer at 55 °C for 2 h.

Raw milk was obtained from a local farm in Cork, Ireland. Skim milk and cream were separated from raw milk using a separator. Composition of milk samples was measured by MilkoScan™ Mars (Foss, Hilleroed, Denmark) and casein content was calculated by multiplying the casein percentage of protein of the milk sample (an estimated percentage of 78% for milk and LHSMP milk and 91.6% for MCC). Control milk was prepared by combining raw milk and skim milk to a casein: fat ratio of 0.7:1. Both reconstituted MCC and reconstituted LHSMP were made up to the same casein level as skim milk. Reconstituted MCC and LHSMP and skim milk were blended with cream and lactose according to the same standardization ratio (casein: fat of 0.70:1.00) and lactose content (~5%) as control milk to obtain three more vats: skim milk with cream (SC), reconstituted MCC milk with cream (MC) and reconstituted LHSMP milk with cream (PC). All milk samples were pasteurized at 72 °C for 15 s (Microthermics Inc., Raleigh, NC, USA) and stored at 4 °C until analysis and cheese-making.

## 2.2 Rennet Coagulation Properties

Dynamic oscillatory analysis (small amplitude oscillatory measurement) of renneted control milk, SC milk, MC milk and PC milk was performed using a Peltier concentric cylinder geometry, which comprised of an aluminium conical rotor [42.01 mm (h) by 28.02 mm (d)], on an AR-G2 controlled stress rheometer (Waters TA Instruments, Leatherhead, Surrey, UK). Aliquots (25 mL) of each sample were pre-warmed in a water bath at 32°C for 15 min, and 80 µL of a 1:10 (v/v) dilution of Maxiren™ (Chr Hansen A/S, Hørsholm, Denmark) was added. The sample was placed immediately in the preheated cup (32 °C), the frequency of oscillation was set at 0.6283 rad s<sup>-1</sup>, and the storage modulus, G' of the sample was recorded continuously as a function of time at a low-amplitude shear strain (0.01) over 90 min. Each sample was analysed in triplicate (Ibáñez et al., 2015).

## 2.3 Cheese Manufacture

Cheddar cheeses were made from 20 L control, SC, MC and PC milks which were prepared, pasteurised and stored at 4 °C overnight before cheese-making. Cheddar-type cheeses were manufactured according to a standard Cheddar cheese manufacture protocol (Fox et al., 2000). An aliquot of 6 g (0.03% w/w) of Cheddar cheese starter culture (R604Y Chr. Hansen Ltd., Little Island, Co. Cork, Ireland) was added into each sample and allowed ripen for 30 min, followed by 0.09% (v/w) of 1 mol L<sup>-1</sup> CaCl<sub>2</sub> and 60 IMCU L<sup>-1</sup> rennet being added to all samples. Whey was drained when the pH dropped to 6.2 and curd was cheddared. When the pH decreased to 5.4,

the curd was milled into small pieces and salted at a level of 2.5% (w/w) NaCl. The curds were then wrapped in cheesecloth and moved to 2-kg circular moulds, which were pressed at a pressure of 2.5 kg cm<sup>-2</sup> for 14 h. The cheeses were then removed, vacuum-packed and ripened at 8 °C for 180 days.

#### *2.4 Compositional analysis*

Gross composition of cheeses was analysed at 14 days old. Moisture was determined by oven-drying method (IDF 1982), protein (%N×6.38) by the Kjeldahl method (IDF 1986) and salt was analysed by titration with AgNO<sub>3</sub> (Fox, 1963). The pH was measured at 14, 30, 60, 120 and 180 days of ripening by using a calibrated pH probe placed in contact with dry grated cheeses. All results were determined in triplicate.

#### *2.5 Microbiological analysis*

Enumeration of starter lactic acid bacteria (LAB) was performed using LM 17 agar plates (Terzaghi and Sandine 1975), incubated for 3 days at 30 °C. Enumeration of non-starter lactic acid bacteria (NSLAB) was performed on Rogosa agar plates (Rogosa and Mitchell 1951), incubated anaerobically for 5 days at 30 °C. Enumeration of LAB and NSLAB were performed in duplicate after 60 and 90 days of ripening.



## 2.6 Analysis of proteolysis

Urea-polyacrylamide gel electrophoresis (urea-PAGE) of cheese samples was used to study the proteolysis of  $\alpha_{S1}$ - and  $\beta$ -casein (CN) during ripening using the method of Andrews (1983) with modifications of Shalabi and Fox (1987). The pH 4.6-soluble and insoluble fraction were prepared (Kuchroo & Fox, 1982), and peptide profiles of pH 4.6-soluble fractions filtered through 0.22- $\mu$ m cellulose acetate filters (Sartorius GmbH, Gottingen, Germany) were analysed by reverse-phase high-performance liquid chromatography (HPLC) using an ultra-performance liquid chromatography (UPLC) Waters Acquity UPLC H-Class Core System (Waters, Milford, MA, USA), with a Waters Acquity UPLC TUV Detector (dual-wavelength; Waters) operated by Empower 3 software (Waters Corp., Milford, MA, USA), following the method of Mane et al. (2019).

For determination of free amino acids (FAA), frozen pH 4.6-soluble fractions were de-proteinised by mixing equal volumes of 24% (w v<sup>-1</sup>) trichloroacetic acid (TCA) and sample and following the method of Fenelon and Guinee (2000).

## 2.7 Plasmin activity

The plasmin activity of cheese samples at 180 days of ripening was measured using the coumarin peptide method (Richardson & Pearce, 1981). A standard curve of the emission intensity at 460 nm was constructed using 7-amido-4-methyl coumarin (AMC), and results expressed in nmol AMC min<sup>-1</sup> mL<sup>-1</sup>, which was defined as one unit of plasmin activity.

## 174 2.8 Residual coagulant assay

175 Grated cheese samples (50 mg) were extracted by dissolving cheese in 1 mL of  
176 0.1 M trisodium citrate, followed by 30 minutes incubation at 37 °C. Fat was  
177 separated by centrifugation at 1000 g (Sigma 1-16K, Harz, Germany) for 1 min, and  
178 the aqueous layer was used for analysis. An aliquot of 70 µL citrate dispersion of  
179 cheese was incubated with 30 µL of 1 mg mL<sup>-1</sup> aqueous solution of a synthetic  
180 heptapeptide substrate (Pro-Thr-Glu-Phe-[NO<sub>2</sub>-Phe]-Arg-Leu) in 400 µL 0.1 M  
181 sodium formate buffer, at 37 °C, pH 3.2, for 24 h. The mixture was heated at 70 °C  
182 for 10 min to stop the reaction, followed by centrifugation at 16,000g for 10 min, and  
183 the supernatant was used for UPLC analysis. Substrate and product levels were  
184 determined using the reversed-phase HPLC system described above, following the  
185 method of Hurley et al. (1999).

## 186 2.9 Texture profile analysis

187 Texture profile analysis was performed using a Texture Analyzer TA-XT2i  
188 (Stable Micro Systems, Godalming, Surrey, UK) at 60, 120, and 180 days of ripening.  
189 Cheese samples were cut into 20 mm height, 20 mm diameter cylinders and kept at  
190 4 °C overnight. Cheese cylinders were compressed to 25% of the initial height in two  
191 continuous compressions with a speed of 1 mm s<sup>-1</sup>. Hardness, cohesiveness and  
192 springiness were measured (Truong et al., 2002), and five cheese samples were  
193 measured for each treatment.

194

## 2.10 Meltability

Meltability was analyzed using the Schreiber meltability test as described by Altan et al. (2005). Cheese samples were heated at 232 °C for 5 min, and meltability was measured as the percentage increase in diameter of the original samples. Analyses were performed in triplicate at 60, 120, 180 days of ripening.

## 2.11 Dynamic small amplitude oscillatory rheology

The rheological properties of meltability of cheese after 180 days ripening were analysed with a controlled stress AR-G2 rheometer (TA Instruments, Waters LLC, Leatherhead, Surrey, UK) according to the method of Ibáñez et al. (2015). Storage modulus ( $G'$ ), loss modulus ( $G''$ ), and loss tangent (LT) were measured during heating. The maximum LT ( $LT_{max}$ ) and the temperature where  $LT=1$ , which are indicators of melting, were also recorded. Each sample was analysed in triplicate.

## 2.12 Colour measurements

Colour values were measured using a Konika-Minolta colourimeter CR400 (Konika-Minolta Optics Inc., Osaka, Japan) at 14, 30, 60, 120, 180 days of ripening. The measurement used the CIELAB system based on illuminant D65 and a visual angle of 2°. Five random readings were taken on fresh-cut cheese at 20°C. The Euclidean distance between the colour of control cheese and that for other treatments was calculated by  $\Delta E^*_{ab} = \sqrt{(\Delta L^*)^2 + (\Delta a^*)^2 + (\Delta b^*)^2}$  where  $\Delta L^* = L^*_{sample} - L^*_{control}$ ,  $\Delta a^* = a^*_{sample} - a^*_{control}$  and  $\Delta b^* = b^*_{sample} - b^*_{control}$  (Mahy et al., 1994).

### 2.13 Statistical analysis

Control, SC, MC and PC cheeses were made in three independent trials,  $4 \times 3$  blocks. Statistical analysis was carried out using one-way analysis of variance (ANOVA) with R project for Windows version i386 3.4.0 to establish significant differences between samples. The probability level used for statistical significance was  $P < 0.05$ .

## 3 Results and Discussion

### 3.1 Rennet coagulation properties

The rennet coagulation properties of milk samples are shown in Table 1. During the first 10 min,  $G'$  value of all the treatments was low and constant, after which the  $G'$  value of all samples increased, with the MC and PC samples exhibiting a slower rate of increase compared to other samples (Fig. 1). The gel strength ( $G'$  value) at 90 min of MC and PC samples was significantly ( $P < 0.05$ ) lower than that of other samples. The gel strength of MC and PC kept increasing and did not reach a constant level during the analysis, while control and SC reached stable values at 45 min. The loss tangent at 90 min of MC sample was significantly ( $P < 0.05$ ) higher than that of the other treatments.

Martin et al. (2010) reported that, during the manufacture of milk protein concentrate (MPC), casein micelles were not damaged; however, the minerals

removed through diafiltration and decreased calcium ionic strength could depress rennet coagulation. The rennet coagulation ability of reconstituted MPC could be restored by adding extra calcium ions. The production of MCC involves two diafiltration steps, which may reduce soluble calcium levels; thus, 0.09% (v/w) of 1 mol L<sup>-1</sup> CaCl<sub>2</sub> was added to all of the batches before cheese manufacture.

### 3.2 Cheese composition and pH

The composition of cheese at 14 days of ripening and pH of cheese samples at 14, 30, 60, 120 and 180 days of ripening are shown in Table 2; the results presented are means of data from the three trials. Guinee et al. (2007) stated that different casein to fat ratio can influence the protein and fat content of cheese. As the casein: fat ratio of milk was adjusted to 0.7 for all samples, no statistically significant differences ( $P>0.05$ ) were found in cheese protein and fat contents between all treatments, and the use of MCC or LHSMP for Cheddar cheese manufacture did not affect these parameters.

The moisture content of PC cheese was significantly ( $P<0.05$ ) higher than that of SC and MC cheese; no significant differences ( $P>0.05$ ) were found for moisture content between control, SC and MC cheese. The salt content of MC cheese was significantly ( $P<0.05$ ) higher than that of control and SC cheese. The pH of MC cheese was significantly ( $P<0.05$ ) higher than that of other treatments, the reason for which is not clear. No significant differences ( $P>0.05$ ) were found for moisture in non-fat substances (MNFS) and fat in dry matter (FDM) for all treatments; MNFS is an

important parameter for Cheddar cheese quality, so it is important that all batches of cheese sample had similar levels of MNFS (Fox et al., 2000; Bogenrief & Olson, 1995).

### 3.3 Proteolysis

Table 2 shows the level of pH 4.6-SN/TN (%) of control, SC, MC and PC cheeses at 180 days of ripening; pH 4.6-SN/TN is an index of proteolysis (Sousa et al., 2001). Significantly ( $P<0.05$ ) higher pH 4.6-SN/TN levels were found in MC cheese than in SC and PC cheese. In addition, the levels of pH 4.6-SN/TN were significantly ( $P<0.05$ ) lower in PC than control cheese. The value (23%) found at 180 days of ripening in this study was comparable with previous studies on Cheddar cheese (Lucey et al., 2005; O'Mahony et al., 2005). The main agent producing pH 4.6-soluble nitrogen is the coagulant (O'Keeffe et al., 1978), while Farkye & Fox (1992) reported that plasmin, the principal indigenous milk proteinase, is also important for primary proteolysis in Cheddar cheese during ripening.

Urea-PAGE electrophoretograms of cheese samples during ripening are shown in Fig. 2. All cheese samples showed break-down of  $\beta$ - and  $\alpha_{S1}$ -CN, but lower levels of intact  $\beta$ - and  $\alpha_{S1}$ -CN were apparent during ripening in MC cheese. Due to the action of plasmin,  $\beta$ -CN is hydrolysed to  $\beta$ -CN (f29-209),  $\beta$ -CN (f106-209) and  $\beta$ -CN (f108-209) (Eigel et al., 1984). The levels of  $\beta$ -CN (f106-209) and  $\beta$ -CN (f108-209) of MC cheese were higher than other treatments during ripening, which suggests higher plasmin activity. Also, in Fig. 2, the MC cheese showed the lowest level of intact  $\alpha_{S1}$ -CN level,

277 followed by PC cheese, control and SC cheese from 60 to 180 days of ripening; at 180  
278 days of ripening, no intact  $\alpha_{S1}$ -CN was found in MC and PC samples. Residual  
279 chymosin in cheese hydrolyses  $\alpha_{S1}$ -CN to  $\alpha_{S1}$ -CN (f24-199) and  $\alpha_{S1}$ -CN (f102-199) and,  
280 as the residual chymosin activity increases, the breakdown of  $\alpha_{S1}$ -CN increases  
281 (Sheehan et al., 2008). The faster breakdown of  $\alpha_{S1}$ -CN of MC and PC samples may  
282 indicate higher residual chymosin activity in these samples.

283 The peptide profiles of the pH 4.6-soluble extracts for control, SC, MC and PC  
284 cheeses at 180 days of ripening were generated by ultra-performance liquid  
285 chromatography (Fig. 3). The highest peak areas were observed at the retention time of  
286 1 to 5 min for all treatments. The peak area of peptides that were eluted later (28-45  
287 min retention time) was higher in MC sample compared to the other treatments.  
288 Peptides in the pH 4.6-soluble extracts reflect the effect of proteinases and peptidases  
289 of starter (Fox & McSweeney, 1997). The number of peptide peaks for all treatments  
290 was similar, which suggests that the proteolysis for all treatments broadly followed  
291 similar pathways, but differences in peak area indicate that there was more extensive  
292 proteolysis in MC cheese compared to the other treatments.

293 The individual FAA levels of all treatments at 120 and 180 days of ripening are  
294 shown in Fig. 4 (A and B). The major FAA determined in cheese were glutamic acid,  
295 valine, leucine, phenylalanine, histidine and lysine; Bansal et al. (2009) also reported  
296 that glutamic acid, valine, leucine, phenylalanine histidine and lysine are the principal  
297 FAA in Cheddar cheese after 180 days of ripening. No significant differences ( $P>0.05$ )  
298 were found between treatments at 120 days of ripening. At 180 days of ripening,

significantly ( $P<0.05$ ) higher concentrations of threonine, serine, glycine, alanine, valine, methionine, isoleucine, phenylalanine, histidine and proline were found in control and SC cheese than that of MC and PC cheese. No significant differences ( $P>0.05$ ) were found between treatments in the levels of aspartic acid, glutamic acid, leucine, tyrosine and arginine. From 120 to 180 days of ripening, the concentrations of glutamic acid, alanine, leucine, tyrosine and phenylalanine increased significantly ( $P<0.05$ ) for all treatments, while only control and SC cheese had a significant increase in the concentration of threonine, glycine, valine and isoleucine. The total FAA levels of all treatments at 120 and 180 days of ripening are shown in Fig. 4 (C). No significant differences ( $P>0.05$ ) were found between all treatments at 120 days of ripening, but significantly higher ( $P<0.05$ ) levels of total FAA were found in control cheese than in MC and PC cheeses at 180 days of ripening. No significant differences ( $P>0.05$ ) were found between control and SC cheese at 180 days of ripening.

Fox and McSweeney (1996) reported that peptidases of starter and non-starter lactic acid bacteria are the principal agents releasing FAA in Cheddar cheese during ripening. No significant ( $P>0.05$ ) differences were found for the numbers of starter bacteria and NSLAB between treatments in this study (result not shown). Therefore, the release of the FAA by starter and NSLAB should not have been an influence in this regard, and so the reasons for this difference is not clear.



#### 320 3.4 Plasmin and residual chymosin activities in cheese

321 The plasmin activity of MC cheese was significantly ( $P<0.05$ ) higher than that of the  
322 other cheese batches (Table 3) which is consistent with the result of urea PAGE  
323 electrophoresis. The plasmin activity of PC cheese was significantly ( $P<0.05$ ) lower  
324 than that of the other cheese samples. As the pH increases, the hydrolysis of  $\beta$ -CN  
325 increases since plasmin has an alkaline optimum pH (Watkinson et al., 2001).

326 Aaltonen and Ollikainen (2011) reported that in diafiltration, with the removal of  
327 whey protein, the concentration of inhibitors of both plasmin and plasminogen  
328 activators decreases, which promotes the conversion from plasminogen to plasmin and  
329 thus increases plasmin activity. In this study, MCC was made using microfiltration,  
330 which may have enhanced the plasmin activity of retentate by removing inhibitors in  
331 the permeate, resulting in the higher plasmin activity and more  $\beta$ -CN breakdown. MC  
332 cheese showed significantly ( $P<0.05$ ) higher residual chymosin activity compared to  
333 the other treatments (Table 3), which is consistent with the faster breakdown of  $\alpha_{S1}$ -CN  
334 in that cheese. No significant differences ( $P>0.05$ ) were found between the residual  
335 chymosin activities of control and SC cheese, so the recombination of fat and skim milk  
336 did not affect the retention of coagulant during cheese manufacture. The residual  
337 chymosin activity of PC cheese was significantly ( $P<0.05$ ) lower than that of control  
338 cheese. The MC cheese apparently retained more coagulant compared to control, SC  
339 and PC cheeses; the percentage of retained chymosin in cheese curd depends on the pH

at the curd-cutting stage, pH at whey drainage, cooking temperature and method (Hurley et al., 1999).

### 3.5 Texture profile analysis

No significant changes ( $P>0.05$ ) were found in the hardness of MC and PC cheeses from 60 to 180 days of ripening (Fig. 5). The hardness of control cheese significantly ( $P<0.05$ ) decreased between 60 and 120 days of ripening, while that of SC cheese significantly ( $P<0.05$ ) increased. At 60 days of ripening, the hardness of control cheese was significantly ( $P<0.05$ ) higher than that of SC and PC cheeses. No significant differences ( $P>0.05$ ) were observed in the hardness of the cheeses made by different treatments at 120 and 180 days of ripening. Chevanan and Muthukumarappan (2007) reported that the contents of calcium, phosphate, and residual lactose affect the texture profile of Cheddar cheese. The cheese-milk of SC, MC and PC samples was reconstituted according to the composition of control cheese-milk, which would not influence lactose content. In this study, extra calcium chloride was added to all treatments, but the addition of calcium chloride does not cause significant changes to the texture of ripened Cheddar cheese (Soodam et al., 2015).

From 60 days to 180 days of ripening, the springiness and cohesiveness of MC cheese decreased, and the springiness and cohesiveness of MC and PC cheeses were significantly ( $P<0.05$ ) higher than that of control and SC cheese at 60 and 180 days of ripening. At 120 days of ripening, the springiness of MC cheese was significantly

( $P < 0.05$ ) higher than that of the other treatments and the cohesiveness of MC and PC cheeses was significantly ( $P < 0.05$ ) higher than that of control and SC cheese. Everard et al. (2006) found that higher pH of Cheddar cheese was associated with increased springiness and cohesiveness; the pH of MC cheese was significantly ( $P < 0.05$ ) higher than that for the other treatments (Table 2), which may explain the increases in springiness and cohesiveness. O'Mahony et al. (2005) reported that, as levels of secondary proteolysis increase, the charged groups released from peptides would associate with free water, which may lead to increased cohesiveness and springiness during ripening.

### 3.6 Meltability

Meltability (percentage increase in diameter) of Cheddar cheeses was determined by the Schreiber method; results are shown in Fig. 6. The four types of cheeses showed increases in meltability between 60 and 180 days. At 60 and 120 days of ripening, no significant differences ( $P > 0.05$ ) for meltability were found between control and SC cheese. Significantly ( $P < 0.05$ ) higher meltability was found in control cheese compared to SC cheese. At 60, 120 and 180 days of ripening, the meltability of MC cheese was significantly higher ( $P < 0.05$ ) than that of cheese from the other treatments. At 120 and 180 days of ripening, the meltability of PC cheese was significantly lower ( $P < 0.05$ ) than that of cheese from the other treatments.

The lower meltability of PC cheese may be caused by the changes of casein in protein during drying of milk powder. Moiseev et al. (2017) reported that Mozzarella

cheese made with reconstituted non-fat milk powder has lower meltability than the control, due to the drying process of non-fat milk powder decreasing the stability and dispersity of casein micelles and promoting demineralization of calcium salts. Dave et al. (2003) found that the meltability of Mozzarella cheese is related to the breakdown of ( $\alpha_{S1}$ -CN and  $\alpha_{S1}$ -CN (f24-199)) and breakdown of intact  $\beta$ -casein, while Bogenrief and Olson (1995) reported that hydrolysis of  $\beta$ -casein increases the meltability of Cheddar cheese. Hydrolysis of  $\beta$ -casein is primarily related to the action of plasmin (Eigel et al., 1984) and, although both MC and PC cheeses were made from powder, and showed a higher breakdown of  $\alpha_{S1}$ -casein and  $\beta$ -casein, the meltability of MC cheese was significantly higher than that of PC.

### *3.7 Dynamic small amplitude oscillatory rheology*

The results of the rheology of meltability of all treatments at 180 days of ripening are shown in Table 4. Higher LT value indicates a higher extent of melting (Lucey et al., 2003). MC cheese showed significantly higher ( $P<0.05$ )  $LT_{max}$  values than PC cheese, followed by control and SC cheese. The temperature at  $LT_{max}$  of MC and PC cheeses was significantly higher ( $P<0.05$ ) than that of control and SC cheese. The  $LT_{max}$  is an indicator of cheese meltability (Mounsey & O’Riordan, 1999); the more thermal energy needed to melt cheese, the higher the  $LT_{max}$  temperature of cheese will be.

In the Schreiber test for cheese meltability, MC cheese also exhibited the highest percentage increase in diameter (Fig. 4). There were differences between

Schreiber melting test and dynamic small amplitude rheology for control, SC and PC cheese. Cooke et al. (2013) stated that differences between results generated by those tests may be linked to the more complete fat melting during the higher temperature of the Schreiber melting test. MC cheese exhibited the lowest temperature when LT=1 compared to the rest of the treatments ( $P<0.05$ ). At the point of LT=1, cheese is considered to start transforming from a solid to a viscous form (Gunasekaran & Ak, 2002). MC cheese thus commenced melting at the lowest temperature and had the highest meltability compared to the other treatments, which may be linked to the more extensive proteolysis of casein.

### *3.8 Cheese colour*

The colour values of samples at 180 days of ripening are shown in Table 5. Whiteness ( $L^*$  values) of control, SC cheese was significantly higher ( $P<0.05$ ) than that of MC cheese at 180 days of ripening. The greenness of MC and PC cheeses was significantly lower (higher  $a^*$  values) ( $P<0.05$ ) than that of control or SC cheese at 180 days of ripening. The yellowness ( $b^*$  values) of MC cheese was significantly lower ( $P<0.05$ ) than that of SC cheese at 180 days of ripening. Ibáñez et al. (2015) reported that the whiteness of cheese may be related to proteolysis; lower whiteness of MC cheese may thus be attributed to higher proteolysis. The  $\Delta E^*_{ab}$  values are also shown in Table 5. Sharma (2003) reported that  $\Delta E^*_{ab}$  of above 2.3 leads to a just noticeable difference (JND) and, on this basis, overall, no JND was found between

control cheese and the other treatments. Overall, no visible difference was found between the cheeses made from protein ingredient and control and SC cheese.

## Conclusions

Cheddar cheeses were made from standardised cheese milk, skim milk with cream, reconstituted MCC with cream and reconstituted LHSMP with cream according to the same casein, fat and lactose content. Only minor differences in composition were found with cheese made from reconstituted MCC with added cream compared to control cheese. The cheese manufactured with MCC had significantly higher ( $P<0.05$ ) levels of pH 4.6-SN/TN than cheese manufactured with skim milk or LHSMP with cream. The level of intact  $\beta$ - and  $\alpha_{S1}$ -CN of MC cheese was lower than rest treatments, consistent with significantly ( $P<0.05$ ) higher plasmin and chymosin activity. No significant difference ( $P>0.05$ ) were found in hardness between all treatments. Significantly higher ( $P<0.05$ ) springiness and cohesiveness were found in cheese manufactured from MCC and LHSMP powder, meltability and maximum loss tangent in MC cheese were significantly ( $P<0.05$ ) higher than that of the other treatments. No overall differences of colour were found between all treatments. Use of novel casein-dominant dairy streams such as MCC has potential for production of Cheddar cheese with tailored functionality. The results of this study suggest that using reconstituted LHSMP with cream for the manufacture of Cheddar cheese may result in changes in

443 functionality, while using reconstituted MCC with cream for the manufacture of  
444 Cheddar cheese may be more feasible.

445

446

#### 447 **Acknowledgements**

448 The authors gratefully acknowledge Department of Agriculture, Food and the  
449 Marine, Ireland and Teagasc, Ireland for providing the samples of MCC for Cheddar  
450 cheese manufacture.

451   **References**

- 452   Aaltonen, T., & Ollikainen, P. (2011). Effect of microfiltration of milk on plasmin  
453   activity. *International Dairy Journal*, 21, 193-197.
- 454   Altan, A., Turhan, M., & Gunasekaran, S. (2005). Short communication: comparison  
455   of covered and uncovered Schreiber test for cheese meltability evaluation. *Journal*  
456   *of Dairy Science*, 88, 857-861.
- 457   Andrews, A. T. (1983). Proteinases in normal bovine milk and their action on  
458   caseins. *Journal of Dairy Research*, 50, 45-55.
- 459   Augustin, M. A., & Margetts, C. L. (2003). Powdered milk | Milk Powders in the  
460   Marketplace. In B. Caballero, P. Finglas & F. Toldra (Eds.), *Encyclopedia of food*  
461   *sciences and nutrition* (2nd edn) (pp. 4694–4702). London, UK: Academic.
- 462   Bansal, N., Drake, M. A., Piraino, P., Broe, M. L., Harboe, M., Fox, P. F., &  
463   McSweeney, P. L. H. (2009). Suitability of recombinant camel (*Camelus*  
464   *dromedarius*) chymosin as a coagulant for Cheddar cheese. *International Dairy*  
465   *Journal*, 19, 510-517.
- 466   Barbano, D. M., & Sherbon, J. W. (1984). Cheddar cheese yields in New  
467   York. *Journal of Dairy Science*, 67, 1873-1883.
- 468   Bogenrief, D. D., and Olson, N. F. (1995). Hydrolysis of  $\beta$ -casein increases Cheddar  
469   cheese meltability. *Milchwissenschaft*, 50, 678-682.
- 470   Boland, M. (2011). Whey proteins. In G. O. Phillips & P. A. Williams (Eds.),  
471   *Handbook of Food Proteins* (pp. 30–55). Cambridge, UK: Woodhead Publishing.
- 472   Bourne, M. C. (1978). Texture profile analysis. *Food Technology*, 32, 62–66,72.



473 Chevanan, N., & Muthukumarappan, K. (2007). Effect of calcium and phosphorus,  
474 residual lactose, and salt-to-moisture ratio on the melting characteristics and  
475 hardness of Cheddar cheese during ripening. *Journal of Food Science*, 72, 168-176.

476 Cooke, D. R., Khosrowshahi, A., & McSweeney, P. L. H. (2013). Effect of gum  
477 tragacanth on the rheological and functional properties of full-fat and half-fat  
478 Cheddar cheese. *Dairy Science & Technology*, 93, 45–62.

479 Crowley, S. V., Burlot, E., Silva, J. V. C., McCarthy, N. A., Wijayanti, H. B.,  
480 Fenelon, M. A., & O'Mahony, J. A. (2018). Rehydration behaviour of spray-dried  
481 micellar casein concentrates produced using microfiltration of skim milk at cold or  
482 warm temperatures. *International Dairy Journal*, 81, 72-79.

483 Dave, R. I., McMahon, D. J., Oberg, C., and Broadbent, J. R. (2003). Influence of  
484 coagulant level on proteolysis and functionality of Mozzarella cheeses made using  
485 direct acidification. *Journal of Dairy Science*, 86, 114-126.

486 Eigel, W. N., Butler, J. E., Ernstrom, C. A., Farrell, H. M., Harwalkar, V. R., Jenness,  
487 R., & Whitney, R. M. (1984). Nomenclature of proteins of cow's milk: Fifth revision.  
488 *Journal of Dairy Science*, 67, 1599-1631.

489 Everard, C. D., O'Callaghan, D. J., Howard, T. V., O'Donnell, C.P., Sheehan, E.M.,  
490 & Delahunty, C.M. (2006). Relationships between sensory and rheological  
491 measurements of texture in maturing commercial Cheddar cheese over a range of  
492 moisture and pH at the point of manufacture. *Journal Texture Studies*, 37, 361-382.

493 FAO. (2019). Dairy Market Review, March 2019. Food and Agriculture  
494 Organisation, Rome.

495 Farkye, N., & Fox, P. F. (1992). Contribution of plasmin to Cheddar cheese ripening:  
496 effect of added plasmin. *Journal of Dairy Research*, 59, 209.

497 Fenelon, M. A., & Guinee, T. P. (2000). Primary proteolysis and textural changes  
498 during ripening in Cheddar cheeses manufactured to different fat contents.  
499 *International Dairy Journal*, 10, 151-158.

500 Fox, P. F. (1963). *Potentiometric determination of salt in cheese*. *Journal of Dairy*  
501 *Science*, 46, 744-745.

502 Fox, P. F., & McSweeney, P. L. H. (1996). Proteolysis in cheese during ripening.  
503 *Food Reviews International*, 12, 457-509.

504 Fox, P. F., & McSweeney, P. L. H. (1997). Rennets: their role in milk coagulation  
505 and cheese ripening. In B.A. Law (Ed.), *Microbiology and Biochemistry of Cheese*  
506 *and Fermented Milk* (second ed.) (pp. 1-49). London, UK: Blackie Academic and  
507 Professional.

508 Fox, P. F., Guinee, T. P., Cogan, T. M., McSweeney, P. L. H. (2000). *Fundamentals*  
509 *of Cheese Science*. MD, USA: Aspen Publishers, Gaithersburg.

510 Freeman, T. R., Bucy, J. L., & Hassan, Z. (1970). Use of nonfat dry milk in the  
511 production of Cheddar cheese. *Journal of Dairy Science*, 53, 727-733.

512 Guinee, T. P., Mulholland, E.O., Kelly, J., & O'Callaghan, D. J. (2007). Effect of  
513 protein-to-fat ratio of milk on the composition, manufacturing efficiency and yield  
514 of Cheddar cheese. *Journal of Dairy Science*, 90, 110-123.

515 Gunasekaran, S., and Ak., M. M. (2002). *Cheese Rheology and Texture*. FL, USA:  
516 CRC Press LLC.

517 Hoogwegt.com. (2019). Market Matters Demand for Whey Products Looks Bright.  
 518 [online] Available at: [https://hoogwegt.com/media/2612/horizons\\_june-2019.pdf](https://hoogwegt.com/media/2612/horizons_june-2019.pdf)  
 519 [Accessed 2 Nov. 2019].

520 Hurley, M. J., O'Driscoll, B. M., Kelly, A. L., & McSweeney, P. L. H. (1999). Novel  
 521 assay for the determination of residual coagulant activity in cheese. *International*  
 522 *Dairy Journal*, 9, 553-558.

523 Ibáñez, R. A., Waldron, D. S., & McSweeney, P. L. H. (2015). Effect of pectin on  
 524 the composition, microbiology, texture, and functionality of reduced-fat Cheddar  
 525 cheese. *Dairy Science & Technology*, 96, 297-316.

526 IDF. (1982). *Determination of the total solids content of cheese and processed cheese.*  
 527 *IDF standard 4A*. Brussels, Belgium: International Dairy Federation.

528 IDF. (1986). *Determination of nitrogen content (Kjeldahl method) and calculation*  
 529 *of crude protein content. IDF standard 20A*. Brussels, Belgium: International Dairy  
 530 Federation.

531 Kelly, P. (2019). Manufacture of whey protein products: concentrates, isolate, whey  
 532 protein fractions and microparticulated. In H. C. Deeth & N. Bansal (Eds.), *Whey*  
 533 *Proteins* (pp. 97–122). London, UK: Elsevier Applied Science.

534 Kuchroo, C. N., & Fox, P. F., (1982). Soluble nitrogen in Cheddar cheese;  
 535 comparison of extraction procedures. *Milchwissenschaft*, 37, 331-335.

536 Kuchtík, J., Šustová, K., Urban, T., & Zapletal, D. (2008). Effect of the stage of  
 537 lactation on milk composition, its properties and the quality of rennet curdling in East  
 538 Friesian ewes. *Czech J. Animal Science*, 53, 55-63.

539 Lucey, J. A., Johnson, M. E., Horne, D. S. (2003). Invited review: perspectives on  
540 the basis of the rheology and texture properties of cheese. *Journal of Dairy Science*,  
541 86, 2725-2743

542 Lucey, J. A., Mishra, R., Hassan, A., & Johnson, M. E. (2005). Rheological and calcium  
543 equilibrium changes during ripening of Cheddar cheese. *International Dairy Journal*,  
544 15, 645-653.

545 Lucey, J., & Fox, P. F. (1992). Rennet coagulation properties of late-lactation milk:  
546 Effect of pH adjustment, addition of  $\text{CaCl}_2$ , variation in rennet level and blending  
547 with mid-lactation milk. *Irish Journal of Agricultural and Food, Research*, 31, 173-  
548 184.

549 Mahy, M., Van Eyckden, L., & Oosterlinck, A., (1994). Evaluation of uniform color  
550 spaces developed after the adoption of CIELAB and CIELUV. *Color Research and*  
551 *Application*, 19, 105-121.

552 Mane, A., Ciocia, F., Beck, T. K., Lillevang, S. K., & McSweeney, P. L. H.  
553 (2019). Proteolysis in Danish Blue cheese during ripening. *International Dairy*  
554 *Journal*, 97, 191-200.

555 Martin, G. J. O., Williams, R. P. W., & Dunstan, D. E. (2010). Effect of manufacture  
556 and reconstitution of milk protein concentrate powder on the size and rennet gelation  
557 behaviour of casein micelles. *International Dairy Journal*, 20, 128-131.

558 Ménard, O., Camier, B., Guyomarc'h, F., (2005). Effect of heat treatment at alkaline  
559 pH on the rennet coagulation properties of skim milk. *Lait*, 85, 515-526.

560 Moiseev, N., Suchkova, E. & Iakovchenko, N. (2017). Possibility of using  
561 reconstituted milk in manufacture of cheese with cheddaring and cheese curd  
562 stretching. *Agronomy Research*, 15, 1358-1368.

563 Mounsey, J. S., and O’Riordan, E. D. (1999). Empirical and dynamic rheological  
564 data correlation to characterize melt characteristics of imitation cheese. *Journal of*  
565 *Food Science*. 64, 701-703.

566 O’Brien, B., Mehra, R., Connolly, J. F., & Harrington, D. (1999). Seasonal variation  
567 in the composition of Irish manufacturing and retail milks. *Irish Journal of*  
568 *Agricultural and Food Research*, 38, 53-64.

569 O’Keeffe, R. B., Fox, P. F., Daly, C. (1978). Proteolysis in Cheddar cheese: role of  
570 coagulant and starter bacteria. *Journal of Dairy Research*, 45, 465–477.

571 O’Mahony, J. A., Lucey, J. A. & McSweeney, P. L. H. (2005). Chymosinmediated  
572 proteolysis, calcium solubilisation, and texture development during the ripening of  
573 Cheddar cheese. *Journal of Dairy Science*, 88, 3101-3114.

574 Richardson, B. C. & Pearce, K. N. (1981). The determination of plasmin in dairy  
575 products. *Journal of Dairy Science and Technology*, 16, 209-220.

576 Rogosa, M., Mitchell, J. A. & Wiseman, R. F. (1951). A selective medium for the  
577 isolation and enumeration of oral and faecal lactobacilli. *Journal of Bacteriology*, 62,  
578 132-133.

579 Shalabi, S. I. & Fox, P. F. (1987). Electrophoretic analysis of cheese, comparison of  
580 methods. *Irish Journal of Food Science and Technology*, 11, 135-151.

581 Sharma, G. & Bala, R. (2003). *Digital Color Imaging Handbook* (1st ed.). FL, USA:  
582 CRC Press LLC.

583 Sheehan, J. J., Wilkinson, M. G. & McSweeney, P. L. H. (2008). Influence of  
584 processing and ripening parameters on starter, non-starter and propionic acid bacteria  
585 and on the ripening characteristics of semi-hard cheese. *International Dairy Journal*,  
586 18, 905-917.

587 Soodam, K., Ong, L., Powell, I. B., Kentish, S. E., & Gras, S. L. (2015). Effect of  
588 calcium chloride addition and draining pH on the microstructure and texture of full  
589 fat Cheddar cheese during ripening. *Food Chemistry*, 181, 111-118.

590 Sousa, M. J., Ardo, Y., & McSweeney, P. L. H. (2001). Advances in the study of  
591 proteolysis during cheese ripening. *International Dairy Journal*, 11, 327-345.

592 Terzaghi, B. E. & Sandine, W. E. (1975). Improved medium for lactic streptococci  
593 and their bacteriophages. *Applied Microbiology*, 29, 807-813.

594 Truong, V. D., Daubert, C. R., Drake, M. A., & Baxter, S. (2002). Vane rheometry  
595 for textural characterization of cheddar cheeses: correlation with other instrumental  
596 and sensory measurements. *LWT - Food Science and Technology*, 35, 305-314.

597 Watkinson, P., Croker, C., Crawford, R., Dodds, C., Johnston, K., McKenna, A., &  
598 White, N. (2001). Effect of cheese pH and ripening time on model cheese textural  
599 properties and proteolysis. *International Dairy Journal*, 11, 455-464.