

Title	High-resolution facies zonation within a cold-water coral mound: The case of the Piddington Mound, Porcupine Seabight, NE Atlantic
Authors	Lim, Aaron;Wheeler, Andrew J.;Arnaubec, Aurélien
Publication date	2017-06-22
Original Citation	Lim, A., Wheeler, A. J. and Arnaubec, A. (2017) 'High-resolution facies zonation within a cold-water coral mound: The case of the Piddington Mound, Porcupine Seabight, NE Atlantic', Marine Geology, 390, pp. 120-130. doi:10.1016/j.margeo.2017.06.009
Type of publication	Article (peer-reviewed)
Link to publisher's version	10.1016/j.margeo.2017.06.009
Rights	© 2017, Elsevier B.V. All rights reserved. This manuscript version is made available under the CC-BY-NC-ND 4.0 license http:// creativecommons.org/licenses/by-nc-nd/4.0
Download date	2025-08-01 00:04:44
Item downloaded from	https://hdl.handle.net/10468/5200



University College Cork, Ireland Coláiste na hOllscoile Corcaigh

1	High-resolution facies zonation within a cold-water coral mound: the case of the
2	Piddington Mound, Porcupine Seabight, NE Atlantic
3	
4	Aaron Lim* <sup>1)</sup> , Andrew J. Wheeler <sup>1)2)</sup> and Aurélien Arnaubec <sup>3)</sup>
5	
6	<sup>1)</sup> School of Biological, Earth and Environmental Sciences, University College Cork, Distillery
7	Fields, North Mall, Cork, Ireland
8	<sup>2)</sup> Irish Centre for Research in Applied Geosciences, University College Cork, Ireland
9	<sup>3)</sup> Unité Systèmes sous-Marins, Centre de Méditerranée, Ifremer, Zone Portuaire de Brégaillon,
10	CS 20330, 83507 La Seyne/Mer Cedex, France
11	
12	Abstract

. . . .

Framework-forming cold-water corals (CWC's) such as Lophelia pertusa and Madrepora 13 oculata generate positive topographic features on the seabed called CWC mounds. In the North 14 East Atlantic, CWC mounds have been studied in detail and reveal heterogeneous spatial on-15 mound organisation of coral patches. Many of these studies are limited by a paucity of 16 17 remotely-sensed and video imagery at an appropriate resolution and coverage. This study is the first attempt to video mosaic an entire CWC mound (the Piddington Mound of the Moira 18 Mounds, Porcupine Seabight, Irish margin). The mosaic is divided into 18,980 0.25 m<sup>2</sup> cells 19 with a manual classification applied to each within a geographic information system (GIS). 20 Geospatial analysis shows that cell distribution is not random but clustered significantly across 21 the mound surface. These clusters of cells make up a ring-like facies pattern. A model for the 22 processes that lead to this facies pattern is suggested based on contemporary environmental 23

controls. Parallels to shallow-water reef atolls are also drawn which subsequently hasimplications for interpreting fossil coral outcrops.

26

27 \*Corresponding author: <u>aaron.lim@ucc.ie</u>

28

Keywords: cold-water coral, habitat mapping, spatial analysis, sediments, facies distribution

### 31 **1. Introduction**

32 Framework-forming, scleractinian cold-water corals (CWCs) are sessile, filter-feeding organisms that can baffle current-suspended sediment and biogenic material between their 33 framework (Roberts et al., 2006). Lophelia pertusa, the most common framework-forming 34 35 CWC in the NE Atlantic, has been found as shallow as 39 m water depth and as deep as 4000 m water depth (Freiwald et al., 2004; Roberts et al., 2006). In general, it occurs in temperatures 36 between 4 and 13°C (Freiwald, 2002) and has proven to be tolerant of a wide range of salinities, 37 38 from 31.7 - 38.8 ‰ (Davies et al., 2008). As the coral framework grows, it baffles sediment which can help to generate topographic features on the seabed called CWC reefs and, through 39 successive periods of reef development, CWC carbonate mounds (Freiwald, 2002; Roberts et 40 al., 2009). Here, we refer to mound-shaped, positive topographic features developed by CWC's 41 as CWC mounds. CWC mounds are common in the NE Atlantic (Wheeler et al., 2007), 42 43 specifically where internal waves concentrate food particles (phytodetritus) which is delivered to CWCs by enhanced bottom currents (Dullo et al., 2008; Mienis et al., 2009; White and 44 45 Dorschel, 2010).

Habitat mapping has proved to be a valuable, efficient and cost-effective tool in understanding 46 the marine environment (e.g. Huang et al., 2011; Lamarche et al., 2011) including CWC 47 48 habitats (Savini et al., 2014). Multibeam Echosounder (MBES) bathymetry and backscatter have been used extensively to characterise current dynamics and their influence on CWC 49 mound morphology and development e.g. in the straits of Florida, W Atlantic (Correa et al., 50 2012a; Correa et al., 2012b) and the midslope Moira Mounds, Porcupine Seabight, NE Atlantic 51 52 (Foubert et al., 2011). Recently, more advanced approaches to MBES surveying have imaged CWC habitats in deep water using ROV-borne MBES (Dolan et al., 2008; Foubert et al., 2011), 53 54 on submarine terraces using AUV-borne MBES (Correa et al., 2012a) and on vertical cliff faces in submarine canyons using forward-facing ROV-borne MBES (Huvenne et al., 2011). 55

In the absence of adequate multibeam data, other studies (e.g. Dorschel et al., 2007; Wheeler 56 et al., 2008) avail of current data, sediment types, video data and/or side scan sonar (SSS) 57 surveying integrated within a Geographical Information System (GIS) to highlight the role of 58 currents and sediment supply on CWC reefs and mounds. Seabed sediment samples are an 59 effective way of studying CWC reefs although limited by the spatial representation of the 60 sample (e.g. Day grab, <0.5 m<sup>2</sup>). Video surveys can discriminate the seabed across substantial 61 62 areas and are widely used in CWC habitat inspections (Foubert et al., 2005; Huvenne et al., 2005; Vertino et al., 2010). Recent advances in underwater imaging have made high-resolution 63 64 underwater imagery with accurate positioning in deep water environments possible (Kocak and Caimi, 2005). As a result, since 2005, the utilisation of ROV-based observations for facies 65 information from CWC mounds has increased dramatically (e.g. De Mol et al., 2007; Foubert 66 et al., 2008; Guinan et al., 2009; Hebbeln et al., 2014; Heindel et al., 2010; Huvenne et al., 67 2016; Purser, 2015; Wienberg et al., 2013). Huvenne et al. (2005) demonstrate the variability 68 between entire mound provinces along the Irish Margin highlighting both the frequency and 69 variability of CWC mounds within the Belgica Mound Province. Later, Dorschel et al. (2007) 70

observed a correlation between living coral and enhanced bottom currents in the Belgica 71 Mound Province and outlined the influence of contour currents, tidal currents and local 72 73 topography on the distinct coral facies distribution across the Galway Mound. Results from other ROV-based facies mapping also highlight the strong relationship between local currents 74 and facies distribution across a mound surface. For example, at the Franken Mound, a CWC 75 mound in a state of "mound retirement" at the western Rockall Bank, shows a distinct facies 76 77 distribution across the mound with living coral dominating the summit region (Wienberg et al., 2008). Many other uses for ROV-based facies observations were realised. Heindel et al. (2010) 78 79 test, for the first time, a method of spatial prediction mapping (maximum likelihood classification) on a CWC ecosystem providing detailed aerial estimates of CWC-typical facies. 80 ROV-based facies observations were utilised to show that L. petusa is restricted to longer term, 81 stable conditions while M. occulata is more tolerant towards environmental fluctuations 82 (Wienberg et al., 2009). Furthermore, its need as a tool in marine reserve designation and 83 implementation is now recognised (Roberts et al., 2005). 84

In addition, advances in image processing has led to the application of video mosaicking to
marine habitat mapping (Rzhanov *et al.*, 2000). For example, Lirman *et al.* (2007) accurately
characterise a tropical coral reef in shallow water using an entire reef-scale video mosaic.
However, no such study has been carried out on an entire CWC reef although small parts of
CWC habitats have been manually photo mosaicked (Wheeler *et al.*, 2011).

90 The need for more local-scale studies and data sets of equal resolution have previously been 91 highlighted (Dolan *et al.*, 2008). This study presents the first attempt of video mosaicking an 92 entire cold-water coral reef and subsequent analyses providing an in-depth facies mapping 93 exercise allowing to discuss facies organisation and potential facies organisational influences.

The Belgica Mound Province (BMP), partly enclosed with a Special Area of Conservation 96 (SAC) designated under the EU Habitats Directive, is located on the eastern flank of the 97 Porcupine Seabight, NE Atlantic (see Fig. 1) (Beyer et al., 2003; Dorschel et al., 2007; 98 Huvenne et al., 2002). It contains an abundance of (giant) CWC carbonate mounds, including 99 100 the well-studied Galway Mound, Thérèse Mound and Challenger Mound (De Mol et al., 2007; Dorschel et al., 2007; Eisele et al., 2008; Huvenne et al., 2009; Kano et al., 2007; Thierens et 101 al., 2010; Titschack et al., 2009; Wheeler et al., 2005). Two distinct CWC carbonate mound 102 chains have been identified, orientated roughly N-S (parallel to depth contours) (Wheeler et 103 al., 2005). Pre-existing bathymetry highlights their slight elongate to conical morphology and 104 105 typical dimensions (approx. 1 km across and 100 m tall) (Beyer et al., 2003). To the east, these large CWC mounds are enclosed by the continental shelf and to the west, the Arwen Channel 106 107 runs through the BMP (Murphy and Wheeler, 2017; Van Rooij, 2004). Towards the south and 108 south-west, the Porcupine Seabight exits out to the abyssal plain (Dorschel et al., 2010) while the north shallows to 500 m, outside the typical depth range for CWC growth on the Irish 109 continental margin (Dullo et al., 2008; White and Dorschel, 2010) 110

The Moira Mounds are small-type CWC reefs (approx. 30 m across and 10 m tall) found 111 throughout the BMP, occurring between 800 - 1100 m water depth (Wheeler et al., 2005). 112 113 While no definitive dating has confirmed their age, it is speculated that they are Holocene features based on their size, seismic profiles and the surrounding seabed substrate (Foubert et 114 al., 2011; Huvenne et al., 2005; Kozachenko, 2005). The Moira Mounds in the BMP can be 115 116 further subdivided into 4 areas or chains of mounds based on the distribution of approx. 256 Moira Mounds; the "up-slope area", the "mid-slope area", the "down-slope area" and the 117 "northern area" (Wheeler et al., 2011) (see Fig. 1). While the number of Moira Mounds in the 118 northern and the up-slope areas are relatively sparse, the main focus of research has been 119

carried out on the mid- and down-slope areas. The mid-slope area occurs between the chain of
large CWC carbonate mound structures. The Moira Mounds here are thought to represent
mound formation under "stressed" conditions due to high sediment input (Foubert *et al.*, 2011).
The down-slope area is unique as it occurs within the Arwen Channel. Unlike the other areas,
the CWC's on the Moira Mounds are predominantly growing, are actively trapping sands
(Wheeler *et al.*, 2011) and occur outside the influence of other large mound structures.

126 <INSERT FIGURE 1>

127

#### 2. Materials and Methods

128 2.1 ROV-borne Multibeam Echosounder (MBES)

ROV-borne MBES data was collected over the Piddington Mound and the surrounding seabed 129 130 during the QuERCi survey (2015) on board RV Celtic Explorer with the Holland 1 ROV (cruise number CE15009: Wheeler et al. (2015). A high-resolution Kongsberg EM2040 MBES was 131 integrated with a sound velocity probe and mounted on the front-bottom of the ROV. Data 132 were acquired at a frequency of 400 kHz while the ROV maintained a height of 30 m above 133 the seabed with a survey speed of approximately 2 knots. This achieved a swath width of ~ 160 134 135 m. Positioning and attitude were obtained using a Kongsberg HAINS inertial navigation system, ultra-short baseline (USBL) system (Sonardyne Ranger 2) and doppler velocity log 136 (DVL). Data acquisition was carried out using SIS software, where calibration values, sensor 137 offsets, navigation and attitude values were incorporated. Two adjacent 170 m long MBES 138 139 lines were collected over Piddington Mound and the surrounding seabed. MBES data were stored as \*.all and \*.wcd files and were processed using CARIS HIPS and SIPS v9.0.14 to 140 141 apply tidal corrections and clean anomalous data spikes. The cleaned data were saved as a single \*.xyz and gridded to a 10 cm ArcView GRID. 142

The 10 cm MBES grid was imported into ArcMap 10.2 and projected in UTM Zone 29N. A 1
m contour \*.shp file was generated using the Arc Toolbox Spatial Analyst Contour tool. Slope
(degrees) and aspect were derived from the bathymetry using the Arc Toolbox Spatial Analyst
tools.

147

148 2.2 ROV-video data collection

ROV-video data was collected over Piddington Mound during the VENTuRE survey (2011) on board *RV Celtic Explorer* with the *Holland 1 ROV* (cruise number CE11009: Wheeler and shipboard party, 2011). A downward-facing, high-definition camera was mounted on the bottom of the ROV. Positioning and navigation were achieved using a USBL (Sonardyne Ranger 2) and RDI Workhouse DVL. The ROV altimeter recorded and logged the height of the ROV from the seabed. The ROV recorded downward-facing HD video during a series of transects across the mound <2 m off the seabed/mound surface.</p>

156

# 157 2.3 Georeferenced video mosaic generation

A georeferenced video mosaic has been generated using the IFREMER in-house Matisse 158 software where the raw video data was imported and from which images have been extracted 159 160 at a rate of 1 per second. Poor quality imagery, possibly due to an excessive fly height, and/or with a poor navigational lock were not included. The associated USBL navigation has been 161 filtered with a sliding median filtering and 2<sup>nd</sup> order polynomial model fit in order to lower 162 trajectory noise. Image and navigation data have been synchronized so that each image has an 163 initial approximate position. This position is refined later by the mosaicking process. For this 164 refinement, the first process consists of feature detection and matching between images using 165 the SIFT (Scale Invariant Feature Transform) algorithm (Lowe, 1999), known for its strong 166

robustness and accuracy. Image matching alone would lead to a drifting mosaic and invalid scale, while USBL navigation alone is not accurate enough for high-quality local-overlapping but gives accurate global positioning. To benefit from the both image matching and USBLnavigation, we merged image and navigation information through a cost function minimization. This developed method is similar to Ferrer *et al.* (2007) except cost function weights are affected according to image and navigation data standard deviations so reprojection errors are minimized in the mosaicking plane rather than the image plane.

After this step, image positions are refined and the mosaic can be drawn. This is carried out
with state of the art seaming and blending techniques after Burt and Adelson (1983) and Kwatra *et al.* (2003).

177 2.4 Seabed classification

As the ROV could not be maintained at a constant height above the seabed in video surveying mode in strong currents, seawater-induced blue-shift varied across the video mosaic. Hence, neither a supervised nor an unsupervised (automatic) seabed classification could be utilized. Instead, a manual classification was applied to each cell of the entire video mosaic. An ArcMap fishnet with a cell size of 0.25 m<sup>2</sup> was overlaid on the video mosaic to manage the supervised classification at the highest resolution possible where each 0.25 m<sup>2</sup> cell can be inspected and classified (totally 18,980 cells).

Classifiers were based on preliminary assessment of the video mosaic and other CWC habitat
ROV-dive-based facies classification from previous work, some of which have similar
objectives or are proximal to the study site (e.g. Dorschel *et al.*, 2007; Dorschel *et al.*, 2009;
Douarin *et al.*, 2014; Heindel *et al.*, 2010; Huvenne *et al.*, 2005; Spezzaferri *et al.*, 2012;
Vertino *et al.*, 2010; Vertino *et al.*, 2015; Wheeler *et al.*, 2005; Wienberg *et al.*, 2008; Wienberg *et al.*, 2009). The facies classifiers are: "hemipelagic sediment with dropstones", "coral

rubble", "hemipelagic sediment", "dead coral framework" and "live coral framework" (see Fig.
2 for examples and results for detailed facies descriptions).

As each individual cell is inspected and a suitable class chosen, a value was assigned to an attribute table. Several preliminary classifications were applied to the video mosaic in different locations to test the individual classifiers and the classification itself. Some areas of the preliminary classification were re-classified and compared to their initial classification to ensure classification accuracy. The entire video mosaic was then classified and saved as a \*.shp file.

199

## 200 2.5 Video-mosaic analysis

201 Spatial dependency is measured by various spatial autocorrelation statistics (e.g. Moran's I: Goodchild, 1986). However, these are unrepresentative when spatial autocorrelation varies 202 203 significantly over the study site. More suitably, Anselin (1995) developed local indicators for 204 spatial association (LISA) which, in contrast to other methods, measures local variation within patterns of spatial dependence that may not be represented in existing techniques. A widely-205 206 used LISA is the G statistic (Getis and Ord, 1992; Ord and Getis, 1995). This measures how concentrated high or low values are for a given study area by calculating a z-score (standard 207 score) and p-value (calculated probability) of a set of geographical values. Here, it is utilised 208 to measure the degree of clustering of the classified cells via the Arc Toolbox spatial statistics 209 tools (High/Low clustering Tool) based on Euclidean distances. 210

The classified \*.shp file was converted to a raster using the conversion tools in Arc Toolbox and imported to the Focal Statistics tool. This calculates, for each input cell location, a statistic of the value that occurs most often (majority) within a specified neighbourhood around it. The specified neighbour was calculated based on a potential navigational error of the ROV (<5 m).

Therefore, the neighbourhood was defined as 2.5 m radius around each cell location.

216

#### 217 **3. Results**

218 3.1 Video mosaic

The video mosaic was generated from approximately 8 hours of high-definition video data. The data was collected by means of a series of over lapping lines covering the mound and immediate off-mound area. The video mosaic is shown in Fig. 2, holes in the mosaic are a result of data rejection. For subsequent display purposes, colours on the mosaic were adjusted manually selecting both a white and black reference within Corel Draw software package.

#### 224 <INSERT FIGURE 2>

3.2 Bathymetry and slope

Piddington Mound exists in an area with several other mounds (Fig. 3A). It has a similar size 226 (11.8 m tall, 60 m x 40 m across) and morphology (slightly elongated conical) as the 227 228 surrounding mounds and it has substantial scouring (1 m - 3 m deep) at its southern limit. Sinuous megaripples (sediment waves) exist across the area around these mounds. They have 229 a wavelength of  $\sim 10$  m and a wave height of 20 cm -75 cm. To the south of each mound, there 230 231 are positive, ridge-shaped features 0.5 m - 2 m tall. These features are unlike the sediment waves as they are larger than the sediment waves, have coral colonising them and biogenic 232 material deposited in their troughs as shown by the oblique video footage (Fig. 4). These ridge 233 and wave features are not included in the facies classification as they are not on the Piddington 234 Mound or in the immediate environment and therefore not covered by the video mosaic. 235

# 236 <INSERT FIGURE 3>

#### 237 <INSERT FIGURE 4>

The deep blue colour represents flat seabed slopes (minimum slope of  $0^{\circ}$ ) while the red colour 238 represents the steep seabed slopes (maximum slope of 55.7°), all relative to this data set (Fig. 239 3B). Note that regional slope angle in this area is  $1.2^{\circ}$ . The steepest areas are found on the 240 flanks of the mounds with slopes dipping as steep as 55°. The summit of each mound is 241 relatively flat  $(0-6^{\circ})$  as are areas away from the mounds  $(0^{\circ} - 6^{\circ})$  (Fig. 3B). This figure also 242 includes the slip faces of mega ripples in the areas away from the mounds. These appear 243 sinuous to cuspate with a wavelength of 7 m - 10 m and wave height of 4 cm - 20 cm with an 244 east-west crest alignment. The larger coral colonised ridge-shaped features (0.5 m - 2 m tall) 245 south of the mounds are morphologically distinct, with short east-west aligned linear crests and 246 a wavelength of approx. 10 m. They diminish in size away from the mounds. 247

248

#### 249 3.3 Seabed classification

#### 250 <INSERT FIGURE 5>

The "hemipelagic sediment with dropstones" class is characterised by the cell being dominated 251 (>50 % cell coverage) by dropstones and sediment (sand or mud). This class occurs most 252 commonly around the mound perimeter. The "coral rubble" class is characterised by a cell 253 dominated (>50% cell coverage) by recognisable biogenic material (i.e. coral rubble, shell 254 fragments) and sediment. This class most commonly occurs in a 10 m wide ring immediately 255 enclosed by the "hemipelagic sediment with dropstones" class. However, minor occurrences 256 257 of this class exist both to the centre of the mound and outside of this ring. This ring is elongate in the same direction of the current (N-S). The "hemipelagic sediment with dropstones" and 258 "coral rubble" classes are the most frequent classified cells in the video mosaic (Fig.5). The 259 "hemipelagic sediment" class is characterised by the cell being dominated (>90% cell 260

coverage) by hemipelagic sediment i.e. sand or mud with no recognisable bioclasts or 261 dropstones. It is the least common class and occurs in small clusters less than 0.75 m wide 262 typically around the edges of the "coral rubble" class. The "dead coral framework" class is 263 characterised by the cell being dominated (>50% cell coverage) coral framework which has no 264 identifiable living parts. This class occurs across the mound but is concentrated in the centre 265 and southern central area as a North-South-elongate ring. The "live coral framework" class is 266 267 characterised by the cell being dominated (>50% cell coverage) by coral framework with identifiable living parts (polyps or mucus-covered frameworks evident) although major 268 269 proportions of the coral framework may be dead. It is found across the mound, but is concentrated in the northern central area. 270

271 3.4 Spatial statistical analysis

The High-Low Clustering (Getis-Ord General G) analysis indicates the concentration of clustering. It firstly applies a null hypothesis to determine if clustering occurs. In this case, with a large Z-score (168.89) and small p-value (<0.001), the null hypothesis can be rejected meaning that clustering occurs. A positive Z-score indicates that high-values (coral frameworks) cluster.

The majority value of cells in the specified neighbourhood (2.5 m radius) (Fig. 6) reveals 4 facies occurring in 5 areas: "Hemipelagic sediment and dropstone" facies, "Coral rubble" facies, "Dead coral framework" facies, and "Live coral framework" facies. This means that in these facies, a particular cell-type dominates, as shown by the focal statistic (majority) based on a 2.5 m radius. The distribution of these facies closely follows the distribution of observed common occurrence of the individual classified cells. These facies have a ring/annulus-like distribution. The "Hemipelagic sediment and dropstone" facies exists in the outer rim of this ring/annulus shape. It is made up of 74.9% "hemipelagic sediment and dropstone" cells, 15.9% "coral rubble" cells, 2.3% "dead coral framework" cells, 1.6% "hemipelagic sediment" cells and 1.3% "live coral framework" cells (Fig. 6b).

Immediately inside the "Hemipelagic sediment with dropstone" facies, the "Coral rubble" facies exists. However, this facies also protrudes 11.5 m through this facies NW of the mound and exists on the highest point of the mound (at -968 m water depth) protruding 11.5 m towards the ENE. It is made up of 60.6% "coral rubble" cells, 16.5% "dead coral framework" cells, 6.9% "live coral framework" cells, 6.3% "hemipelagic sediment and dropstone" cells and 1.6% "hemipelagic sediment" cells (Fig. 6b).

The "Dead coral framework" facies exists immediately inside the main body of the "Coral rubble" facies ring. It has a U-shape where the centre of the "U" is broadly based around the highest point on the mound. This facies is made up of 53.1% "dead coral framework" cells, 24.3% "coral rubble" cells, 17.4% "live coral framework" cells, 0.6% "hemipelagic sediment" cells and 0.1% "hemipelagic sediment and dropstone" cells (Fig. 6b).

Finally, the "Live coral framework" facies occurs 4.8 m north of the highest point on the mound, where it completes the U-shape of the "Dead coral framework zone" to make a complete ring, encircling the central "Coral rubble" facies. This zone has an irregular morphology where the zone is intruded by minor occurrences of the "Dead coral framework" facies. It is made up of 47.8% "live coral framework" cells, 32.1% "dead coral framework" cells, 13.6% "coral rubble" cells, 0.2% "hemipelagic sediment" cells and 0% "hemipelagic sediment and dropstone" cells (Fig. 6b).

The "hemipelagic sediment" cells were not common enough in any neighbourhood (2.5 m) to develop its own facies in the focal statistics tool. However, their distribution in relation to these facies is noted.

309 <INSERT FIGURE 6>

310 **4. Discussion** 

311 4.1 Local sedimentary environment

East-west orientated sediment waves surrounding Piddington Mound and its neighbouring 312 mounds suggest a high-energy, bottom current-influenced, sedimentary environment (Foubert 313 et al., 2011). The cross-profile morphology (steepened lee slope) of the sediment waves suggest 314 a south to north prevailing current across the area (Fig. 3B). The current-orientated, elongate 315 316 morphology may be owed to the dominance of currents here. In fact, substantial scouring around the base of the mounds (Fig. 3A) highlights the intensity of these currents. The larger 317 ridge-form features (see Fig. 3A) to the south of the mounds may no longer be active sediment 318 transport bedforms as the presence of coral on them suggest stability since the growth of the 319 corals seen on the surface (see Fig. 4). However, as the internal composition is unknown 320 321 nothing about their origin can be said with certainty. De Mol et al. (2007) and Wheeler et al. (2007) describe similar coral-topped features at the base of the Thérèse Mound. 322

323

### 324 4.2 Piddington Mound spatial organisation

The off-mound area of the Moira Mounds has been previously described (see Foubert *et al.*, 2011; Wheeler *et al.*, 2005). Off-mound areas are typically less spatially heterogeneous than the on-mound areas (e.g. Vertino *et al.*, 2010). The video mosaic covers the entire Piddington Mound surface and the immediate off-mound area, allowing inclusion of the full-mound surface in the analyses. Further, the chosen resolution of the analyses is both commensurate with typical box core footprints used to study other CWC mounds and is of sufficient resolutionto map facies change and distribution.

332 The G-statistic reveals that, despite the heterogeneous nature of the facies on Piddington Mound observed from the video mosaic at a  $0.25 \text{ m}^2$  scale, there is a statistically significant 333 clustered pattern. The focal statistics layer shows the geographic distribution of cell-type 334 335 dominance (Fig. 6a). The ring-like pattern of these dominating cell-types is clearly focussed around the summit of Piddington Mound observed in the bathymetric data (Fig. 3A). This ring-336 like clustered pattern is made up of facies typical of CWC mounds ("Live coral framework", 337 "Dead coral framework" and "Coral rubble"). Similar facies have been identified at the nearby 338 Galway Mound (Dorschel et al., 2007), other Moira Mounds (Foubert et al., 2011; Wheeler et 339 al., 2005), Atlantis Mound and Yellow Chain at the Santa Maria di Leuca Coral Province in 340 the Mediterranean (Vertino et al., 2010), Franken Mound (Wienberg et al., 2008), Propeller 341 Mound (Heindel et al., 2010) and various mound provinces within the Porcupine Seabight 342 (Huvenne et al., 2005). The distribution of these mound-typical facies closely follows the 343 mound depth contours. Where the seabed becomes relatively flat, surrounding Piddington 344 Mound (Fig. 3B), it is characterised by "Hemipelagic sediment with dropstones" facies which 345 is the typical background facies present in the area (Foubert et al., 2011). It is therefore likely 346 that the on-mound facies distribution can be attributed to the presence of corals where each 347 348 facies may represent the result of a dominant process or set of processes at that part of the mound. The "Live coral framework" area slightly to the north of the Piddington Mound summit 349 is the area on the mound where coral growth dominates. The "Coral rubble" areas on the 350 mound, found near the summit and on the fringes of the reef, represent the areas where notable 351 accumulations of eroded (physical or biological) coral and other biogenic material has been 352 deposited. The "Dead coral framework" area found mainly on the flanks of the reef and are 353 typically found where the "Coral rubble" and "Live coral framework" intersect. This area 354

represents an area where live coral growth no longer dominates and older framework remainsexposed.

357

358 4.3 Influences on facies distribution

The "Hemipelagic sediment and dropstone" facies dominates around the Piddington Mound 359 within the surrounding scour pits (Fig. 3A). Initial deposition of this dropstone-rich layer and 360 posterior re-exposure by erosion is rather unlikely as coring and drilling attempts in the vicinity 361 362 of the Belgica Mound Province CWC mounds never observed such a layer (e.g. IODP 307 Expedition scientists, 2005). Alternatively, perhaps the dropstones were scattered in 363 hemipelagic sediment in which winnowing of this sediment up-concentrated the dropstones. 364 365 This would also explain the preferential occurrence of the hemipelagic sediment with dropstone facies within the scour pit. 366

367 The "hemipelagic sediment" cell-type is uncommon, existing in very small patches scattered across Piddington Mound. However, its distribution is noted (Fig. 5) occurring in 2 different 368 settings. The first setting is next to coral framework cell-types ("Dead coral framework" and 369 370 "Live coral framework") where the coral frameworks may act as a physical barrier to erosion for the already deposited sediment from a former lower hydrodynamic energy regime. The 371 second setting is at the boundary between the "hemipelagic sediment and dropstone" cell-types 372 and "coral rubble" cell-types in the current-facing (southern) side of the mound. This 373 deposition of "hemipelagic sediment" may be explained by a sudden change in seabed rugosity 374 slowing the current between these two cell-types, allowing the deposition of current-suspended 375 sediment. 376

Wilson (1979) proposes a ring-like CWC growth (Wilson Ring model), particularly focussed
around the coppice stage (*sensu* Squires, 1964). Wilson (1979) describe CWC reef

development in optimum conditions, with limited sediment supply, where autogenic (biogenic)
processes dominate; an isolated colony grows and successive rings of coral debris and colonies
develop around this initial point (colony). This forms a circular CWC coppice made up of
alternating rings of colonies and debris where, presumably, the centre is older than the edges
of the structure. However, at the reef-scale, this process of ring-like development seems rather
unlikely.

385 <INSERT FIGURE 7>

386 A more likely explanation for the ring-like distribution of facies across the Piddington Mound is based on benthic environmental conditions. Currents are well-known to influence facies 387 distribution across CWC mounds (Dorschel et al., 2007). In fact, previous research show that 388 389 even local-scale (bio)zonation may be primarily controlled by current flow over them (Messing 390 et al., 1990). Here, currents are considered as the dominant (allogenic) control on the distribution of facies across Piddington Mound. CWC's preferentially settle on elevated 391 392 structures where they have access to faster flowing water and enhanced food supply, undiluted by benthic sediment transport processes (Freiwald, 2002; Roberts et al., 2003; Roberts et al., 393 2006). F1 and F2 (Figure 7) occur relatively high (several meters above the surrounding 394 seabed) on the Piddington Mound. This is also the steepest part of the mound  $(45-50^\circ)$  which 395 may be a result of coral trapping sediment and generating topography (Wheeler *et al.*, 2005; 396 397 Wheeler et al., 2008). Such steepened slopes may cause biologically and physically eroded bioclasts to roll onto the flatter parts of the mound (predominately the downslope R2). Thus, 398 F1 and F2 are likely the source of the bioclasts that make up the "Coral rubble" facies in R2. 399 400 The central "Coral rubble" facies at the mound summit (R1) occurs upslope and sheltered by a ring of coral framework and therefore its distribution cannot be explained by the same process. 401 Here, this central area is fully encircled by dense coral framework which locally cuts it off from 402 surrounding currents and subsequent food supply. Despite its elevated occurrence, a reduction 403

in food supply and current velocity may result in a less favourable environment at this central 404 area. This may explain the degradation of these corals to become an area of coral rubble. As 405 406 such, 3 comparisons can be drawn between Piddington Mound and shallow-water reef atolls where a) a ring of coral shelters/cuts off the centre, creating a micro-environment, markedly 407 different from the surrounding environment akin to the central lagoon area of a shallow-water 408 reef atoll, b) the slope of the Piddington Mound is similar to that of shallow-water reef atolls 409 410 (45°) (e.g. Mururoa Atoll) (Chevalier, 1968) and subsequently c) the reef material is transported down its slope (Woodroffe and Biribo, 2011). 411

On Piddington Mound, the location and distribution of the "Live coral framework" area is 412 interesting as live coral framework is typically found at or near the summit of mounds 413 (Rüggeberg et al., 2011). However, the main loci for live coral growth on different mounds 414 varies from setting to setting: on the summit (e.g. Galway Mound: Dorschel et al. (2007), the 415 lee side of the mound structure (e.g. Giant Mound and the Hedge Mounds: Dorschel et al., 416 2009), and to the current-facing side of the mound structure (e.g. CWC reefs at Hola, offshore 417 Norway: (Buhl-Mortenson et al., 2015). In the case of the Piddington Mound, it occurs just 418 north of the summit (Fig. 6) on the lee side of the mound. In areas of excessively high current 419 420 flow on the Porcupine Bank, CWC have been found flourishing on the lee-side of mounds (Dorschel et al., 2009). However, this is unlike other mounds in the eastern Porcupine Seabight 421 422 (i.e. Galway Mound: Dorschel et al., 2007), and here the summit of Piddington Mound is dominated by Coral rubble facies. Fig. 3C maps the facies types across the Piddington Mound 423 bathymetric cross-profile. Lim (2017) measure a current speed here of 39 - 41 cm s<sup>-1</sup> in this 424 area, the highest noted for the western Moira Mounds chain. It may be that the live corals in 425 F2 thrive in the shelter provided by the mound which acts as a physical barrier slowing the 426 currents. Purser et al. (2010) and Orejas et al. (2016) show that Lophelia capture food more 427 effectively in lower flow velocities. However, whilst these are laboratory-based studies, a 428

detailed study from the Mingulay Reef Complex may also explain the distribution of live corals
on the Piddington Mound where mound topography-induced turbulence may account for
enhanced delivery of food particles suspended in deeper water (see Davies *et al.*, 2009).

The facies zonation observed on the Piddington Mound has various implications for the 432 interpretation of coral deposits in fossil outcrop. Coral diversity, coral morphology, sediment-433 434 type (hydrodynamic energy), presence of algae, abundance of pelagic or planktonic components, microborings, surrounding facies or geochemistry are among the numerous 435 recognition criteria to distinguish between shallow- or deep-water occurring CWC mound 436 deposits (Mullins et al., 1981). With more studies available on modern CWC mounds, Hebbeln 437 and Samankassou (2015) show that misinterpretation of some of these distinguishing criteria 438 are possible and, in fact, for many ancient carbonate mounds, formation in deep water appear 439 just as likely as formation in shallow shelf seas. In line with this argument, various comparisons 440 have been outlined herein between the Piddington Mound and shallow water reef atolls. In 441 442 addition, facies zonations appear common in both modern, shallow-water coral reef build ups (Janßen et al., 2017) and in outcrops (Gischler, 1995; Reolid et al., 2014). Here, the first deep-443 water CWC mound video mosaic also exhibits a reef-scale facies zonation. Thus, these 444 similarities must also be taken into consideration when interpreting fossil coral outcrops. 445

446

## 447 **5.** Conclusion

Although video mosaicking an entire CWC reef at this resolution is time consuming in data acquisition, processing and analysis, our preliminary investigation offers new, unique and detailed insights to CWC reef organisation and facies distribution. Despite mound surface heterogeneity, areas of the Piddington Mound surface (as small as 0.25 m<sup>2</sup>) cluster significantly in a ring-like pattern focussed around the mound summit. While ring-like observations have been reported at other CWC mounds (e.g. Henriet *et al.*, 1998; Wilson, 1979), the facies pattern
here is physically dissimilar. A definitive explanation cannot be put forward for the existence
of this ring-like facies pattern however, a scenario is put forward based on the influences of the
surrounding environment drawing parallels to shallow-water coral atolls. The existence of a
facies zonation implies that deep-water CWC mounds are similar to shallow-water coral buildups and this should be taken into consideration when making interpretations from fossil coral
outcrops.

Looking to the future, this data set may be used to either compare supervised and unsupervised
classification techniques or to ground-truth high-resolution facies distribution prediction
models.

463

## 464 **6.** Acknowledgements

The authors would like to thank Dr. Jürgen Titschack and an anonymous reviewer for their time and effort in extremely useful and constructive reviews. Further, the authors would like to thank the crew and officers of ROV Holland 1 and RV Celtic Explorer for assistance in collecting high resolution, accurately positioned data. Ship time on the RV Celtic Explorer was funded by the Marine Institute under the 2011 and 2015 Ship Time Programme of the National Development Plan. Aaron Lim is funded by the Irish Research Council Graduate of Ireland Scholarship IRC (2015 - 2016).

472

# 473 **7. References**

Anselin, L., 1995. Local Indicators of Spatial Association—LISA. Geographical Analysis 27
(2), 93-115, 10.1111/j.1538-4632.1995.tb00338.x.

- 476 Beyer, A., Schenke, H.W., Klenke, M., Niederjasper, F., 2003. High resolution bathymetry of
- the eastern slope of the Porcupine Seabight. Marine Geology 198 (1-2), 27-54,
   <a href="http://dx.doi.org/10.1016/S0025-3227(03)00093-8">http://dx.doi.org/10.1016/S0025-3227(03)00093-8</a>.
- Buhl-Mortenson, L., Hodnesdal, H., Thorsnes, T., 2015. The Norwegian Sea Floor.
  MAREANO 2015,
- Burt, P.J., Adelson, E.H., 1983. A Multiresolution Spline with Application to Image Mosaics.
  Acm Transactions on Graphics 2 (4), 217-236, Doi 10.1145/245.247.
- 483 Chevalier, J.-P., 1968. Etude géomorphologique et bionomique de l'atoll de Mururoa
  484 (Tuamotu),
- 485 Correa, T.B.S., Eberli, G.P., Grasmueck, M., Reed, J.K., Correa, A.M.S., 2012a. Genesis and
  486 morphology of cold-water coral ridges in a unidirectional current regime. Marine Geology
  487 326, 14-27, 10.1016/j.margeo.2012.06.008.
- 488 Correa, T.B.S., Grasmueck, M., Eberli, G.P., Reed, J.K., Verwer, K., Purkis, S.A.M., 2012b.
  489 Variability of cold-water coral mounds in a high sediment input and tidal current regime,
- 490 Straits of Florida. Sedimentology 59 (4), 1278-1304, 10.1111/j.1365-3091.2011.01306.x.
- 491 Coughlan, M., Wheeler, A.J., Dorschel, B., Lordan, C., Boer, W., Gaever, P.v., Haas, H.d.,
- 492 Mörz, T., 2015. Record of anthropogenic impact on the Western Irish Sea mud belt.
- 493 Anthropocene 9, 56-69, <u>http://dx.doi.org/10.1016/j.ancene.2015.06.001</u>.
- 494 Davies, A.J., Duineveld, G.C., Lavaleye, M.S., Bergman, M.J., van Haren, H., Roberts, J.M.,
- 495 2009. Downwelling and deep-water bottom currents as food supply mechanisms to the cold-
- 496 water coral Lophelia pertusa (Scleractinia) at the Mingulay Reef complex. Limnology and
- 497 Oceanography 54 (2), 620, DOI 10.4319/lo.2009.54.2.0620.
- 498 Davies, A.J., Wisshak, M., Orr, J.C., Murray Roberts, J., 2008. Predicting suitable habitat for
- the cold-water coral Lophelia pertusa (Scleractinia). Deep Sea Research Part I:
- 500 Oceanographic Research Papers 55 (8), 1048-1062, 10.1016/j.dsr.2008.04.010.
- 501 De Mol, B., Kozachenko, M., Wheeler, A.J., Alvares, H., Henriet, J.-P., Olu-Le Roy, K.,
- 502 2007. Thérèse Mound: a case study of coral bank development in the Belgica Mound
- Province, Porcupine Seabight. International Journal of Earth Sciences 96 (1), 103-120,
   <u>http://dx.doi.org/10.1007/s00531-005-0496-x</u>.
- Dolan, M.F.J., Grehan, A.J., Guinan, J.C., Brown, C., 2008. Modelling the local distribution
  of cold-water corals in relation to bathymetric variables: Adding spatial context to deep-sea
  video data. Deep-Sea Research I 55 (11), 1564-1579,
- 508 <u>http://dx.doi.org/10.1016/j.dsr.2008.06.010</u>.
- 509 Dorschel, B., Hebbeln, D., Foubert, A., White, M., Wheeler, A.J., 2007. Hydrodynamics and
- 510 cold-water coral facies distribution related to recent sedimentary processes at Galway Mound
- 511 west of Ireland. Marine Geology 244 (1–4), 184-195,
- 512 <u>http://dx.doi.org/10.1016/j.margeo.2007.06.010</u>.
- 513 Dorschel, B., Wheeler, A.J., Huvenne, V.A.I., de Haas, H., 2009. Cold-water coral mounds in
- an erosive environmental setting: TOBI side-scan sonar data and ROV video footage from

- the northwest Porcupine Bank, NE Atlantic. Marine Geology 264 (3-4), 218-229,
- 516 10.1016/j.margeo.2009.06.005.
- 517 Dorschel, B., Wheeler, A.J., Monteys, X., Verbruggen, K., 2010. Atlas of the Deep-water
  518 Seabed: Ireland. Springer, Dordrecht Heidelberg London New York DOI 10.1007/978-90-
- 519 481-9376-1.
- 520 Douarin, M., Sinclair, D.J., Elliot, M., Henry, L.-A., Long, D., Mitchison, F., Roberts, J.M.,
- 521 2014. Changes in fossil assemblage in sediment cores from Mingulay Reef Complex (NE
- 522 Atlantic): Implications for coral reef build-up. Deep Sea Research Part II: Topical Studies in
- 523 Oceanography 99, 286-296, <u>http://dx.doi.org/10.1016/j.dsr2.2013.07.022</u>.
- 524 Dullo, C., Flögel, S., Rüggeberg, A., 2008. Cold-water coral growth in relation to the
- build, e., 110ge, e., 100ge, e., 100ge, e., 2000 e.e. and 100 e.e. and 100 e.e.
  hydrography of the Celtic and Nordic European continental margin. Marine Ecology Progress
  Series 371, 165-176, 10.3354/meps07623.
- 527 Eisele, M., Hebbeln, D., Wienberg, C., 2008. Growth history of a cold-water coral covered
- 528 carbonate mound Galway Mound, Porcupine Seabight, NE-Atlantic. Marine Geology 253
- 529 (3-4), 160-169, 10.1016/j.margeo.2008.05.006.
- 530 Ferrer, J., Elibol, A., Delaunoy, O., Gracias, N., Garcia, R., 2007. Large-area photo-mosaics
- 531 using global alignment and navigation data. MTS/IEEE OCEANS Conference, Vancouver,
- 532 Canada, pp. 1-9<sup>,</sup> <u>http://dx.doi.org/10.1109/OCEANS.2007.4449367</u>.
- 533 Foubert, A.T.G., Beck, T., Wheeler, A.J., Opderbecke, J., Grehan, A., Klages, M., Thiede, J.,
- Henriet, J.-P., Polarstern ARK-XIX/3a shipboard party, 2005. New view of the Belgica
- 535 Mounds, Porcupine Seabight, NE Atlantic: preliminary results from the Polarstern ARK-
- 536 XIX/3a ROV cruise. In: Freiwald, A., Roberts, J.M. (Eds.), Deep-water corals and
- 537 Ecosystems, . Springer-Verlag, Berlin Heidelberg, pp. 403-415,
- 538 Foubert, A.T.G., Depreiter, D., Beck, T., Maignien, L., Pannemans, B., Frank, N., Blamart,
- 539 D., Henriet, J.-P., 2008. Carbonate mounds in a mud volcano province off north-west
- 540 Morocco: Key to processes and controls. Marine Geology 248, 74-96,
- 541 Foubert, A.T.G., Huvenne, V.A.I., Wheeler, A.J., Kozachenko, M., Opderbecke, J., Henriet,
- 542 J.-P., 2011. The Moira Mounds, small cold-water coral mounds in the Porcupine Seabight,
- 543 NE Atlantic: Part B Evaluating the impact of sediment dynamics through high-resolution
- 544 ROV-borne bathymetric mapping Marine Geology 282 (1-2), 65-78,
- 545 doi:10.1016/j.margeo.2011.02.008
- 546 Freiwald, A., 2002. Reef-Forming Cold-Water Corals. In: Wefer, G., Billett, D.S.M.,
- Hebbeln, D., Jørgensen, B.B., van Weering, T.C.E. (Eds.), Ocean Margin Systems. Springer,
- 548 Berlin Heidelberg New York, pp. 365-385, <u>http://dx.doi.org/10.1007/978-3-662-05127-6\_23</u>.
- Freiwald, A., Fosså, J.H., Grehan, A., Koslow, T., Roberts, J.M., 2004. Cold-water coral
  reefs. UNEP-WCMC, Cambridge, UK 84,
- Getis, A., Ord, J.K., 1992. The Analysis of Spatial Association by Use of Distance Statistics.
  Geographical Analysis 24 (3), 189-206, 10.1111/j.1538-4632.1992.tb00261.x.
- Gischler, E., 1995. Current and Wind Induced Facies Patterns in a Devonian Atoll: Iberg
- 554 Reef, Harz Mts., Germany. Palaios 10 (2), 180-189, 10.2307/3515181.

- 555 Goodchild, M., 1986. Spatial Autocorrelation. Concepts and Techniques in Modern
- 556 Geography 47. Norwich, UK: Geo Books,
- 557 <u>https://alexsingleton.files.wordpress.com/2014/09/47-spatial-aurocorrelation.pdf</u>
- Guinan, J., Grehan, A.J., Dolan, M.F.J., Brown, C., 2009. Quantifying relationships between
  video observations of cold-water coral cover and seafloor features in Rockall Trough, west of
- Ireland. Marine Ecology Progress Series 375, 125-138, 10.3354/meps07739.
- 561 Hebbeln, D., Samankassou, E., 2015. Where did ancient carbonate mounds grow In
- bathyal depths or in shallow shelf waters? Earth-Science Reviews 145, 56-65,
- 563 <u>http://dx.doi.org/10.1016/j.earscirev.2015.03.001</u>.
- Hebbeln, D., Wienberg, C., Wintersteller, P., Freiwald, A., Becker, M., Beuck, L., Dullo, C.,
  Eberli, G.P., Glogowski, S., Matos, L., 2014. Environmental forcing of the Campeche cold-
- 566water coral province, southern Gulf of Mexico.
- 567 Heindel, K., Titschack, J., Dorschel, B., Huvenne, V.A.I., Freiwald, A., 2010. The sediment
- 568 composition and predictive mapping of facies on the Propeller Mound—A cold-water coral
- mound (Porcupine Seabight, NE Atlantic). Continental Shelf Research 30 (17), 1814-1829,
- 570 doi:10.1016/j.csr.2010.08.007.
- Henriet, J.-P., De Mol, B., Pillen, S., Vanneste, M., Van Rooij, D., Versteeg, W., Croker,
- P.F., Shannon, P.M., Unnithan, V., Bouriak, S., Chachkine, P., BELGICA 97 Shipboard
  scientific crew, 1998. Gas hydrate crystals may help build reefs. Nature 391 (6668), 648-649,
  Doi 10.1038/35530.
- Huang, Z., Brooke, B.P., Harris, P.T., 2011. A new approach to mapping marine benthic
- habitats using physical environmental data. Continental Shelf Research 31 (2, Supplement),
  S4-S16, <u>http://dx.doi.org/10.1016/j.csr.2010.03.012</u>.
- 578 Huvenne, V.A., Tyler, P.A., Masson, D.G., Fisher, E.H., Hauton, C., Huhnerbach, V., Le
- Bas, T.P., Wolff, G.A., 2011. A picture on the wall: innovative mapping reveals cold-water
- coral refuge in submarine canyon. PLoS ONE 6 (12), e28755, 10.1371/journal.pone.0028755.
- Huvenne, V.A.I., Bett, B.J., Masson, D.G., Le Bas, T.P., Wheeler, A.J., 2016. Effectiveness
- 582 of a deep-sea cold-water coral Marine Protected Area, following eight years of fisheries
- closure. Biological Conservation 200, 60-69, 10.1016/j.biocon.2016.05.030.
- Huvenne, V.A.I., Beyer, A., de Haas, H., Dekindt, K., Henriet, J.-P., Kozachenko, M., Olu-
- Le Roy, K., Wheeler, A.J., participants, T.P.c., participants, C.c., 2005. The seabed
- appearance of different coral bank provinces in the Porcupine Seabight, NE Atlantic: results
- from sidescan sonar and ROV seabed mapping. In: Freiwald, A., Roberts, J.M. (Eds.), Cold-
- water Corals and Ecosystems. Springer-Verlag, Berlin Heidelberg, pp. 535-569,
- 589 <u>http://dx.doi.org/10.1007/3-540-27673-4\_27</u>.
- 590 Huvenne, V.A.I., Blondel, P., Henriet, J.-P., 2002. Textural analyses of sidescan sonar
- imagery from two mound provinces in the Porcupine Seabight. Marine Geology 189 (3-4),
  323-341, Pii Doi 10.1016/S0025-3227(02)00420-6.
- Huvenne, V.A.I., Van Rooij, D., De Mol, B., Thierens, M., O'Donnell, R., Foubert, A.T.G.,
  2009. Sediment dynamics and palaeo-environmental context at key stages in the Challenger

- cold-water coral mound formation: Clues from sediment deposits at the mound base. Deep
  Sea Research Part I 56 (12), 2263-2280, 10.1016/j.dsr.2009.08.003.
- 597 IODP 307 Expedition scientists, 2005. Modern carbonate mounds: Porcupine drilling. IODP
   598 Prel. Rept., 307. p. 58, doi:10.2204/iodp.pr.307.2005
- Janßen, A., Wizemann, A., Klicpera, A., Satari, D.Y., Westphal, H., Mann, T., 2017.
- 600 Sediment Composition and Facies of Coral Reef Islands in the Spermonde Archipelago,
- Indonesia. Frontiers in Marine Science 4 (144), 10.3389/fmars.2017.00144.
- 602 Kano, A., Ferdelman, T.G., Williams, T., Henriet, J.-P., Ishikawa, T., Kawagoe, N.,
- Takashima, C., Kakizaki, Y., Abe, K., Sakai, S., Browning, E.L., Li, X., Andres, M.S.,
- Bjerager, M., Cragg, B.A., De Mol, B., Dorschel, B., Foubert, A.T.G., Frank, T.D., Fuwa, Y.,
- 605 Gaillot, P., Gharib, J.J., Gregg, J.M., Huvenne, V.A.I., Léonide, P., Mangelsdorf, K.,
- 606 Monteys, X., Novosel, I., O'Donnell, R., Rüggeberg, A., Samarkin, V.A., Sasaki, K.,
- 607 Spivack, A.J., Tanaka, A., Titschack, J., van Rooij, D., Wheeler, A.J., 2007. Age constraints
- 608 on the origin and growth history of a deep-water coral mound in the northeast Atlantic drilled 609 during Integrated Ocean Drilling Program Expedition 307. Geology 35 (11), 1051-1054, doi:
- 609 during integrated Ocean Drining Program Expedition 507. Geology 55 (11), 1051-1054,
   610 10.1130/G23917A.
- 611 Kocak, D.M., Caimi, F.M., 2005. The Current Art of Underwater Imaging With a
- 612 Glimpse of the Past and Vision of the Future. Marine Technology Society Journal 39 (3), 5-
- **613 26**, 10.4031/002533205787442576.
- Kozachenko, M., 2005. Present and Past Environments of the Belgica Mounds (deep-water
- 615 coral carbonate mounds) Eastern Porcupine Seabight, North East Atlantic (unpublished PhD
   616 Thesis), University College Cork, Cork,
- 617 Kwatra, V., Schödl, A., Essa, I., Turk, G., Bobick, A., 2003. Graphcut textures: image and
- Kwatra, V., Schodi, A., Essa, I., Turk, G., Bobick, A., 2003. Graphcut textures: image and
  video synthesis using graph cuts. ACM Transactions on Graphics (TOG). ACM, pp. 277286<sup>^</sup>, http://dx.doi.org/10.1145/882262.882264.
- 620 Lamarche, G., Lurton, X., Verdier, A.-L., Augustin, J.-M., 2011. Quantitative
- 621 characterisation of seafloor substrate and bedforms using advanced processing of multibeam
- backscatter—Application to Cook Strait, New Zealand. Continental Shelf Research 31 (2,
- 623 Supplement), S93-S109, <u>http://dx.doi.org/10.1016/j.csr.2010.06.001</u>.
- Lim, A., 2017. Spatio-temporal patterns and controls on cold-water coral reef development: the Moira Mounds, Porcupine Seabight, NE Atlantic, University College Cork, Cork Open
- 626 Research Archive, <u>http://hdl.handle.net/10468/4031</u>.
- 627 Lirman, D., Gracias, N.R., Gintert, B.E., Gleason, A.C.R., Reid, R.P., Negahdaripour, S.,
- Kramer, P., 2007. Development and application of a video-mosaic survey technology to
- 629 document the status of coral reef communities. Environmental Monitoring and Assessment
- 630 125 (1), 59-73, 10.1007/s10661-006-9239-0.
- 631 Lowe, D.G., 1999. Object recognition from local scale-invariant features. Computer vision,
- 1999. The proceedings of the seventh IEEE international conference on Computer Vision.
  Ieee, pp. 1150-1157<sup>^</sup>, http://dx.doi.org/10.1109/ICCV.1999.790410.
- 634 Messing, C.G., Neumann, A.C., Lang, J.C., 1990. Biozonation of deep-water lithoherms and 635 associated hardgrounds in the northeastern Straits of Florida. Palaios 5, 15-33,

- Mienis, F., de Stigter, H.C., de Haas, H., van Weering, T.C.E., 2009. Near-bed particle
- 637 deposition and resuspension in a cold-water coral mound area at the Southwest Rockall
- Trough margin, NE Atlantic. Deep Sea Research Part I: Oceanographic Research Papers 56
- 639 (6), 1026-1038, <u>http://dx.doi.org/10.1016/j.dsr.2009.01.006</u>.
- Mullins, H.T., Newton, C.R., Heath, K., Van Buren, H.M., 1981. Modern deep-water coral
- mounds north of Little Bahama Bank; criteria for recognition of deep-water coral bioherms in
  the rock record. Journal of sedimentary research 51 (3), 999-1013, 10.1306/212f7dfb-2b24-
- 643 11d7-8648000102c1865d.
- Murphy, P., Wheeler, A.J., 2017. A GIS-based application of drainage basin analysis and
- 645 geomorphometry in the submarine environment: The Gollum Canyon System, North-east
- 646 Atlantic. In: Bartlett, D., Celliers, L. (Eds.), Geoinformatics for Marine and Coastal
- 647 Management. CRC Press, Taylor & Francis Group, Boca Raton, USA,
- 648 <u>http://dx.doi.org/10.1201/9781315181523-4</u>.
- 649 Ord, J.K., Getis, A., 1995. Local Spatial Autocorrelation Statistics: Distributional Issues and
- 650 an Application. Geographical Analysis 27 (4), 286-306, 10.1111/j.1538-
- 651 4632.1995.tb00912.x.
- 652 Orejas, C., Gori, A., Rad-Menendez, C., Last, K.S., Davies, A.J., Beveridge, C.M., Sadd, D.,
- Kiriakoulakis, K., Witte, U., Roberts, J.M., 2016. The effect of flow speed and food size on
- the capture efficiency and feeding behaviour of the cold-water coral Lophelia pertusa. Journal
   of Experimental Marine Biology and Ecology 481, 34-40, 10.1016/j.jembe.2016.04.002.
- Purser, A., 2015. A Time Series Study of Lophelia pertusa and Reef Megafauna Responses to
- Drill Cuttings Exposure on the Norwegian Margin. PLoS ONE 10 (7), e0134076,
- 658 10.1371/journal.pone.0134076.
- Purser, A., Larsson, A.I., Thomsen, L., van Oevelen, D., 2010. The influence of flow velocity
   and food concentration on *Lophelia pertusa* (Scleractinia) zooplankton capture rates. Journal
- of Experimental Marine Biology and Ecology 395 (1-2), 55-62,
- 662 10.1016/j.jembe.2010.08.013.
- Reolid, J., Betzler, C., Braga, J.C., Martín, J.M., Lindhorst, S., Reijmer, J.J.G., 2014. Reef
  slope geometries and facies distribution: controlling factors (Messinian, SE Spain). Facies 60
  (3), 737-753, 10.1007/s10347-014-0406-4.
- Roberts, J.M., Brown, C.J., Long, D., Bates, C.R., 2005. Acoustic mapping using a
  multibeam echosounder reveals cold-water coral reefs and surrounding habitats. Coral Reefs
  24 (4), 654-669, 10.1007/s00338-005-0049-6.
- 669 Roberts, J.M., Long, D., Wilson, J.B., Mortensen, P.B., Gage, J.D., 2003. The cold-water
- 670 coral Lophelia pertusa (Scleractinia) and enigmatic seabed mounds along the north-east
- 671 Atlantic margin: are they related? Marine Pollution Bulletin 46 (1), 7-20,
- 672 <u>http://dx.doi.org/10.1016/S0025-326X(02)00259-X</u>.
- Roberts, J.M., Wheeler, A., Freiwald, A., Cairns, S., 2009. Cold-Water Corals: The Biology
- and Geology of Deep-Sea Coral Habitats. Cambridge University Press,
- 675 <u>http://books.google.ie/books?id=6WA3YIH27t0C</u>

- Roberts, J.M., Wheeler, A.J., Freiwald, A., 2006. Reefs of the Deep: The Biology and
- 677 Geology of Cold-Water Coral Ecosystems. Science 312 (5773), 543-547,
- 678 <u>http://dx.doi.org/10.1126/science.1119861</u>.
- Rüggeberg, A., Flögel, S., Dullo, W.-C., Hissmann, K., Freiwald, A., 2011. Water mass
- 680 characteristics and sill dynamics in a subpolar cold-water coral reef setting at Stjernsund,
- 681 northern Norway. Marine Geology 282 (1–2), 5-12,
- 682 <u>http://dx.doi.org/10.1016/j.margeo.2010.05.009</u>.
- Rzhanov, Y., Linnett, L.M., Forbes, R., 2000. Underwater video mosaicing for seabed
  mapping. Image Processing, 2000. Proceedings. 2000 International Conference on, pp. 224227 vol.221<sup>^</sup>, 10.1109/ICIP.2000.900935.
- Savini, A., Vertino, A., Marchese, F., Beuck, L., Freiwald, A., 2014. Mapping cold-water
  coral habitats at different scales within the Northern Ionian Sea (Central Mediterranean): an
  assessment of coral coverage and associated vulnerability. PLoS ONE 9 (1), e87108,
  10.1371/journal.pone.0087108.
- 690 Spezzaferri, S., Vertino, A., Addamo, A.M., Backers, J., Baratti, V., Constandache, M.,
- 691 Camozzi, O., El Kateb, A., Gennari, G., Leuzinger, L., McGrath, M., Mc Hugh, S.K., Naudts,
- L., Savini, A., Stalder, C., 2012. E-CWC Moira (2012/16) *RV Belgica*: Cold-water coral
- 693 ecosystems from the Moira Mounds (NE Atlantic): affinities and differences with modern
- and Pleistocene Mediterranean counterparts. University Freibourg, Switzerland, UnpublishedCruise Report,
- Squires, D.F., 1964. Fossil coral thickets in Wairarapa, New Zealand. Journal Paleontol. 38
  (5), 904-915, <u>http://www.jstor.org/stable/1301611</u>
- Thierens, M., Titschack, J., Dorschel, B., Huvenne, V.A.I., Wheeler, A.J., Stuut, J.-B.W.,
- 699 O'Donnell, R., 2010. The 2.6 Ma depositional sequence from the Challenger cold-water coral
- carbonate mound (IODP Exp. 307): sediment contributors and hydrodynamic palaeo-
- rot environments. Marine Geology 271 (3-4), 260-277, doi:10.1016/j.margeo.2010.02.021.
- 702 Titschack, J., Thierens, M., Dorschel, B., Schulbert, C., Freiwald, A., Kano, A., Takashima,
- C., Kawagoe, N., Li, X., 2009. Carbonate budget of a cold-water coral mound (Challenger
- Mound, IODP Exp. 307). Marine Geology 259 (1-4), 36-46, 10.1016/j.margeo.2008.12.007.
- Van Rooij, D., 2004. An integrated study of Quaternary sedimentary processes on the eastern
   slope of the Porcupine Seabight, SW of Ireland, Ghent University,
   http://dx.doi.org/1854/10815
- 707 <u>http://dx.doi.org/1854/10815</u>.
- Vertino, A., Savini, A., Rosso, A., Di Geronimo, I., Mastrototaro, F., Sanfilippo, R., Gay, G.,
- Etiope, G., 2010. Benthic habitat characterization and distribution from two representative
- sites of the deep-water SML Coral Province (Mediterranean). Deep-Sea Research Part Ii-
- 711 Topical Studies in Oceanography 57 (5-6), 380-396, 10.1016/j.dsr2.2009.08.023.
- 712 Vertino, A., Spezzaferri, S., Ruggeberg, A., Stalder, C., Wheeler, A.J., Party, t.E.C.-M.c.S.,
- 713 2015. An Overview on Cold-Water Coral Ecosystems and Facies. In: Spezzaferri, S.,
- Ruggeberg, A., Stalder, C. (Eds.), Atlas of benthic foraminifera from cold-water coral reefs.
- 715 Cushman Foundation for Foraminiferal Research, pp. 12-19,

- 716 Wheeler, A.J., Beyer, A., Freiwald, A., de Haas, H., Huvenne, V.A.I., Kozachenko, M., Olu-
- Le Roy, K., Opderbecke, J., 2007. Morphology and environment of cold-water coral
- carbonate mounds on the NW European margin. International Journal of Earth Sciences 96
- 719 (1), 37-56, <u>http://dx.doi.org/10.1007/s00531-006-0130-6</u>.
- 720 Wheeler, A.J., Capocci, R., Crippa, L., Connolly, N., Hogan, R., Lim, A., McCarthy, E.,
- 721 McGonigle, C., O' Donnell, E., O' Sullivan, K., Power, K., Ryan, G., Vertino, A., Holland 1
- 722 ROV Technical Team, Officers and Crew of the RV Celtic Explorer, 2015. Cruise Report:
- 723 Quantifying Environmental Controls on Cold-Water coral Reef Growth (QuERCi).
- 724 University College Cork, Ireland,
- 725 Wheeler, A.J., Kozachenko, M., Beyer, A., Foubert, A.T.G., Huvenne, V.A.I., Klages, M.,
- Masson, D.G., Olu-Le Roy, K., Thiede, J., 2005. Sedimentary processes and carbonate
- mounds in the Belgica Mound province, Porcupine Seabight, NE Atlantic. In: Freiwald, A.,
- Roberts, J.M. (Eds.), Cold-water Corals and Ecosystems. Springer-Verlag, Berlin Heidelberg,
- 729 pp. 533-564, <u>http://dx.doi.org/10.1007/3-540-27673-4\_28</u>.
- 730 Wheeler, A.J., Kozachenko, M., Henry, L.A., Foubert, A., de Haas, H., Huvenne, V.A.I.,
- Masson, D.G., Olu, K., 2011. The Moira Mounds, small cold-water coral banks in the
- 732 Porcupine Seabight, NE Atlantic: Part A—an early stage growth phase for future coral
- carbonate mounds? Marine Geology 282 (1-2), 53-64, 10.1016/j.margeo.2010.08.006.
- Wheeler, A.J., Kozachenko, M., Masson, D.G., Huvenne, V.A.I., 2008. Influence of benthic
  sediment transport on cold-water coral bank morphology and growth: the example of the
  Darwin Mounds, north-east Atlantic. Sedimentology 55 (6), 1875-1887, 10.1111/j.13653091.2008.00970.x.
- Wheeler, A.J., shipboard party, 2011. Vents & Reefs deep-sea ecosystem study of the 45°
  North MAR hydrothermal vent field and the cold-water coral Moira Mounds, Porcupine
  Seabight. Cruise report, p. 160,
- White, M., Dorschel, B., 2010. The importance of the permanent thermocline to the cold
  water coral carbonate mound distribution in the NE Atlantic. Earth and Planetary Science
  Letters 296, 395-402,
- 744 Wienberg, C., Beuck, L., Heidkamp, S., Hebbeln, D., Freiwald, A., Pfannkuche, O., Monteys,
- F.X., 2008. Franken Mound: facies and biocoenoses on a newly-discovered "carbonate
  mound" on the western Rockall Bank, NE Atlantic. Facies 54 (1), 1-24, 10.1007/s10347-007-
- 747 0118-0.
  - 748 Wienberg, C., Hebbeln, D., Fink, H.G., Mienis, F., Dorschel, B., Vertino, A., López Correa,
- 749 M., Freiwald, A., 2009. Scleractinian cold-water corals in the Gulf of Cádiz-first clues about
- their spatial and temporal distribution. Deep Sea Research Part I 56 (10), 1873-1893,
  doi:10.1016/j.dsr.2009.05.016.
- Wienberg, C., Wintersteller, P., Beuck, L., Hebbeln, D., 2013. Coral Patch seamount (NE
- Atlantic) a sedimentological and megafaunal reconnaissance based on video and
- 754 hydroacoustic surveys. Biogeosciences 10, 3421-3443, 10.5194/bg-10-3421-2013.

Wilson, J.B., 1979. "Patch" development of the deep-water coral *Lophelia pertusa* (L.) on the

- Rockall bank. Journal of the Marine Biological Association of the United Kingdom 59 (1),
- 757 165-177, http://dx.doi.org/10.1017/S0025315400046257.

761

- 758 Woodroffe, C.D., Biribo, N., 2011. Atolls. In: Hopley, D. (Ed.), Encyclopedia of Modern
- Coral Reefs: Structure, Form and Process. Springer Netherlands, Dordrecht, pp. 51-71,
  10.1007/978-90-481-2639-2 4.



Figure 1 A) Map showing the Belgica Mound Province (BMP) Special Area of Conservation
(SAC) (red box) within the Porcupine Seabight relative to Ireland; B) the subdivision of Moira
Mounds with the position of the Piddington Mound (yellow star) with side scan sonar data in
the background (white tones are high backscatter, darker tones are low backscatter), Red

dots= Moira Mounds identified on 30 kHz side-scan sonar data; blue dots=Moira Mounds
identified on 100 kHz side-scan sonar data, green dots= Moira Mounds identified on 410 kHz
side-scan sonar data; yellow dots= Moira Mounds groundtruthed by ROV and identified on



100 and 410 kHz side-scan sonar after Wheeler et al. (2011); C) the subdivision of Moira

770 Mounds after Wheeler et al. (2011); D) bathymetry of the Piddington Mound area.

Figure 2 A) Map of georeferenced video mosaic with relative position of close-up images
(yellow boxes) with 1 m contours superimposed; B) example close-up image of area
dominated by "hemipelagic sediment and dropstone" cells with ArcGIS fishnet overlaid; C)
example close-up image of area dominated by "hemipelagic sediment" cells with ArcGIS

fishnet overlaid; D) example close-up image of area dominated by "coral rubble" cells with
ArcGIS fishnet overlaid; E) example close-up image of area dominated by "dead coral
framework" cell with ArcGIS fishnet overlaid; F) example close-up image of area dominated
by "live coral framework" cells with ArcGIS fishnet overlaid.



Figure 3 A) Bathymetry of the Piddington Mound (centre) area with coral ridge feature position
identified (black box); B) ArcGIS-generated slope of the Piddington Mound area. Relatively

- flat areas in blue, relatively steep areas in red and line X-Y (Fig. 3C) indicated; C) Cross profile
- of Piddington Mound with facies type plotted across the surface.



Figure 4 Image of ridge-form feature, covered with corals to the south of Piddington Mound.
View to the north. Laser scalers = 11 cm. Ripple marks on the camera-facing side of the ridge
indicate a northerly-directed current.



Figure 5 Classified fishnet \*.shp file with contours superimposed showing the distribution of
facies types across the Piddington Mound mosaic (red = hemipelagic sediment with dropstones,
orange = hemipelagic sediment, blue = coral rubble, purple = dead coral framework, and pink
= live coral framework).



Figure 6 A) Map showing the focal statistics layer which shows the 4 identified facies occurring
in 5 areas. Each facies represents where a cell-type is most dominant within a 2.5 m radius
neighbourhood (red = "Hemipelagic sediment with dropstones" facies, blue = "Coral rubble"
facies, dark pink = "Dead coral framework" facies, light pink = "Live coral framework" facies);
B) Pie charts showing the relative proportion and percentages of cells that define each facies.



805

Figure 7 A) map showing the distribution of facies across Piddington Mound with 1 m contours overlaid (red = hemipelagic sediment and dropstone facies, blue = coral rubble facies, purple = dead coral framework facies, pink = live coral framework facies) B) Simplified drawing of Piddington Mound with differentiation between live and dead coral frameworks (red = hemipelagic sediment and dropstone facies, blue = coral rubble facies, purple = combined coral framework facies, R1 & R2 = coral rubble facies 2, F1 = dead coral framework facies, F2 = live coral framework facies, arrow = current direction).