

Title	Food for thought: formulating away the food effect - a PEARRL review
Authors	O'Shea, Joseph P.;Holm, René;O'Driscoll, Caitríona M.;Griffin, Brendan T.
Publication date	2018-06-28
Original Citation	O'Shea, J. P., Holm, R., O'Driscoll, C. M. and Griffin, B. T. 'Food for thought: formulating away the food effect – a PEARRL review', Journal of Pharmacy and Pharmacology, 71(4), pp.510-535 doi: 10.1111/jphp.12957
Type of publication	Article (peer-reviewed)
Link to publisher's version	https://onlinelibrary.wiley.com/doi/abs/10.1111/jphp.12957 - 10.1111/jphp.12957
Rights	© 2018 Royal Pharmaceutical Society. This is the peer reviewed version of the following article: O'Shea, J. P., Holm, R., O'Driscoll, C. M. and Griffin, B. T. (2018), Food for thought: formulating away the food effect – a PEARRL review. J Pharm Pharmacol., which has been published in final form at https://doi.org/10.1111/jphp.12957 . This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Self-Archiving.
Download date	2026-09-25 18:27:02
Item downloaded from	https://hdl.handle.net/10468/7108



UCC

University College Cork, Ireland
Coláiste na hOllscoile Corcaigh

Food for thought: formulating away the food effect- a PEARRL Review

Joseph P. O'Shea¹, René Holm², Caitriona M. O'Driscoll¹, Brendan T. Griffin^{1,*}

¹ *School of Pharmacy, University College Cork; Ireland*

² *Drug Product Development, Janssen Research and Development, Johnson & Johnson, Turnhoutseweg 30, 2340 Beerse, Belgium*

*Correspondence to:

Abstract

Objectives: Co-ingestion of oral dosage forms with meals can cause substantial changes in the rate and extent of drug absorption relative to the fasted state. Food mediated effects on bioavailability can have significant consequences in drug development, regulatory and clinical settings. To date the primary focus of research and analysis has focused on the ability to mechanistically understand the causes and predict the occurrence of these effects.

Key findings: The current review describes the mechanisms underpinning the occurrence of food effects, sheds new insights on the relative frequency that these effects occur in new medicinal products and describes the various methods by which they can be overcome. Analysis of new oral medicines licensed by either the EMA or FDA since 2010 revealed that over 40% of new medicinal products display significant food effects. Due to altered bioavailability, these medicines are often required to be dosed, rather restrictively, in either the fed or the fasted state, which can hinder clinical usefulness.

Conclusions: There are clinical and commercial advantages to predicting the presence of food effects early in the drug development process, in order to mitigate this risk of variable food effect bioavailability. Formulation approaches aimed at reducing variable food-dependent bioavailability, through the use of bio-enabling formulations, are an essential tool in addressing this challenge and the latest state-of-the-art in this field are summarised here

Key words

Food effect; bioavailability; bio-enabling formulations; food-drug interactions; formulation screening; gastrointestinal physiology

Introduction

The concomitant administration of oral dosage forms with food can have a significant impact on drug pharmacokinetics and bioavailability relative to the fasted state. With oral drug delivery continuing to be the method of choice for drug administration, understanding the effects food has on the biopharmaceutical aspects of drug delivery is key to the drug development process as well as the effective and rational use of medicines in the clinical setting ^[1, 2]. Oral medicines are generally required to be repeatedly administered, often chronically and in multiple daily dosings, so it is inevitable that drugs will be administered in different prandial states. The understanding of the effects food has on pharmacokinetics is consequently a critical factor in assessing the clinical potential of new medicines and designing a food effect resistant formulation early in drug development can both provide a commercial advantage and prevent costly reformulation later in the product lifecycle ^[1, 3].

It is just over 40 years since the publication of the first major review focusing on the manner by which food affects drug absorption ^[4], and the topic has been subject to extensive research and review in the interim ^[5-9, 3, 10, 11]. Despite the abundance of studies examining the predictability, mechanistic understanding, and ability to overcome the effects of food on bioavailability, a universal approach to quantitatively predict food effect does not exist, nor is it a likely prospect. Significant progress has, however, been made in identifying potential drug candidates and drug products that display food effect bioavailability, understanding the mechanisms by which food effects occurs and developing formulations to overcome this effect.

While there has been comprehensive review and analysis of the mechanisms underlying the food effect, and more recently of current approaches to predict food effect (Pentafragka *et al.*, this issue)^[12], to date there has been limited analysis of the relative abundance of medicines which display food effect and the systematic approaches utilised to eliminate food mediated changes in bioavailability. The aims of the current review are, therefore, to briefly summarise the main causes of food mediated changes in bioavailability, discuss the clinical and regulatory impact with regard to the types of and abundance of preparations which display significant food effects and to describe the various formulation approaches currently implemented to overcome the food effect. To our knowledge this is the first review to focus primarily on the use of enabling formulations to overcome

59 food effects on bioavailability in clinical and pre-clinical studies, while it also provides an updated
60 compilation of recently licensed medicines which demonstrate significant food effect.

61

Food Effects; causes and clinical consequences

What is a food-effect?

In its simplest terms, food effects on drug absorption are observed when the rate and/or extent of drug bioavailability is altered when a drug or drug product is administered in fed state, compared to the fasted state. The clinical effects and significance of food effects on absorption are generally assessed with regard to the rate and extent of bioavailability – as measured by peak plasma concentrations ($C_{\max}$), time to peak plasma concentration ($T_{\max}$) and the total extent of bioavailability (area under the curve; AUC) ^[1]. Welling classified food drug interactions into five categories causing ^[5],

- Reduced extent of bioavailability
- Delayed rate of absorption
- Increased extent of bioavailability
- Accelerated rate of absorption
- No effect

With regard to clinical significance, the most crucial aspect of food effect is generally considered to be the extent of bioavailability change, and the terms ‘positive food effect’ and ‘negative food effect’ have been coined to describe either an increase or decrease in the overall extent of bioavailability, respectively^[1]. While some variation in bioavailability is tolerated, larger deviations in the fed, relative to the fasted state can have clinical implications. It is, thus, necessary to have some guidance on defining what exactly constitutes a significant food effect. Accordingly, the FDA have provided guidelines on how to design clinical trials to investigate food effects, recommending dosing in both fasted and fed states. The FDA guidance defines that a food-effect is established if the 90% confidence intervals for the ratio of population geometric means, based on log-transformed data, for either $AUC_{0 \rightarrow \infty}$ or $C_{\max}$ fall outside the 80-125% bioequivalence limits relative to the reference, i.e. the same formulation administered in the fasted state ^[13]. The fed state represents dosing post ingestion of a high fat, FDA standard breakfast, containing 800 – 1000 kcal with approximately 50% of total calories coming from fat, to maximise potential for demonstrating a food effect ^[13].

Figure 1 illustrates the key steps in drug absorption and bioavailability and indicates how food influences these processes, while the underlying mechanisms of these processes are described in subsequent sections of this review.




Figure 1 Schematic diagram of critical steps in drug absorption and influence of food and food components; FPM: first pass metabolism

Mechanisms underlying the food effect

Drug absorption via the oral route is a function of the interplay of various complex biopharmaceutical processes, namely (i) drug molecular and physicochemical properties, (ii) formulation characteristics, (iii) the physiological changes of the gastrointestinal tract induced in the fed state and (iv) the physical chemical changes in the composition of the gastrointestinal fluid ^[1]. The Biopharmaceutical Classification System (BCS) and Biopharmaceutical Drug Disposition Classification system (BDDCS) provide a useful predictor of potential food effects based on drug physicochemical properties, as summarised in figure 2 ^[14, 15]. The anticipated effects are predicted by the most likely limiting factor for bioavailability, namely solubility or dissolution for BCS/BDDCS class II compounds, permeability for class III compounds, or a combination thereof for BCS class IV compounds. An overall delay in T_{max} and reduced C_{max} for highly bioavailable compounds can be associated with a delayed gastric emptying ^[16]. While this tool does not capture all the potential effects of food, it is the most widely utilised simple tool to predict food effect behaviour, and is estimated to be accurate in approximately 70% of cases ^[17].

	High Solubility	Low Solubility
High Permeability/Metabolism	Class I $F_{\text{extent}} \leftrightarrow$ $T_{\text{max}} \uparrow$ $C_{\text{max}} \downarrow$	Class II $F_{\text{extent}} \uparrow$ $T_{\text{max}} \uparrow \downarrow \leftrightarrow$ $C_{\text{max}} \uparrow$
Low Permeability/Metabolism	Class III $F_{\text{extent}} \downarrow$ $T_{\text{max}} \uparrow$ $C_{\text{max}} \downarrow$	Class IV $F_{\text{extent}} \uparrow \downarrow \leftrightarrow$ $T_{\text{max}} \uparrow \downarrow \leftrightarrow$ $C_{\text{max}} \uparrow \downarrow \leftrightarrow$

Figure 2 Predicted effect of high fat meals by BCS/BDDCS class. Adapted from Custodio *et al.* (2008)

Drug absorption is inherently variable, owing to both inter- and intra-individual variability in the physiology of the GIT. When considering the gut physiology McConnell *et al.* have stated that there is ‘no such thing as an average person’^[18], and despite regulatory guidance, equally there is no such thing as a standard meal^[13]. The purpose of FDA guidance is to provide a standard for bioavailability and bioequivalence studies, where the likelihood of observing a food effect is maximised. However, this is not always reflective of the fed state for patients, which adds further to the variability and complexity of absorption and drug product performance.

In the fed state the physicochemical composition of the gastrointestinal fluid, including its volume, pH, osmolality, surface tension, hydrodynamics and overall composition change. These changes have been extensively reviewed by Pentafragka *et al.* in the current issue^[12]. The reader is directed here for greater detail, specifically with regard to the intraluminal environment after intake of meals similar in composition to that suggested by the FDA and EMA for food effect and fed state bioequivalence studies i.e. a high-fat, high-caloric meal^[13, 19]. There are a number of additional factors that may influence GIT absorption from oral dosage forms and the most pertinent aspects are described below, and summarised in figure 3.

Food induced changes on drug absorption

Gastrointestinal fluid composition

With regard to drug absorption of immediate release dosage forms, it is the characteristics of the stomach and upper intestine which are generally most crucial for drug absorption. Relative to the fasted state, the most pertinent changes in the intraluminal environment include the increase in gastric pH to 5 or higher, along with a corresponding increase in buffer capacity^[20-22]. The intragastric fluid volume also increases significantly in the fed state, with increases in the presence of dietary lipids and their digestion products along with increased viscosity of the luminal contents^[23-26]. The most significant changes in the small intestinal luminal fluid composition is the increase in bile salt concentrations and the presence of lipid digestion products^[27, 24, 28-30]. The extensive absorption in the small intestine means that despite the fluid ingested with a meal and significant gastrointestinal secretions, the overall volume of fluid in the small intestine actually decreases in the fed state^[23, 25].

Gastrointestinal motility and its impact on transit time of dosage forms

The interplay between GIT motility and intestinal transit of dosage forms can be complex and affected by numerous factors. In the fasted state, emptying of liquid formulations will occur quite rapidly, whereas emptying of solids can be delayed by up to 2 hours^[31-34, 10]. In the fed state, liquids and smaller particles (<3-4mm) will empty with food, at a rate controlled by the caloric density of the food, but which is invariably slower than the fasted state^[31, 35]. Larger particles (>7mm) can be retained in the fed state, displaying a significant lag time^[36]. Fadda *et al.* have estimated the gastric transit of a non-disintegrating tablet in the fasted, fed and pre-fed state with a median (IQR) gastric emptying time of 37 (19-74) minutes, 149 (119-171) minutes and 39 (25-169) minutes in each state, respectively^[37]. Small intestinal transit time appears to be remarkably independent of fed state, and mean values are consistently reported to be 3-4 hours^[38, 37]. However, this mean value masks considerable inter- and intra-individual variation.

Metabolism and transporter effects

Both dietary components, including monoglycerides, and bile salts have been shown to have an inhibitory effect on both uptake and efflux transporters *in vitro*, regularly leading to suggestions that high fat meals may lead to inhibition of intestinal enzymes, as well as efflux and uptake transporters^[16, 39]. While many enteric enzymes are responsible for drug metabolism, it is the cytochrome P450 (CYP450) family, in particular the CYP3A and CYP2C subfamilies, which are most widely implicated in such interactions as they are the most abundant family and play a crucial role in bioavailability of a wide range of molecules^[40, 39]. The most commonly implicated transporters are

the organic anion transporter polypeptides (OATP) and P-glycoprotein (p-GP) efflux transporters^[41, 42, 16, 43, 39]. However, despite the widely cited assertion that food, generally, inhibits intestinal transporter and enzyme function, to date the clinically important interactions of note regarding enzyme and/or transporter inhibition involve specific food components. Most notable among these is the inhibition of CYP450 enzymes by grapefruit juice, while other foods rich in phytochemicals, such as fruits, herbs and red wine have also been implicated^[42, 44, 16, 39]. This constitutes a specific, and well characterised phenomenon involving an individual food component, which is not what is typically considered when discussing food effect, which usually refers to the effects of meals generally rather than the impact of individual components.




Figure 3 Summary of human physiological changes in the fed state (adapted from Varum *et al.* [3])

Mechanisms of food effect

As has been described above, food has a complex and significant effect on the physiology of the gastrointestinal tract and the physicochemical properties of gastrointestinal fluid, which in turn can have a significant effect on drug absorption. These effects are dependent on both the physicochemical properties of the drug, principally solubility, pKa and LogP/logD, and formulation characteristics, including release and disintegration of solid dosage forms [45, 46, 8]. For the purposes of this review, the focus will predominantly be on immediate release and bio-enabling formulations, the mechanisms by which food causes these changes in bioavailability are discussed here and summarised in table 1. The effect of food on modified release dosage forms can be significantly different to that of immediate release preparations, notably with regard to disintegration and release and the potential for dose dumping. The effects of food on modified release formulations have recently been reviewed by Varum *et al.* (2013)^[3], Yasuji *et al.* (2012)^[9] and Abuhelwa *et al.* (2017)^[10].

Positive food effects

The principal cause of positive food effects is the increase in dissolution and solubilisation of poorly water soluble drugs (PWSD) in the fed state. The release of bile salts and the presence of exogenous solubilising species, such as ingested lipids and their digestion products serve to enhance solubilising capacity of gastrointestinal fluid [28, 47, 30, 48-50]. For drugs which are dissolution rate, rather than solubility limited, the increased gastric residence time also can improve bioavailability, while the increase in gastric pH may result in improved solubility and dissolution of weak acids. In practical terms, it is difficult to isolate the impact of any one of these factors, which work synergistically to increase solubility and dissolution of PWSD.

The inhibition of intestinal transporters can play a role in enhancing bioavailability of certain drugs. Wu and Benet have demonstrated that for BCS class II compounds efflux transporters predominate, and that for these compounds transporter inhibition is likely to improve bioavailability [51, 15, 16, 52].

Reduction in first pass metabolism in the fed state can also lead to increases in bioavailability and this can occur through numerous mechanisms including altered blood flow, increased lymphatic uptake and reduced enteric metabolism. Food intake is associated with an increase in splanchnic blood flow by as much as 60% depending on the volume and nature of the meal. This allows drug to bypass the liver, while the increase in hepatic blood flow may also reduce the first pass effect for drugs which display low to moderate clearance [53-55]. Co-administering lipophilic drugs with food allows efficient absorption of these molecules with dietary lipids, via lipid absorption pathways, while particularly lipophilic drugs (logP>5) can also show significant lymphatic uptake [56, 30, 57]. This

208 can increase the systemic absorption by both increasing the fraction escaping the gastrointestinal
209 lumen and reducing the first pass effect.

210 The inhibitory effect of meal components on CYP3A4 is also a significant contributor to the reduction
211 of enteric drug metabolism and increased bioavailability in the fed state. Inhibition of CYP3A
212 metabolism by grapefruit juice has been widely associated with increases in bioavailability and
213 subsequent increases in adverse events for a wide range of pharmacologically diverse compounds ^{[40,}
214 ^{16]}. While both dietary monoglycerides and bile salts have been demonstrated to have an inhibitory
215 effect on enzymatic activity *in vitro*, and it has regularly been asserted that this inhibition leads to
216 clinically relevant enzyme inhibition by high fat meals such interactions have not yet been
217 extensively characterised *in vivo* ^[42, 16, 39, 3].

218 *Negative food effects*

219 Negative food effects encompass both reduced and delayed drug absorption. With regard to delayed
220 absorption in the fed state, this often occurs for immediate release preparations without a
221 corresponding reduction in overall bioavailability. The main mechanism by which this occurs is
222 delayed gastric transit in the fed state. This manifests itself as a prolonged T_{max} , which may or may
223 not be accompanied by a reduction in C_{max} or a significant lag time. For medicines which are
224 chronically dosed and where overall exposure, rather than peak plasma levels, mediate
225 pharmacodynamic action, this is unlikely to result in clinically meaningful effects ^[7].

226 Decreased absorption in the fed state results in a reduction in AUC, along with a reduction in C_{max} ,
227 and can lead to sub-therapeutic plasma levels and loss of efficacy. The most common causes of
228 reduced bioavailability in the fed state are direct physicochemical interactions between drugs, or
229 drug products, and food. One potential cause of this effect is the reduced diffusivity of drug in the
230 viscous postprandial upper GIT. The increased viscosity can result in either inhibition of
231 disintegration of a dosage form, preventing drug release, or hindering diffusion of drug to the
232 absorptive membranes of the GIT ^[58-61]. This can be problematic for poorly permeable drugs,
233 particularly those with narrow absorption windows, as by the time viscosity has reduced in the distal
234 gut, the absorption window has been transited and absorption will be reduced ^[62-64]. A second direct
235 mechanism by which food can hinder drug absorption is by binding of drug with food components
236 ^[65, 7]. This is prevalent in the case of polyvalent cations, which are abundant in dairy products ^{[66-68, 7,}
237 ^{69]}.

238 Physiological factors can also play a role in negative food effects, especially in the case of drugs
239 displaying instability and possibly acid lability in the GIT. Prolonged gastric residence can result in
240 increased degradation of these molecules, though in the case of acid labile drugs the effect may be

241 somewhat mitigated by the increase in gastric pH ^[70]. Food can also result in alterations in
242 absorption through altering both passive permeability and active transport. The presence of
243 increased lipids and bile salts in the fed state can result in micellar entrapment, with the consequent
244 decrease in free drug causing a reduction in permeability ^[45, 71-73, 49]. While for poorly soluble drugs,
245 this is generally more than compensated for by increases in solubility, highly soluble and poorly
246 permeable compounds may display reduced absorption in this case.

247 The inhibition of uptake transporters may also result in negative food effects. For poorly permeable
248 drugs, the inhibition of these transporters may result in a reduction in absorption, as these
249 compounds are often reliant on the action of uptake transporters. The general inhibition of
250 intestinal transporters observed in the fed state is therefore likely to reduce the bioavailability of
251 BCS class III compounds. Care is needed, however, when applying this rule of thumb, as class III
252 compounds may be candidates for both uptake and efflux transporters and the relative inhibition of
253 either uptake or efflux transporters, or the extent to which a specific molecule will be a substrate for
254 each particular class can determine the overall effect of bioavailability ^[16]. Fexofenadine is a BCS
255 class III compound which displays a negative food effect, as predicted by its BDDCS class.
256 Fexofenadine is a substrate for both OATP uptake transporters and P-gp efflux transporters. In the
257 fed state, principally when taken with fruit juices, the inhibition of OATP transporters predominates
258 and absorption is decreased ^[74-77]. Fruit juice related inhibition of OATP uptake has also been
259 implicated in a reduction in AUC for other drugs, including aliskiren and celiprolol^[74, 78-80]. The
260 inhibition of PAT1 has been suggested as a possible reason for the reduced rate of absorption of
261 vigabatrin, though this is most likely due to a reduction in the rate of gastric emptying ^[81, 82].

262 The events described here are summarised and examples of drugs affected by the various
263 mechanisms are provided in table 1.

Table 1 Summary of physiological mechanisms and biopharmaceutical aspects underpinning the food effect

Physiological mechanism	Biopharmaceutical aspects	Effect on Drug exposure	Example(s)
Increased pH in stomach	Solubility and dissolution of ionisable compounds can be altered	Increases AUC and C _{max} for weak acids Decreases AUC and C _{max} for weak bases	Cefuroxime Dipyridamole, indinavir
Increased concentration of solubilising species e.g. bile salts, lipid digestion products	Solubilisation of poorly water soluble drugs increases	Increases AUC and C _{max}	Fenofibrate Alectinib Danazol
Increased splanchnic blood flow	Saturation of liver enzymes and avoidance of FPM	Increases AUC and C _{max}	Propranolol Tacrine Dronedarone
Inhibition of gastrointestinal Cytochrome P450 – e.g. with Grapefruit juice	Fraction of drug escaping gut metabolism increases	Increases AUC and C _{max}	Felodipine Ciclosporin Atorvastatin
Inhibition of intestinal absorptive and efflux transporters	Fraction of drug subject to either absorptive or efflux transport is reduced	Increases AUC and C _{max} for drugs subject to efflux Decreases AUC and C _{max} for drugs which require uptake transporters	Ganciclovir Fexofenadine Talinolol

Delayed gastric emptying	Presence of food in stomach delays transit of drug to small intestine	Increases T_{max} , can decrease C_{max} , may cause T_{lag}	Widespread NSAIDs Paracetamol
Increase in viscosity of intestinal fluid	Reduction in water diffusivity, increase in luminal viscosity, slower water penetration of dosage form, increased disintegration time	Increases T_{max} , may reduce C_{max} and F , may cause T_{lag}	Chlorothiazide, Metformin

Clinical significance

Most medicines contain instructions to take the medication with a glass of water and often gives specific instructions to either take with food, occasionally specifying the size or content of the meal, or in the fasted state. These recommendations are generally designed either to improve safety and tolerability or to maximise the oral absorption. For example, it is recommended to take Famprya® (fampridine) without food in order to reduce the risk of adverse events as ‘there is a clear relationship between C_{max} and dose related adverse reactions’, and taking Famprya® with food is associated a 15-23% increase in C_{max} ^[83]. Similarly, the reduction in C_{max} of trifluridine observed in the fed state when taking Lonsurf® may prevent a reduction in neutrophils^[84]. Conversely, it is recommended to take both Orkambi® and Kalydeco® with fat containing meals to improve bioavailability and clinical efficacy, as there a 2-4 fold increase in exposure of both lumacaftor and ivacaftor are anticipated when these medicines are administered with fat containing food^[85, 86]. Other considerations may include the slower rate of absorption widely observed in the fed state. While there is no overall effect on bioavailability, the orexin receptor antagonist suvorexant (Belsomra®) should not be administered with or soon after a meal, as this may delay sleep onset^[87].

However, occasionally there may be contradictory advice or a lack of evidence for justifying these recommendations and occasionally the justification can seem counter-intuitive. For example, it is often recommended to take non-steroidal anti-inflammatory drugs (NSAIDs) with food, with the justification that this can reduce the incidence of gastric side effects, though the extent to which this is effective is questionable. Rainsford and Bjarnason (2011) have stated that ‘there are no specifically claimed benefits from these recommendations and their origins have not been made clear’, while Moore *et al.* (2015) have said that the evidence that taking NSAIDs with food achieves its objectives is non-existent^[88, 89]. Taking NSAIDs with food has been shown to delay T_{max} and reduce C_{max} with no overall effect on bioavailability. Considering that early, high plasma drug concentrations produce better and longer lasting analgesia in acute pain, and reduce the frequency of re-medication, it appears that the recommendation to take NSAIDs with food is misguided^[90, 89].

Another implication of significant food effect is the potential implications for the clinical efficacy. Ziprasidone (Geodon®) is an orally active atypical antipsychotic used in the treatment of bipolar affective disorder, which displays non-linear pharmacokinetics in the fasted state, while its absorption is approximately doubled by taking with a meal containing at least 500 kcal. Despite the significant food effect observed and label instructions to take Geodon® with food, about 40% of patients do not consistently take the medication with sufficient food and physicians have suggested that it is less effective in patients displaying poor compliance to the dose instructions^[91].

The impact of food on drug bioavailability is pertinent for new molecularly targeted therapies in oncology, particularly in the case of the kinase inhibitors^[92]. While FDA drug label instructions are generally designed to maximise the bioavailability of the drug, there is a distinct reversal of this situation for oncological preparations, where there is a noticeable trend towards label instructions to take medication in the fasted state despite significant increases in bioavailability in the fed state^[93]. This appears to run contrary to established understanding of basic biopharmaceutical principles, which would suggest that bioavailability may be enhanced, while variability can be reduced by co-administering these drugs in the fed state^[94]. This has resulted in suggestions of wastefulness, with some clinicians proposing that by ignoring the label recommendations and administering some of these antineoplastic agents with food that significant savings may be made such as \$1,700 per month in the case of lapatinib or \$3,750 per month in the case of abiraterone acetate^[95, 96]. While the case for taking these medicines with food in an off-label manner seems to stand to reason, other factors are important and warrant consideration, not least of which is conditions under which the drug product is licensed. While food effect studies are most often carried out as single dose studies in healthy subjects, the pivotal phase 3 clinical studies, which establish safety and efficacy in patients, may have been initiated in different prandial conditions, leading to a licensed dosing recommendation which reflects that of the relevant clinical study^[97, 98]. Dosing in differing prandial conditions runs to that recommended in the drug product label constitutes off-label administration, and risks administration under conditions which have not definitively been demonstrated as safe and effective, while patient adherence to label recommendations is a major concern for oncologists and their ability to manage dosage regimens^[99, 98]. The role of inter- and intra-individual variation, regarding meal composition and timing of taking medication with food is also a pertinent consideration^[100, 98]. For example, while lapatinib exposure can be increased greater than four-fold when taken with a high fat meal relative to the fasted state, this increase is only two fold when administered with a low fat meal. Considering the high fat and caloric content of the FDA high fat breakfast, it is not realistic to replicate the controlled environment of a food effect trial in the clinical setting, and variations in meal composition from day to day can result in large intra-individual variation^[101, 102]. In these cases it may be more reproducible, easily understood and easier to promote patient and clinician adherence where medicines are dosed in the fasted state^[97].

In cases where a specific type of meal is explicitly detailed, this can add further to the complexity. With regard to the multi-kinase inhibitor regorafenib (Stivarga®), the type of meal is particularly important, not only for the magnitude but, in fact, for the direction of the food effect. A significant increase in bioavailability was observed with a low fat breakfast, while a high fat meal causes a reduction in bioavailability with the resultant recommendation to 'take Stivarga® with food (a low-

fat breakfast)' ^[84]. Specifying a particular meal further adds to the risk associated with clinical use of medicines which display significant food mediated alterations in bioavailability, and risks reducing compliance with dosage regimens. It must be acknowledged that, with regard to oncological products, there may be specific challenges for fed state administration when considering the side effect profiles, such as nausea and vomiting, along with reduced appetite of patients undergoing certain chemotherapeutic regimens. While the debate continues as to whether these medications are best administered in the fed or fasted state, one thing which is abundantly clear is that a method of delivering these drugs in a reproducible, bio-enhanced manner, independent of prandial state would be advantageous.

Overall, establishing the clinical implications of food effect can be difficult, from the point of view of development scientists, clinicians and, indeed, regulators. The FDA bioequivalence criteria are deliberately conservative, ensuring maximal opportunity to observe a food effect, and do not take into account the variability and therapeutic window of the drug being assessed. A modest increase or decrease in bioavailability in the fed state will mean that bioequivalence is not demonstrated, however, if this drug displays large variability in bioavailability and/or possesses a wide therapeutic window, a modest change in variability, such as the 30% increase in exposure observed for gefitinib, is unlikely to be clinically significant ^[98]. This is notable among some of the recently licensed polymerase and protease inhibitors licensed in the treatment of the hepatitis C virus (HCV), in both individual and combination products, including Zepatier® (elbasvir and grazoprevir), Epclusa® (sofosbuvir and velpatasvir) and Daklinza® (daclatasvir). These products all display modest variations in bioavailability in the fed compared to the fasted state, though these minor changes are not deemed clinically relevant, allowing dosing independent of meal intake (table 2).

Another factor to consider is that while food effect studies are most often single dose studies, often this effect is lessened with multiple dosing, where variability in the patient population pharmacokinetics and the therapeutic window of the drug in question are important considerations. Such an example is that of Syndros® (dronabinol) where, despite a 2.5 fold, increase in exposure in a single dose fed state pharmacokinetic study, only the first dose is recommended to be taken with food, with subsequent dosing taken without regard to meals (table 2).

Yan *et al.* have recently identified numerous cases where there are label differences with regard to food effect between the US product information (PI) and European summary of product characteristics (SPC) ^[11]. This demonstrates the difficulties in interpretation of food effect data, which may be subjective and not entirely dependent on pharmacokinetic considerations, but also on the clinical pharmacodynamic response. It is also interesting to note that there does not appear to

be a consistent trend in these differences and of the products they identified, two-thirds displayed significant pharmacokinetic food effects. In the case of the anticoagulant Xarelto® (rivaroxaban), the clinical recommendation in fact varies between product strength, where the 10mg and 15mg preparations can be taken with or without a meal, while the 20mg strength should be taken with a meal^[103]. It is easy to envisage difficulties for clinicians in advising patients where dosing instructions vary for the same products between jurisdictions, but also between dose strengths.

Table 2 provides a summary of the food effects of newly licensed drugs or formulations approved in the US and/or EU over the last seven years which have, demonstrated significant food effect, or have been designated with a label restriction with regard to the administration of drug with regard to food. A food effect was considered significant if the ratio of AUC and/or C_{max} in the fed and fasted states fell outside 80-125%. We have also included products with a specific label claim regarding dosing with food. In cases where it was stated that a product showed no change or a non-significant change in either AUC or C_{max} in the fed or fasted state, but no values or ratios were obtained, a value of 1 was assigned.

Interestingly, our estimates have suggested that approximately 40% (67 of 157 products identified; 42.68%) of medicines licensed by the EMA and FDA since January 1st 2010 display a significant food effect or have been licensed with a label restriction with regard to dosing with or without food. Included in this analysis were new chemical entities, new combination products and previously marketed active pharmaceutical ingredients which have been reformulated. Excluded from our analysis were generic medicines/ abbreviated new drug applications (ANDAs), parenteral, topical, transdermal and other non-oral preparations (including buccal and sublingual preparations and orally disintegrating tablets), extended/ controlled release preparations and oral medicines designed for local administration within the GIT, i.e. those not subject to appreciable levels of absorption. In a competitive market place, the ability to take a medicine without regard to the timing of meals presents a clear commercial advantage for developing dosage forms that can be administered independent of food^[104].

393 Table 2 Recently licenced medicines displaying significant food effect and/or food specific dosage instructions
394 Data obtained from FDA Drug Label (from Drugs@FDA database) or European Summary of Pharmaceutical Characteristics (SPC) unless otherwise stated

Year licensed	Drug Name	Commercial Name	Clinical Recommendation regarding timing of food	Food Effect	AUC _{Fed} /AUC _{Fasted}	C _{max} _{Fed} /C _{max} _{Fasted}
2017	Spironalactone	Carospir®	CAROSPIR can be taken with or without food, but should be taken consistently with respect to food	Positive	1.9	-
2017	Glecarprevir Pibrentasvir	Mavyret®	Take orally once daily with food	Positive	1.83-2.63 ^a 1.4-1.53 ^a	-
2017	Sofosbuvir Velpatasvir Voxilaprevir	Vosevi®	Taken orally once daily with food	Positive	1.64-2.44 ^b 1.4-2.66 ^b 2.12-5.35 ^b	-
2017	Deutetrabenazine	Austedo®	Administer with food	Positive	1 ^c	1.5
2017	Betrixaban	Bevyxxa®	Take at the same time each day with food	Negative	0.39 ^d 0.52 ^e	0.3 ^d 0.5 ^e
2017	Telotristat ethyl	Xermelo®	Take with food	Positive	3.64 ^f 1.33 ^g	2.12 ^f 1.47 ^g
2016	Tenofovir alafenamide	Vemlidy®	Take with food	Positive	1.65	-
2016	Cabozatinib	Cabometyx®	Take at least 2 hours before and at least one hour after food	Positive	1.57	1.41
2016	Elbasvir Grazoprevir	Zepatier®	Taken with or without food	Neutral; Elbasvir Positive; Grazoprevir	0.89 1.5	0.85 2.8
2016	Migalastat hydrochloride	Galafold®	Take on an empty stomach, at least 2 hours before or after food intake	Negative	0.6	-
2016	Dronabinol	Syndros®	Because food delays the absorption of SYNDROS, administer the first dose on an empty stomach at least 30 minutes before eating. Subsequent doses can be taken without regard to meals.	Positive	2.5	0.8
2016	Emtricitabine Rilpivirine	Odefsey®	Take with a meal	Positive	Emtricitabine 0.91 ^h , 0.88 ^e	-

Tenofovir alafenamide

Rilpivirine
1.13^h, 1.72^e
Tenofovir
alafenamide
1.45^h, 1.53^e

2016	Rucaparib	Rubraca®	Take with or without food	Positive	1.38	1.2
2016	Sofosbuvir Velpatasvir	Epclusa®	Take with or without food	Positive;	1.6 ^h	-
				sofosbuvir	1.78 ^e	
				Positive;	1.21 ^h	-
				velpatasvir	1.34 ^e	
2016	Venetoclax	Venclexta®	Take with a meal and water	Positive	3.4 ^d 5.1 ^e	-
2015	Elvitegravir Cobicistat Emtricitbine Tenofovir alafenamide	Genvoya®	Take Once Daily with Food	Positive	Elvitegravir	
					1.34 ^d	-
					1.87 ^e	
2015	Alectinib	Alecensa®	Administer with food	Positive	3.1	-
2015	Daclatasvir	Daklinza®	With or without food, with sofosbuvir	Negative	0.77	0.72
2015	Eluxadoline	Viberzi®	Taken twice daily with food	Negative	0.4	0.5
2015	Flibanserin	Addyi®	No instructions with regard to food intake	Positive	1.18 ^d	1.02 ^d
					1.43 ^h	1.13 ^h
					1.56 ^e	1.15 ^e
2015	Idebenone	Raxone®	Administer with food	Positive	5-7	-
2015	Ivabradine	Corlanor®/ Procoralan®	Take with meals	Positive	1.2-1.4	-
2015	Ixazomib	Ninlaro®	Taken at least one hour before or 2 hours after food	Negative	0.72	0.31
2015	lumacaftor ivacaftor	Orkambi®	Take with fat containing food	Positive;	2	-
				Lumacaftor		
				Positive; Ivacaftor	3	-
2015	Palbociclib	Ibrance®	Take with food	Positive	1.12 ^d	1.27 ^d
					1.13 ^h	1.24 ^h

					1.21 ^e	1.38 ^e
2015	Sonidegib	Odomzo [®]	Take on an empty stomach	Positive	7.4-7.8	7.4-7.8
2015	Tasimelteon	Hetlioz [®]	Taken without food	Negative	-	0.56
2015	Trifluridine Tipiracil	Lonsurf [®]	Within One hour after completion of meal	Negative	1 ^c	0.6
			Take with food, as decrease in Trifluridine C _{max} can prevent decrease in neutrophils	Negative	0.6	0.6
2014	Ceritinib	Zykadia [®]	Administer ZYKADIA on an empty stomach (i.e., do not administer within 2 hours of a meal)	Positive	1.58 ^d 1.73 ^e	1.43 ^d 1.41 ^e
2014	Delamanid	Delytba [®]	Delamanid should be taken with food	Positive	2.7	-
2014	Droxidopa	Northera [®]	Take consistently with or without food	Negative	0.8	0.65
2014	Idelalisib	Zydelig [®]	Zydelig can be taken with or without food	Positive	1.4	-
2014	Ledipasvir	Harvoni [®]	Taken daily with or without food	None; ledipasvir	1 ^c	1 ^c
	Sofosbuvir			Positive; sofosbuvir	2	1 ^c
2014	Naloxegol	Movantik [®]	Take on an empty stomach at least 1 hour prior to the first meal of the day or 2 hours after the meal	Positive	1.45	1.3
2014	Nintedanib	Ofev [®]	Take with food	Positive	1.2	-
2014	Pirfenidone	Esbriet [®]	Three times daily taken with food.	Negative	0.84	0.51
2014	Suvorexant	Belsomra [®]	Suvorexant may be taken with or without food; however for faster sleep onset, suvorexant should not be administered with or soon after a meal	None	1 ^c	1 ^c
2014	Tasimelteon	Hetlioz [®]	Take without food	Negative	-	0.56
2013	Afatinib	Gilotrif [®]	Take at least 1 hour before or 2 hours after a meal	Negative	0.61	0.5
2013	Dabrafenib	Tafinlar [®]	Taken at least 1 hour before or at least 2 hours after a meal	Negative	0.69	0.49
2013	Dimethyl fumarate	Tecfidera [®]	Take TECFIDERA with or without food	Negative	1 ^c	0.6

2013	Dolutegravir	Tivicay®	May be taken without regard to meals	Positive	1.33 ^d 1.41 ^h 1.66 ^e	1.46 ^d 1.52 ^h 1.67 ^e
2013	Obrutinib	Imbruvica®	No instructions with regard to food intake	Positive	2	-
2013	Nalmefene hydrochloride dihydrate	Selincro®	Selincro can be taken with or without food	Positive	1.3	1.5
2013	Ospemifene	Osphena®	One tablet taken orally once daily with food	Positive	1.7	2.3
2013	Simeprevir	Olysio®	One 150 mg capsule taken once daily with food	Positive	1.69 ^e	-
2013	Trametinib	Mekinist®	Take at least 1 hour before or at least 2 hours after a meal	Negative	0.76	0.3
2012	Bedaquiline	Sirturo®	Bedaquiline should be taken with food to enhance its oral bioavailability	Positive	2	-
2012	Bosutinib	Bosulif®	Taken once daily with food.	Positive	1.7	1.8
2012	Cabozantinib	Cometriq®	Instruct patients not to eat for at least 2 hours before and at least 1 hour after taking COMETRIQ.	Positive	1.57	1.41
2012	Elvitegravir	Stribild®	The recommended dose of STRIBILD is one tablet taken orally once daily with food	Positive	1.34 ^d 1.87 ^e	-
	Cobicistat			Neutral	1 ^c	-
	Emtricitabine			Neutral	1 ^c	-
	Tenofovir disoproxil fumarate			Positive	1.24 ^d 1.23 ^e	-
2012	Ivacaftor	Kalydeco®	Taken orally every 12 hours with fat-containing food	Positive	2 to 4	-
2012	Mirabegron	Myrbetriq®/ Betmiga®	Taken once daily with or without food	Negative	0.49 ^d 0.83 ^e	0.25 ^d 0.55 ^e
2012	Regorafenib	Stivarga®	Take Stivarga with food (a low-fat breakfast)	Positive	1.48 ^e 0.8 ^{e,g} 0.49 ^{e,g} 1.36 ^d	-

					1.4 ^{d,g} 1.23 ^{d,g}	
2012	Isotretinoin	Absorica®/ Epuris®	Recommended dosage of 0.5 to 1 mg/kg/day given in two divided doses without regards to meals for 15 to 20 weeks	Positive	1.5	1.26
2011	Abiraterone acetate	Zytiga®	ZYTIGA must be taken on an empty stomach	Positive	5 ^d 10 ^e	7 ^d 17 ^e
2011	Boceprevir	Victrelis®	800 mg administered orally three times daily (every 7 - 9 hours) with food (a meal or light snack).	Positive	1.65	-
2011	Gabapentin enacarbil	Horizant®	Once daily taken with food at about 5 PM	Positive	1.24 ^d 1.34 ^h 1.44 ^e	-
2011	Piperaquine tetraphosphate Arteminol	Eurartesim®	Eurartesim should be administered with water no less than 3 hours after the last food intake, and no food should be taken within 3 hours after each dose	Positive	Piperaquine; 3 Arteminol; 1.43	-
2011	Rilpivirine hydrochloride	Edurant®	Taken once daily with a meal	Positive	1.666667	-
2011	Rivaroxaban	Xarelto®	10mg and 15mg; With or without food 20mg; Take with food	10mg and 15mg; Neutral 20mg Positive	1 ^c (10mg) 1.39 (20mg)	-
2011	Telaprevir	Incivek®	INCIVEK tablets is 750 mg (two 375-mg tablets) taken orally 3 times a day (7-9 hours apart) with food (not low fat)	Positive	2.17 ^d 3.37 ^h 4.3 ^e	-
2011	Vemurafenib	Zelboraf®	Take with or without a meal	Positive	4.6-5.1	2.5
2011	Vilazodone hydrochloride	Viibryd®	VIIBRYD should be taken with food.	Positive	1.64 - 1.85	2.47 - 2.6
2010	Dronedarone	Multaq®	Take twice a day with morning and evening meals	Positive	3.75	-
2010	Fampridine	Fampyra®	Since there is a clear relationship between Cmax and dose related adverse reactions,	Positive	0.93-0.98	1.15-1.23

it is recommended to take Fampyra
without food

	2010	Lurasidone HCl	Latuda®	Latuda should be taken with food	Positive	2	3
395	^a Mean systemic exposures with moderate to high fat meals						
396	^b Values refer to geometric mean systemic exposure.						
397	^c Where no numerical values for food effect were obtained but no significant food effect was observed a value of 1 was assigned						
398	^d Low fat fed						
399	^e High fat fed						
400	^f Parent compound						
401	^g Active metabolite						
402	^h Moderate fat meal						
403							

Strategies to overcome food effect bioavailability in drug development

Where food effects are identified, there is generally three choices facing drug development and/or regulatory scientists: (1) consider an alternative lead drug molecule that will not display food effects, (2) apply specific instructions for how a medicine is taken with regard to food or (3) design a formulation which overcomes to food effect. With an increasing desire to improve R&D efficiency in drug development, and the 'quick win, fast fail approach' now favoured in drug development, there is an increasing desire to predict food effects earlier in the drug development process^[105, 106]. This will also allow potential to identify a food-independent formulation, approaches to which are described here. The primary focus is on the clinical performance observed with such formulation approaches, while notable studies in preclinical studies, principally in beagle dogs are also reviewed. While pre-clinical animal models, including the dog model, are not always representative of human bioavailability, they remain a cornerstone of pre-clinical formulation development, particularly with regard to food effect^[105, 107]. Dogs are indeed the most widely characterised animal model in food effect studies, and dog specific food effect models are widely available, with a general tendency to be over-predictive of human food effect^[108, 109].

Lead candidate modification and optimisation

Once a potential lead compound has been identified during the drug development process, the final drug discovery phase involves modifying the molecular structure or physicochemical properties of the potential drug candidate to improve biopharmaceutical performance. The two guiding principles are the maintenance of favourable properties in lead compounds, retaining the motifs identified as crucial to the structure activity ratio (SAR), while also improving deficiencies in drug structure, often with the aim of improving the deliverability of the drug^[110]. While it is unlikely that development scientists will specifically focus on food effect at this stage, identification and selection of appropriate lead candidates can lead to a reduction in food effect bioavailability later in the development process. While studies focussing on modifying the structure and physiochemical properties of a lead candidates specifically with the aim of reducing the impact of food are sparse, there are numerous examples of marketed drugs with related chemical and clinical properties, but differing food effects.

Pithavala *et al.* examined the effect that crystal habit may have on absorption and food-effect, and demonstrated the importance of screening drug polymorphs. Initial first in human (FIH) trials suggested a negative food effect for axitinib form IV in a film-coated, immediate release tablet. A 23% reduction in absorption in the fed state was demonstrated. Subsequent investigations identified

a more stable polymorph, form XLI. Food effect studies carried out with form XLI demonstrated an increase in the overall bioavailability of 19% with a high fat meal, and a 10% reduction with a moderate fat meal compared to fasting, which were not considered to be clinically significant changes ^[111].

Modifying the chemical structure by means of producing prodrugs can also be used to eliminate food effect, as demonstrated by the development of fosamprenavir, a prodrug of the previously marketed protease inhibitor amprenavir. Due to its poor solubility, amprenavir was originally formulated as a lipid based formulation (Agenerase[®]) which demonstrated reduced bioavailability (AUC decreased by 23%) when taken with a high fat meal. Fosamprenavir, a phosphate ester prodrug with improved solubility, was originally developed with a view to reduce the significant daily pill burden associated with Agenerase[®] (eight capsules, twice daily). Successful formulation of fosamprenavir (Telzir[®]) not only reduced the dosing schedule to one tablet twice daily, but also eliminated the negative food effect seen with amprenavir, allowing dosing independent of prandial state ^[112].

The prodrug approach has also been used to produce the gabapentin ester, gabapentin enacarbil. The original aim of such an approach was to increase the poor and saturable bioavailability of gabapentin. The ester prodrug is completely hydrolysed to gabapentin by esterase enzymes in the gut and liver ^[113, 114]. While gabapentin bioavailability is greater from the prodrug when dosed in equimolar concentrations, Horizant[®] (gabapentin enacarbil) is required to be dosed with a meal, while Neurontin[®] (gabapentin) can be dosed with or without food. Numerous studies have demonstrated increases in exposure to gabapentin after oral dosing as gabapentin enacarbil ^[113, 115, 114]. Mean increases in AUC_{inf} of 23% for low fat, 31% for moderate fat and 40% for high fat meals have been observed in one study ^[115]. Meanwhile exposure to gabapentin from Neurontin[®] is not significantly different in the fasted and fed states with an increase of 14% in AUC and C_{max} in the fed state ^[116]. Direct comparison in these cases is, however, difficult as gabapentin enacarbil is only utilised in extended release preparations, while gabapentin is an immediate release formulation and both compounds are utilised for different indications. A similar approach, using an ester prodrug, has also been demonstrated to improve the bioavailability and eliminate the food effect for the direct thrombin inhibitor melagatran ^[117].

Formulation approaches to enhance bioavailability

Numerous formulation approaches have been utilised to overcome food effects on bioavailability and the type of formulation chosen will depend on the nature and mechanism of the food effect, the drugs physiochemical properties and the intended therapeutic profile. To date, the majority of

studies aimed at overcoming food effect have focused on poorly water soluble, BCS class II compounds. This is both due to these molecules being the most commonly observed class in drug development pipelines, and the fact that these molecules are the most amenable to formulation approaches designed to overcome their biopharmaceutical limitations. This has provided a focus for the development of bio-enabling formulations to improve dissolution and bioavailability, ultimately with the aim of ensuring BCS class II compounds will behave more like BCS class I compounds *in vivo*. It is widely stated that by maximising dissolution *in vivo* in the fasted state it may also be possible to prevent the postprandial increases in solubilisation and mitigate or eliminate a positive food effect entirely, though as we will discuss below, this may be an oversimplification. While each of the formulations discussed in this article have indeed been well characterised elsewhere, they are discussed here specifically in the context of their use in eliminating food effects on bioavailability.

Nanosized preparations

The term nanocrystal has emerged to describe drug particles with a crystalline structure in the nanoscale range ^[118]. Nanosizing refers to the reduction of API particle size to the sub-micron range, typically <500nm, and with modern production techniques it is possible to achieve particle size in the 100-200nm range ^[119]. The reduction in particle size leads to an increase in surface area available for solvation and increases the rate of dissolution. The formation of nanoparticles may not only enhance dissolution, but evidence exists that solubility may also be increased through changes in the particle curvature and introduction of defects into the crystal lattice, while the thickness of the diffusion layer surrounding the particle may also be reduced ^[120-122, 118]. Nanonisation of API has proven useful in enhancing the bioavailability of PWSD, and numerous commercial examples exist, and many of these commercial preparations have been shown to eliminate a positive food effect previously seen with marketed preparations or in the drug development process.

Fenofibrate has been widely investigated as a model PWSD displaying positive food effect bioavailability. Originally marketed as a co-micronized capsule, with an API particle size of 5-15 µm, which required dosing with food to achieve maximal absorption of a 200mg dose, it has repeatedly been reformulated using different bio-enabling approaches. Two nanonized preparations of fenofibrate have so far reached market, namely Tricor® (also marketed as Lipantil® Supra; prepared using NanoCrystal® milling technique developed by Elan Nanosystems) and Triglide™ (prepared via high pressure homogenisation). Comparison of absorption from 145 mg nanosized Tricor® formulation in the fasted and fed state to that of the 160 mg microcoated tablets demonstrated similar exposure in the fed state, while absorption from the nanonized tablet was increased in the fasted state and resulted in the elimination of a food effect ^[123, 124].

Aprepitant is a BCS class IV compound which was formulated as a nanoparticle, using NanoCrystal® technology, during drug development to enhance fasted state dissolution. The final preparation was marketed as EMEND® and was found to improve fasted state exposure and eliminate the positive food effect seen with early tablet formulations in clinical development [125, 126].

Megestrol acetate is a steroidal progestin which is licensed for use as an appetite stimulant in anorexia and cachexia. Thus, the positive food effect seen with the original Megace® oral suspension, along with the 800mg dose in the relatively large volume of 20 mL suspension was seen as problematic in patients with decreased appetite. Reformulation as the nanocrystalline Megace® ES demonstrated a reduction of food effect, but also allowed dose reduction to 625mg administered in 5 mL of the new formulation [127, 128].

The advantages of nanonized API compared to other methods of particle size reduction, specifically micronization through hammer- or jet-milling was demonstrated by Jinno *et al.* [129]. Here, a spray dried nanocrystalline suspension of cilostazol not only improved bioavailability approximately 5 fold in fasted beagle dogs relative to two different micronized preparations, but also eliminated the positive food effect seen with the micronized formulations. This was attributed to improved dissolution, as demonstrated in biorelevant FaSSIF media [129].

Several other nanocrystalline preparations have also demonstrated enhanced fasted state bioavailability in the fasted state and elimination of food effect in pre-clinical animal models, including ziprasidone [91, 130], lurasidone [131] and the novel gamma secretase inhibitor ELND006 [132]. Table 4 contains numerous examples of commercially available nanocrystalline preparations where food effect has been studied. In all cases a significant food effect observed with previous formulations has been mitigated or eliminated, demonstrating that nanosizing is an effective approach to eliminating food effect bioavailability.

Amorphisation and solid dispersion

The term solid dispersion describes a wide range of different, but related formulations which are designed to maintain drug in an amorphous or phase-separated crystalline state [133]. By reducing the drug particle size to the molecular level rapid dissolution can be facilitated, and production of an amorphous form will improve the apparent solubility, while solid dispersion can also confer improved wettability, increased porosity and, ultimately, improved biopharmaceutical performance [134-136]. Solid dispersions are being used increasingly often as bio-enabling formulations for PWSD to enhance oral bioavailability and numerous commercial preparations exist [137]. These preparations most often exist as amorphous drug dispersed in an inert carrier matrix, and this narrow definition has been used to describe their behaviour [138].

One such example is that of Kaletra[®], a combination product of lopinavir and ritonavir produced using solid dispersion technology, specifically hot melt extrusion, using PVP/VA as a carrier ^[139]. Having originally been formulated as a soft gelatin capsule containing lipid excipients, the capsule formulation of Kaletra[®] were required to be taken with food, with a 48% increase in bioavailability observed in the fed state. The poor solubility of the API also meant that the capsule dose was limited to an 80mg/20mg strength capsule. Reformulation as a solid dispersion allowed production of a 200mg/50mg tablet, reducing the pill burden from 10 capsules daily to four tablets daily. The amorphous solid dispersion formulation also displayed only insignificant changes in bioavailability in the fed compared to the fasted state, allowing food independent dosing ^[139].

Similarly, Lynparza[®] (olaparib) has been reformulated from a lipid-based, crystalline solid dispersion of micronised olaparib in Gelucire[®], to a hot-melt extrusion based dispersion using copovidone as a carrier ^[140]. While the original formulation was developed after significant pre-clinical development, and displayed a 2-fold increase relative to a standard immediate release tablet, relatively low drug loading (10%) led to a significant pill burden for patients (16 capsules daily) ^[141]. The development of the melt extrusion tablet formulation both increased olaparib bioavailability and drug loading, allowing a dose reduction from 400mg to 300mg twice daily, and reduced the pill burden to four tablets daily. The food effect was also reduced, with a 20% increase in exposure observed for the capsule compared to a 9% increase with the tablet formulation ^[142, 143, 140].

Banarjee *et al.* developed a ziprasidone solid dispersion via hot melt extrusion, which retained crystalline characteristics of ziprasidone while suspending the drug in a hydrophilic matrix to improve wettability and dissolution, resulting in a nearly 10-fold increase in solubility. The enhanced dissolution translated to improved bioavailability in fasted healthy volunteers, while simultaneously eliminating the positive food effect observed with the commercial Zeldox[®] formulation ^[144].

However, while these approaches have successfully reduced food effect using solid dispersion technology, table 4 contains numerous examples where this is not the case. For the marketed solid dispersion formulations and amorphous drug preparations for which food effect data could be found (n = 21) almost half (n = 10) display positive food effect, while four preparations displayed a negative food effect. The fact that two thirds of these bio-enabled formulations display food effect bioavailability suggests that while solid dispersions may well improve dissolution in the fasted state, quite often solubility limitations remain.

Lipid Based Formulations

The original rationale for the investigation of the use of lipid-based formulations (LBF) to increase bioavailability of PWSD was the observation of positive food effects for many of these compounds

^[145]. The ability of food to enhance the absorption of PWSD has long been attributed to the ability of meal components, and in particular lipids, to enhance drug solubilisation, dissolution and absorption. Thus, the addition of exogenous lipids to pharmaceutical preparations was proposed and investigated as a viable option to enhance the bioavailability of PWSD. LBFs have thus become one of the most widely investigated and characterised formulation types for bioavailability enhancement and the elimination of a positive food effect, and have become 'renowned for their potential to reduce the impact of food on drug absorption' ^[146]. One of the earliest studies to specifically focus on the utility of LBF to eliminate food effect bioavailability was that of Charman *et al.* (1993). This study demonstrated that the approximately 3 fold increase in C_{max} and AUC observed for a commercial danazol capsule formulation (Danocrine®) was eliminated using a lipid emulsion of danazol in glycerol mono-oleate ^[147].

However, eliminating food effect using LBF is not always straightforward, and can require significant formulation development, as is the case for lipid-based formulations of cyclosporine. The commercial success of self-emulsifying drug delivery system (SEDDS) formulation of Neoral® owing principally to its elimination of the food effects and reducing inter-subject variability relative to the crude lipid emulsion formulation of Sandimmune® ^[148, 149]. Delivery as a crude emulsion was not sufficient to overcome the food effect, which required a more elaborate SEDDS formulation.

Similarly, Roaccutane® is a soft gelatin capsule, which contains isotretinoin solubilised in water-insoluble solvents, namely beeswax, soya bean oil and hydrogenated soya bean oil, which displays an approximately 2.7 fold increase in bioavailability in the fed compared to fasted state. Absorica® is a novel isotretinoin formulation developed using Lidose® technology, which enhanced the fasting state bioavailability and reduced the food effect to a 1.5 fold increase, which is not considered to be clinically significant and allows food independent dosing ^[150].

While the use of LBF to eliminate food effect has been widely acknowledged, it is interesting to look more critically at this claim. The use of LBF to enhance the fasted state bioavailability has been the major focus of formulation development over the last five decades, and it is a logical inference that by enhancing the solubility limited bioavailability in the fasted state, the post-prandial increase in absorption mediated by increased solubility can be reduced or avoided. However, as presented in table 4, of the 29 LBFs for which food effect data was gathered, 17 of these formulations displayed significant positive food effect, while only 9 formulations demonstrating truly food independent dosing.

This can be exemplified by the case study of Fortovase®, a SEDDS formulation designed to improve the oral bioavailability of saquinavir, relative to the conventional capsule formulation, Invirase®.

While bioavailability was enhanced approximately three fold by Fortovase®, a significant food effect was still evident, with a similar increase in the fed state to that observed with Invirase® (6.7 fold increase).

Perlman *et al.* have examined the food effect of torcetrapib in dogs using a range of different SEDDS formulations, finding that the composition of the formulation can be crucial in determining the food effect, with a food effect ranging from complete absence to 3.8 fold increase in beagle dogs. It should be noted, however, that all formulations reduced the food effect seen with aqueous (18 fold increase) and lipid (5 fold increase) suspensions used in early drug development ^[151]. Thus, while simply administering PWSD with lipids may reduce the magnitude of the food effect, true elimination may require more extensive formulation optimisation ^[8]. Christiansen *et al.* have similarly demonstrated a reduction of food effect for cinnarizine tablets when co-administered with placebo SNEDDS, relative to administration without this placebo lipid formulation in healthy volunteers, though complete elimination of food effect was not possible ^[152].

Various other lipid-based formulations have also been investigated in pre-clinical species with varying levels of success, and with a general trend towards reducing rather than eliminating food effects ^[153-163]. While the assertion that LBF are excellent candidates to eliminate food effect is widespread, the evidence from the literature, and from product literature in particular is underwhelming and a more systematic investigations are required to fully elucidate the potential for LBF to overcome food effects ^[164].

Cyclodextrins

Cyclodextrins have been widely used to enhance the oral bioavailability of lipophilic and poorly water-soluble drugs, in both pre-clinical animals and in humans. The bioavailability enhancing effects are mainly due to enhanced dissolution kinetics, increased solubility and potential reduction in degradation as well as increased permeability ^[165]. Experience with cyclodextrins with a specific focus on elimination of food effect is, however, limited.

Sporanox® (itraconazole) has been formulated both as an amorphous solid dispersion, which displayed significant, positive food effect and as an oral solution solubilised by hydroxypropyl-β-cyclodextrin inclusion complex. Sporanox® cyclodextrin solution has demonstrated higher bioavailability than Sporanox capsules in the fed state, and has also been demonstrated to show enhanced bioavailability in the fasted state, eliminating the need for fed state dosing ^[166-168].

Along with the commercial itraconazole preparation, cyclodextrin complexes have also been investigated in pre-clinical food effect studies. Thombre *et al.* have demonstrated that a sulfobutyl

633 ether- β -cyclodextrin (SBE- β -CD) inclusion complex enhanced fasted bioavailability and eliminated
634 food effect for ziprasidone relative to the commercial preparation (Geodon®) in beagle dogs ^[91, 169].
635 Wang *et al.* have recently demonstrated similar results with an SBE- β -CD complex of amiodarone
636 ^[170].

637 While the experience with cyclodextrin preparations for eliminating food effect is limited, these
638 examples show promise for this formulation method to eliminate food effect, though overall clinical
639 acceptability may be limited as the relatively large intake volume (up to 20 mL for Sporanox®
640 solution) may be problematic for some patients.

641

Guiding formulation selection

With the range of formulations available, identification of the biopharmaceutical risks for a particular drug candidate is essential in order to ensure the most appropriate formulation is chosen. While it may be possible that different types of formulation may achieve improved solubilisation for a particular drug candidate, identification of the most beneficial formulation can be advantageous in the industrial setting where developing parallel formulation portfolios can put a strain on the limited timelines and resources. Kuentz *et al.* have recently reviewed the various methods by which formulations are selected in the pharmaceutical industry^[171]. The key elements of formulation screening involve identifying the critical physiochemical and biopharmaceutical properties that are likely to play a role in drug bioavailability and generating a target product profile (TPP). Ideally, formulation decision trees based on a basic set of drug properties, such as that proposed by Rabinow *et al.*^[172], or those based on identifying the biopharmaceutical limitations, such as the BCS based decision trees suggested by Ku *et al.*^[173], can be implemented and will provide a relative simple strategy to formulation choice. There is an abundance of such decision trees in the literature, with focus on enhancing bioavailability and manufacturability of drug candidates. However, given the lack of a clear consensus on the appropriate prediction of food effect bioavailability and the contradictory evidence of the various formulation options at eliminating food effect, it is no surprise that no decision tree exists specifically focus on eliminating the food effect. In the absence of a specific decision tree, choice is best guided by analysing BCS/BDDCS class assignment and utilisation of existing decision trees for each appropriate class. Formulation approaches suitable for each class are summarised in figure 4.

While significant experience, no doubt, exists within the pharmaceutical industry, this data is often not published in the literature. The paucity of data regarding the systematic and structured development of formulations with the intention of eliminating food effect means that, initially, the decision to focus on one particular formulation approach to eliminate a food effect remains largely empiric. More thorough formulation development and characterisation, however, can benefit from a more mechanistic approach, using a range of *in vitro*, *in vivo* and *in silico* tools. A recent example of such an approach is that of Pandey *et al.*^[174]. In this work, the group first identified a large food effect in an early stage clinical trial in healthy volunteers. This food effect was subsequently investigated mechanistically using biorelevant *in vitro* screening tools along with *in silico* modelling using the GastroPlus® ACAT model. This biorelevant screening identified that the key mechanism governing the observed positive food effect was the enhanced solubilisation by dietary lipids, while *in silico* modelling suggested that other contributory factors were involved, including the changing pH and the impact of gastrointestinal transporters and metabolism. The integration of the early

clinical data with the biorelevant measurements with the *in silico* model allowed development of a bio-predictive *in vitro* dissolution method, which enabled rapid formulation screening. Formulation screening led to the development of a surfactant containing, wet-granulated tablet formulation. The approximately 3.5-fold increase in fed state bioavailability observed for the dry granulated formulation was reduced to an approximately 1.5-fold increase, which was deemed not to be clinically significant for this compound ^[174].

	High Solubility	Low Solubility
High Permeability/Metabolism	Class I Conventional immediate release dosage forms	Class II <u>Bio-enabling formulations</u> Micronised API Nanonised API Lipid Based Formulations Amorphisation Solid Dispersions Crystal modification Cyclodextrin complexation
Low Permeability/Metabolism	Class III Permeation enhancers	Class IV Combination of II and III

Figure 4 Approaches to formulation design based on BCS/BDDCS classification

685 Table 3 Approved bio-enabling formulations with clinical food effect data
686 Data obtained from FDA Drug Label (from Drugs@FDA database) or European Summary of Pharmaceutical
687 Characteristics (SPC) unless otherwise stated

	Trade name	API	Clinical recommendation	AUC
Amorphous solid dispersions	Cesamet®	Nabilone	No specific instructions	
	Sporanox® (capsule)/ Onmel®	Itraconazole	Sporanox is for oral administration and must be taken immediately after a meal for maximal absorption.	
	Prograf®	Tacrolimus	Administer capsules consistently with or without food	
	Kaletra®	Ritonaivr/lopinavir	Kaletra tablets can be taken with or without food.	
	Intelence®	Etravirine	Taken following a meal	
	Zortress®/ Certican®	Everolimus	Administer consistently with or without food at the same time as cyclosporine.	
	Norvir®	Ritonavir	Take Norvir with meals	
	Zelboraf®	Vemurafenib	Administer ZELBORAF approximately 12 hours apart with or without a meal	
	Incivek®	Telaprevir	Take with food (not low fat)	
	Kalydeco®	Ivacaftor	With fat containing food	
		Ombitasvir		
	Viekirax®	Paritaprevir Ritonavir	Viekirax tablets should be taken with food, without regard to fat and calorie content	
	Fenoglide®	Fenofibrate	Should be taken with meals	
	Rezulin®	Troglitazone	Rezulin should be taken with a meal	1.
	Noxafil®	Posaconazole	Taken with food	
Pure Amorphous Drug	Ceftin®	Cefuroxime axetil	Administer tablets with or without food	
	Viracept®	Nelfinavir mesylate	VIRACEPT should be taken with a meal	

Lipid Based Formulations	Accupril®	Quinapril HCl	No specific instructions
	Victrelis®	boceprevir	Take three times daily with food (a meal or light snack)
	Crestor®	Rosuvastatin Calcium	CRESTOR can be administered as a single dose at any time of day, with or without food
	Zepatier®	elbasavir grazoprevir	One tablet taken orally once daily with or without food
	Accolate®	Zafirlukast	ACCOLATE should be taken at least 1 hour before or 2 hours after meals.
	Agenerase®	Amprenavir	AGENERASE may be taken with or without food, however, a high fat meal decreases the absorption of amprenavir and should be avoided
	Neoral®	Cyclosporin A/I	Neoral be administered on a consistent schedule with regard to time of day and relation to meals.
	Accutane®/ Roaccutane®	Isotretinoin	Accutane should be administered with a meal
	Kaletra®	Lopinavir Ritonavir	Kaletra capsules must be taken with food
	Norvir®	Ritonavir	Take with food
	Restandol®/ Andriol® Testocaps	Testosterone undecanoate	Restandol Testocaps must be taken with a normal meal
	Targretin®	Bexarotene	Targretin® capsules should be taken as a single oral daily dose with a meal.
	Lamprene®	Clofazimine	Administer 100 mg LAMPRENE daily with meals
	Sandimmune®	Cyclosporin A	To be administered on a consistent schedule with regard to time of day and relation to meals.
	Marinol®	Dronabinol	Take twice daily , before lunch and supper
	Avodart®	Dutasteride	May be administered with or without food
	Procardia®	Nifedipine	No specific instructions

	Rapamune®	Sirolimus	Oral solution; Administer once daily by mouth, consistently with or without food
	Fortovase®	Saquinavir	FORTOVASE should be taken within 2 hours after a full meal
	Amitiza®	Lubiprostone	Take with food to reduce nausea
	Aptivus®	Tipranavir	Always take APTIVUS with food.
	Hycamtin®	Topotecan HCl	May be administered with or without food
	Akynzeo®	Netupitant Palonsetron	Can be taken with or without food
	Prometrium®	Progesterone	No specific instructions
	Utrogestan®	Progesterone	Utrogestan 100mg Capsules should not be taken with food and should be taken at bedtime.
	Absorica®	Isotretinoin	Recommended dosage of 0.5 to 1 mg/kg/day given in two divided doses without regards to meals for 15 to 20 weeks
	Lipofen®	Fenofibrate	LIPOFEN™ capsules should be given with meals, thereby optimizing the absorption of the medication.
	Ofev®	Nintedanib	Take with food
	Navelbine®	vinorelbine tartrate	Administer the capsule with some food, as this has also been shown to reduce the incidence of nausea and vomiting
	Zemlar®	Paricalcitol	Without regard to meals
	Toctino®	Alitretinoin	With a main meal
	Vyndaqel®	Tafamidis meglumine	With or without food
	Rayaldee®	calcifediol	Once Daily at bedtime
	Xtandi®	Enzalutamide	XTANDI can be taken with or without food
Nanocrystalline	Lipantil Supra®	Fenofibrate	Without regard to meals
	Megace® ES	megestrol acetate	No specific instructions

	Emend®	aprepitant	With or without food
	Triglide®	Fenofibrate	TRIGLIDE may be administered with or without food
	Rapamune®	sirolimus	To minimise variability, Rapamune should consistently be taken either with or without food
Cyclodextrin	Sporanox® (Oral solution)	Itraconazole	Oral solution; Taking SPORANOX® Oral Solution under fasted conditions improves the systemic availability of itraconazole. Instruct patients to take SPORANOX® Oral Solution without food, if possible
Crystalline Solid Dispersion	Lynparza® (capsule)	Olaparib micronised in Gelucire; manufactured as a suspension of drug in molten excipient	No specific instructions
	Lynparza® (tablet)	Hot melt extruded olaparib in co-povidone carrier matrix	Taken orally twice daily with or without food

^a Where no numerical values for food effect were obtained but no significant food effect was observed a value of 1 was assigned

^b Moderate fat meal

^c High fat meal

^d Low fat meal

^e With Breakfast

^f 2 hours post breakfast

^g 4 hours post breakfast

^h Active metabolite

Conclusion

This review has investigated the causes and impact of food mediated changes in drug bioavailability. While our mechanistic understanding of the causes of food effects, and our ability to predict when and why it might occur have developed significantly since Welling first reviewed this topic 40 years ago, food effects still pose significant problems with regard to both development and regulatory scientists.

Despite the increased awareness of the negative clinical impact of food effects on bioavailability and the strict regulatory guidance regarding the appropriate testing of new medicinal products in the fed and fasted states there appears to be an ever-increasing challenge of food mediated alterations in drug bioavailability, likely reflecting the increasing prevalence of PWSD in drug development

707 pipelines. While there has been increasing understanding and development of improved drug
708 delivery technologies, there remains an overall lack of appreciation of the scale of the food effect
709 challenge, as evidenced by the fact that over 40% of new medicines display significant food effects,
710 or possess a label claim in respect of dosing with regard to food intake. This has had a knock-on
711 effect in the clinic, where the success or commercial advantage of compounds can be affected,
712 particularly with antipsychotic and oncological preparations.

713 While this review has summarised the various formulation approaches that have been utilised to
714 mitigate food effect, it is still difficult to definitively suggest a method of choice for formulating new
715 compounds to overcome significant food effects. The major focus of formulation approaches to
716 mitigate food effects to date has focused on compounds displaying positive food effects mediated
717 by poor dissolution or solubility, while relatively limited approaches exist for drugs displaying
718 negative food effects, where permeability, diffusivity or metabolism related limitations occur. It is
719 interesting to note that despite significant improvements in formulation design and characterisation
720 with regard to supersaturable and bioenabling formulations that many of these marketed
721 formulations still appear to behave sub-optimally *in vivo*, specifically with regard to food effects.

722 Formulating compounds to overcome food effect remains largely empirically driven, with only
723 sporadic case studies for individual compounds published. While the presence or absence of food
724 effects is unlikely to be a key driving factor in early formulation development, it can be a critical
725 factor when entering the clinic. In the absence of large databanks of formulation design studies in
726 easily obtainable literature, greater use of mechanistic and *in silico* approaches will be central to
727 enhancing our ability to discriminate between formulations likely to overcome food-mediated
728 alterations in drug bioavailability.

729 Acknowledgements

730 This work was supported by the European Union's Horizon 2020 Research and Innovation
731 Programme under grant agreement No 674909 (PEARRL)

732 References

- 733 1. Fleisher D et al. Drug, meal and formulation interactions influencing drug absorption after
734 oral administration - Clinical implications. *Clinical Pharmacokinetics* 1999; 3: 233-254.
- 735 2. Aulton ME, Taylor KM. *Aulton's Pharmaceutics: The Design and Manufacture of Medicines*:
736 Elsevier Health Sciences, 2017.
- 737 3. Varum FJO et al. Food, physiology and drug delivery. *International Journal of Pharmaceutics*
738 2013; 2: 446-460.
- 739 4. Welling PG. Influence of food and diet on gastrointestinal drug absorption: A review. *Journal*
740 *of Pharmacokinetics and Biopharmaceutics* 1977; 4: 291-334.
- 741 5. Welling PG. Effects of Food on Drug Absorption. *Pharmacology & Therapeutics* 1989; 3: 425-
742 441.
- 743 6. Winstanley PA, Orme MLE. THE EFFECTS OF FOOD ON DRUG BIOAVAILABILITY. *British*
744 *Journal of Clinical Pharmacology* 1989; 6: 621-628.
- 745 7. Schmidt LE, Dalhoff K. Food-drug interactions. *Drugs* 2002; 10: 1481-1502.
- 746 8. Mullertz A. Food Effects on Drug Absorption and Dosage Form Performance. In: Dressman, J.
747 B., Reppas, C. eds. *Oral Drug Absorption: Prediction and Assessment, 2nd Edition*. 2010: 90-
748 107.
- 749 9. Yasuji T et al. The effect of food on the oral bioavailability of drugs: a review of current
750 developments and pharmaceutical technologies for pharmacokinetic control. *Therapeutic*
751 *delivery* 2012; 1: 81-90.
- 752 10. Abuhelwa AY et al. Food, gastrointestinal pH, and models of oral drug absorption. *European*
753 *Journal of Pharmaceutics and Biopharmaceutics* 2017: 234-248.
- 754 11. Yan JH. Food Effect on Oral Bioavailability: Old and New Questions. *Clin Pharmacol Drug Dev*
755 2017.
- 756 12. C. Pentafragka MS, M. McAllister, J. Dressman, M. Vertzoni, C. Reppas. The Impact Of Food
757 Intake On The Luminal Environment And Performance Of Oral Drug Products With A View To
758 In Vitro And In Silico Simulations. *Journal of Pharmacy and Pharmacology* 2017; This issue.
- 759 13. FDA. Food-effect bioavailability and fed bioequivalence studies: guidance for industry. Food-
760 effect bioavailability and fed bioequivalence studies: guidance for industry
761 <http://www.fda.gov/downloads/RegulatoryInformation/Guidances/UCM126833.pdf>, 2002.
- 762 14. Amidon GL et al. A theoretical basis for a biopharmaceutic drug classification: the correlation
763 of in vitro drug product dissolution and in vivo bioavailability. *Pharm Res* 1995; 3: 413-20.
- 764 15. Wu C-Y, Benet LZ. Predicting Drug Disposition via Application of BCS:
765 Transport/Absorption/Elimination Interplay and Development of a Biopharmaceutics
766 DrugDisposition Classification System. *Pharmaceutical Research* 2005; 1: 11-23.
- 767 16. Custodio JM et al. Predicting drug disposition, absorption/elimination/transporter interplay
768 and the role of food on drug absorption. *Advanced Drug Delivery Reviews* 2008; 6: 717-733.
- 769 17. Benet LZ. The role of BCS (biopharmaceutics classification system) and BDDCS
770 (biopharmaceutics drug disposition classification system) in drug development. *J Pharm Sci*
771 2013; 1: 34-42.
- 772 18. McConnell EL et al. Gut instincts: Explorations in intestinal physiology and drug delivery.
773 *International Journal of Pharmaceutics* 2008; 2: 213-226.

- 774 19. European Medicines Agency. Guideline on the investigation of bioequivalence London, UK,
775 2010.
- 776 20. McLauchlan G et al. Comparison of gastric body and antral pH: a 24 hour ambulatory study
777 in healthy volunteers. *Gut* 1989; 5: 573-8.
- 778 21. Simonian HP et al. Regional postprandial differences in pH within the stomach and
779 gastroesophageal junction. *Dig Dis Sci* 2005; 12: 2276-85.
- 780 22. Abuhelwa AY et al. A Quantitative Review and Meta-Models of the Variability and Factors
781 Affecting Oral Drug Absorption-Part I: Gastrointestinal pH. *Aaps j* 2016; 5: 1309-21.
- 782 23. Schiller C et al. Intestinal fluid volumes and transit of dosage forms as assessed by magnetic
783 resonance imaging. *Alimentary Pharmacology & Therapeutics* 2005; 10: 971-979.
- 784 24. Kalantzi L et al. Characterization of the human upper gastrointestinal contents under
785 conditions simulating bioavailability/bioequivalence studies. *Pharmaceutical Research* 2006;
786 1: 165-176.
- 787 25. Koziolk M et al. Intra gastric Volume Changes after Intake of a High-Caloric, High-Fat
788 Standard Breakfast in Healthy Human Subjects Investigated by MRI. *Molecular*
789 *Pharmaceutics* 2014; 5: 1632-1639.
- 790 26. Koziolk M et al. Navigating the human gastrointestinal tract for oral drug delivery:
791 Uncharted waters and new frontiers. *Advanced Drug Delivery Reviews* 2016; Supplement C:
792 75-88.
- 793 27. Camilleri M. Integrated upper gastrointestinal response to food intake. *Gastroenterology*
794 2006; 2: 640-58.
- 795 28. Clarysse S et al. Postprandial Changes in Solubilizing Capacity of Human Intestinal Fluids for
796 BCS Class II Drugs. *Pharmaceutical Research* 2009; 6: 1456-1466.
- 797 29. Clarysse S et al. Postprandial Evolution in Composition and Characteristics of Human
798 Duodenal Fluids in Different Nutritional States. *Journal of Pharmaceutical Sciences* 2009; 3:
799 1177-1192.
- 800 30. Di Maio S, Carrier RL. Gastrointestinal contents in fasted state and post-lipid ingestion: In
801 vivo measurements and in vitro models for studying oral drug delivery. *Journal of Controlled*
802 *Release* 2011; 2: 110-122.
- 803 31. Hunt JN, Stubbs DF. The volume and energy content of meals as determinants of gastric
804 emptying. *The Journal of Physiology* 1975; 1: 209-225.
- 805 32. Siegel JA et al. Biphasic nature of gastric emptying. *Gut* 1988; 1: 85-89.
- 806 33. Oberle RL et al. The influence of the interdigestive migrating myoelectric complex on the
807 gastric emptying of liquids. *Gastroenterology* 1990; 5: 1275-82.
- 808 34. Mudie DM et al. Quantification of Gastrointestinal Liquid Volumes and Distribution
809 Following a 240 mL Dose of Water in the Fasted State. *Molecular Pharmaceutics* 2014; 9:
810 3039-3047.
- 811 35. Sunesen VH et al. Effect of liquid volume and food intake on the absolute bioavailability of
812 danazol, a poorly soluble drug. *European Journal of Pharmaceutical Sciences* 2005; 4: 297-
813 303.
- 814 36. Stotzer, Abrahamsson. Human postprandial gastric emptying of indigestible solids can occur
815 unrelated to antral phase III. *Neurogastroenterology & Motility* 2000; 5: 415-419.
- 816 37. Fadda HM et al. Meal-Induced Acceleration of Tablet Transit Through the Human Small
817 Intestine. *Pharmaceutical Research* 2009; 2: 356-360.
- 818 38. Davis SS et al. Transit of pharmaceutical dosage forms through the small intestine. *Gut* 1986;
819 8: 886-892.
- 820 39. Won CS et al. Mechanisms underlying food-drug interactions: Inhibition of intestinal
821 metabolism and transport. *Pharmacology & Therapeutics* 2012; 2: 186-201.
- 822 40. Gibbs MA, Hosea NA. Factors affecting the clinical development of cytochrome p450 3A
823 substrates. *Clin Pharmacokinet* 2003; 11: 969-84.

- 824 41. Deferme S, Augustijns P. The effect of food components on the absorption of P-gp
825 substrates: a review. *Journal of Pharmacy and Pharmacology* 2003; 2: 153-162.
- 826 42. Harris RZ et al. Dietary Effects on Drug Metabolism and Transport. *Clinical Pharmacokinetics*
827 2003; 13: 1071-1088.
- 828 43. Won CS et al. Influence of Dietary Substances on Intestinal Drug Metabolism and Transport.
829 *Current Drug Metabolism* 2010; 9: 778-792.
- 830 44. Jang GR, Harris RZ. Drug interactions involving ethanol and alcoholic beverages. *Expert Opin*
831 *Drug Metab Toxicol* 2007; 5: 719-31.
- 832 45. Singh BN. A quantitative approach to probe the dependence and correlation of food-effect
833 with aqueous solubility, dose/solubility ratio, and partition coefficient (Log P) for orally
834 active drugs administered as immediate-release formulations. *Drug Development Research*
835 2005; 2: 55-75.
- 836 46. Gu C-H et al. Predicting effect of food on extent of drug absorption based on
837 physicochemical properties. *Pharmaceutical Research* 2007; 6: 1118-1130.
- 838 47. Clarysse S et al. Intestinal drug solubility estimation based on simulated intestinal fluids:
839 Comparison with solubility in human intestinal fluids. *European Journal of Pharmaceutical*
840 *Sciences* 2011; 4: 260-269.
- 841 48. Augustijns P et al. A review of drug solubility in human intestinal fluids: Implications for the
842 prediction of oral absorption. *European Journal of Pharmaceutical Sciences* 2014: 322-332.
- 843 49. Stappaerts J et al. Human and simulated intestinal fluids as solvent systems to explore food
844 effects on intestinal solubility and permeability. *European Journal of Pharmaceutical*
845 *Sciences* 2014: 178-186.
- 846 50. Geboers S et al. The Effect of Food on the Intraluminal Behavior of Abiraterone Acetate in
847 Man. *Journal of Pharmaceutical Sciences* 2016; 9: 2974-2981.
- 848 51. Benet LZ et al. Unmasking the dynamic interplay between efflux transporters and metabolic
849 enzymes. *Int J Pharm* 2004; 1-2: 3-9.
- 850 52. Elgart A et al. Improved Oral Bioavailability of BCS Class 2 Compounds by Self Nano-
851 Emulsifying Drug Delivery Systems (SNEDDS): The Underlying Mechanisms for Amiodarone
852 and Talinolol. *Pharmaceutical Research* 2013; 12: 3029-3044.
- 853 53. Melander A, McLean A. Influence of Food Intake on Presystemic Clearance of Drugs. *Clinical*
854 *Pharmacokinetics* 1983; 4: 286-296.
- 855 54. Liedholm H et al. MECHANISMS AND VARIATIONS IN THE FOOD EFFECT ON PROPRANOLOL
856 BIOAVAILABILITY. *European Journal of Clinical Pharmacology* 1990; 5: 469-475.
- 857 55. Marasanapalle VP et al. Correlation between the systemic clearance of drugs and their food
858 effects in humans. *Drug Development and Industrial Pharmacy* 2011; 11: 1311-1317.
- 859 56. Charman WNA, Stella VJ. Estimating the maximal potential for intestinal lymphatic transport
860 of lipophilic drug molecules. *International Journal of Pharmaceutics* 1986; 1: 175-178.
- 861 57. Lawless E et al. Exploring the Impact of Drug Properties on the Extent of Intestinal Lymphatic
862 Transport - In Vitro and In Vivo Studies. *Pharmaceutical Research* 2015; 5: 1817-1829.
- 863 58. Kelly K et al. Comparison of the rates of disintegration, gastric emptying, and drug
864 absorption following administration of a new and a conventional paracetamol formulation,
865 using gamma scintigraphy. *Pharm Res* 2003; 10: 1668-73.
- 866 59. Cole ET et al. In vitro and in vivo pharmacoscintigraphic evaluation of ibuprofen
867 hypromellose and gelatin capsules. *Pharm Res* 2004; 5: 793-8.
- 868 60. Radwan A et al. Bio-predictive tablet disintegration: Effect of water diffusivity, fluid flow,
869 food composition and test conditions. *European Journal of Pharmaceutical Sciences* 2014:
870 273-279.
- 871 61. Yildiz HM et al. Food-associated stimuli enhance barrier properties of gastrointestinal
872 mucus. *Biomaterials* 2015: 1-8.

- 873 62. Radwan A et al. Mechanistic investigation of food effect on disintegration and dissolution of
874 BCS class III compound solid formulations: the importance of viscosity. *Biopharmaceutics &*
875 *Drug Disposition* 2012; 7: 403-416.
- 876 63. Radwan A et al. Mechanistic Understanding of Food Effects: Water Diffusivity in
877 Gastrointestinal Tract Is an Important Parameter for the Prediction of Disintegration of Solid
878 Oral Dosage Forms. *Molecular Pharmaceutics* 2013; 6: 2283-2290.
- 879 64. Radwan A et al. Food effect: The combined effect of media pH and viscosity on the
880 gastrointestinal absorption of ciprofloxacin tablet. *Eur J Pharm Sci* 2017: 100-106.
- 881 65. Gertz BJ et al. Studies of the oral bioavailability of alendronate. *Clin Pharmacol Ther* 1995; 3:
882 288-98.
- 883 66. Leyden JJ. Absorption of minocycline hydrochloride and tetracycline hydrochloride: Effect of
884 food, milk, and iron. *J Am Acad Dermatol* 1985; 2, Part 1: 308-312.
- 885 67. Polk RE. Drug-drug interactions with ciprofloxacin and other fluoroquinolones. *Am J Med*
886 1989; 5a: 76s-81s.
- 887 68. Neuvonen PJ et al. Interference of dairy products with the absorption of ciprofloxacin. *Clin*
888 *Pharmacol Ther* 1991; 5 Pt 1: 498-502.
- 889 69. Wallace AW, Amsden GW. Is it really OK to take this with food? Old interactions with a new
890 twist. *Journal of Clinical Pharmacology* 2002; 4: 437-443.
- 891 70. Jones HM et al. Predicting pharmacokinetic food effects using biorelevant solubility media
892 and physiologically based modelling. *Clinical Pharmacokinetics* 2006; 12: 1213-1226.
- 893 71. Sugano K et al. Prediction of food effect by bile micelles on oral drug absorption considering
894 free fraction in intestinal fluid. *European Journal of Pharmaceutical Sciences* 2010; 2: 118-
895 124.
- 896 72. Kataoka M et al. Application of Dissolution/Permeation System for Evaluation of Formulation
897 Effect on Oral Absorption of Poorly Water-Soluble Drugs in Drug Development.
898 *Pharmaceutical Research* 2012; 6: 1485-1494.
- 899 73. Holmstock N et al. Exploring food effects on indinavir absorption with human intestinal
900 fluids in the mouse intestine. *European Journal of Pharmaceutical Sciences* 2013; 1: 27-32.
- 901 74. Dresser GK et al. Fruit juices inhibit organic anion transporting polypeptide-mediated drug
902 uptake to decrease the oral availability of fexofenadine. *Clin Pharmacol Ther* 2002; 1: 11-20.
- 903 75. Dresser GK, Bailey DG. The effects of fruit juices on drug disposition: a new model for drug
904 interactions. *Eur J Clin Invest* 2003: 10-6.
- 905 76. Dresser GK et al. Effect of grapefruit juice volume on the reduction of fexofenadine
906 bioavailability: Possible role of organic anion transporting polypeptides. *Clinical*
907 *Pharmacology & Therapeutics* 2005; 3: 170-177.
- 908 77. Glaeser H et al. Intestinal drug transporter expression and the impact of grapefruit juice in
909 humans. *Clin Pharmacol Ther* 2007; 3: 362-70.
- 910 78. Greenblatt DJ. Analysis of drug interactions involving fruit beverages and organic anion-
911 transporting polypeptides. *J Clin Pharmacol* 2009; 12: 1403-7.
- 912 79. Kato Y et al. Involvement of influx and efflux transport systems in gastrointestinal absorption
913 of celiprolol. *J Pharm Sci* 2009; 7: 2529-39.
- 914 80. Bailey DG. Fruit juice inhibition of uptake transport: a new type of food-drug interaction. *Br J*
915 *Clin Pharmacol* 2010; 5: 645-55.
- 916 81. Frisk-Holmberg M et al. Effect of food on the absorption of vigabatrin. *British Journal of*
917 *Clinical Pharmacology* 1989; Suppl 1: 23S-25S.
- 918 82. Nøhr MK et al. Is oral absorption of vigabatrin carrier-mediated? *European Journal of*
919 *Pharmaceutical Sciences* 2015: 10-18.
- 920 83. Biogen Idec Ltd. Fampyra(R) [Summary of Product Characteristics]. Biogen Idec Ltd.,
921 Berkshire, UK; May 2017. 2017.
- 922 84. Bayer HealthCare Pharmaceuticals Inc. Stivarga(R) [Highlights of Prescribing Information].
923 Bayer HealthCare Pharmaceuticals Inc., Wayne, NJ; September 2012. 2012.

924 85. Vertex Pharmaceuticals (Europe) Ltd. Orkambi(R) [Summary of Product Characteristics].
925 Vertex Pharmaceuticals (Europe) Ltd., London, UK; November 2015. 2015.

926 86. Vertex Pharmaceuticals (Europe) Ltd. Kalydeco(R) [Summary of Product Characteristics].
927 Vertex Pharmaceuticals (Europe) Ltd., London, UK; April 2017. 2017.

928 87. Merck Sharp & Dohme Corp. Belsomra(R) [Highlights of Prescribing Information]. Merck
929 Sharp & Dohme Corp., Whitehouse Station, NJ; 2014. 2014.

930 88. Rainsford KD, Bjarnason I. NSAIDs: take with food or after fasting? *Journal of Pharmacy and*
931 *Pharmacology* 2012; 4: 465-469.

932 89. Moore RA et al. Effects of food on pharmacokinetics of immediate release oral formulations
933 of aspirin, dipyrrone, paracetamol and NSAIDs - a systematic review. *British Journal of Clinical*
934 *Pharmacology* 2015; 3: 381-388.

935 90. Moore RA et al. Validating speed of onset as a key component of good analgesic response in
936 acute pain. *Eur J Pain* 2015; 2: 187-92.

937 91. Thombre AG et al. Improved Ziprasidone Formulations with Enhanced Bioavailability in the
938 Fasted State and a Reduced Food Effect. *Pharmaceutical Research* 2011; 12: 3159-3170.

939 92. Szmulewitz RZ, Ratain MJ. Playing Russian Roulette With Tyrosine Kinase Inhibitors. *Clinical*
940 *Pharmacology & Therapeutics* 2013; 3: 242-244.

941 93. Kang SP, Ratain MJ. Inconsistent labeling of food effect for oral agents across therapeutic
942 areas: differences between oncology and non-oncology products. *Clin Cancer Res* 2010; 17:
943 4446-51.

944 94. Singh BN, Malhotra BK. Effects of food on the clinical pharmacokinetics of anticancer agents
945 - Underlying mechanisms and implications for oral chemotherapy. *Clinical Pharmacokinetics*
946 2004; 15: 1127-1156.

947 95. Ratain MJ, Cohen EE. The value meal: How to save \$1,700 per month or more on lapatinib.
948 *Journal of Clinical Oncology* 2007; 23: 3397-3398.

949 96. Ratain MJ. Flushing Oral Oncology Drugs Down the Toilet. *Journal of Clinical Oncology* 2011;
950 30: 3958-3959.

951 97. Todd M et al. Fast and Flawed or Scientifically Sound: The Argument for Administering Oral
952 Oncology Drugs During Fasting. *Journal of Clinical Oncology* 2012; 8: 888-889.

953 98. Deng J et al. To Take or Not to Take With Meals? Unraveling Issues Related to Food Effects
954 Labeling for Oral Antineoplastic Drugs. *Clin Pharmacol Drug Dev* 2017.

955 99. Halfdanarson TR, Jatoi A. Oral cancer chemotherapy: the critical interplay between patient
956 education and patient safety. *Curr Oncol Rep* 2010; 4: 247-52.

957 100. Koch KM et al. Reply to I.F. Tannock. *Journal of Clinical Oncology* 2009; 22: e43-e43.

958 101. Rahman A et al. The Value Meal: Effect of Food on Lapatinib Bioavailability. *Journal of*
959 *Clinical Oncology* 2007; 33: 5333-5334.

960 102. Jain RK et al. Food and oral antineoplastics: more than meets the eye. *Clin Cancer Res* 2010;
961 17: 4305-7.

962 103. Janssen Pharmaceuticals Inc. Xarelto(R) [Highlights of Prescribing Information]. Janssen
963 Pharmaceuticals Inc., Titusville, NJ., 2011.

964 104. Raman S, Polli JE. Prediction of positive food effect: Bioavailability enhancement of BCS class
965 II drugs. *International Journal of Pharmaceutics* 2016; 1-2: 110-115.

966 105. Lentz KA. Current methods for predicting human food effect. *Aaps Journal* 2008; 2: 282-288.

967 106. Paul SM et al. How to improve R&D productivity: the pharmaceutical industry's grand
968 challenge. *Nat Rev Drug Discov* 2010; 3: 203-14.

969 107. J. HL et al. The pig as a preclinical model for predicting oral bioavailability and in vivo
970 performance of pharmaceutical oral dosage forms: a PEARRL review. *Journal of Pharmacy*
971 *and Pharmacology* 2018; 0.

972 108. Lentz KA et al. Development and validation of a preclinical food effect model. *Journal of*
973 *Pharmaceutical Sciences* 2007; 2: 459-472.

974 109. Mathias N et al. Food Effect in Humans: Predicting the Risk Through In Vitro Dissolution and
975 In Vivo Pharmacokinetic Models. *Aaps Journal* 2015; 4: 988-998.

976 110. Hughes JP et al. Principles of early drug discovery. *British Journal of Pharmacology* 2011; 6:
977 1239-1249.

978 111. Pithavala YK et al. Evaluation of the effect of food on the pharmacokinetics of axitinib in
979 healthy volunteers. *Cancer Chemother Pharmacol* 2012; 1: 103-12.

980 112. Falcoz C et al. Pharmacokinetics of GW433908, a prodrug of amprenavir, in healthy male
981 volunteers. *J Clin Pharmacol* 2002; 8: 887-98.

982 113. Cundy KC et al. Clinical Pharmacokinetics of XP13512, a Novel Transported Prodrug of
983 Gabapentin. *The Journal of Clinical Pharmacology* 2008; 12: 1378-1388.

984 114. Anderson GD, Saneto RP. Modified-Release Formulations of Second-Generation Antiepileptic
985 Drugs: Pharmacokinetic and Clinical Aspects. *Cns Drugs* 2015; 8: 669-81.

986 115. Lal R et al. The effect of food with varying fat content on the clinical pharmacokinetics of
987 gabapentin after oral administration of gabapentin enacarbil. *International Journal of Clinical*
988 *Pharmacology and Therapeutics* 2010; 2: 120-128.

989 116. Pfizer Inc. Neurontin(R) [Highlights of Prescribing Information]. Pfizer Inc., New York, NY;
990 September 2015. 2015.

991 117. Gustafsson D et al. The Direct Thrombin Inhibitor Melagatran and Its Oral Prodrug H 376/95:
992 Intestinal Absorption Properties, Biochemical and Pharmacodynamic Effects. *Thrombosis*
993 *Research* 2001; 3: 171-181.

994 118. Chen M-L et al. Development Considerations for Nanocrystal Drug Products. *Aaps j* 2017: 1-
995 10.

996 119. Kesisoglou F et al. Nanosizing - Oral formulation development and biopharmaceutical
997 evaluation. *Advanced Drug Delivery Reviews* 2007; 7: 631-644.

998 120. Möschwitzer JP. Drug nanocrystals in the commercial pharmaceutical development process.
999 *International Journal of Pharmaceutics* 2013; 1: 142-156.

1000 121. Williams HD et al. Strategies to Address Low Drug Solubility in Discovery and Development.
1001 *Pharmacological Reviews* 2013; 1: 315-499.

1002 122. Shah DA et al. A Review: Pharmaceutical and Pharmacokinetic Aspect of Nanocrystalline
1003 Suspensions. *Journal of Pharmaceutical Sciences* 2016; 1: 10-24.

1004 123. Guivarc'h PH et al. A new fenofibrate formulation: Results of six single-dose, clinical studies
1005 of bioavailability under fed and fasting conditions. *Clinical Therapeutics* 2004; 9: 1456-1469.

1006 124. Sauron R et al. Absence of a food effect with a 145 mg nanoparticle fenofibrate tablet
1007 formulation. *International Journal of Clinical Pharmacology and Therapeutics* 2006; 2: 64-70.

1008 125. Wu Y et al. The role of biopharmaceutics in the development of a clinical nanoparticle
1009 formulation of MK-0869: a Beagle dog model predicts improved bioavailability and
1010 diminished food effect on absorption in human. *International Journal of Pharmaceutics*
1011 2004; 1-2: 135-146.

1012 126. Kesisoglou F, Wu Y. Understanding the Effect of API Properties on Bioavailability Through
1013 Absorption Modeling. *Aaps Journal* 2008; 4: 516-525.

1014 127. Femia RA, Goyette RE. The science of megestrol acetate delivery: potential to improve
1015 outcomes in cachexia. *BioDrugs* 2005; 3: 179-87.

1016 128. Deschamps B et al. Food effect on the bioavailability of two distinct formulations of
1017 megestrol acetate oral suspension. *Int J Nanomedicine* 2009: 185-92.

1018 129. Jinno J et al. Effect of particle size reduction on dissolution and oral absorption of a poorly
1019 water-soluble drug, cilostazol, in beagle dogs. *Journal of Controlled Release* 2006; 1-2: 56-64.

1020 130. Thombre AG et al. Solid Nanocrystalline Dispersions of Ziprasidone with Enhanced
1021 Bioavailability in the Fasted State. *Molecular Pharmaceutics* 2012; 12: 3526-3534.

1022 131. Yu P et al. Enhanced oral bioavailability and diminished food effect of lurasidone
1023 hydrochloride nanosuspensions prepared by facile nanoprecipitation based on dilution.
1024 *Powder Technology* 2017: 11-20.

- 1025 132. Quinn K et al. A formulation strategy for gamma secretase inhibitor ELND006, a BCS class II
1026 compound: Development of a nanosuspension formulation with improved oral
1027 bioavailability and reduced food effects in dogs. *Journal of Pharmaceutical Sciences* 2012; 4:
1028 1462-1474.
- 1029 133. Williams HD et al. Strategies to address low drug solubility in discovery and development.
1030 *Pharmacol Rev* 2013; 1: 315-499.
- 1031 134. Leuner C, Dressman J. Improving drug solubility for oral delivery using solid dispersions.
1032 *European Journal of Pharmaceutics and Biopharmaceutics* 2000; 1: 47-60.
- 1033 135. Vo CL-N et al. Current trends and future perspectives of solid dispersions containing poorly
1034 water-soluble drugs. *European Journal of Pharmaceutics and Biopharmaceutics* 2013; 3, Part
1035 B: 799-813.
- 1036 136. Mishra DK et al. Amorphous solid dispersion technique for improved drug delivery: basics to
1037 clinical applications. *Drug Deliv Transl Res* 2015; 6: 552-65.
- 1038 137. Wyttenbach N, Kuentz M. Glass-forming ability of compounds in marketed amorphous drug
1039 products. *European Journal of Pharmaceutics and Biopharmaceutics* 2017: 204-208.
- 1040 138. Baghel S et al. Polymeric Amorphous Solid Dispersions: A Review of Amorphization,
1041 Crystallization, Stabilization, Solid-State Characterization, and Aqueous Solubilization of
1042 Biopharmaceutical Classification System Class II Drugs. *Journal of Pharmaceutical Sciences*
1043 2016; 9: 2527-2544.
- 1044 139. Klein CE et al. The Tablet Formulation of Lopinavir/Ritonavir Provides Similar Bioavailability
1045 to the Soft-Gelatin Capsule Formulation With Less Pharmacokinetic Variability and
1046 Diminished Food Effect. *JAIDS Journal of Acquired Immune Deficiency Syndromes* 2007; 4:
1047 401-410.
- 1048 140. Hughes DL. Patent Review of Manufacturing Routes to Recently Approved PARP Inhibitors:
1049 Olaparib, Rucaparib, and Niraparib. *Organic Process Research & Development* 2017; 9: 1227-
1050 1244.
- 1051 141. European Medicines Agency. Lynparza [CHMP assessment report]. 2014.
- 1052 142. Plummer R et al. Olaparib tablet formulation: effect of food on the pharmacokinetics after
1053 oral dosing in patients with advanced solid tumours. *Cancer Chemother Pharmacol* 2015; 4:
1054 723-9.
- 1055 143. Rolfo C et al. Effect of Food on the Pharmacokinetics of Olaparib after Oral Dosing of the
1056 Capsule Formulation in Patients with Advanced Solid Tumors. *Adv Ther* 2015; 6: 510-22.
- 1057 144. Banerjee S et al. Formulation development and systematic optimization of stabilized
1058 ziprasidone hydrochloride capsules devoid of any food effect. *Pharmaceutical Development
1059 and Technology* 2016; 7: 775-786.
- 1060 145. Feeney OM et al. 50 years of oral lipid-based formulations: Provenance, progress and future
1061 perspectives. *Advanced Drug Delivery Reviews* 2016: 167-194.
- 1062 146. Dening TJ et al. Oral nanomedicine approaches for the treatment of psychiatric illnesses.
1063 *Journal of Controlled Release* 2016: 137-156.
- 1064 147. Charman WN et al. Effect of food and a monoglyceride emulsion formulation on danazol
1065 bioavailability. *J Clin Pharmacol* 1993; 4: 381-6.
- 1066 148. Mueller EA et al. Influence of a fat-rich meal on the pharmacokinetics of a new oral
1067 formulation of cyclosporine in a crossover comparison with the market formulation. *Pharm
1068 Res* 1994; 1: 151-5.
- 1069 149. Mueller EA et al. Improved dose linearity of cyclosporine pharmacokinetics from a
1070 microemulsion formulation. *Pharm Res* 1994; 2: 301-4.
- 1071 150. Webster GF et al. Comparative pharmacokinetic profiles of a novel isotretinoin formulation
1072 (isotretinoin-Lidose) and the innovator isotretinoin formulation: a randomized, 4-treatment,
1073 crossover study. *J Am Acad Dermatol* 2013; 5: 762-7.
- 1074 151. Perlman ME et al. Development of a self-emulsifying formulation that reduces the food
1075 effect for torcetrapib. *International Journal of Pharmaceutics* 2008; 1-2: 15-22.

- 1076 152. Christiansen ML et al. Effect of food intake and co-administration of placebo self-
1077 nanoemulsifying drug delivery systems on the absorption of cinnarizine in healthy human
1078 volunteers. *European Journal of Pharmaceutical Sciences* 2016; 77-82.
- 1079 153. Hong JY et al. A new self-emulsifying formulation of itraconazole with improved dissolution
1080 and oral absorption. *Journal of Controlled Release* 2006; 2: 332-338.
- 1081 154. Grove M et al. Bioavailability of seocalcitol: III. Administration of lipid-based formulations to
1082 minipigs in the fasted and fed state. *European Journal of Pharmaceutical Sciences* 2007; 1: 8-
1083 15.
- 1084 155. Nielsen FS et al. Bioavailability of probucol from lipid and surfactant based formulations in
1085 minipigs: Influence of droplet size and dietary state. *European Journal of Pharmaceutics and*
1086 *Biopharmaceutics* 2008; 2: 553-562.
- 1087 156. Tan A et al. Hybrid Nanomaterials that Mimic the Food Effect: Controlling Enzymatic
1088 Digestion for Enhanced Oral Drug Absorption. *Angewandte Chemie-International Edition*
1089 2012; 22: 5475-5479.
- 1090 157. Christiansen ML et al. Cinnarizine food-effects in beagle dogs can be avoided by
1091 administration in a Self Nano Emulsifying Drug Delivery System (SNEDDS). *European Journal*
1092 *of Pharmaceutical Sciences* 2014: 164-172.
- 1093 158. Han SD et al. Reduced Food-Effect on Intestinal Absorption of Dronedarone by Self-
1094 microemulsifying Drug Delivery System (SMEDDS). *Biological & Pharmaceutical Bulletin*
1095 2015; 7: 1026-1032.
- 1096 159. Li Y et al. Nanomulsion of megestrol acetate for improved oral bioavailability and reduced
1097 food effect. *Archives of Pharmacol Research* 2015; 10: 1850-1856.
- 1098 160. O'Shea JP et al. Lipidic dispersion to reduce food dependent oral bioavailability of
1099 fenofibrate: In vitro, in vivo and in silico assessments. *European Journal of Pharmaceutics*
1100 *and Biopharmaceutics* 2015: 207-216.
- 1101 161. Dening TJ et al. Silica encapsulated lipid-based drug delivery systems for reducing the
1102 fed/fasted variations of ziprasidone in vitro. *European Journal of Pharmaceutics and*
1103 *Biopharmaceutics* 2016: 33-42.
- 1104 162. Kamboj S, Rana V. Quality-by-design based development of a self-microemulsifying drug
1105 delivery system to reduce the effect of food on Nelfinavir mesylate. *International Journal of*
1106 *Pharmaceutics* 2016; 1-2: 311-325.
- 1107 163. Miao YF et al. Enhanced oral bioavailability of lurasidone by self-nanoemulsifying drug
1108 delivery system in fasted state. *Drug Development and Industrial Pharmacy* 2016; 8: 1234-
1109 1240.
- 1110 164. Rao S et al. Perspective and potential of oral lipid-based delivery to optimize
1111 pharmacological therapies against cardiovascular diseases. *Journal of Controlled Release*
1112 2014: 174-187.
- 1113 165. Carrier RL et al. The utility of cyclodextrins for enhancing oral bioavailability. *Journal of*
1114 *Controlled Release* 2007; 2: 78-99.
- 1115 166. Van de Velde VJ et al. Effect of food on the pharmacokinetics of a new hydroxypropyl-beta-
1116 cyclodextrin formulation of itraconazole. *Pharmacotherapy* 1996; 3: 424-8.
- 1117 167. Barone JA et al. Food interaction and steady-state pharmacokinetics of itraconazole oral
1118 solution in healthy volunteers. *Pharmacotherapy* 1998; 2: 295-301.
- 1119 168. Barone JA et al. Enhanced bioavailability of itraconazole in hydroxypropyl-beta-cyclodextrin
1120 solution versus capsules in healthy volunteers. *Antimicrob Agents Chemother* 1998; 7: 1862-
1121 5.
- 1122 169. Thombre AG et al. In vitro and in vivo characterization of amorphous, nanocrystalline, and
1123 crystalline ziprasidone formulations. *International Journal of Pharmaceutics* 2012; 1-2: 8-17.
- 1124 170. Wang D et al. Preparation and Characterization of the Sulfobutylether- β -Cyclodextrin
1125 Inclusion Complex of Amiodarone Hydrochloride with Enhanced Oral Bioavailability in Fasted
1126 State. *Aaps Pharmscitech* 2016: 1-10.

1127 171. Kuentz M et al. Methodology of oral formulation selection in the pharmaceutical industry.
1128 *European Journal of Pharmaceutical Sciences* 2016; 136-163.

1129 172. Rabinow BE. Nanosuspensions in drug delivery. *Nat Rev Drug Discov* 2004; 9: 785-796.

1130 173. Ku MS. Use of the Biopharmaceutical Classification System in Early Drug Development. *Aaps*
1131 *j* 2008; 1: 208-212.

1132 174. Pandey P et al. From Bench to Humans: Formulation Development of a Poorly Water Soluble
1133 Drug to Mitigate Food Effect. *Aaps Pharmscitech* 2014; 2: 407-416.

1134 175. Van Peer A et al. The effects of food and dose on the oral systemic availability of
1135 itraconazole in healthy subjects. *Eur J Clin Pharmacol* 1989; 4: 423-6.

1136 176. Woo JS et al. Reduced food-effect and enhanced bioavailability of a self-microemulsifying
1137 formulation of itraconazole in healthy volunteers. *European Journal of Pharmaceutical*
1138 *Sciences* 2008; 2: 159-165.

1139 177. Ribas A et al. The Effects of a High-Fat Meal on Single-Dose Vemurafenib Pharmacokinetics.
1140 *Journal of Clinical Pharmacology* 2014; 4: 368-374.

1141 178. Finn A et al. Effect of dose and food on the bioavailability of cefuroxime axetil. *Biopharm*
1142 *Drug Dispos* 1987; 6: 519-26.

1143 179. Bagchus WM et al. Important effect of food on the bioavailability of oral testosterone
1144 undecanoate. *Pharmacotherapy* 2003; 3: 319-325.

1145 180. Nix DE et al. Pharmacokinetics and relative bioavailability of clofazimine in relation to food,
1146 orange juice and antacid. *Tuberculosis (Edinb)* 2004; 6: 365-73.

1147 181. Schaad-Lanyi Z et al. Pharmacokinetics of clofazimine in healthy volunteers. *Int J Lepr Other*
1148 *Mycobact Dis* 1987; 1: 9-15.

1149 182. Oh DA et al. Effect of food on the pharmacokinetics of dronabinol oral solution versus
1150 dronabinol capsules in healthy volunteers. *Clinical Pharmacology-Advances and Applications*
1151 2017: 9-17.

1152 183. Reitberg DP et al. Effect of food on nifedipine pharmacokinetics. *Clin Pharmacol Ther* 1987;
1153 1: 72-5.

1154 184. Schering Corporation. Prometrium(TM) Clinical Pharmacology and Biopharmaceutics
1155 Review; NDA 20-843. 1997.

1156 185. Simon JA et al. The absorption of oral micronized progesterone: the effect of food, dose
1157 proportionality, and comparison with intramuscular progesterone. *Fertil Steril* 1993; 1: 26-
1158 33.

1159 186. Gibbons JA et al. Clinical Pharmacokinetic Studies of Enzalutamide. *Clinical Pharmacokinetics*
1160 2015; 10: 1043-1055.

1161

1162