

Title	Pharmacotherapy for neonatal seizures: current knowledge and future perspectives
Authors	Donovan, Maria D.;Griffin, Brendan T.;Kharoshankaya, Liudmila;Cryan, John F.;Boylan, Geraldine B.
Publication date	2016-03-04
Original Citation	Donovan, M. D., Griffin, B. T., Kharoshankaya, L., Cryan, J. F. and Boylan, G. B. [2016] 'Pharmacotherapy for Neonatal Seizures: Current Knowledge and Future Perspectives', <i>Drugs</i> , 76(6), pp. 647-661. doi:10.1007/s40265-016-0554-7
Type of publication	Article (peer-reviewed)
Link to publisher's version	10.1007/s40265-016-0554-7
Rights	© Springer International Publishing Switzerland 2016. The final publication is available at Springer via http://dx.doi.org/10.1007/s40265-016-0554-7
Download date	2026-09-19 02:46:24
Item downloaded from	https://hdl.handle.net/10468/3537



UCC

University College Cork, Ireland
 Coláiste na hOllscoile Corcaigh

Pharmacotherapy for neonatal seizures: current knowledge and future perspectives

Maria D. Donovan^{1, 2}, Brendan T. Griffin¹, Liudmila Kharoshankaya^{3, 4}, John F. Cryan^{2, 5},
Geraldine B. Boylan^{3, 4}.

¹Pharmacodelivery group, School of Pharmacy, University College Cork, Cork, Ireland

²Department of Anatomy and Neuroscience, University College Cork, Cork, Ireland

³Irish Centre for Fetal and Neonatal Translational Research, University College Cork and Cork University Maternity Hospital, Cork, Ireland.

⁴Department of Paediatrics and Child Health, University College Cork, Cork, Ireland.

⁵Alimentary Pharmabiotic Centre, University College Cork, Cork, Ireland.

Corresponding author: Geraldine Boylan; Email: g.boylan@ucc.ie; Telephone: +353 21 4205040; Fax: +353 21

Running head: Neonatal seizures: current and future treatment strategies

Abstract

Seizures are the most common neurological emergencies in the neonatal period and are associated with poor neurodevelopmental outcomes. Seizures affect up to five per 1000 term births and population based studies suggest that they occur even more frequently in premature infants. Seizures are a sign of an underlying cerebral pathology, the most common of which is hypoxic-ischemic encephalopathy in term infants. Due to a growing body of evidence that seizures exacerbate cerebral injury, effective diagnosis and treatment of neonatal seizures is of paramount importance to reduce long-term adverse outcomes. Electroencephalography is essential for the diagnosis of seizures in neonates due to their subtle clinical expression, non-specific neurological presentation and a high frequency of electro-clinical uncoupling in the neonatal period. Hypoxic-ischaemic encephalopathy may require neuroprotective therapeutic hypothermia, accompanying sedation with opioids, anticonvulsant drugs or a combination of all of these. The efficacy, safety, tolerability and pharmacokinetics of seven anticonvulsant drugs (phenobarbital, phenytoin, levetiracetam, lidocaine, midazolam, topiramate and bumetanide) are reviewed. This review is focused only on studies reporting electrographically confirmed seizures and highlights the knowledge gaps that exist in optimal treatment regimens for neonatal seizures. Randomised controlled trials are needed to establish a safe and effective treatment protocol for neonatal seizures.

Key points:

- The optimal treatment protocol for neonatal seizures remains elusive. Phenobarbital remains the first-line antiepileptic of choice, despite suboptimal efficacy and altered pharmacodynamic effects in neonates. There is currently no consensus regarding second-line drug choice, which often varies between phenytoin, lidocaine, levetiracetam or benzodiazepines.

- Hypothermia is the current standard of care for neuroprotection in HIE and many novel neuroprotective drugs are also emerging. Drug-drug interactions as well as drug-hypothermia interactions between antiepileptic drugs, novel neuroprotectants and hypothermia need to be investigated prior to administration in neonates, due to the potential for both pharmacokinetic and pharmacodynamic interactions.
- Continuous EEG monitoring is essential as a measure of antiepileptic efficacy.
- Randomised, controlled trials are required to establish a safe and effective treatment regimen for neonatal seizures.

1. Neonatal seizures-an overview

Neonatal seizures affect between one and five full-term neonates per 1000 live births, and are the most common neonatal neurological emergencies [1]. Moreover, the incidence of seizures increases in very low birth weight infants [2]. Phenobarbital remains the first line drug for treatment of neonatal seizures, despite having only around 50% efficacy [3]. There is little consensus about the best second-line treatment for neonatal seizures and there is considerable off-label use of antiepileptic medications with sparse efficacy data in the neonatal period.

1.1 Aetiology of neonatal seizures

Seizures are a hallmark of neurological injury and approximately 60% of all neonatal seizures are attributable to hypoxic-ischaemic encephalopathy (HIE) [4]. In Europe, HIE is the third most common cause of neonatal mortality, accounting for 9% of all deaths and 21% of term deaths, while globally it is estimated to cause approximately one million neonatal deaths each year [5, 6]. For the survivors of HIE, there is significant secondary morbidity, including cerebral palsy (29%), cognitive delay (45%), seizure disorders (12%), sensorineural deafness (9%), and visual loss (26%) [7]. Seizures in HIE may exacerbate the underlying cerebral injury and increase the risk of detrimental neurodevelopmental consequences [8-11]. Perinatal arterial

ischaemic stroke is the second most common cause of seizures in term neonates and accounts for 7.5%-20% of neonatal seizures [12, 13]. Seizures also arise from intracranial haemorrhage (7%-18%), congenital cerebral malformations (3%-17%), infection (2%-14%), metabolic causes (3%-5%), electrolyte imbalances (1%-4%) and other less common causes [14, 15]. There is little evidence on which to base recommendations for antiepileptic protocols regardless of seizure aetiology, although a prudent approach would be to treat the underlying cause and administer antiepileptic treatment according to hospital protocols.

1.2 Pathophysiological aspects of neonatal seizures

Seizures are the result of excessive electrical firing of neurons in the brain [16]. The immature brain is more susceptible to seizures, primed by the early development of excitatory neurotransmitters, delayed inhibitory function of gamma-amino butyric acid (GABA) and an excess of excitatory glutamatergic neurons which are composed of more excitable subunits than the equivalent in adults [17, 18].

The binding of GABA agonists/modulators to the GABA_A receptor triggers either an influx or an efflux of chloride ions, depending on the neuronal equilibrium potential for chloride [19]. It has been shown that the expression of inward sodium-potassium-chloride cotransporter (NKCC1) in human cortex is increased at birth compared to one year of age, whereas expression of outward potassium-chloride cotransporter (KCC2) increases from birth onwards [17]. This leads to an accumulation of intracellular chloride in immature neurons through NKCC1, thus the equilibrium potential for chloride becomes positive in relation to the resting membrane potential [19]. In immature neurons, activation of the GABA_A receptor results in chloride efflux and neuronal depolarisation [19]. Furthermore, birth injuries such as ischaemia increases NKCC1 and decrease KCC2 expression, whereas hypoxic-ischaemic injury increases NKCC1 alone [20]. For these reasons, treatment of neonatal seizures with GABA_A agonists, such as phenobarbital or benzodiazepines, may be suboptimal. Moreover, GABA_A receptors

are expressed at low levels in human and rodent cortex and contain less α_1 subunits than their adult counterparts, decreasing their sensitivity to modulation by benzodiazepines [21]. Female rats have increased levels of outward potassium-chloride transporter (KCC2), which translates to an inhibitory GABA action emerging earlier in females than in males [22]. Indeed, sex differences have been noted in mice, rats and humans with regards to susceptibility to brain injury, mechanisms of brain injury and response to treatment [23]. The increased seizure susceptibility due to developmental peculiarities of immature brain and excitatory GABA function might suggest that a class of antiepileptic drug (AED) other than GABA modulators should be considered as a first-line treatment for neonatal seizures. Furthermore, the sex differences observed in cotransporter expression raise questions regarding a differential approach to seizure treatment in male and female subjects.

1.3 Diagnosis of neonatal seizures

The diagnosis of neonatal seizures is challenging. Clinical seizure detection may lead to both over- and under-diagnosis [24]. Apart from classical tonic, clonic and myoclonic seizures, neonates may exhibit a wide variety of subtle seizure presentations including eye deviations, blinking, staring, chewing, sucking, cycling and boxing limb movements, apnoea and blood pressure changes [25]. Only a small portion of suspected neonatal clinical seizures are confirmed by electroencephalography (EEG) [24], while clinical signs may be absent in up to 80-90% of electrographic seizures [26, 27]. The only randomised controlled trial comparing the effect of treatment of electrographic-only seizures to clinical-only seizures in neonates with HIE using a traditional AED protocol (phenobarbital up to 40 mg/kg, followed by fosphenytoin 20 mg/kg and third line midazolam bolus or infusion) demonstrated significantly reduced seizure burden in neonates treated based on electrographic seizure activity [28].

The absence or cessation of clinical correlates when electrographic seizures are confirmed is called electro-clinical uncoupling [29, 30]. The subtle clinical seizure presentation in neonates

and the phenomenon of electro-clinical uncoupling may be at least partly explained by the incomplete axonal dendritic and synaptic development, as well as incomplete myelination in the immature brain. Synaptic connectivity continues to increase until 2 years of age [31, 32]. Clinical seizures can become even more difficult to detect following the administration of anticonvulsants or sedative agents, during hypothermia treatment or in neonates in critical condition [30, 33]. Both phenobarbital and phenytoin produced equal rates of uncoupling, with 58% of neonates exhibiting only or mostly electrographic evidence of seizures after drug administration [30]. Differential maturation of transporters that control intracellular chloride levels in different regions of the brain could be the mechanism underlying AED-induced uncoupling. Phenobarbital, a GABA_A agonist, reduced epileptiform power in slices taken from the ventroposterior thalamus of postnatal day 9/10 rat pups, but had no such effect on slices of neocortex from the same animals, suggesting that GABA signalling is inhibitory in the ventroposterior thalamus, but may be excitatory in the neocortex at this age [29].

A simplified and compressed version of multichannel EEG called amplitude-integrated EEG (aEEG) uses fewer channels than traditional EEG and requires less expertise for interpretation. It is often used for diagnosis of neonatal seizures in the neonatal intensive care unit (NICU) [1]. However, some seizures may be missed using this technology, as it struggles to identify low amplitude, short duration (< 1 minute) and infrequent seizures [27, 34, 35]. In addition, neonatal seizures often remain focal and do not generalise [27]. Therefore, focal seizures in the regions beyond aEEG electrode placement sites may remain undetected. Furthermore, artefacts that mimic seizure activity on aEEG may cause additional complications and lead to false positive readings [36]. Experience is required for reliable interpretation of both clinical and electrographic seizures, and studies have shown that non-expert users perform poorly in aEEG seizure detection [37]. There has been considerable effort in recent years to develop an automated neonatal seizure detection system to aid in clinical decision support in the NICU

144 and one such algorithm is currently undergoing a clinical trial across Europe (NCT02160171)
145 [38, 39].

146 Reliable diagnosis of neonatal seizures can only be performed using continuous EEG (cEEG)
147 monitoring which is considered the gold standard for the diagnosis of all neonatal seizures and
148 for the assessment of anticonvulsant efficacy [24]. The role of cEEG monitoring extends to the
149 differential diagnosis of seizure aetiology, particularly for HIE, stroke, infantile
150 encephalopathy and congenital metabolic diseases [40, 41]. Multichannel cEEG monitoring of
151 neonates at risk of seizures or suspected clinical seizures should be implemented rapidly to
152 confirm diagnosis and optimise outcomes [42]. Laboratory tests and magnetic resonance
153 imaging are also required to determine the underlying seizure pathology [43]. A protocol for
154 laboratory workup in seizures is detailed in a previously published review [25].

155

2. Treatment strategies

Once neonatal seizures are suspected, the neonate should be rapidly assessed for treatable underlying causes, such as hypoglycaemia or electrolyte disturbances [44]. AEDs are then administered according to clinical preference, independent of seizure cause. AEDs should only be initiated once seizure activity is confirmed, due to a lack of evidence for any positive outcomes if they are administered in the absence of seizures [3, 45].

As HIE is responsible for the majority of neonatal seizures and seizures are treated with the same AEDs regardless of underlying injury, the various treatments available for HIE-induced seizures are reviewed here. Neuro-protective strategies, currently led by therapeutic hypothermia, are initiated during the latent phase of HIE and may interact with AEDs that are administered during the secondary phase of HIE, and are therefore briefly mentioned in this context (Section 3).

2.1 Drug treatment for neonatal seizures

Neonatal seizures are neurological emergencies and must be treated promptly since seizures, particularly high seizure burden, may exacerbate neuronal injury in the immature brain and contribute to pathogenesis of later cerebral palsy and epilepsy [10, 46, 47]. In neonates with HIE who do not receive therapeutic hypothermia, there is a peak in seizure burden shortly after seizure onset (within six hours) [48]. AEDs should ideally be administered within the time period prior to the peak seizure burden. However, current AEDs are sub-optimal in terms of effectiveness, safety and long-term outcomes [3, 49] and a systematic review has shown that the use of AEDs following perinatal asphyxia in the absence of confirmed seizures are of little benefit with no improvement in survival or neurodevelopmental outcome [45]. AEDs used in neonates act through a variety of mechanisms to reduce excitability in the brain, thereby suppressing the seizure. The mode of action of neonatal AEDs is illustrated in Fig. 1.

180 **Insert Figure 1 here**

181

2.2 Antiepileptic Drugs: Efficacy, Safety and Tolerability

The most frequently used AEDs in both term and preterm babies include phenobarbital, phenytoin, midazolam, lorazepam, clonazepam, and lidocaine [54]. Current recommendations suggest initiating anticonvulsant therapies in neonates with phenobarbital, adding either a benzodiazepine, phenytoin or lidocaine as a second-line agent if seizures continue [3] (Table 1). In a treatment protocol designed by Slaughter *et al.*, a similar treatment regimen is proposed starting with phenobarbital, followed by levetiracetam, phenytoin or lidocaine, and finally the addition of a benzodiazepine as a third-line agent [55]. In other studies, if seizures were not controlled by phenobarbital and/or phenytoin, drugs such as midazolam, clonazepam, lidocaine, levetiracetam and topiramate have been used [42, 55-57]. A survey of clinicians in USA found that a majority (73%) would use levetiracetam and/or topiramate despite limited knowledge about the pharmacokinetics of these drugs in newborn infants [58]. However, topiramate was shown to exacerbate cell apoptosis caused by phenytoin in rat pups, despite the absence of neurodegenerative properties when administered as monotherapy [59]. Thus, certain AED combinations may be detrimental to neurodevelopment. While the use of other AEDs (carbamazepine, paraldehyde, sodium valproate, vigabatrin, lamotrigine) in the treatment of neonatal seizures has been described in case reports [60, 61] and recent animal studies have shown a beneficial anti-seizure effect of potassium channel opener flupirtine in a hypoxia-model of neonatal seizures [62], we will focus on AEDs that have been recommended in neonatal treatment protocols and that have been studied in conjunction with EEG monitoring. AED efficacy is defined differently in many of the studies cited in Sections 2.2, 2.3 and Table 1, but the vast majority state that efficacy is an 80% reduction in seizure severity or complete seizure cessation, with one notable exception that defined 50% seizure reduction as efficacious [33]. However, further work is required to define AED efficacy optimally using EEG criteria

in view of the well described natural evolution of acute seizures in neonates, particularly those with HIE [48, 63, 64].

2.2.1 Phenobarbital and phenytoin

Phenobarbital remains the first choice of AED in neonatal seizures, due to an extensive history of its use in this population [3]. Phenobarbital acts by increasing GABA_A mediated inhibition [51]. Neonates with persistent seizures are likely to have more severe brain damage and poor neurodevelopmental outcomes; thus half of the babies on two AEDs and a staggering 95% of babies on three AEDs were reported to have poor outcomes [47, 65]. Phenytoin, an antiepileptic that reduces excitatory neurotransmission by blocking a voltage-gated sodium channel, is often administered second-line to phenobarbital [51]. A Cochrane review found that there was very little supportive evidence for the main AEDs currently used in the neonatal period, as even with a combination treatment with phenobarbital and phenytoin, seizures remained in up to 50% of babies, as confirmed by cEEG [42, 49, 66, 67].

2.2.2 Lidocaine

Lidocaine acts by inhibiting voltage-gated sodium channels, thereby preventing depolarisation [50]. Lidocaine is a promising AED in neonatal seizures administered either second-line or third line with efficacy rates as high as 78%, based on aEEG assessment [68-70]. A very recent retrospective study of aEEG data has found that lidocaine as a second- or third-line AED had a good (seizure control for at least four hours) or intermediate (seizure control for at least two hours) antiepileptic effect in 71.4% of neonates, both term and preterm [70]. An earlier study demonstrated the lower efficacy rate of 60% with lidocaine, supported by cEEG [42]. One of the main challenges of using lidocaine in neonates is the risk of adverse events, particularly with plasma concentrations >9mg/L, including both bradycardia and ventricular tachycardia [68, 71]. As with many AEDs, a tailored neonatal dosage regimen is needed, as cardio-toxic levels were found in the majority of neonates treated with a standard lidocaine infusion [68].

A neonate-specific regimen was designed using pharmacokinetic modelling, and optimal lidocaine plasma levels were achieved in the majority of treated full-term neonates [68]. Furthermore, lidocaine dosing was studied in both term and preterm neonates, and it was found that both cohorts of neonates should receive approximately 50% of the previously recommended dose i.e. a 1 kg neonate should receive 52mg as opposed to 110mg [72]. However, lidocaine demonstrated a good safety profile in neonates [73].

2.2.3 Benzodiazepines

Benzodiazepines have had varied success as second- and third-line agents in the treatment of neonatal seizures. Benzodiazepines allosterically modulate the chloride channel in the GABA_A receptor to increase inhibitory neurotransmission [51]. Midazolam response rates vary from 0-100%, with both 0% and 100% efficacy being observed using cEEG monitoring (see Table 1) [42, 67]. Efficacy rates measured by aEEG are reported as 50% when midazolam is used as a second-line AED, increasing to 73-100% when administered as a third-line AED [69, 74]. Midazolam appears to be less effective than lidocaine at treating persistent seizures, particularly those caused by the most severe form of HIE [69, 75].

The evidence for the effect of other benzodiazepines used in neonatal seizures is less convincing [55]. Clonazepam did not abolish any seizures as a second-line AED in three neonates monitored by cEEG [42]. The support for lorazepam as an AED is sparse, with less than half of the studied neonates monitored by cEEG [76, 77]. Seizure control rates were as high as 86% and 100% in two studies, but these results are unreliable due to the absence of cEEG monitoring [76, 77].

2.2.4 Levetiracetam

Levetiracetam is a relatively new AED which is proposed to act through synaptic vesicle glycoprotein 2A (SV2A) which is a protein thought to be involved in the release of

neurotransmitters [78]. Levetiracetam is efficacious in treating various seizures in both adults and children. In addition, levetiracetam has a very favourable pharmacokinetic and safety profile in neonates [79, 80]. Levetiracetam has demonstrated some efficacy as a neonatal and paediatric AED, according to cEEG findings which show 35-64% efficacy within 24 hours, rising to improvements in 52-100% of patients in 72 hours [33, 56]. Levetiracetam was initiated as a second- or third- line AED in the majority of recorded cases [33]. Evidence from randomised-controlled trials is needed to endorse levetiracetam as a safe and effective AED. A trial is ongoing in America looking at the safety, efficacy and pharmacokinetic profile of levetiracetam in neonates (NCT01720667), with more efficacy/safety trials planned in France (NCT02229123) and China (NCT02550028) [38].

2.2.5 Topiramate

Topiramate reduces the frequency of action potential firing by altering GABA neurotransmission, blocking voltage-gated sodium channels and by weakly blocking AMPA glutamate receptors [81]. Similar to levetiracetam, pharmacokinetic and safety profiles are favourable, but little is known about the safety, efficacy or pharmacokinetics in a critically-ill newborn population [57]. In a small, retrospective study, topiramate was considered an effective add-on agent in neonatal seizures in four out of six neonates, and no major safety concerns were highlighted [57]. However, this study was limited by the lack of EEG monitoring [57].

2.3 Potential adjunct antiepileptics

2.3.1 Bumetanide

Bumetanide is a potential adjunct to AED treatments for neonatal seizures [82]. A number of years ago, bumetanide was observed to have antiepileptic effects in kainic acid-induced seizures *in vivo* [83]. This was believed to be due to its ability to block ion cotransporters in neurons and glia of the central nervous system, which in turn affected GABA signalling [83].

280 Bumetanide blocks NKCC co-transporters, NKCC1 and NKCC2, which both move chloride
281 into cells [84]. Bumetanide was originally developed as a loop diuretic, which reduces oedema
282 by inhibiting the reabsorption of sodium, potassium and chloride through NKCC2 in the thick
283 ascending loop of Henle of the kidney [84]. Bumetanide also inhibits NKCC1, an isoform of
284 the NKCC cotransporter that is widely expressed, including on neurons in the brain [84].
285 GABA is excitatory in immature neurons due to the accumulation of chloride through NKCC1
286 [85]. By preventing intracellular chloride build-up, bumetanide is thought to decrease or even
287 reverse the excitatory action of GABA, thus presenting a potentially useful combination
288 therapy with GABAergic anticonvulsants [64, 82]. There are gaps in our knowledge of this
289 potential adjunct to AEDs for the treatment of neonatal seizures, namely the dose at which it
290 acts in the brain, the human blood-brain barrier permeability/transport of bumetanide as well
291 as its effect on development of the central nervous system (CNS). Two clinical trials were
292 initiated to establish the safety and efficacy of bumetanide in neonatal seizures, one in Europe
293 (NCT01434225) and one in the USA (NCT00830531) [38]. In the European dose-finding
294 clinical study, bumetanide was administered according to a bivariate Bayesian sequential dose-
295 escalation design, in which participants were treated with four doses of bumetanide
296 (0.05mg/kg-0.3mg/kg) each given twelve hours apart, with the first dose given in conjunction
297 with phenobarbital [64]. However, the trial was concluded early as the benefit: risk ratio was
298 no longer favourable and the efficacy endpoint was not achieved in any of the trial participants
299 [64, 86]. It has been suggested that this is partially due to a poor CNS effect of bumetanide at
300 the doses used and evidence to corroborate this have come from animal studies that indicate a
301 poor brain permeability of bumetanide [87]. Many studies are examining novel ways to
302 enhance brain levels of bumetanide in an effort to overcome the pharmacokinetic issues
303 hindering its therapeutic success [87-90].

Table 1: Drug treatments of neonatal seizures-efficacy, safety and tolerability

Drug	Place in treatment protocol	Efficacy	Safety	Tolerability
Phenobarbital	First-line.	Effective in 43% of neonates in a randomised controlled trial (n = 30) [66]. Phenobarbital achieved seizure control in 50% of neonates (n = 22) [42], and 47% of neonates in a further study (n = 32) [67]. Cost-effective AED [91].	May impair neurodevelopment and increase apoptosis of neurons [92]. Potential for drug-drug interactions.	Many CNS side-effects i.e. sedation, impaired cognition, depressed mood [91].
Phenytoin	Second-line.	Response in 45% of neonates to a dose that achieves a free plasma concentration of 3µg/mL (n = 29) [66].	Concerns about potential detrimental effect on developing neurons [60]. Potential for drug-drug interactions [55].	No changes in cardiac or respiratory function observed [66]
Levetiracetam	Emerging. Second- or third-line [3, 55]. Effect may be additive with other AEDs [93].	Effective with twice daily dosing. Efficacious in 82% of preterm neonates with seizures (n=11) [94]. Achieves full control of seizures in 33% (n = 18) [95], 35% (n = 23) [33] and between 32% and 100% of cases, depending on the treatment duration (n = 22) [56].	Does not cause neuronal apoptosis in rat pups [93].	Side effects in infants and children: somnolence and irritability. Well tolerated [79].
Lidocaine	Second- or third-line.	Response rate varies from 60% (n = 5) [42], to 71.4% (n = 413) [70], to 76% (n = 20) [68] and 77% (n = 22) [69]. Optimised dosing regimen achieved seizure control in 78% neonates (n = 15) [68].	Cardiac toxicity i.e. bradycardia-increased risk following other cardio-toxic agents e.g. phenytoin.	Risk of arrhythmias in 5% patients [68].
Midazolam	Second-line.	Reported response to treatment shows a wide variability from no response (n = 3) [42] to 50% (n = 8) [69] to 73% (n = 15) [74] to 100% response (n = 13) [67]. Improved neurodevelopment at 1 year of age compared to non-responders [67].	Higher doses or combination treatment with hypothermia may cause cardiac depression [69, 75]. Short-term drowsiness observed [67].	Well tolerated, no serious adverse effects noted [67].
Topiramate	Emerging.	Efficacy studies in neonates ongoing [81]. Efficacy of 67% in one small, retrospective study (n = 6), but no EEG monitoring so unreliable [57]. Hypothesised to have synergistic neuro-protective effects in neonates; reduces brain injury in animal models of HIE [96].	Seems safe- no increase in risk of death, short-term detrimental effects or gross brain pathology [97].	Well-tolerated, no adverse effects noted [96]

Bumetanide	Emerging.	Low efficacy rates of ~36% (n = 14) with research protocol doses of between 0.05 mg/kg and 0.3 mg/kg given 12 hours apart for a total of four doses-rescue AEDs were required by most neonates and efficacy endpoint not met by any trial participants [64, 86].	Potential risk of ototoxicity, especially if given concomitantly with other ototoxic drugs [64]. Other side effects include dehydration.	Well tolerated up to 0.1mg/kg dose [64].
------------	-----------	--	--	--

2.4 Pharmacokinetic properties of AEDs

There are a variety of physiological differences between neonates and adults. These variations in physiology affect all pharmacokinetic processes in the neonate, including absorption, distribution, metabolism and elimination [60]. These variations are detailed in a review by Alcorn *et al.*, but the salient changes are noted in Table 2. Key pharmacokinetic parameters, including volume of distribution (V_d), fraction unbound in plasma (f_u), clearance (Cl) and elimination half-life ($t_{1/2}$), are different in neonates compared to adults. Moreover, there is wide variability in these pharmacokinetic parameters within the neonatal population, as can be seen by the ranges reported (Table 3).

316 **Table 2: Physiological differences between neonates and adults**

Stage	Pharmacokinetic parameter	Neonate	Adult
Absorption	Gastric pH	6-8	2
	Gastric emptying time	Reduced rate	-
	Intestinal Motility	Reduced rate	-
Distribution	Body composition		
	❖ Water	74%	55-60%
	❖ Fat	14%	~20%
Metabolism	Plasma proteins		
	❖ Albumin	~75%	100%
	❖ α 1 acid glycoprotein	~25%	100%
Metabolism	Enzyme expression		
	❖ Foetal	0-10%	100%
	o ST, GST		
	❖ Early Neonatal	25-37.5%	100%
	o UGT, NAT		
	Cytochrome P activity		
	❖ Foetal		
	o 3A7	500-600%	100%
	❖ Early neonatal		
	o 2D6	4-24%	100%
Excretion	o 2E1	21-36%	100%
	❖ Neonatal		
	o 1A2	5-20%	100%
	o 2C	2-42%	100%
	o 3A4	3-29%	100%
	Bacterial flora	Very limited	100%
	Glomerular filtration rate	~90%	100%
	Tubular secretion	~50%	100%
	Renal bloodflow	~65%	100%

317

318

319

320 **Table 3: Drug treatments of neonatal seizures from published studies-pharmacokinetics**

AED	Dose	t _{1/2} (h)	f _u (%)	Cl (L/hr/kg)	V _d (L/kg)	Ref
Phenobarbital	D _L : 20mg/kg (twice if required) D _M : 5mg/kg/day	73.9-154.5	57-64	0.0053-0.0141	0.64-1.17	[55, 66, 99-102]
Phenytoin	D _L : 20mg/kg D _M : 5mg/kg/day	Wk 1: 20.7 ± 11.6 Wks 2-4: 7.6 ± 3.5	19.8 ± 2.6	0.00151-0.139	0.8 ± 0.26	[55, 103, 104]
Levetiracetam	D _L : 10-50 mg/kg D _M : 10-80 mg/kg/day (mean 45 mg/kg/day)	Day 1: 18.5 ± 7.1 Day 7: 9 ± 2	~100	0.042-0.078	0.89-1.01	[33, 55, 56, 79, 95]
Lidocaine	D _L : 2mg/kg D _M (normothermia): 5-7 mg/kg/h for 4 h, 2.5-3.5 mg/kg/h for 6-12 h, 1.25-1.75 mg/kg/h for 12 h *Altered D _M recommended in hypothermia-see ref. 54	5.2-5.4	20-40	0.462-1.68	3-3.2	[54, 60, 68, 72]
Midazolam	D _L : 0.05-0.15mg/kg D _M : 0.06-0.4 mg/kg/h	6.9	3.1	0.124	1-1.7	[55, 105]
Topiramate	D _M : 5mg/kg-10mg/kg daily	35.6 ± 19.3	~85	0.0156 ± 0.0048	0.6-1	[60, 81, 96, 102]

321 ^a t_{1/2}: half-life; fu: fraction unbound; Cl: clearance; V_d: volume of distribution; D_L: Loading dose; D_M: Maintenance dose; Wk: week.

3. Combining therapeutic strategies

3.1 Adjunct therapies in HIE with potential for interaction with antiepileptics

3.1.1 Hypothermia

Hypothermia has demonstrated neuro-protective properties in neonates with moderate to severe HIE [107-109]. Since the introduction of therapeutic hypothermia, the composite risk of death and major disability has been reduced by approximately 25% [108]. Neurological outcomes in cooled neonates with HIE improved at both 18 months and six-seven years of age [108, 109]. Hypothermia significantly reduces seizure burden, as measured by cEEG, in neonates with HIE [110, 111]. Seizure burden during hypothermia is characterised by a more even distribution over time (as opposed to the accumulation seen at seizure-onset in normothermia) and de-novo seizures may occur after re-warming [10, 48, 63]. It has been proposed that hypothermia should also be tested as a therapeutic strategy in late premature neonates with HIE and neonatal stroke, both of which can also result in seizures [112].

3.1.2 Emerging neuro-protective treatments

Additional neuro-protective strategies that are emerging include xenon, erythropoietin, melatonin, allopurinol and sevoflurane [113-119]. Thus far, the authors have found no reports of combination treatment with AEDs and emerging neuro-protective drugs. However, the combination of these novel neuro-protective agents, hypothermia and AEDs are a definite possibility in the future. Briefly, xenon protects the brain from excitatory injury by antagonising the N-methyl-D-aspartate (NMDA) glutamate receptor reducing total neurotransmission and is currently under investigation in a Phase 2 trial [120]. Erythropoietin has anti-inflammatory properties and is also anti-apoptotic [121, 122]. It has been shown to reduce detrimental neurodevelopmental outcomes in neonates with moderate-severe HIE [123]. Melatonin reduces oxidative stress through a variety of mechanisms, such as scavenging

oxygen free radicals and has been shown to augment neuroprotection by hypothermia in a piglet model of HIE [117]. Sevoflurane reduced hippocampal apoptosis in a rat model of intrauterine perinatal asphyxia and thus may be neuroprotective [113]. Allopurinol was found to have anti-oxidant properties [116].

3.1.3 Sedation

Intravenous morphine is commonly used as a sedative during hypothermia, as it reduces pain and stress, allows the patient to tolerate hypothermia and can be titrated to optimal response [124]. In a preclinical model of HIE, hypothermia without sedation lacked neuro-protective properties [125]. In a small group of term and preterm neonates without underlying brain injury, morphine infusion at a rate of 10-20 mcg/kg/hour was found to be associated with excessive epileptiform activity on cEEG [126].

It is known that morphine clearance is decreased during hypothermia, resulting in an increased concentration of morphine in both cerebrospinal fluid and plasma [127, 128]. In terms of pharmacodynamic considerations, the affinity of morphine for its receptor appears reduced in hypothermia, but the incidence of hypotension is increased [127, 129]. Neonates with HIE who are sedated with opioids show less brain injury and display better outcomes [130]. Little is known about drug-drug interactions with AEDs, but it is advised that barbiturates such as phenobarbital may increase the sedating effect of opioids [131].

3.2 Antiepileptics and hypothermia

It is thought that synergistic therapy including a traditional AED and hypothermia may augment neuro-protective properties of either treatment given alone [132]. Combination treatments need to be explored further to complement this claim. However, caution needs to be exercised as hypothermia may alter pharmacokinetics of AEDs in neonates by decreasing absorption, distribution or metabolism/clearance [100, 133-135]. Moreover, as multi-organ

dysfunction is frequently a characteristic of HIE, the combination of therapeutic hypothermia and organ impairment, particularly renal and hepatic, may have additive detrimental effects on fundamental pharmacokinetic processes [136]. The rewarming phase following hypothermia is another period of pharmacokinetic and pharmacodynamic uncertainty and is likely to be a window of time in which serious toxicity and adverse reactions could occur, due to a lag time between the return of normal metabolic enzyme and transporter function [135]. There have been reports of seizures re-occurring during the rewarming phase, but the affected infants were not receiving regular AEDs [10, 137]. Thus, combination treatment with hypothermia and AEDs may be useful, but must be approached with caution due to uncertainties regarding the effect of hypothermia on efficacy, safety and pharmacokinetics of such medications. It is important to identify AEDs, doses and dosage intervals that are suitable for neonates during and after hypothermia.

3.2.1 Phenobarbital and hypothermia

Positive synergism of first-line AED phenobarbital and hypothermia was observed in a rodent model of HIE, with both early and late assessment of neuropathology and sensorimotor performance demonstrating improvements [138]. However, current evidence suggests that this combination has not translated to a reduced risk of death or brain damage in neonates [139, 140]. Seizures were detected using aEEG, and a 66% reduction in seizures was demonstrated for neonates treated with hypothermia and with plasma concentrations of phenobarbital above 20mg/L [140].

The pharmacokinetics of phenobarbital were examined in hypothermic critically-ill neonates [100, 101]. It was found that minimum, maximum and average plasma concentrations were all larger in cooled neonates versus normothermia [100]. However, V_d and clearance remained unchanged [101]. It was concluded the alterations in pharmacokinetics of phenobarbital during hypothermia in neonates were not clinically significant, and that a total maximum dose of

40mg/kg can be safely administered in hypothermia prior to initiation of second-line AED [140]. In contrast, metabolism of phenobarbital via CYP2C19 was significantly reduced when it was administered to critically injured children who were cooled under more severe hypothermic conditions to 30-31°C [135, 141]. Therapeutic drug monitoring of phenobarbital allows for tight control of AED concentrations, which may be particularly important during hypothermia.

3.2.2 Lidocaine and hypothermia

Lidocaine was administered as a third-line anticonvulsant to neonates undergoing hypothermia treatment for asphyxia-induced seizures with aEEG monitoring. An impressive 91% of these patients responded to lidocaine [134]. This is a similar response rate to that observed in normothermic babies [42].

The pharmacokinetics of lidocaine are altered by hypothermia. Clearance of lidocaine is reduced by 24% as hepatic blood flow is reduced during hypothermia [134]. Despite these changes, no cardiotoxicity was observed in hypothermic neonates when an altered dosing regimen, equating to 70% of the total lidocaine dose given to normothermic neonates, was administered [134].

3.2.3 Topiramate and hypothermia

Animal studies suggested that the combination of topiramate and hypothermia improved motor and brain tissue damage in a model of HIE, where neither drug alone conferred any neuroprotection [142]. In neonates, there were no statistically significant changes in survival rate or brain damage observed when topiramate was given in combination with hypothermia when compared to hypothermia alone [97]. A randomised-controlled trial of topiramate and hypothermia in combination is underway, which will examine efficacy of seizure control with this treatment strategy (NCT01765218) [38, 81].

The pharmacokinetic profile of topiramate is altered when administered during hypothermia treatment: maximum, minimum and average concentrations, $t_{1/2}$ and area under the concentration-time curve are significantly higher in hypothermia [96]. However, these pharmacokinetic variations are not clinically significant, and the majority of neonates are observed to have topiramate concentrations within the safe, effective concentration range [96].

3.2.4 Midazolam and hypothermia

The efficacy of midazolam as a second-line AED in seizing neonates undergoing hypothermia treatment is modest, achieving seizure control in only 23% of neonates, confirmed using aEEG monitoring [75].

The pharmacokinetic profile of midazolam in neonates were not significantly changed by hypothermia [75, 143]. However, the incidence of midazolam-induced hypotension increased in neonates undergoing therapeutic hypothermia [75]. Midazolam levels in the serum of asphyxiated infants (both normothermic and hypothermic) were found to be highly variable and unpredictable, due to renal/hepatic impairment caused by the initial injury [143]. Furthermore, it is worth noting that the combination of midazolam and hypothermia in adults with disorders of the CNS gave rise to significant decreases in clearance and increases in V_d of midazolam compared to midazolam treatment alone [144].

3.2.5 Bumetanide and hypothermia

Bumetanide pharmacokinetics including clearance and V_d were calculated in a neonatal population [64]. These patients were also receiving hypothermia treatment and phenobarbital. Clearance values appear to generally be in agreement with values previously reported in a neonatal population [145, 146]. The combination of phenobarbital, bumetanide and hypothermia in a neonatal population with HIE-induced seizures was not effective, as none of

the neonates achieved the requisite 80% seizure reduction without the need for rescue AEDs [64, 86].

3.2.6 Phenytoin and hypothermia

There are no data from neonates with seizures on the efficacy or safety of phenytoin and hypothermia together. However, there are reports from trials on the use of phenytoin and hypothermia in older children aged 2-16 years, as well as adult patients, for the treatment of traumatic brain injury [147, 148]. In these populations, decreases in metabolism by CYP2C9 and CYP2C19 resulted in reduced clearance compared to values obtained after rewarming [135, 147, 148]. In children, it was also found that increased phenytoin levels are present both during and after rewarming which increased the risk of drug toxicity even after hypothermia had finished [147]. Moreover, a case report has described an additive bradycardic effect of therapeutic hypothermia and phenytoin [149]. Hypothermia in this case occurred during surgery and was not controlled. The authors hypothesise that the cardiac depressant effects of both treatments acted synergistically and that extreme caution should be exercised when co-administration is necessary [149].

4. Knowledge gaps

There is an urgent need for more randomised controlled trials in neonates to validate a treatment algorithm for seizures, especially when used in combination with hypothermia. Due to a lack of evidence from clinical studies, seizure treatments consist of older generation drugs that have more side-effects than newer drugs [3]. In general, efficacy rates of treatments are underwhelming (Table 1). Furthermore, there is a need to observe long-term neurodevelopmental outcomes following each of the proposed treatments, and to define the optimal length of time to continue with AED therapy given the concern regarding their effect on long-term brain development [16, 150].

There is a paucity of data on the pharmacokinetics and efficacy of many AEDs used in neonates, including levetiracetam, lidocaine and topiramate [151]. There are also major gaps in our knowledge about the efficacy and safety of most anticonvulsant drugs, particularly in preterm neonates.

In HIE it is imperative that the pharmacokinetics of AEDs during both active therapeutic hypothermia and the rewarming phase in neonates with seizures are elucidated, particularly with regards to phenytoin, a popular second-line AED [135]. Furthermore, drug-drug interactions are significantly under-investigated, especially co-administration of AEDs with novel neuro-protective drugs such as xenon, allopurinol, melatonin and erythropoietin.

Multichannel EEG is essential to accurately measure the efficacy of AEDs [24]. The optimal algorithm for detecting seizures remains to be developed, to enhance bedside recognition of seizures by non-EEG experts.

In the last decade there has been an increased interest in developing safe and effective drugs for neonates. The European Commission through the FP7 Framework promoted research on the safe and effective use of medicine in children by specifically supporting applications for

the neonatal age group [152]. More recently, the International Neonatal Consortium was launched in May 2015. This is a consortium of stakeholders focused on the development of effective medicines for neonates, including the Food and Drug Administration (FDA), European Medicines Agency (EMA), the pharmaceutical industry, academia, patient research groups, and family advocates. The consortium aims to align priorities and initiate collaborations to accelerate the development of safe and effective treatments for neonates. One of the first topics that this consortium has prioritised for further development is the treatment of neonatal seizures.

5. Conclusion

The treatment of neonatal seizures remains sub-optimal. Treatment algorithms are based on minimal trial data on older generation drugs. Phenobarbital remains the first-line antiepileptic of choice, despite suboptimal efficacy and altered underlying pharmacodynamics in the immature brain. However, there is no consensus on a replacement first-line drug or even on the most efficacious and suitable second-line AED. A lack of randomised controlled trials to guide treatment regimens in neonatal seizures is halting progress in the field. There are a multitude of drug-drug and drug-hypothermia interactions that remain to be elucidated, including the efficacy/safety of antiepileptic polypharmacy in neonates. These knowledge gaps have been identified and urgently need to be bridged by designing and conducting high quality clinical trials in neonates. Until the pharmacokinetic/pharmacodynamic profiles of antiepileptic medications in hypothermia are sufficiently investigated, therapeutic drug monitoring of serum antiepileptic levels is encouraged. The efficacy of antiepileptic treatment protocols should always be measured using cEEG monitoring.

Acknowledgments

The authors would like to extend their gratitude to Dr Roman Stilling for providing expert help with artwork.

Conflict of Interest

Maria Donovan, Brendan Griffin, Liudmila Kharoshankaya, John Cryan and Geraldine Boylan declare that they have no conflict of interest.

Compliance with Ethical Standards

Funding: MDD is in receipt of an Irish Research Council for Science Engineering and Technology scholarship. GBB was supported under European Community's Seventh Framework Programme (FP7/2007-2013) under grant agreement n° 241479 and by Science Foundation Ireland in the form of a centre grant (INFANT SFI/12/RC/2272). JFC is supported in part by Science Foundation Ireland in the form of a centre grant (Alimentary Pharmabiotic Centre Grant Number SFI/12/RC/2273), by the Health Research Board of Ireland (Grant Numbers HRA_POR/2011/23 and HRA_POR/2012/32) and by the European Community's Seventh Framework Programme, Grant no. FP7/2007-2013, Grant Agreement number 278948 (TACTICS – Translational Adolescent and Childhood Therapeutic Interventions in Compulsive Syndrome). No funding was specifically received for the publication of this article.

References

1. Glass HC. Neonatal seizures: advances in mechanisms and management. Clin Perinatol. 2014;41(1):177-90.
2. Lanska MJ, Lanska DJ. Neonatal seizures in the United States: results of the National Hospital Discharge Survey, 1980-1991. Neuroepidemiology. 1996;15(3):117-25.
3. World Health Organisation. Guideline on Neonatal Seizures. 2011.
http://apps.who.int/mental_health/publications/guidelines_neonatal_seizures/en/ Accessed 17th April 2015.
4. Volpe J. Neonatal Seizures. In: Volpe J, editor. Neurology of the Newborn. 5th ed. Philadelphia: WB Saunders; 2008. p. 203-44.
5. Lee AC, Kozuki N, Blencowe H, Vos T, Bahalim A, Darmstadt GL, Niermeyer S, Ellis M, Robertson NJ, Cousens S, Lawn JE. Intrapartum-related neonatal encephalopathy incidence and impairment at regional and global levels for 2010 with trends from 1990. Pediatr Res. 2013;74 Suppl 1:50-72.
6. Lawn JE, Cousens S, Zupan J. 4 million neonatal deaths: when? Where? Why? Lancet. 2005;365(9462):891-900.
7. Mwaniki MK, Atieno M, Lawn JE, Newton CR. Long-term neurodevelopmental outcomes after intrauterine and neonatal insults: a systematic review. Lancet. 2012;379(9814):445-52.
8. Miller SP, Weiss J, Barnwell A, Ferriero DM, Latal-Hajnal B, Ferrer-Rogers A, Newton N, Partridge JC, Glidden DV, Vigneron DB, Barkovich AJ. Seizure-associated brain injury in term newborns with perinatal asphyxia. Neurology. 2002;58(4):542-8.
9. Glass HC, Glidden D, Jeremy RJ, Barkovich AJ, Ferriero DM, Miller SP. Clinical Neonatal Seizures are Independently Associated with Outcome in Infants at Risk for Hypoxic-Ischemic Brain Injury. J Pediatr. 2009;155(3):318-23.

552 10. Shah DK, Wusthoff CJ, Clarke P, Wyatt JS, Ramaiah SM, Dias RJ, Becher JC, Kapellou
553 O, Boardman JP. Electrographic seizures are associated with brain injury in newborns
554 undergoing therapeutic hypothermia. *Arch Dis Child Fetal Neonatal Ed.* 2014;99(3):F219-24.

555 11. McBride MC, Laroia N, Guillet R. Electrographic seizures in neonates correlate with
556 poor neurodevelopmental outcome. *Neurology.* 2000;55(4):506-13.

557 12. Wu YW, Lynch JK, Nelson KB. Perinatal arterial stroke: understanding mechanisms and
558 outcomes. *Semin Neurol.* 2005;25(4):424-34.

559 13. Kirton A, Armstrong-Wells J, Chang T, Deveber G, Rivkin MJ, Hernandez M, Carpenter
560 J, Yager JY, Lynch JK, Ferriero DM. Symptomatic neonatal arterial ischemic stroke: the
561 International Pediatric Stroke Study. *Pediatrics.* 2011;128(6):e1402-10.

562 14. Vasudevan C, Levene M. Epidemiology and aetiology of neonatal seizures. *Semin Fetal*
563 *Neonatal Med.* 2013;18(4):185-91.

564 15. Levene MI, Trounce JQ. Cause of neonatal convulsions. Towards more precise diagnosis.
565 *Arch Dis Child.* 1986;61(1):78-9.

566 16. Beaulieu MJ. Levetiracetam. *Neonatal Netw.* 2013;32(4):285-8.

567 17. Dzhalal VI, Talos DM, Sdrulla DA, Brumback AC, Mathews GC, Benke TA, Delpire E,
568 Jensen FE, Staley KJ. NKCC1 transporter facilitates seizures in the developing brain. *Nat*
569 *Med.* 2005;11(11):1205-13.

570 18. D'Souza SW, Slater P. Excitatory amino acids in neonatal brain: contributions to
571 pathology and therapeutic strategies. *Arch Dis Child Fetal Neonatal Ed.* 1995;72(3):F147-50.

572 19. Nardou R, Ferrari DC, Ben-Ari Y. Mechanisms and effects of seizures in the immature
573 brain. *Semin Fetal Neonatal Med.* 2013;18(4):175-84.

574 20. Kang S, Kadam S. Pre-Clinical Models of Acquired Neonatal Seizures: Differential
575 Effects of Injury on Function of Chloride Co-Transporters. *Austin J Cerebrovasc Dis Stroke.*
576 2014;1(6):1026.

577 21. Silverstein FS, Jensen FE, Inder T, Hellstrom-Westas L, Hirtz D, Ferriero DM. Improving
578 the treatment of neonatal seizures: National Institute of Neurological Disorders and Stroke
579 workshop report. *J Pediatr*. 2008;153(1):12-5.

580 22. McGoldrick MK, Galanopoulou AS. Developmental pharmacology of benzodiazepines
581 under normal and pathological conditions. *Epileptic Disord*. 2014;16 Suppl 1:59-68.

582 23. Fatemi A, Wilson MA, Johnston MV. Hypoxic Ischemic Encephalopathy in the Term
583 Infant. *Clinics in perinatology*. 2009;36(4):835-58.

584 24. Murray DM, Boylan GB, Ali I, Ryan CA, Murphy BP, Connolly S. Defining the gap
585 between electrographic seizure burden, clinical expression and staff recognition of neonatal
586 seizures. *Arch Dis Child Fetal Neonatal Ed*. 2008;93(3):F187-91.

587 25. Hallberg B, Blennow M. Investigations for neonatal seizures. *Semin Fetal Neonatal Med*.
588 2013;18(4):196-201.

589 26. Abend NS, Wusthoff CJ, Goldberg EM, Dlugos DJ. Electrographic seizures and status
590 epilepticus in critically ill children and neonates with encephalopathy. *Lancet Neurol*.
591 2013;12(12):1170-9.

592 27. Bye AM, Flanagan D. Spatial and temporal characteristics of neonatal seizures. *Epilepsia*.
593 1995;36(10):1009-16.

594 28. Srinivasakumar P, Zempel J, Trivedi S, Wallendorf M, Rao R, Smith B, Inder T, Mathur
595 AM. Treating EEG Seizures in Hypoxic Ischemic Encephalopathy: A Randomized
596 Controlled Trial. *Pediatrics*. 2015;136(5):e1302-9.

597 29. Glykys J, Dzhala VI, Kuchibhotla KV, Feng G, Kuner T, Augustine G, Bacskai BJ,
598 Staley KJ. Differences in cortical versus subcortical GABAergic signaling: a candidate
599 mechanism of electroclinical uncoupling of neonatal seizures. *Neuron*. 2009;63(5):657-72.

600 30. Scher MS, Alvin J, Gaus L, Minnigh B, Painter MJ. Uncoupling of EEG-clinical neonatal
601 seizures after antiepileptic drug use. *Pediatr Neurol*. 2003;28(4):277-80.

602 31. Haynes RL, Borenstein NS, Desilva TM, Folkerth RD, Liu LG, Volpe JJ, Kinney HC.
603 Axonal development in the cerebral white matter of the human fetus and infant. *J Comp*
604 *Neurol.* 2005;484(2):156-67.

605 32. van den Heuvel MP, Kersbergen KJ, de Reus MA, Keunen K, Kahn RS, Groenendaal F,
606 de Vries LS, Benders MJNL. The Neonatal Connectome During Preterm Brain Development.
607 *Cerebral Cortex* (New York, NY). 2015;25(9):3000-13.

608 33. Abend NS, Gutierrez-Colina AM, Monk HM, Dlugos DJ, Clancy RR. Levetiracetam for
609 treatment of neonatal seizures. *J Child Neurol.* 2011;26(4):465-70.

610 34. Shah DK, Boylan GB, Rennie JM. Monitoring of seizures in the newborn. *Arch Dis Child*
611 *Fetal Neonatal Ed.* 2012;97(1):F65-9.

612 35. Shellhaas RA, Soaita AI, Clancy RR. Sensitivity of amplitude-integrated
613 electroencephalography for neonatal seizure detection. *Pediatrics.* 2007;120(4):770-7.

614 36. Shah DK, Mackay MT, Lavery S, Watson S, Harvey AS, Zempel J, Mathur A, Inder TE.
615 Accuracy of bedside electroencephalographic monitoring in comparison with simultaneous
616 continuous conventional electroencephalography for seizure detection in term infants.
617 *Pediatrics.* 2008;121(6):1146-54.

618 37. Rennie JM, Chorley G, Boylan GB, Pressler R, Nguyen Y, Hooper R. Non-expert use of
619 the cerebral function monitor for neonatal seizure detection. *Arch Dis Child Fetal Neonatal*
620 *Ed.* 2004;89(1):F37-40.

621 38. US National Institutes of Health. ClinicalTrials.gov. 2015. <http://clinicaltrials.gov>
622 Accessed 13th July 2015.

623 39. Mathieson SR, Stevenson NJ, Low E, Marnane WP, Rennie JM, Temko A, Lightbody G,
624 Boylan GB. Validation of an automated seizure detection algorithm for term neonates. *Clin*
625 *Neurophysiol.* 2016;127(1):156-158.

626 40. Low E, Mathieson SR, Stevenson NJ, Livingstone V, Ryan CA, Bogue CO, Rennie JM,
627 Boylan GB. Early postnatal EEG features of perinatal arterial ischaemic stroke with seizures.
628 PLoS One. 2014;9(7):e100973.

629 41. Pressler R, Binnie, C.D., Cooper, R., Robinson, R. (editors). Neonatal and Paediatric
630 Clinical Neurophysiology. London: Churchill Livingstone; 2007.

631 42. Boylan GB, Rennie JM, Chorley G, Pressler RM, Fox GF, Farrer K, Morton M, Binnie
632 CD. Second-line anticonvulsant treatment of neonatal seizures: a video-EEG monitoring
633 study. Neurology. 2004;62(3):486-8.

634 43. Weeke LC, Groenendaal F, Toet MC, Benders MJ, Nievelstein RA, van Rooij LG, de
635 Vries LS. The aetiology of neonatal seizures and the diagnostic contribution of neonatal
636 cerebral magnetic resonance imaging. Dev Med Child Neurol. 2015;57(3):248-56.

637 44. Glass HC, Sullivan JE. Neonatal seizures. Curr Treat Options Neurol. 2009;11(6):405-13.

638 45. Evans DJ, Levene MI, Tsakmakis M. Anticonvulsants for preventing mortality and
639 morbidity in full term newborns with perinatal asphyxia. Cochrane Database Syst Rev.
640 2007(3):CD001240.

641 46. van Rooij LG, Toet MC, van Huffelen AC, Groenendaal F, Laan W, Zecic A, de Haan T,
642 van Straaten IL, Vrancken S, van Wezel G, van der Sluijs J, Ter Horst H, Gavilanes D,
643 Laroche S, Naulaers G, de Vries LS. Effect of treatment of subclinical neonatal seizures
644 detected with aEEG: randomized, controlled trial. Pediatrics. 2010;125(2):e358-66.

645 47. Glass HC, Nash KB, Bonifacio SL, Barkovich AJ, Ferriero DM, Sullivan JE, Cilio MR.
646 Seizures and magnetic resonance imaging-detected brain injury in newborns cooled for
647 hypoxic-ischemic encephalopathy. J Pediatr. 2011;159(5):731-5.

648 48. Lynch NE, Stevenson NJ, Livingstone V, Murphy BP, Rennie JM, Boylan GB. The
649 temporal evolution of electrographic seizure burden in neonatal hypoxic ischemic
650 encephalopathy. Epilepsia. 2012;53(3):549-57.

651 49. Booth D, Evans DJ. Anticonvulsants for neonates with seizures. Cochrane Database Syst
652 Rev. 2004(4):CD004218.

653 50. Borowicz KK, Banach M. Antiarrhythmic drugs and epilepsy. Pharmacological Reports.
654 2014;66(4):545-51.

655 51. Bialer M, White HS. Key factors in the discovery and development of new antiepileptic
656 drugs. Nat Rev Drug Discov. 2010;9(1):68-82.

657 52. Irish Medicines Board. Summaries of Product Characteristics-Levetiracetam [Online].
658 2013. www.medicines.ie Accessed 5th January 2016.

659 53. Nardou R, Yamamoto S, Bhar A, Burnashev N, Ben-Ari Y, Khalilov I. Phenobarbital but
660 Not Diazepam Reduces AMPA/kainate Receptor Mediated Currents and Exerts Opposite
661 Actions on Initial Seizures in the Neonatal Rat Hippocampus. Front Cell Neurosci.
662 2011;5:16.

663 54. van Rooij LG, Hellstrom-Westas L, de Vries LS. Treatment of neonatal seizures. Semin
664 Fetal Neonatal Med. 2013;18(4):209-15.

665 55. Slaughter LA, Patel AD, Slaughter JL. Pharmacological treatment of neonatal seizures: a
666 systematic review. J Child Neurol. 2013;28(3):351-64.

667 56. Khan O, Chang E, Cipriani C, Wright C, Crisp E, Kirmani B. Use of intravenous
668 levetiracetam for management of acute seizures in neonates. Pediatr Neurol. 2011;44(4):265-
669 9.

670 57. Glass HC, Poulin C, Shevell MI. Topiramate for the treatment of neonatal seizures.
671 Pediatr Neurol. 2011;44(6):439-42.

672 58. Silverstein FS, Ferriero DM. Off-label use of antiepileptic drugs for the treatment of
673 neonatal seizures. Pediatr Neurol. 2008;39(2):77-9.

674 59. Kim J, Kondratyev A, Gale K. Antiepileptic drug-induced neuronal cell death in the
675 immature brain: effects of carbamazepine, topiramate, and levetiracetam as monotherapy
676 versus polytherapy. *J Pharmacol Exp Ther.* 2007;323(1):165-73.

677 60. Tulloch JK, Carr RR, Ensom MH. A systematic review of the pharmacokinetics of
678 antiepileptic drugs in neonates with refractory seizures. *J Pediatr Pharmacol Ther.*
679 2012;17(1):31-44.

680 61. Mikati MA, Fayad M, Koleilat M, Mounla N, Hussein R, Kazma A, Yunis K. Efficacy,
681 tolerability, and kinetics of lamotrigine in infants. *J Pediatr.* 2002;141(1):31-5.

682 62. Sampath D, Shmueli D, White AM, Raol YH. Flupirtine effectively prevents
683 development of acute neonatal seizures in an animal model of global hypoxia. *Neurosci Lett.*
684 2015;607:46-51.

685 63. Lynch NE, Stevenson NJ, Livingstone V, Mathieson S, Murphy BP, Rennie JM, Boylan
686 GB. The temporal characteristics of seizures in neonatal hypoxic ischemic encephalopathy
687 treated with hypothermia. *Seizure.* 2015;33:60-5.

688 64. Pressler RM, Boylan GB, Marlow N, Blennow M, Chiron C, Cross JH, de Vries LS,
689 Hallberg B, Hellstrom-Westas L, Jullien V, Livingstone V, Mangum B, Murphy B, Murray
690 D, Pons G, Rennie J, Swarte R, Toet MC, Vanhatalo S, Zohar S. Bumetanide for the
691 treatment of seizures in newborn babies with hypoxic ischaemic encephalopathy (NEMO): an
692 open-label, dose finding, and feasibility phase 1/2 trial. *Lancet Neurol.* 2015;14(5):469-77.

693 65. Maartens IA, Wassenberg T, Buijs J, Bok L, de Kleine MJ, Katgert T, Andriessen P.
694 Neurodevelopmental outcome in full-term newborns with refractory neonatal seizures. *Acta*
695 *Paediatr.* 2012;101(4):e173-8.

696 66. Painter MJ, Scher MS, Stein AD, Armatti S, Wang Z, Gardiner JC, Paneth N, Minnigh B,
697 Alvin J. Phenobarbital compared with phenytoin for the treatment of neonatal seizures. *N*
698 *Engl J Med.* 1999;341(7):485-9.

699 67. Castro Conde JR, Hernandez Borges AA, Domenech Martinez E, Gonzalez Campo C,
700 Perera Soler R. Midazolam in neonatal seizures with no response to phenobarbital.
701 Neurology. 2005;64(5):876-9.

702 68. Malingre MM, Van Rooij LG, Rademaker CM, Toet MC, Ververs TF, van Kesteren C, de
703 Vries LS. Development of an optimal lidocaine infusion strategy for neonatal seizures. Eur J
704 Pediatr. 2006;165(9):598-604.

705 69. Shany E, Benzaqen O, Watemberg N. Comparison of continuous drip of midazolam or
706 lidocaine in the treatment of intractable neonatal seizures. J Child Neurol. 2007;22(3):255-9.

707 70. Weeke LC, Toet MC, van Rooij LG, Groenendaal F, Boylan GB, Pressler RM,
708 Hellstrom-Westas L, van den Broek MP, de Vries LS. Lidocaine response rate in aEEG-
709 confirmed neonatal seizures: Retrospective study of 413 full-term and preterm infants.
710 Epilepsia. 2015 [Epub ahead of print].

711 71. van Rooij LG, Toet MC, Rademaker KM, Groenendaal F, de Vries LS. Cardiac
712 arrhythmias in neonates receiving lidocaine as anticonvulsive treatment. Eur J Pediatr.
713 2004;163(11):637-41.

714 72. van den Broek MP, Huitema AD, van Hasselt JG, Groenendaal F, Toet MC, Egberts TC,
715 de Vries LS, Rademaker CM. Lidocaine (lignocaine) dosing regimen based upon a
716 population pharmacokinetic model for preterm and term neonates with seizures. Clin
717 Pharmacokinet. 2011;50(7):461-9.

718 73. Lundqvist M, Agren J, Hellstrom-Westas L, Flink R, Wickstrom R. Efficacy and safety of
719 lidocaine for treatment of neonatal seizures. Acta Paediatr. 2013;102(9):863-7.

720 74. van Leuven K, Groenendaal F, Toet MC, Schobben AF, Bos SA, de Vries LS, Rademaker
721 CM. Midazolam and amplitude-integrated EEG in asphyxiated full-term neonates. Acta
722 Paediatr. 2004;93(9):1221-7.

723 75. van den Broek MP, van Straaten HL, Huitema AD, Egberts T, Toet MC, de Vries LS,
724 Rademaker K, Groenendaal F. Anticonvulsant effectiveness and hemodynamic safety of
725 midazolam in full-term infants treated with hypothermia. *Neonatology*. 2015;107(2):150-6.

726 76. Deshmukh A, Wittert W, Schnitzler E, Mangurten HH. Lorazepam in the treatment of
727 refractory neonatal seizures. A pilot study. *Am J Dis Child*. 1986;140(10):1042-4.

728 77. Maytal J, Novak GP, King KC. Lorazepam in the treatment of refractory neonatal
729 seizures. *J Child Neurol*. 1991;6(4):319-23.

730 78. Talos DM, Chang M, Kosaras B, Fitzgerald E, Murphy A, Folkerth RD, Jensen FE.
731 Antiepileptic effects of levetiracetam in a rodent neonatal seizure model. *Pediatr Res*.
732 2013;73(1):24-30.

733 79. Merhar SL, Schibler KR, Sherwin CM, Meinzen-Derr J, Shi J, Balmakund T, Vinks AA.
734 Pharmacokinetics of levetiracetam in neonates with seizures. *J Pediatr*. 2011;159(1):152-4.e3.

735 80. Briggs DE, French JA. Levetiracetam safety profiles and tolerability in epilepsy patients.
736 *Expert Opin Drug Saf*. 2004;3(5):415-24.

737 81. Filippi L, Fiorini P, Daniotti M, Catarzi S, Savelli S, Fonda C, Bartalena L, Boldrini A,
738 Giampietri M, Scaramuzzo R, Papoff P, Del Balzo F, Spalice A, la Marca G, Malvagia S,
739 Della Bona ML, Donzelli G, Tinelli F, Cioni G, Pisano T, Falchi M, Guerrini R. Safety and
740 efficacy of topiramate in neonates with hypoxic ischemic encephalopathy treated with
741 hypothermia (NeoNATI). *BMC Pediatr*. 2012;12:144.

742 82. Dzhalal VI, Brumback AC, Staley KJ. Bumetanide enhances phenobarbital efficacy in a
743 neonatal seizure model. *Ann Neurol*. 2008;63(2):222-35.

744 83. Schwartzkroin PA, Baraban SC, Hochman DW. Osmolarity, ionic flux, and changes in
745 brain excitability. *Epilepsy Res*. 1998;32(1-2):275-85.

746 84. Haas M, Forbush B, 3rd. The Na-K-Cl cotransporters. *J Bioenerg Biomembr*.
747 1998;30(2):161-72.

748 85. Ben-Ari Y. Excitatory actions of gaba during development: the nature of the nurture. Nat
749 Rev Neurosci. 2002;3(9):728-39.

750 86. Pressler RM, Boylan GB, Marlow N, de Vries LS, Blennow M, Chiron C, Cross JH,
751 Hallberg B, Hellstrom-Westas L, Jullien V, Mangum B, Murphy B, Murray D, Pons G,
752 Rennie J, Toet MC, Zohar S. Bumetanide for neonatal seizures-back from the cotside. Nat
753 Rev Neurol. 2015;11(24):724.

754 87. Donovan MD, O'Brien FE, Boylan GB, Cryan JF, Griffin BT. The effect of organic anion
755 transporter 3 inhibitor probenecid on bumetanide levels in the brain: an integrated in vivo
756 microdialysis study in the rat. J Pharm Pharmacol. 2015;67(4):501-10.

757 88. Topfer M, Tollner K, Brandt C, Twele F, Broer S, Loscher W. Consequences of
758 inhibition of bumetanide metabolism in rodents on brain penetration and effects of
759 bumetanide in chronic models of epilepsy. Eur J Neurosci. 2014;39(4):673-87.

760 89. Tollner K, Brandt C, Topfer M, Brunhofer G, Erker T, Gabriel M, Feit PW, Lindfors J,
761 Kaila K, Loscher W. A novel prodrug-based strategy to increase effects of bumetanide in
762 epilepsy. Ann Neurol. 2014;75(4):550-62.

763 90. Tollner K, Brandt C, Romermann K, Loscher W. The organic anion transport inhibitor
764 probenecid increases brain concentrations of the NKCC1 inhibitor bumetanide. Eur J
765 Pharmacol. 2014;746c:167-73.

766 91. Brodie MJ, Kwan P. Current position of phenobarbital in epilepsy and its future.
767 Epilepsia. 2012;53 Suppl 8:40-6.

768 92. Bittigau P, Sifringer M, Ikonomidou C. Antiepileptic drugs and apoptosis in the
769 developing brain. Ann N Y Acad Sci. 2003;993:103-14.

770 93. Loiacono G, Masci M, Zaccara G, Verrotti A. The treatment of neonatal seizures: focus
771 on Levetiracetam. J Matern Fetal Neonatal Med. 2016;29(1):69-74.

772 94. Khan O, Cipriani C, Wright C, Crisp E, Kirmani B. Role of intravenous levetiracetam for
773 acute seizure management in preterm neonates. *Pediatr Neurol.* 2013;49(5):340-3.

774 95. Sharpe CM, Capparelli EV, Mower A, Farrell MJ, Soldin SJ, Haas RH. A seven-day
775 study of the pharmacokinetics of intravenous levetiracetam in neonates: marked changes in
776 pharmacokinetics occur during the first week of life. *Pediatr Res.* 2012;72(1):43-9.

777 96. Filippi L, la Marca G, Fiorini P, Poggi C, Cavallaro G, Malvagia S, Pellegrini-Giampietro
778 DE, Guerrini R. Topiramate concentrations in neonates treated with prolonged whole body
779 hypothermia for hypoxic ischemic encephalopathy. *Epilepsia.* 2009;50(11):2355-61.

780 97. Filippi L, Poggi C, la Marca G, Furlanetto S, Fiorini P, Cavallaro G, Plantulli A, Donzelli
781 G, Guerrini R. Oral topiramate in neonates with hypoxic ischemic encephalopathy treated
782 with hypothermia: a safety study. *J Pediatr.* 2010;157(3):361-6.

783 98. Alcorn J, McNamara PJ. Pharmacokinetics in the newborn. *Adv Drug Deliv Rev.*
784 2003;55(5):667-86.

785 99. Marsot A, Brevaut-Malaty V, Vialet R, Boulamery A, Bruguerolle B, Simon N.
786 Pharmacokinetics and absolute bioavailability of phenobarbital in neonates and young
787 infants, a population pharmacokinetic modelling approach. *Fundam Clin Pharmacol.*
788 2014;28(4):465-71.

789 100. Filippi L, la Marca G, Cavallaro G, Fiorini P, Favelli F, Malvagia S, Donzelli G,
790 Guerrini R. Phenobarbital for neonatal seizures in hypoxic ischemic encephalopathy: a
791 pharmacokinetic study during whole body hypothermia. *Epilepsia.* 2011;52(4):794-801.

792 101. Shellhaas RA, Ng CM, Dillon CH, Barks JD, Bhatt-Mehta V. Population
793 pharmacokinetics of phenobarbital in infants with neonatal encephalopathy treated with
794 therapeutic hypothermia. *Pediatr Crit Care Med.* 2013;14(2):194-202.

795 102. Patsalos PN, Berry DJ, Bourgeois BF, Cloyd JC, Glauser TA, Johannessen SI, Leppik
796 IE, Tomson T, Perucca E. Antiepileptic drugs--best practice guidelines for therapeutic drug

797 monitoring: a position paper by the subcommission on therapeutic drug monitoring, ILAE
 798 Commission on Therapeutic Strategies. *Epilepsia*. 2008;49(7):1239-76.

799 103. Loughnan PM, Greenwald A, Purton WW, Aranda JV, Watters G, Neims AH.
 800 Pharmacokinetic observations of phenytoin disposition in the newborn and young infant.
 801 *Archives of Disease in Childhood*. 1977;52(4):302-9.

802 104. Al Za'abi M, Lanner A, Xiaonian X, Donovan T, Charles B. Application of routine
 803 monitoring data for determination of the population pharmacokinetics and enteral
 804 bioavailability of phenytoin in neonates and infants with seizures. *Ther Drug Monit*.
 805 2006;28(6):793-9.

806 105. Bjorkman S. Prediction of drug disposition in infants and children by means of
 807 physiologically based pharmacokinetic (PBPK) modelling: theophylline and midazolam as
 808 model drugs. *Br J Clin Pharmacol*. 2005;59(6):691-704.

809 106. Jullien V, Pressler RM, Boylan G, Blennow M, Marlow N, Chiron C, Pons G. Pilot
 810 evaluation of the population pharmacokinetics of bumetanide in term newborn infants with
 811 seizures. *J Clin Pharmacol*. 2015 [Epub ahead of print].

812 107. Azzopardi DV, Strohm B, Edwards AD, Dyet L, Halliday HL, Juszczak E, Kapellou O,
 813 Levene M, Marlow N, Porter E, Thoresen M, Whitelaw A, Brocklehurst P. Moderate
 814 hypothermia to treat perinatal asphyxial encephalopathy. *N Engl J Med*. 2009;361(14):1349-
 815 58.

816 108. Tagin MA, Woolcott CG, Vincer MJ, Whyte RK, Stinson DA. Hypothermia for
 817 neonatal hypoxic ischemic encephalopathy: an updated systematic review and meta-analysis.
 818 *Arch Pediatr Adolesc Med*. 2012;166(6):558-66.

819 109. Azzopardi D, Strohm B, Marlow N, Brocklehurst P, Deierl A, Eddama O, Goodwin J,
 820 Halliday HL, Juszczak E, Kapellou O, Levene M, Linsell L, Omar O, Thoresen M, Tusor N,

821 Whitelaw A, Edwards AD. Effects of hypothermia for perinatal asphyxia on childhood
822 outcomes. *N Engl J Med.* 2014;371(2):140-9.

823 110. Srinivasakumar P, Zempel J, Wallendorf M, Lawrence R, Inder T, Mathur A.
824 Therapeutic hypothermia in neonatal hypoxic ischemic encephalopathy: electrographic
825 seizures and magnetic resonance imaging evidence of injury. *J Pediatr.* 2013;163(2):465-70.

826 111. Low E, Boylan GB, Mathieson SR, Murray DM, Korotchikova I, Stevenson NJ,
827 Livingstone V, Rennie JM. Cooling and seizure burden in term neonates: an observational
828 study. *Arch Dis Child Fetal Neonatal Ed.* 2012;97(4):F267-72.

829 112. Gancia P, Pomero G. Therapeutic hypothermia in the prevention of hypoxic-ischaemic
830 encephalopathy: new categories to be enrolled. *J Matern Fetal Neonatal Med.* 2012;25 Suppl
831 4:94-6.

832 113. Yang T, Zhuang L, Rei Fidalgo AM, Petrides E, Terrando N, Wu X, Sanders RD,
833 Robertson NJ, Johnson MR, Maze M, Ma D. Xenon and sevoflurane provide analgesia
834 during labor and fetal brain protection in a perinatal rat model of hypoxia-ischemia. *PLoS*
835 *One.* 2012;7(5):e37020.

836 114. Thoresen M, Hobbs CE, Wood T, Chakkarapani E, Dingley J. Cooling combined with
837 immediate or delayed xenon inhalation provides equivalent long-term neuroprotection after
838 neonatal hypoxia-ischemia. *J Cereb Blood Flow Metab.* 2009;29(4):707-14.

839 115. Palmer C, Towfighi J, Roberts RL, Heitjan DF. Allopurinol administered after inducing
840 hypoxia-ischemia reduces brain injury in 7-day-old rats. *Pediatr Res.* 1993;33(4 Pt 1):405-11.

841 116. Kaandorp JJ, Benders MJ, Rademaker CM, Torrance HL, Oudijk MA, de Haan TR,
842 Bloemenkamp KW, Rijken M, van Pampus MG, Bos AF, Porath MM, Oetomo SB, Willekes
843 C, Gavilanes AW, Wouters MG, van Elburg RM, Huisjes AJ, Bakker SC, van Meir CA, von
844 Lindern J, Boon J, de Boer IP, Rijnders RJ, Jacobs CJ, Uiterwaal CS, Mol BW, Visser GH,
845 van Bel F, Derks JB. Antenatal allopurinol for reduction of birth asphyxia induced brain

846 damage (ALLO-Trial); a randomized double blind placebo controlled multicenter study.
847 BMC Pregnancy Childbirth. 2010;10:8.

848 117. Robertson NJ, Faulkner S, Fleiss B, Bainbridge A, Andorka C, Price D, Powell E,
849 Lecky-Thompson L, Thei L, Chandrasekaran M, Hristova M, Cady EB, Gressens P, Golay X,
850 Raivich G. Melatonin augments hypothermic neuroprotection in a perinatal asphyxia model.
851 Brain. 2013;136(Pt 1):90-105.

852 118. Mazur M, Miller RH, Robinson S. Postnatal erythropoietin treatment mitigates neural
853 cell loss after systemic prenatal hypoxic-ischemic injury. J Neurosurg Pediatr. 2010;6(3):206-
854 21.

855 119. Rangarajan V, Juul SE. Erythropoietin: emerging role of erythropoietin in neonatal
856 neuroprotection. Pediatr Neurol. 2014;51(4):481-8.

857 120. Dingley J, Tooley J, Liu X, Scull-Brown E, Elstad M, Chakkarapani E, Sabir H,
858 Thoresen M. Xenon ventilation during therapeutic hypothermia in neonatal encephalopathy: a
859 feasibility study. Pediatrics. 2014;133(5):809-18.

860 121. Traudt CM, McPherson RJ, Bauer LA, Richards TL, Burbacher TM, McAdams RM,
861 Juul SE. Concurrent erythropoietin and hypothermia treatment improve outcomes in a term
862 nonhuman primate model of perinatal asphyxia. Dev Neurosci. 2013;35(6):491-503.

863 122. Traudt CM, Juul SE. Erythropoietin as a neuroprotectant for neonatal brain injury:
864 animal models. Methods Mol Biol. 2013;982:113-26.

865 123. Zhu C, Kang W, Xu F, Cheng X, Zhang Z, Jia L, Ji L, Guo X, Xiong H, Simbruner G,
866 Blomgren K, Wang X. Erythropoietin improved neurologic outcomes in newborns with
867 hypoxic-ischemic encephalopathy. Pediatrics. 2009;124(2):e218-26.

868 124. Gill H, Thoresen M, Smit E, Davis J, Liu X, Dingley J, Elstad M. Sedation management
869 during therapeutic hypothermia for neonatal encephalopathy: atropine premedication for

870 endotracheal intubation causes a prolonged increase in heart rate. Resuscitation.
871 2014;85(10):1394-8.

872 125. Thoresen M, Satas S, Loberg EM, Whitelaw A, Acolet D, Lindgren C, Penrice J,
873 Robertson N, Haug E, Steen PA. Twenty-four hours of mild hypothermia in unsedated
874 newborn pigs starting after a severe global hypoxic-ischemic insult is not neuroprotective.
875 *Pediatr Res*. 2001;50(3):405-11.

876 126. Young GB, da Silva OP. Effects of morphine on the electroencephalograms of neonates:
877 a prospective, observational study. *Clin Neurophysiol*. 2000;111(11):1955-60.

878 127. Bansinath M, Turndorf H, Puig MM. Influence of hypo and hyperthermia on disposition
879 of morphine. *J Clin Pharmacol*. 1988;28(9):860-4.

880 128. Roka A, Melinda KT, Vasarhelyi B, Machay T, Azzopardi D, Szabo M. Elevated
881 morphine concentrations in neonates treated with morphine and prolonged hypothermia for
882 hypoxic ischemic encephalopathy. *Pediatrics*. 2008;121(4):e844-9.

883 129. Puig MM, Warner W, Tang CK, Laorden ML, Turndorf H. Effects of temperature on the
884 interaction of morphine with opioid receptors. *Br J Anaesth*. 1987;59(11):1459-64.

885 130. Angeles DM, Ashwal S, Wycliffe ND, Ebner C, Fayard E, Sowers L, Holshouser BA.
886 Relationship between opioid therapy, tissue-damaging procedures, and brain metabolites as
887 measured by proton MRS in asphyxiated term neonates. *Pediatr Res*. 2007;61(5 Pt 1):614-21.

888 131. Stockley's Drug Interactions. In: Medicines Complete Database. 2009.
889 www.medicinescomplete.com Accessed: 23rd July 2015.

890 132. Cilio MR, Ferriero DM. Synergistic neuroprotective therapies with hypothermia. *Semin*
891 *Fetal Neonatal Med*. 2010;15(5):293-8.

892 133. Tortorici MA, Kochanek PM, Poloyac SM. Effects of hypothermia on drug disposition,
893 metabolism, and response: A focus of hypothermia-mediated alterations on the cytochrome
894 P450 enzyme system. *Crit Care Med*. 2007;35(9):2196-204.

895 134. van den Broek MP, Rademaker CM, van Straaten HL, Huitema AD, Toet MC, de Vries
896 LS, Egberts AC, Groenendaal F. Anticonvulsant treatment of asphyxiated newborns under
897 hypothermia with lidocaine: efficacy, safety and dosing. *Arch Dis Child Fetal Neonatal Ed.*
898 2013;98(4):F341-5.

899 135. Zhou J, Poloyac SM. The effect of therapeutic hypothermia on drug metabolism and
900 response: cellular mechanisms to organ function. *Expert Opin Drug Metab Toxicol.*
901 2011;7(7):803-16.

902 136. Shah P, Riphagen S, Beyene J, Perlman M. Multiorgan dysfunction in infants with post-
903 asphyxial hypoxic-ischaemic encephalopathy. *Arch Dis Child Fetal Neonatal Ed.*
904 2004;89(2):F152-5.

905 137. Kendall GS, Mathieson S, Meek J, Rennie JM. Recooling for rebound seizures after
906 rewarming in neonatal encephalopathy. *Pediatrics.* 2012;130(2):e451-5.

907 138. Barks JD, Liu YQ, Shangguan Y, Silverstein FS. Phenobarbital augments hypothermic
908 neuroprotection. *Pediatr Res.* 2010;67(5):532-7.

909 139. Meyn DF, Jr., Ness J, Ambalavanan N, Carlo WA. Prophylactic phenobarbital and
910 whole-body cooling for neonatal hypoxic-ischemic encephalopathy. *J Pediatr.*
911 2010;157(2):334-6.

912 140. van den Broek MP, Groenendaal F, Toet MC, van Straaten HL, van Hasselt JG, Huitema
913 AD, de Vries LS, Egberts AC, Rademaker CM. Pharmacokinetics and clinical efficacy of
914 phenobarbital in asphyxiated newborns treated with hypothermia: a thermopharmacological
915 approach. *Clin Pharmacokinet.* 2012;51(10):671-9.

916 141. Kadar D, Tang BK, Conn AW. The fate of phenobarbitone in children in hypothermia
917 and at normal body temperature. *Can Anaesth Soc J.* 1982;29(1):16-23.

918 142. Liu Y, Barks JD, Xu G, Silverstein FS. Topiramate extends the therapeutic window for
919 hypothermia-mediated neuroprotection after stroke in neonatal rats. *Stroke*. 2004;35(6):1460-
920 5.

921 143. Welzing L, Junghaenel S, Weiss V, Roth B, Mueller C, Wiesen MH. Disposition of
922 midazolam in asphyxiated neonates receiving therapeutic hypothermia--a pilot study. *Klin*
923 *Padiatr*. 2013;225(7):398-404.

924 144. Fukuoka N, Aibiki M, Tsukamoto T, Seki K, Morita S. Biphasic concentration change
925 during continuous midazolam administration in brain-injured patients undergoing therapeutic
926 moderate hypothermia. *Resuscitation*. 2004;60(2):225-30.

927 145. Lopez-Samblas AM, Adams JA, Goldberg RN, Modi MW. The pharmacokinetics of
928 bumetanide in the newborn infant. *Biol Neonate*. 1997;72(5):265-72.

929 146. Pacifici GM. Clinical pharmacology of the loop diuretics furosemide and bumetanide in
930 neonates and infants. *Paediatr Drugs*. 2012;14(4):233-46.

931 147. Empey PE, de Mendizabal NV, Bell MJ, Bies RR, Anderson KB, Kochanek PM,
932 Adelson PD, Poloyac SM. Therapeutic hypothermia decreases phenytoin elimination in
933 children with traumatic brain injury. *Crit Care Med*. 2013;41(10):2379-87.

934 148. Iida Y, Nishi S, Asada A. Effect of mild therapeutic hypothermia on phenytoin
935 pharmacokinetics. *Ther Drug Monit*. 2001;23(3):192-7.

936 149. Bhagat H, Bithal PK, Chouhan RS, Arora R. Is phenytoin administration safe in a
937 hypothermic child? *J Clin Neurosci*. 2006;13(9):953-5.

938 150. Jensen FE. Neonatal seizures: an update on mechanisms and management. *Clin*
939 *Perinatol*. 2009;36(4):881-900.

940 151. Kanhere S. Recent advances in neonatal seizures. *Indian J Pediatr*. 2014;81(9):917-25.

941 152. Donnelly F. EU initiatives for research involving children. *Eur J Pediatr*.
942 2008;167(7):837-8.

Fig. 1 Mode of action of neonatal AEDs

Many drugs act by reducing excitatory neurotransmission (glutamatergic synapse). Phenytoin, lidocaine and topiramate prevent depolarisation by inhibiting voltage-gated sodium channels [50, 51]. Levetiracetam prevents calcium influx through N-type calcium channels which in turn reduces exocytosis and reduces the release of glutamate from intracellular vesicles by modulating synaptic vesicle protein 2A (SV2A) [51, 52]. On the postsynaptic terminal, phenobarbital and topiramate reduce excitatory neurotransmission via the AMPA/kainate glutamate receptor [51, 53]. Anticonvulsants including phenobarbital, benzodiazepines and topiramate work by enhancing inhibitory neurotransmission via the GABA_A receptor (GABAergic synapse) [51]. Bumetanide can alter the action of GABAergic agents by preventing intracellular accumulation of chloride through NKCC1 [17].

(AMPA = α -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid; NMDA = N-methyl-D-aspartate; GABA = γ -amino butyric acid; GAD = glutamic acid decarboxylase; SV2A = synaptic vesicle protein 2A)