

Title	Associations between the gut microbiome, inflammation, and cardiovascular profiles in people with human immunodeficiency virus
Authors	MacCann, Rachel;Li, Junhui;Leon, Alejandro Abner Garcia;Negi, Riya;Alalwan, Dana;Tinago, Willard;Mcgettrick, Padraig;Cotter, Aoife G;Landay, Alan;Sabin, Caroline;O'toole, Paul W;Mallon, Patrick W. G;Barco, Elena Alvarez;Leon, Alejandro Garcia;Mcdermott, Aoife;Mcginty, Tara;Macken, Alan;Kavanagh, Eoin;Mccarthy, Geraldine;Sheehan, Gerard;Lambert, John;Powderly, William;Compston, Juliet
Publication date	2025
Original Citation	MacCann, R., Li, J., Leon, A. A. G., Negi, R., Alalwan, D., Tinago, W., McGettrick, P., Cotter, A. G., Landay, A., Sabin, C. and O'Toole, P. W. (2025) 'Associations between the gut microbiome, inflammation, and cardiovascular profiles in people with human immunodeficiency virus', The Journal of Infectious Diseases, 231(4), pp.e781-e791. DOI: 10.1093/infdis/jiaf043
Type of publication	Article (peer reviewed)
Link to publisher's version	10.1093/infdis/jiaf043
Rights	© 2025, the Author(s). Published by Oxford University Press on behalf of Infectious Diseases Society of America. This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs licence (https://creativecommons.org/licenses/by-nc-nd/4.0/), which permits non-commercial reproduction and distribution of the work, in any medium, provided the original work is not altered or transformed in any way, and that the work is properly cited. - https://creativecommons.org/licenses/by-nc-nd/4.0/
Download date	2026-09-17 00:51:31
Item downloaded from	https://hdl.handle.net/10468/17332



University College Cork, Ireland
Coláiste na hOllscoile Corcaigh

Associations Between the Gut Microbiome, Inflammation, and Cardiovascular Profiles in People With Human Immunodeficiency Virus

Rachel MacCann,^{1,2,3,*} Junhui Li,⁴ Alejandro Abner Garcia Leon,³ Riya Negi,³ Dana Alalwan,^{3,*} Willard Tinago,^{3,*} Padraig McGettrick,^{1,3,5} Aoife G. Cotter,^{3,5} Alan Landay,⁶ Caroline Sabin,⁷ Paul W. O'Toole,⁴ and Patrick W. G. Mallon^{1,2,3}; for the Understanding the Pathology of Comorbid Disease in HIV-Infected Individuals (HIV UPBEAT) Study Group^a

¹School of Medicine, University College Dublin; ²Department of Infectious Diseases, St Vincent's University Hospital; ³Centre for Experimental Pathogen Host Research, University College Dublin; ⁴School of Microbiology and APC Microbiome Ireland, University College Cork; ⁵Department of Infectious Diseases, Mater Misericordiae University Hospital, Dublin, Ireland; ⁶Department of Internal Medicine, University of Texas Medical Branch, Galveston; and ⁷Institute for Global Health, University College London, United Kingdom

Background. Inflammation and innate immune activation are associated with chronic human immunodeficiency virus (HIV) infection, despite effective treatment. Although gut microbiota alterations are linked to systemic inflammation, their relationship with HIV infection the relationships between the gut microbiome, inflammation, and HIV remains unclear.

Methods. The HIV UPBEAT Coronary Artery Disease sub-study evaluated cardiovascular disease (CVD) in people with and without HIV. Subclinical CVD was assessed using coronary computed tomography angiography (CCTA). Thirty-four biomarkers were measured using quantitative immunoassays. Stool samples underwent 16S rRNA sequencing. Differentially abundant species were identified by analysis of compositions of microbiomes with bias correction (ANCOM-BC) and correlated to biomarkers, diet, and CCTA outcomes using Spearman correlation.

Results. Among 81 participants (median age, 51 years; 73% male), people with HIV ($n = 44$) had higher rates of hypercholesterolemia ($P < .025$). Gut microbiome β -diversity differed significantly by HIV status. Enriched *Bifidobacterium pseudocatenulatum*, *Megamonas hypermegale*, and *Selenomonas ruminantium* correlated with lower plaque burden, while depleted *Ruminococcus bromii* correlated with higher plaque burden and fat intake. Depleted *Bacteroides* spp and *Alistipes* spp correlated with elevated biomarkers (D-dimer, CD40 ligand, C-reactive protein, and interferon- γ).

Conclusions. Gut microbiota differences in people with HIV were linked to subclinical CVD, diet, and inflammation, highlighting the microbiome's role in cardiovascular risk in HIV infection.

Keywords. HIV; cardiovascular disease; inflammation; microbiome; ageing.

Since the introduction of antiretroviral therapy (ART), many people with human immunodeficiency virus (HIV) now achieve a life expectancy approaching that of the general population [1]. However, some experience an accentuated aging phenotype, with increased risk of developing age-related noncommunicable diseases, such as cardiovascular disease (CVD), at a younger age than those without HIV [2]. CVD is now a leading cause of death

in those receiving ART [1], with traditional CVD risk factors, although prevalent in this group, not fully accounting for this excess risk [3]. Initially, prolonged ART exposure was suspected, but recent data suggest its impact on CVD is relatively small [4].

Unresolved chronic inflammation is a major contributor to CVD pathogenesis, involving endothelial activation, vascular inflammation, and type 1 T-helper (Th1) cell activation [5], all implicated in coronary artery plaque (CAP) development [5]. Innate immune activation and systemic inflammation are also associated with subclinical coronary artery disease (CAD) in those with HIV, with markers like high-sensitivity C-reactive protein (hsCRP) and interleukin 6 predicting worse outcomes [6].

HIV infection has been associated with gut dysbiosis due to alteration of the gut epithelial barrier and depletion of CD4⁺ T cells in the gut-associated lymphoid tissue [7]. This leads to changes in gut microbial diversity and community compositions. Moreover, commensal bacteria crucial for a healthy gut are replaced by taxa that may contribute to chronic inflammation and immune dysfunction [8]. Important commensal bacteria include short chain fatty acid (SCFA)-producing organisms, which produce butyrate, propionate, and acetate as metabolic

Received 07 November 2024; editorial decision 15 January 2025; published online 24 January 2025

^aMembers of the HIV UPBEAT Study Group are listed in the Acknowledgments.

Correspondence: Rachel MacCann, PhD, Centre for Experimental Pathogen Host Research, University College Dublin, Belfield, Dublin 4, Ireland (rachelmaccann@svhg.ie).

The Journal of Infectious Diseases® 2025;231:e781–91

© The Author(s) 2025. Published by Oxford University Press on behalf of Infectious Diseases Society of America.

This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs licence (<https://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial reproduction and distribution of the work, in any medium, provided the original work is not altered or transformed in any way, and that the work is properly cited. For commercial re-use, please contact reprints@oup.com for reprints and translation rights for reprints. All other permissions can be obtained through our RightsLink service via the Permissions link on the article page on our site—for further information please contact journals.permissions@oup.com.

<https://doi.org/10.1093/infdis/jiaf043>

products through the fermentation of dietary fibers and other indigestible carbohydrates [9]. *Firmicutes* and *Bacteroides* are the main butyrate-producing phyla [9]. Depletion of butyrate-producing bacteria has been associated with CVD, including atherosclerosis and dyslipidemia [10, 11].

In the context of HIV, decreased abundance of SCFA-producing bacteria has been associated with lower CD4⁺ T-cell counts and immune activation [12]. A reduction of butyrate-producing bacteria in particular has been observed in people with HIV with metabolic disorders such as hyperglycemia, hypertension, and dyslipidemia [13]. HIV infection also increases microbial translocation (MT), with movement of bacteria (or bacterial products) from the gastrointestinal tract into the systemic circulation, driving inflammation [14, 15]. Markers used for MT include lipopolysaccharide, soluble C14 (sCD14), and intestinal fatty acid-binding protein (I-FABP), a marker of enterocyte damage. Reductions in butyrate-producing bacteria have been associated with MT as well as immune activation and vascular inflammation in people with HIV [15, 16].

A previous study of this cohort identified distinct biomarker-derived inflammatory patterns associated with both subclinical CAD and prevalent CVD events [17]. One particular inflammatory pattern displayed higher markers of gut epithelial dysfunction (I-FABP), T-cell stimulation, and systemic inflammation and was distinguished from the other clusters by being significantly associated with higher CAP and prevalent clinical CVD, independent of HIV status [17]. Similarly, in the Pharmacokinetic and Clinical Observations in People Over Fifty (POPPY) study, the systemically inflamed/gut epithelial dysfunction group showed higher estimated CVD risk [18]. These studies suggest a distinct inflammatory profile in persons with HIV that is associated with a high clinical CVD risk, with biomarkers linked to gut dysfunction implicated.

Despite these significant findings, the potential role of the gut microbiome on influencing this high-risk inflammatory phenotype remains unclear. Understanding the role of the gut microbiome in host inflammatory responses in HIV infection could provide valuable insights into the pathogenesis of premature aging and help identify individuals at higher risk of noncommunicable diseases like CVD, aiding earlier therapeutic or preventive strategies to reduce CVD risk. Our study aim was to investigate the interactions between host gut microbiome compositions and systemic inflammatory responses and to correlate these with subclinical CVD outcomes.

MATERIALS AND METHODS

Study Cohort

The Understanding the Pathology of Comorbid Disease in HIV-Infected Individuals Coronary Artery Disease (HIV UPBEAT CAD) study is a substudy of the HIV UPBEAT study, a single-center, prospective, observational cohort that enrolled

people with and without HIV from similar demographic backgrounds. Participants with HIV were recruited from the infectious diseases clinics at the Mater Misericordiae University Hospital, Dublin, Ireland, whereas people without HIV were recruited from the local geographical area. Participants were included if they were >40 years old, with no known history of CVD. Those with and without HIV were propensity score matched for traditional CVD risk factors to controls, ensuring an even distribution of estimated CVD risk across both groups.

Participants underwent coronary computed tomography angiography (CCTA) to assess subclinical CAD, as described previously [17]. Participants also provided clinical/medical history, in addition to stool samples and fasting bloods, from which serum and plasma were derived and stored at −80°C. The study was approved by the Mater Misericordiae University Hospital and Mater Private Hospital Institutional Review Board and all participants provided written informed consent.

Nucleic Acid Purification and 16S rRNA Amplification

DNA was extracted from thawed stool samples using the repeated bead-beating Qiagen DNA extraction method as previously described [19]. A part of the 16S rRNA gene was amplified from isolated DNA using primers for the V3–V4 region according to the Illumina 16S Metagenomic Sequencing Protocol. Library DNA concentration was quantified using a Qubit fluorometer (Invitrogen) using the high-sensitivity assay, and samples were pooled at a standardized concentration. The uniquely barcoded amplicons were sequenced on an Illumina MiSeq platform utilizing 2 × 300 bp chemistry.

Processing and Quality Control

Resulting sequence data were quality-filtered using TrimGalore v0.6.7 [20]. Read de-replication, learning of the error rates, and sample sequence variant inference with pooled samples followed by the construction of amplicon sequence variant (ASV) table and removal of chimeras were performed using LotuS2 pipeline (with dada2 clustering) at the ASV level for α -diversity analysis [21]. To ensure fair comparison across samples, we applied rarefaction, randomly subsampling reads to a uniform depth across all samples for α -diversity estimation, thereby controlling for differences in sequencing depth. Annotate of the 16S sequences at the species level was constructed using SPINGO [22]. In addition, we also applied a quality filtering step removing rare species from the analysis, keeping those present in 10% or more of samples. Absolute counts were transformed to relative abundance to normalize the data and focus on the proportional representation of taxa across samples.

Measurement of Inflammatory Biomarkers

The biomarkers analyzed in this study have been previously assessed using quantitative immunoassays as previously described (Supplementary Table 1) [17].

Collection of Dietary Data

Dietary data were collected using a semi-quantitative food frequency questionnaire (FFQ) weighted by 10 consumptive frequencies according to the method used by Claesson et al [23]. Participants were asked to recall dietary intakes, including nutritional supplements, over the previous 4 weeks to provide an average daily nutritional intake. FFQ coding, data cleaning, and data checks were conducted by a single, trained individual to ensure consistency of data. The Healthy Food Diversity (HFD) index is based on dietary proportions derived from FFQ data and associated health values from nutritional guidelines [23]. It considers diversity of food in conjunction with using a weighted health value to circumnavigate many of the traditional problems of measures of dietary diversity. The HFD provides a score between 0 and 1, where a higher value indicates a more diverse diet.

Statistical Analysis

For clinical factors, continuous and categorical variables were summarized using median with interquartile range (IQR) and frequency (%), respectively. Categorical and continuous variables were compared using χ^2 and Kruskal-Wallis or Wilcoxon rank-sum tests, respectively.

Kruskal-Wallis and Mann-Whitney tests were used to explore significant differences in clinical and biochemical measures and α -diversity (Chao1 and Shannon indices). The permutational multivariate analysis of variance (PERMANOVA) was performed with Bray-Curtis dissimilarity, Jaccard dissimilarity, and binary Jaccard dissimilarity at the species and genus level, respectively, through vegan adonis2 function. The covariates of sex, age, comorbidities, and medications (antihypertensives, statins, and ART) were included in the model, and the marginal sums of squares was applied to avoid issues related to the order of factors in adonis2. To identify differentially abundant species and genera between people with and those without HIV, we used analysis of compositions of microbiomes with bias correction (ANCOM-BC; version 2.2.0) [24]. ANCOM-BC incorporates a centered log-ratio transformation to address the compositional nature of microbiome data and reports bias-corrected log fold changes in abundance between groups, adjusting for covariates (ie, sex, age, comorbidities, and the use of statins and ART). The *P* values were adjusted using the Benjamini-Hochberg procedure for multiple hypothesis testing [25]. We used nonparametric Spearman correlation to test associations between biomarkers, the CCTA outcome data, and the dietary data with the differential abundance of the identified microbiome species and genera. Prior to formation of the correlation matrix, the biomarkers were log-transformed (for approximate normality) and scaled to ensure that biomarkers with larger intrinsic variation did not dominate the correlation matrix.

Table 1. Baseline Demographics and Coronary Artery Disease (CAD) Risk Factors of Participants of the HIV UPBEAT CAD Study, Stratified by Human Immunodeficiency Virus Status

Characteristic	People Without HIV (n = 37)	People With HIV (n = 44)	<i>P</i> Value ^a
Age, y, median (IQR)	50 (46–56)	50 (45–57)	.67
Sex			.63
Female	11 (30)	11 (25)	
Male	26 (70)	33 (75)	
Race/ethnicity			>.99
African	8 (22)	10 (23)	
Asian	1 (2.7)	2 (4.5)	
Caucasian	28 (76)	32 (73)	
Smoking history			.98
Current smoker	7 (19)	9 (20)	
Former smoker	11 (30)	13 (30)	
Never smoker	19 (51)	22 (50)	
Statin use	4 (11)	22 (50)	<.001
History of hypertension	7 (19)	16 (36)	.083
Diabetes	1 (2.7)	3 (6.8)	.62
High cholesterol (TC level in mmol/L)	11 (30)	24 (55)	.025
BMI, kg/m ² , median (IQR)	27.2 (25.0–29.3)	29.4 (24.5–32.2)	.38
ART duration, y, median (IQR)	...	9.5 (7.2–13.9)	

Data are presented as No. (%) unless otherwise indicated.

Abbreviations: ART, antiretroviral therapy; BMI, body mass index; HIV, human immunodeficiency virus; IQR, interquartile range; TC, total cholesterol.

^aWilcoxon rank-sum test, Pearson χ^2 test, Fisher exact test, or Wilcoxon rank-sum exact test.

RESULTS

Clinical Characteristics of the Study Population

Eighty-one participants with available stool samples were included in this analysis, of whom 44 were people with HIV and 37 were people without HIV (Table 1). The median age was 51 (IQR, 46–56) years, 73% were male, and 74% were Caucasian. Although estimated CVD risk was similar between the 2 groups (as a result of the propensity score matching), those with HIV had a higher prevalence of hypercholesterolemia (*P* < .025) and were more likely to be on statins (*P* < .001).

Microbiome Diversity in People With and Without HIV

While there was no significant difference in α -diversity at the species and genus level (using Chao1 and Shannon indices) between people with and without HIV (Figure 1), a significant separation in β -diversity was observed between the 2 groups, using the Bray-Curtis dissimilarity (PERMANOVA, *P* < .001, *R*² = 0.034) and Jaccard dissimilarity (PERMANOVA, *P* < .001, *R*² = 0.027). ANCOM-BC differential abundance analysis identified 42 species (Figure 2 and Supplementary Figure 1) and 10 genera (Supplementary Figure 2) that were significantly enriched or depleted in people with HIV compared to people without HIV. In this group, *Olsenella uli* was the most significantly enriched of these species, followed by *Anaerovibrio*




Figure 1. Alpha-diversity at the amplicon sequence variant level for species (*A*) and genus (*B*) for people with and without human immunodeficiency virus (HIV). Beta-diversity with Bray-Curtis dissimilarity and Jaccard similarity at the species level (*C* and *D*) and genus level (*E* and *F*), respectively. Permutational multivariate analysis of variance was performed, and the effect size for Axis 1 in the β -diversity results was interpreted using the 95% confidence interval.

lipolyticus and *Slackia exigua*, respectively. The most significantly depleted species, primarily SCFA producers, were *Odoribacter splanchnicus*, followed by *Alistipes shahii*, *Bacteroides uniformis*, and *Barnesiella intestinihominis*. *Barnesiella intestinihominis* is a taxa that has been positively associated with healthy aging, and its reduced

abundance has been linked to an increased burden of cerebral small vessel disease [26]. *Olsenella* was the most significantly enriched genus, followed by *Megasphaera*, *Mitsuokella*, and *Flavonifractor*, respectively. The most significantly depleted genera were *Bacteroides*, followed by *Faecalibacterium* and *Fusicatenibacter*.




Figure 2. Differential abundance (DA) analysis of 42 species by in people with human immunodeficiency virus (HIV) compared to those without HIV. DA was assessed by analysis of compositions of microbiomes with bias correction while adjusting for the covariates based on filtered species abundance (present in at least 10% of samples and with maximum reads ≥ 10). False discovery rate < 0.1 was considered as DA species.

Comparison of Microbiome Taxa With Subclinical Cardiovascular Disease

Overall, 30 (37%) participants had evidence of subclinical CAD on CCTA. Focusing on Agatston score and CAP burden as measures of subclinical CAD, CAP burden was similar in both groups, but those without HIV were more likely to be on a statin ($n = 22$ vs $n = 4$, $P < .001$) and have an Agatston score > 100 ($n = 9$ vs $n = 3$, $P < .05$) (Table 2). In people with HIV, an increased abundance of the bacterium *Bifidobacterium pseudocatenulatum*, a potentially probiotic microbe that is associated with SCFA production and bile acid metabolism, correlated with a lower Agatston score and overall CAP burden (Supplementary Figure 3) [27]. Similarly, an increased abundance of *Megamonas hypermegale* and *Selenomonas ruminantium*, both SCFA-producing bacteria involved in carbohydrate fermentation, was linked to a lower CAP burden in people with HIV. These findings suggest a role of SCFA-producing bacteria in modulating inflammation and maintaining cardiovascular health. No significant associations at the genus level were associated with Agatston score or CAP burden (Supplementary Figure 4).

Dietary Data

Dietary data were collected from 81 participants (Table 3). There was no significant difference in daily nutrient intake or the HFD index between the groups with and without HIV. However, dietary patterns correlated with several microbiome changes relevant to CVD risk. For example, higher intake of saturated and trans-unsaturated fatty acids was associated with an enrichment of *Ruminococcus bromii*, a SCFA producer, and depletion of

key gut commensals such as *Pyramidobacter pisciolens* and *Bacteroides* spp (Figure 3). Additionally, higher intake of sugars and calories correlated with depletion of SCFA-producing *Bacteroides* spp and *Alistipes* spp (Supplementary Figure 5). These important gut commensals are involved in the breakdown of complex dietary fibers and polysaccharides, contributing to digestion. Importantly, the observed depletion of SCFA producers is consistent with findings from the CCTA analysis, further emphasizing their potential role in modulating cardiovascular outcomes.

Associations Between Microbial and Inflammatory Biomarkers

Building on the links between diet, microbiota, and cardiovascular outcomes, we explored associations between microbial taxa and 38 inflammatory biomarkers. Previous work within this cohort identified a distinct inflammatory phenotype defined by elevated circulating levels of I-FABP, T-cell stimulation, and systemic inflammation markers that were associated with subclinical CAD and prevalent CVD [17]. In this study, we found an increased abundance of several SCFA bacteria, including *Prevotella salivae*, *Megasphaera elsdenii*, *Enterorhabdus mucosicola*, and *Succinivibrio dextrinosolvens*, that correlated with elevated circulating I-FABP levels (Figure 4). Similarly, levels of interleukin 18 (IL-18), a potent proinflammatory cytokine that facilitates type 1 inflammatory responses [28], were positively associated with increased abundance of several gram-negative microbes (*P salivae*, *Prevotella albensis*, *M hypermegale*, *Burkholderia andropogonis*) and lipid-metabolizing bacteria, such as *A lipolyticus*. An overabundance of lipid-metabolizing bacteria may contribute

Table 2. Subclinical Coronary Artery Disease (CAD) as Measured by Coronary Computed Tomography Angiography in HIV UPBEAT CAD Substudy Participants

Characteristic	HIV Negative (n = 37)	People With HIV (n = 44)	P Value ^a
History of statin use	4 (11)	22 (50)	<.001
Any coronary plaque	16 (43)	14 (32)	.29
Calcified plaque	15 (41)	13 (30)	.30
Noncalcified plaque	3 (8.1)	6 (14)	.50
Partially calcified plaque	9 (24)	8 (18)	.50
Worst stenosis			.30
<25% stenosis	4 (11)	7 (16)	
26%–50% stenosis	9 (24)	4 (9.1)	
>50% stenosis	3 (8.1)	3 (6.8)	
No significant stenosis	21 (57)	30 (68)	
Agatston score, median (IQR)	0.00 (0.00–4.20)	0.00 (0.00–2.10)	.12
Agatston score >100	9 (24)	3 (6.8)	.027
Calcium volume score, median (IQR)	0.00 (0.00–3.73)	0.00 (0.00–1.87)	.15
Calcium volume score >100	6 (16)	2 (4.5)	.13

Data are presented as No. (%) unless otherwise indicated.

Abbreviations: HIV, human immunodeficiency virus; IQR, interquartile range.

^aPearson χ^2 test, Fisher exact test, or Wilcoxon rank-sum test.

to an imbalance in lipid metabolism, potentially leading to adverse metabolic outcomes [29].

In people with HIV, a depletion of SCFA-producing bacterial species, including *Bacteroides* spp, *Alistipes* spp, and *R. bromii* (Figure 4), as well as the genus *Succinivibrio* (Supplementary Figure 6), correlated with upregulated inflammatory markers (hsCRP, D-dimer, CD40 ligand, interferon- γ , and lipopolysaccharide-binding protein [LBP]) and endothelial dysfunction markers (Vascular cell adhesion protein 1 [VCAM-1], E-selectin). These inflammatory patterns align with the microbiota associations seen in the CCTA and dietary findings, further connecting changes in the gut microbiome with cardiovascular risk.

DISCUSSION

This study describes the differences in fecal microbiota compositions associated with CVD risk in people with HIV. We identified decreased abundance of SCFA-producing bacteria, linked to subclinical CVD findings on CCTA, elevated inflammatory biomarkers, and higher dietary intake of fatty acids and sugars. Additionally, an increased abundance of proinflammatory bacteria correlated with elevated levels of I-FABP and IL-18, markers previously associated with CVD in this cohort [17]. While the cross-sectional design precludes causal inference, these findings suggest that the gut microbiome may influence inflammation and CVD risk in HIV.

Compared to people without HIV, an enrichment of *B. pseudocatenulatum* was associated with a lower CAP burden and Agatston score in people with HIV, indicating a more

Table 3. Median Daily Dietary Nutritional Intake and Healthy Food Diversity Index in People With and Without Human Immunodeficiency Virus

Characteristic	People Without HIV (n = 35)	People With HIV (n = 41)	P Value ^a
Energy, kcal	7.75 (7.42–7.98)	7.71 (7.52–8.05)	.66
Carbohydrates, g	5.28 (5.05–5.82)	5.51 (5.11–5.94)	.24
Sugars, g	4.63 (4.15–4.88)	4.71 (4.31–5.14)	.14
Fiber, g	3.34 (2.90–3.63)	3.33 (3.07–3.66)	.51
Saturated fatty acids, g	3.37 (3.09–3.52)	3.48 (3.15–3.80)	.36
Polyunsaturated fatty acids, g	2.53 (2.18–2.81)	2.51 (2.38–3.01)	.32
Trans-saturated fatty acids, g	0.28 (0.09–0.39)	0.36 (0.01–0.65)	.43
Cholesterol, mg	5.80 (5.40–6.05)	5.75 (5.54–6.08)	>.99
Protein, g	4.85 (4.53–5.18)	4.58 (4.45–4.95)	.15
Alcohol, g	2.18 (1.38–2.55)	1.38 (0.00–2.60)	.18
Caffeine, mg	7.16 (4.79–9.42)	6.12 (5.47–7.09)	.17
HFD index	0.49 (0.46–0.51)	0.49 (0.47–0.52)	.58

Data are presented as median (interquartile range) unless otherwise indicated.

Abbreviations: HFD, Healthy Food Diversity; HIV, human immunodeficiency virus.

^aWilcoxon rank-sum exact test, Wilcoxon rank-sum test, Pearson χ^2 test, or Fisher exact test.

favorable cardiovascular profile. Conversely, in this group, we found an increased abundance of bacteria known to be associated with CVD, including *Megasphaera micronuciformis*, *Streptococcus* spp, and *Prevotella* spp [15]. Additionally, in people with HIV, a depletion of *Bacteroides plebeius*, *Bacteroides barnesi*, *Alistipes* spp, the butyrate-producing *Eubacterium ventriosum*, and the keystone species *R. bromii* was associated with greater CAP burden and elevated Agatston scores. Although these are not the typical major butyrate producers, these other SCFA-producing microbes may have a role to play in CVD outcomes. Previous studies have shown that SCFA-producing microbes may influence immune regulation and can exert positive effects on blood pressure, CAD, and atherosclerosis [30]. The depletion of SCFA-producing organisms has been associated with atherosclerosis [9] and heart failure [31] in the general population.

A reduction of SCFA microbes may lead to increased systemic inflammation through multiple interconnected mechanisms. SCFAs play a key role in maintaining intestinal barrier integrity by serving as an energy source for colonocytes and promoting tight junction function [32]. Reduced SCFA production compromises this barrier, leading to increased gut permeability and MT [33]. Furthermore, SCFAs modulate inflammation through anti-inflammatory pathways, including the activation of regulatory T cells and inhibition of histone deacetylases, both of which help suppress excessive immune responses [32].

While studies investigating the role of SCFA-producing microbes in relation to CVD outcomes in people with HIV are limited, a reduction in production of a key SCFA, butyrate, has been found to be associated with diabetes in women with HIV [34]. Similarly, reductions in butyrate-producing




Figure 3. Heatmap showing the correlation of the differentially abundant (DA) species with mean nutrient intake and the Healthy Food Diversity (HFD) index. DA was assessed by analysis of compositions of microbiomes with bias correction while adjusting for the covariates based on filtered species abundance (present in at least 10% of samples and with maximum reads ≥ 10). False discovery rate < 0.1 was considered as DA species. Green stars indicate * $P < .05$; ** $P < .01$; *** $P < .001$.

microbes have been observed in people with HIV with hyperglycemia, hypertension, and dyslipidemia [13]. Our findings are in line with these previous studies and suggest that alterations in gut microbiota, particularly a reduction in SCFA-producing bacteria like butyrate producers, may be linked to cardiovascular and metabolic conditions in people with HIV.

In keeping with previous studies, we found no significant differences in nutrient intake or HFD index between people with and without HIV [35]. However, a depletion of SCFA-producing species was linked to higher intake of fats and sugars in both groups, affecting gut microbiota and CVD risk. Several studies have shown that a high-fat diet has a negative impact on gut microbiota, with a reduction in SCFA-producing bacteria affecting host metabolism and linked with obesity, insulin sensitivity, and lipid metabolism in the general population [9]. In the context of HIV, studies in simian immunodeficiency virus-infected macaques have shown that a high-fat diet exacerbates inflammation, increasing levels of proinflammatory cytokines such as IL-18 [36]. Consistent with these findings, we observed elevated circulating IL-18, alongside other proinflammatory biomarkers, which were associated with the depletion of SCFA-producing species. IL-18 has been previously

reported to be elevated in obesity and linked with lipodystrophy in people with HIV, underscoring its relevance to metabolic and inflammatory dysregulation in this population [37, 38].

An increased abundance of *Megasphaera* and *Prevotella* spp, along with *A. lipolyticus*, was associated with elevated I-FABP and IL-18 levels in people with HIV. I-FABP, a marker of endothelial damage, has been associated with CVD [39] and hypertension [40] in the general population. In HIV groups, I-FABP has been associated with increased adiposity, suggesting that intestinal damage is linked with nutrient malabsorption and inflammation [41]. The elevation of I-FABP likely reflects increased enterocyte turnover due to systemic metabolic derangements rather than, in the absence of suggestive markers (sCD14, LBP), gut MT. Together, these findings suggest an inflammatory state driven by reduced SCFA-producing microbes and elevated proinflammatory biomarkers, contributing to CVD risk in HIV infection.

Of note, a number of oral microbes were identified in people with HIV in this study, including the species *O. uli* and *S. exigua*. The oral microbiome plays a significant role not only in maintaining oral health but also in normal homeostasis. The translocation of oral microbes to the gut is particularly evident in inflammatory bowel diseases, sporadic colorectal cancer, and




Figure 4. Heatmap showing the correlation of the differentially abundant (DA) species with the inflammatory cytokines and T-cell surface markers for people with human immunodeficiency virus (HIV) and those without HIV. DA was assessed by analysis of compositions of microbiomes with bias correction (ANCOM-BC) while adjusting for the covariates based on filtered species abundance (present in at least 10% of samples and with maximum reads ≥ 10). False discovery rate < 0.1 was considered as DA species. Green stars indicate $*P < .05$; $**P < .01$; $***P < .001$. actCD8, activated CD8 T cells; CD14, cluster of differentiation 14; ExhCD4, exhausted CD4 T cells; ExhCD8, exhausted CD8 T cells; hsCRP, high sensitivity C Reactive Protein; I-FABP, intestinal fatty acid binding protein; IFN, interferon; IL, interleukin; IL1b, interleukin 1 beta; IL1RA, IL-1 receptor antagonist; LBP, LPS binding protein; Lp-PLA2, lipoprotein-associated phospholipase A2; MCP-1, monocyte chemoattractant protein; MIP-1, macrophage inflammatory protein; sCD163, soluble cluster of differentiation 163; sCD40L, soluble CD40 ligand; s-ICAM, soluble intercellular adhesion molecule; sensCD8, sensitised CD8 T cells; TNFR, tumor necrosis factor receptor; Tregs, regulatory T cells; TSLP, thymic stromal lymphopoietin; VCAM1, vascular cell adhesion molecule; vwf, von Willebrand factor.

liver disease [42–44]. Furthermore, HIV infection can significantly alter the composition of the oral microbiome, potentially contributing to dysbiosis and increasing susceptibility to systemic diseases [45]. Meta-analysis also shows that increased abundance of oral microbes is associated with unhealthy aging, suggesting that microbial shifts in the oral cavity can contribute to inflammation and chronic diseases associated with aging [46]. Additionally, a number of rumen microbes were identified in our results, including *M. elsdenii*, *S. dextrinosolvens*, and *S. ruminantium*. The presence of these microbes could be attributed to diet or animal exposure, but these species have also been identified in studies of the human gut [47] and in human disease states [47–50].

The strengths of this study include (1) a well-matched CVD risk cohort of participants representing a diverse population of both men and women from different ethnicities; (2) a comprehensive comparison of diet, inflammatory phenotypes, and gut

microbiome composition in the context of both subclinical and clinical CVD risk; and (3) a direct comparison of people with and without HIV, allowing for the identification of unique differences attributable to HIV status.

Limitations of this study include its cross-sectional design and relatively small sample size. This highlights the need for replication in larger, longitudinal cohorts with diverse demographics. Although SPINGO provides the highest accuracy at the species level compared to other tools, the annotation of 16S amplicon sequences at the species level is still limited. We included genus-level classification to aid a more reliable microbial community analysis. While the FFQ offers easy administration, low burden on participants, and cost-efficiency, it is subject to recall and social desirability bias. Inclusion of other dietary assessment tools, such as dietary records and 24-hour food recall, may enhance dietary data accuracy. Finally, while propensity scores were used to match groups, unmeasured factors, including sexual orientation, which

was not assessed in this cohort but is known to influence microbiome changes, may still affect outcomes. However, as both study groups were recruited from similar socioeconomic backgrounds, this likely helps mitigate some of these unmeasured behavioral and socioeconomic factors. Future research should explore the functional capacities of specific microbial species and examine SCFA associations through targeted metabolomics to assess how gut microbiome interventions, such as probiotic use or dietary interventions, may impact inflammatory phenotypes in people with HIV.

CONCLUSIONS

In conclusion, we describe how gut microbiome alterations, particularly the depletion of SCFA-producing microbes, is associated with diet and systemic inflammation, contributing to clinical and subclinical CVD in people with HIV. These findings underscore the importance of dietary counseling and targeted interventions to address gut dysbiosis, as well as the potential for monitoring inflammatory markers to enhance CVD risk prediction and prevention. This evidence supports a precision medicine approach to managing HIV-associated comorbidities and improving CVD outcomes.

Supplementary Data

Supplementary materials are available at *The Journal of Infectious Diseases* online (<http://jid.oxfordjournals.org/>). **Supplementary materials** consist of data provided by the author that are published to benefit the reader. The posted materials are not copyedited. The contents of all **supplementary data** are the sole responsibility of the authors. Questions or messages regarding errors should be addressed to the author.

Notes

Acknowledgments. The authors wish to thank the HIV UPBEAT CAD substudy participants and their families for their involvement in this study. HIV UPBEAT study group members are as follows: Dr Padraig McGettrick, Dr Elena Alvarez Barco, Dr Willard Tinago, Mr Alejandro Garcia Leon, Ms Aoife McDermott, Dr Tara McGinty, Dr Aoife G. Cotter, Mr Alan Macken, and Prof Patrick W. G. Mallon (Centre for Experimental Pathogen Host Research, University College Dublin, Ireland); Dr Eoin Kavanagh, Prof Geraldine McCarthy, Dr Gerard Sheehan, and Dr John Lambert (Mater Misericordiae University Hospital, Dublin, Ireland); Prof William Powderly (Institute for Public Health, Washington University School of Medicine, St Louis, Missouri); Prof Juliet Compston (Department of Medicine, Cambridge Biomedical Campus, Cambridge, United Kingdom); and Prof Caroline Sabin (Institute for Global Health, University College London, United Kingdom).

Author contributions. All authors contributed substantially to the development of this study and contributed significantly and equally to this manuscript.

Disclaimer. The funders had no role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Financial support. The HIV UPBEAT Cohort Study has received funding from the Irish Health Research Board (award number HRA_POR/2010/66 to P. W. M. as principal investigator). R. M. C. was supported by the Irish Clinical Academic Training program, supported by the Wellcome Trust and the Irish Health Research Board (grant number 203930/B/16/Z), the Health Service Executive, National Doctors Training and Planning and the Health and Social Care, Research and Development Division, Northern Ireland. Work in P. W. O.'s laboratory is supported by Science Foundation Ireland through a Centre Award to APC Microbiome Ireland (12/RC/2273_P2) and by the European Union's Horizon Europe research and innovation program under grant agreement number 101079777 with the MicrobAIome consortium.

Potential conflicts of interest. P. W. M. has received honoraria and/or travel grants from Gilead Sciences, MSD, Bristol-Myers Squibb, and ViiV Healthcare. C. S. has received honoraria and funding to support participation in data safety and advisory groups, for preparation of educational materials, and for participation in speaker's panels from Gilead Sciences, ViiV Healthcare, and MSD. A. L. has consulted for Gilead and Abbott. All other authors report no potential conflicts.

All authors have submitted the ICMJE Form for Disclosure of Potential Conflicts of Interest. Conflicts that the editors consider relevant to the content of the manuscript have been disclosed.

References

1. Antiretroviral Therapy Cohort Collaboration. Survival of HIV-positive patients starting antiretroviral therapy between 1996 and 2013: a collaborative analysis of cohort studies. *Lancet HIV* **2017**; 4:e349–56.
2. Hawkins KL, Brown TT, Margolick JB, Erlandson KM. Geriatric syndromes: new frontiers in HIV and sarcopenia. *AIDS* **2017**; 31(Suppl 2):S137–46.
3. McGettrick P, Mallon PWG, Sabin CA. Cardiovascular disease in HIV patients: recent advances in predicting and managing risk. *Expert Rev Anti Infect Ther* **2020**; 18:677–88.
4. Smit M, Van Zoest RA, Nichols BE, et al. Cardiovascular disease prevention policy in human immunodeficiency virus: recommendations from a modeling study. *Clin Infect Dis* **2018**; 66:743–50.
5. Lorenzatti AJ, Servato ML. New evidence on the role of inflammation in CVD risk. *Curr Opin Cardiol* **2019**; 34: 418–23.

6. Nordell AD, McKenna M, Borges ÁH, et al. Severity of cardiovascular disease outcomes among patients with HIV is related to markers of inflammation and coagulation. *J Am Heart Assoc* **2014**; 3:e000844.
7. Li SX, Armstrong A, Neff CP, Shaffer M, Lozupone CA, Palmer BE. Complexities of gut microbiome dysbiosis in the context of HIV infection and antiretroviral therapy. *Clin Pharmacol Ther* **2016**; 99:600–11.
8. Vujkovic-Cvijin I, Somsouk M. HIV and the gut microbiota: composition, consequences, and avenues for amelioration. *Curr HIV/AIDS Rep* **2019**; 16:204–13.
9. Nogal A, Valdes AM, Menni C. The role of short-chain fatty acids in the interplay between gut microbiota and diet in cardio-metabolic health. *Gut Microbes* **2021**; 13:1–24.
10. Chen W, Zhang S, Wu J, et al. Butyrate-producing bacteria and the gut-heart axis in atherosclerosis. *Clin Chim Acta* **2020**; 507:236–41.
11. Jie Z, Xia H, Zhong SL, et al. The gut microbiome in atherosclerotic cardiovascular disease. *Nat Commun* **2017**; 8:845.
12. Lu W, Feng Y, Jing F, et al. Association between gut microbiota and CD4 recovery in HIV-1 infected patients. *Front Microbiol* **2018**; 9:1451.
13. Villanueva-Millán MJ, Pérez-Matute P, Recio-Fernández E, Lezana Rosales JM, Oteo JA. Characterization of gut microbiota composition in HIV-infected patients with metabolic syndrome. *J Physiol Biochem* **2019**; 75:299–309.
14. Sandler NG, Douek DC. Microbial translocation in HIV infection: causes, consequences and treatment opportunities. *Nat Rev Microbiol* **2012**; 10:655–66.
15. Wang Z, Peters BA, Usyk M, et al. Gut microbiota, plasma metabolomic profiles, and carotid artery atherosclerosis in HIV infection. *Arterioscler Thromb Vasc Biol* **2022**; 42:1081–93.
16. Dillon SM, Kibbie J, Lee EJ, et al. Low abundance of colonic butyrate-producing bacteria in HIV infection is associated with microbial translocation and immune activation. *AIDS* **2017**; 31:511–21.
17. McGettrick P, Tinago W, O'Brien J, et al. Distinct inflammatory phenotypes are associated with subclinical and clinical cardiovascular disease in people with human immunodeficiency virus. *J Infect Dis* **2024**; 230:e616–21.
18. Sukumaran L, Kunisaki KM, Bakewell N, et al. Association between inflammatory biomarker profiles and cardiovascular risk in individuals with and without HIV. *AIDS* **2023**; 37:595–603.
19. Yu Z, Morrison M. Improved extraction of PCR-quality community DNA from digesta and fecal samples. *BioTechniques* **2004**; 36:808–12.
20. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* **2014**; 30:2114–20.
21. Özkurt E, Fritscher J, Soranzo N, et al. Lotus2: an ultrafast and highly accurate tool for amplicon sequencing analysis. *Microbiome* **2022**; 10:176.
22. Allard G, Ryan FJ, Jeffery IB, Claesson MJ. SPINGO: a rapid species-classifier for microbial amplicon sequences. *BMC Bioinformatics* **2015**; 16:324.
23. Claesson MJ, Jeffery IB, Conde S, et al. Gut microbiota composition correlates with diet and health in the elderly. *Nature* **2012**; 488:178–84.
24. Lin H, Peddada SD. Analysis of compositions of microbiomes with bias correction. *Nat Commun* **2020**; 11:3514.
25. Benjamini Y, Hochberg Y. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *J R Stat Soc Series B Stat Methodol* **1995**; 57:289–300.
26. Fongang B, Satizabal C, Kautz TF, et al. Cerebral small vessel disease burden is associated with decreased abundance of gut *Barnesiella intestinihominis* bacterium in the Framingham Heart Study. *Sci Rep* **2023**; 13:13622.
27. Al-Sheraji SH, Ismail A, Manap MY, Mustafa S, Yusof RM, Hassan FA. Hypocholesterolaemic effect of yoghurt containing *Bifidobacterium pseudocatenulatum* G4 or *Bifidobacterium longum* BB536. *Food Chem* **2012**; 135: 356–61.
28. El-Far M, Durand M, Turcotte I, et al. Upregulated IL-32 expression and reduced gut short chain fatty acid caproic acid in people living with HIV with subclinical atherosclerosis. *Front Immunol* **2021**; 12:664371.
29. Schoeler M, Caesar R. Dietary lipids, gut microbiota and lipid metabolism. *Rev Endocr Metab Disord* **2019**; 20: 461–72.
30. den Besten G, van Eunen K, Groen AK, Venema K, Reijngoud DJ, Bakker BM. The role of short-chain fatty acids in the interplay between diet, gut microbiota, and host energy metabolism. *J Lipid Res* **2013**; 54:2325–40.
31. Modrego J, Ortega-Hernández A, Goirigolzarri J, et al. Gut microbiota and derived short-chain fatty acids are linked to evolution of heart failure patients. *Int J Mol Sci* **2023**; 24:13892.
32. Vinolo MAR, Rodrigues HG, Nachbar RT, Curi R. Regulation of inflammation by short chain fatty acids. *Nutrients* **2011**; 3: 858–76.
33. Lewis K, Lutgendorff F, Phan V, Söderholm JD, Sherman PM, McKay DM. Enhanced translocation of bacteria across metabolically stressed epithelia is reduced by butyrate. *Inflamm Bowel Dis* **2010**; 16:1138–48.
34. Moon JY, Zolnik CP, Wang Z, et al. Gut microbiota and plasma metabolites associated with diabetes in women with, or at high risk for, HIV infection. *EBioMedicine* **2018**; 37:392–400.
35. Dillon SM, Lee EJ, Kotter CV, et al. An altered intestinal mucosal microbiome in HIV-1 infection is associated with mucosal and systemic immune activation and endotoxemia. *Mucosal Immunol* **2014**; 7:983–94.

36. Mansfield KG, Carville A, Wachtman L, et al. A diet high in saturated fat and cholesterol accelerates simian immunodeficiency virus disease progression. *J Infect Dis* **2007**; 196:1202–10.
37. Lindegaard B, Hansen AB, Pilegaard H, Keller P, Gerstoft J, Pedersen BK. Adipose tissue expression of IL-18 and HIV-associated lipodystrophy. *AIDS* **2004**; 18:1956–8.
38. Jia X, Buckley L, Sun C, et al. Association of interleukin-6 and interleukin-18 with cardiovascular disease in older adults: atherosclerosis risk in communities study. *Eur J Prev Cardiol* **2023**; 30:1731–40.
39. Isnard S, Fombuena B, Sadouni M, et al. Circulating β -d-glucan as a marker of subclinical coronary plaque in antiretroviral therapy-treated people with human immunodeficiency virus. *Open Forum Infect Dis* **2021**; 8: ofab109.
40. Kim S, Goel R, Kumar A, et al. Imbalance of gut microbiome and intestinal epithelial barrier dysfunction in patients with high blood pressure. *Clin Sci (Lond)* **2018**; 132:701–18.
41. Dirajlal-Fargo S, Moser C, Rodriguez K, et al. Changes in the fungal marker β -D-glucan after antiretroviral therapy and association with adiposity. *Open Forum Infect Dis* **2019**; 6:ofz434.
42. Li J, O'Toole PW. Disease-associated microbiome signature species in the gut. *PNAS Nexus* **2024**; 3:pgae352.
43. Read E, Curtis MA, Neves JF. The role of oral bacteria in inflammatory bowel disease. *Nat Rev Gastroenterol Hepatol* **2021**; 18:731–42.
44. Wirbel J, Pyl PT, Kartal E, et al. Meta-analysis of fecal metagenomes reveals global microbial signatures that are specific for colorectal cancer. *Nat Med* **2019**; 25:679–89.
45. Li S, Su B, He QS, Wu H, Zhang T. Alterations in the oral microbiome in HIV infection: causes, effects and potential interventions. *Chin Med J (Engl)* **2021**; 134:2788–98.
46. Ghosh TS, Das M, Jeffery IB, O'Toole PW. Adjusting for age improves identification of gut microbiome alterations in multiple diseases. *Elife* **2020**; 9:e50240.
47. Pomeroy C, Shanholtzer CJ, Peterson LR. *Selenomonas* bacteraemia—case report and review of the literature. *J Infect* **1987**; 15:237–42.
48. Horiba K, Aso S, Oda R, Tateishi Y, Ura K, Kuroda M. Complete genome sequence of *Selenomonas species* strain TAMA-11512, isolated from blood culture of a septic patient. *Microbiol Resour Announc* **2024**; 13:e0007024.
49. Tecer D, Gogus F, Kalkanci A, et al. Succinivibrionaceae is dominant family in fecal microbiota of Behçet's Syndrome patients with uveitis. *PloS one* **2020**; 15:e0241691.
50. Cabral LDS, Weimer PJ. *Megasphaera elsdenii*: Its Role in Ruminant Nutrition and Its Potential Industrial Application for Organic Acid Biosynthesis. *Microorganisms* **2024**; 12.