


Title	CT in the detection of latent tuberculosis: a systematic review
Authors	Moore, Niamh;Maher, M.;Murphy, G.;O'Callaghan Maher, M.;O'Connor, O. J.;McEntee, Mark F.
Publication date	2023-05-13
Original Citation	Moore, N., Maher, M., Murphy, G., Maher, M. C., O'Connor, O. J. and McEntee, M. F. (2023) 'CT in the detection of latent tuberculosis: a systematic review', Clinical Radiology, 78(8), pp.568-575. https://doi.org/10.1016/j.crad.2023.04.014
Type of publication	Article (peer-reviewed)
Link to publisher's version	10.1016/j.crad.2023.04.014
Rights	© 2023, Published by Elsevier Ltd on behalf of The Royal College of Radiologists. All rights reserved. This manuscript version is made available under the CC BY-NC-ND 4.0 license. - https://creativecommons.org/licenses/by-nc-nd/4.0/
Download date	2026-09-20 14:27:19
Item downloaded from	https://hdl.handle.net/10468/14476

Journal Pre-proof

CT in the detection of latent tuberculosis: a systematic review

N. Moore, M. Maher, G. Murphy, M. O'Callaghan Maher, O.J. O'Connor, M.F. McEntee



PII: S0009-9260(23)00180-0

DOI: <https://doi.org/10.1016/j.crad.2023.04.014>

Reference: YCRAD 6282

To appear in: *Clinical Radiology*

Received Date: 29 November 2022

Revised Date: 7 April 2023

Accepted Date: 23 April 2023

Please cite this article as: Moore N, Maher M, Murphy G, O'Callaghan Maher M, O'Connor OJ, McEntee MF, CT in the detection of latent tuberculosis: a systematic review, *Clinical Radiology*, <https://doi.org/10.1016/j.crad.2023.04.014>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2023 Published by Elsevier Ltd on behalf of The Royal College of Radiologists.

Purpose:

The accurate detection of latent tuberculosis can prevent the disease progressing to active tuberculosis. Currently there is no gold standard for the diagnosis of latent TB. Diagnosis is commonly based on a combination of clinical history, immune response to antigens and radiological assessments. Chest x-rays are most commonly used, but lack diagnostic accuracy. CT is an alternative but is associated with a higher radiation dose. This review aims to evaluate the use of CT and low dose CT in the detection of latent TB.

Method:

A systematic search of literature in adherence with the PRISMA guidelines was carried out. Quality assessment of the included studies was conducted.

Results:

The search strategy identified a total of 4621 studies. Sixteen studies were considered eligible and included in the review. There was high heterogeneity among all studies. CT was identified as much more sensitive for the detection of latent TB in all studies despite chest x-ray often being recommended in guidelines to assess patients for latent TB. Low Dose CT showed promising results in four of the studies, however, these results were limited due to small sample sizes.

Conclusion

CT is much superior to chest x-ray consistently identifying additional cases of latent TB. There are limited high-quality publications available using low dose CT but findings thus far suggest low dose CT could be used as an alternative to standard-dose CT for the detection of latent TB. It is recommended that a randomised control trial investigating Low Dose CT should be carried out.

CT in the detection of latent tuberculosis: a systematic review

N. Moore^{a,*}, M. Maher^{b,c}, G. Murphy^d, M. O'Callaghan Maher^b, O. J. O'Connor^{b,c}, M. F. McEntee^a

^aMedical Imaging and Radiation Therapy, University College Cork, Ireland

^bDepartment of Radiology, University College Cork, Cork Ireland

^cDepartment of Radiology, Cork University Hospital, Wilton, Cork, Ireland

^dDepartment of Rheumatology, Cork University Hospital, Cork, Ireland

*Guarantor and correspondent: N. Moore, College of Medicine and Health, University College Cork, Cork, Ireland. Tel.: 00353 868599976

E-mail address: niamh.moore@ucc.ie

ABSTRACT

AIM: To evaluate the use of computed tomography (CT) and low-dose CT in the detection of latent tuberculosis (TB).

MATERIALS AND METHODS: A systematic search of literature in adherence with the PRISMA guidelines was carried out. Quality assessment of the included studies was conducted.

RESULTS: The search strategy identified a total of 4,621 studies. Sixteen studies were considered eligible and included in the review. There was high heterogeneity among all studies. CT was identified as much more sensitive for the detection of latent TB in all studies despite chest radiography often being recommended in guidelines to

assess patients for latent TB. Low-dose CT showed promising results in four of the studies; however, these results were limited due to small sample sizes.

CONCLUSION: CT is much superior to chest radiography consistently identifying additional cases of latent TB. There are limited high-quality publications available using low-dose CT but findings thus far suggest low-dose CT could be used as an alternative to standard-dose CT for the detection of latent TB. It is recommended that a randomised controlled trial investigating low-dose CT should be carried out.

INTRODUCTION

Tuberculosis (TB) continues to be one of the top 10 causes of death worldwide [1], caused by *Mycobacterium tuberculosis*. Latent tuberculosis infection (LTBI) is an immune response against *M. tuberculosis* without clinical manifestations or radiological evidence of active TB [1][2]. Approximately 22.5% of the global population (1.78 billion) are said to have LTBI [3][4]. This chronic inactive infection is controlled by the cellular immunity of the infected person. Detection of LTBI is crucial as the risk is that it might become active TB. This risk is heightened in particular groups, including young children, immunodeficient patients, individuals with a high-risk of exposure, recently infected individuals, and intravenous drug users [5][6] [7][8]. LTBI is a term that includes latent TB and previous (inactive)TB.

Although the incidence of active TB has declined recently in most high-income countries, the prevalence of LTBI has remained stable [2]. Despite the historic decline, the incidence of active TB has reversed in the UK in the last 30 years and has become an area of concern for public health with the increase predominantly among the foreign born people in the UK [9]. Appropriate screening therefore remains important because

of the rising number of people travelling from high endemic areas and also because of the increasing use of immunosuppressive therapies[10]. The magnitude of the reactivation of TB risk varies among high-risk categories; however, early diagnosis of LTBI may prevent the development of active TB [11][12]. A systematic approach to managing LTBI in cohorts at high-risk reactivation is a critical component of the World Health Organization (WHO) End TB Strategy [13][14]. Consequently, many countries have introduced LTBI control programmes targeting high-risk groups and screening to identify those with LTBI as well as TB [15]. Nonetheless, the WHO has expressed concerns over the effect of COVID-19 on TB prevention control programmes, noting a sharp decrease in TB notifications in 2020[1]. Regardless, there is consensus that screening for LTBI is indicated in high-risk groups but little consensus on how this screening is done.

There is no reference standard test for the diagnosis of LTBI. The diagnosis is commonly based on immune response against *M. tuberculosis* antigens using the tuberculin skin test (TST) or the more recent *M. tuberculosis*-specific interferon-gamma release assays (IGRA)[16][13] commercially available as Quantiferon-TB (Qiagen, Hilden, Germany) and T.Spot TB (Oxford Immunotec, Abington, UK); however, neither can differentiate between active TB and LTBI, which is problematic for establishing appropriate treatments[17][8]. Consequently, differentiation often relies on additional radiological assessments to increase diagnostic sensitivity.

Clinical practice guidelines regarding LTBI are inconsistent in their recommendations[11]. The National Institute for Health and Care Excellence (NICE) guidelines recommend carrying out TST followed by IGRA and additional work-up with

chest radiography (CR)[17]. The findings of each test dictating the next test required. All of the international guidelines deem an individual at risk if either TST or IGRA are positive[18]. CR is usually recommended as part of the screening process in addition to immunological tests[5][14][13][11]. Despite this, CR lacks diagnostic accuracy with limited sensitivity for the detection of LTBI[11][19]. A recent meta-analysis reported that the sensitivity of a CR for detection of LTBI is as low as 15%[20]. CR has limited value for LTBI diagnosis in high-risk individuals[21] and in patients with other underlying diseases[22]. Therefore, there is a risk of undertreating TB due to the insensitivity of a CR and overtreating LTBI due to the lack of specificity of TST and IGRA.

Chest CT is more accurate than CR[3], but is associated with much higher radiation doses. CT is often considered a supplementary technique to CR for screening high-risk individuals[23] and is sometimes used for serial screening in high endemic regions[24][25]. Given the associated high radiation doses, there has been a range of dose-reduction strategies trialled. The most successful are iterative reconstruction algorithms, which allow substantial reductions in radiation dose by removing image noise and artefacts [26]. These have led to the development of low-dose (LDCT) or ultra-low-dose CT (ULDCT) protocols, which are already accepted in clinical practice for lung nodule follow-up and in lung cancer screening programmes[27]. Low radiation doses in the thorax have been reported to deliver doses similar to conventional chest radiography, which are 0.05–0.24 mSv[27].

The primary aim of this review was to evaluate the use of CT to detect radiological abnormalities suggestive of LTBI and evaluate its use as part of a screening protocol.

A secondary aim is to identify if LDCT could be used as an alternative to standard-dose CT to do this.

MATERIALS AND METHODS

As a systematic review was carried out no ethical approval was required. A systematic search of the literature was conducted in adherence to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses guidelines for reporting systematic reviews [28]. The Population, Intervention, Comparison, Outcomes and Study design tool[29] was the framework used to formulate the eligibility criteria (Table 1). The search terms “latent tuberculosis” and “computed tomography” and “screening” and “biologic therapies” with their synonyms, acronyms, and spelling alternatives were generated.

An extensive literature search was performed using PubMed, Scopus, Web of Science and Embase databases to identify relevant English language literature. Over 10 years from 1 September 2011 to 1 September 2021. This timeframe was considered appropriate due to the technical developments in multidetector CT.

Selection process

All identified records were screened on title and abstract, the remaining studies were assessed for eligibility. Full texts were obtained for abstracts that met the selection criteria. Reference lists of retrieved studies were also reviewed for relevant studies.

Quality assessment

The quality assessment of each of the included studies was conducted using a Critical Appraisal Skills Programme (CASP)[30]. When the validity of results, clarity of results, and the clinical relevance of results were absent or unclear, they were excluded from the review. Internal validity of the studies was assessed using the Quality Assessment of Diagnostic Accuracy Studies-2 tool (QUADAS-2)[31] to determine bias in four critical domains patient selection, index test(s), reference standard, and flow and timing. For this review, any study containing high or unclear risk of bias in all four domains was eliminated.

A data extraction form was developed and was used on studies that met the reviewers' inclusion criteria. Data extracted included study design, clinical history, diagnostic methods, radiological imaging for LTBI diagnosis. These headings were used for thematic analysis (Electronic Supplementary Material Table S1).

RESULTS

The Preferred Reporting Items for Systematic Reviews and Meta-Analyses flowchart (Fig.1) summarises the search results and the review process. The search strategy identified 4,621 studies, 4,385 after removal of duplicates. The titles and abstracts were screened, and 4,266 were removed. A full-text review was completed on the remaining 119 studies. Sixteen of these studies were considered eligible for inclusion. These 16 potential studies were appraised for inclusion using the modified QUADAS-2 ROB tool[31], and none were considered "high risk" or "unclear risk" in all of the domains (Table 2). The 16 studies were also appraised using the CASP diagnostic checklist[30], which did not identify any low-quality studies (Table 2).

All included studies evaluated high-risk individuals. There was high heterogeneity in the literature with variations in the combination of diagnostic tools used to diagnose LTBI. There was variation in the groups of participants investigated between high-risk participants undergoing immunosuppressive therapy, close contact with a person with TB, and participants working in high endemic regions. The retrospective studies, in particular, were often ambiguous on clinical histories such as BCG (*Bacillus Calmette–Guérin*) vaccination and also lacked detail on the diagnostic methods being reviewed, e.g., methodology (Electronic Supplementary Material Table S1). Variation exists across all studies between the use of TST, IGRA, CR, and CT for the diagnosis of LTBI (Table 2). The criteria implemented to determine positivity for each test within a screening protocol varied in many of the studies included. These variations included radiological appearances of LTBI and the cut-off points to determine immunological test positivity.

Radiological imaging for the diagnosis of LTBI

CT was used as the reference standard in only one study [32]. Four publications [21][24][32][22] included CT as part of a screening protocol for all patients and three used CT for most of its participants. Tannus et al. carried out CT in 34/37 participants[33], Lee et al. used LDCT in six out of seven participants[25] and 104/107 participants in Allwood et al. underwent both CT and CR[34] as the authors attempted to identify the optimal approach between structured questionnaires, CT, and CR for ruling out previous pulmonary TB. The remaining nine studies [6][8][35][36][37][38][39][40][41] reported CT as either an additional screening tool, as part of routine practice[6], or for a reason other than screening for LTBI, e.g., metastatic work-up[39].

The typical radiological appearances suggestive of LTBI on CT (Table 3) varied in the literature, with many listing pleural thickening, interstitial granuloma, fibrotic scarring, nodules, or lymphadenopathy as the predominant appearances [6][21][22][24][33][32][34][36][37][38][39][40]. Examples of some typical appearances can be seen in Fig. 2. The remaining four studies[8][25][35][41] provided no detail on radiological appearances. In combination with results from other tests some reported CT positive for LTBI [8][41] when no radiological signs of active TB were classified. Yoon et al. employed serial LDCT findings to determine diagnosis in patients with some non-calcified nodules disappearing in two contacts[40].

CRs were not included as part of the screening protocol in six of the studies [24][25][32][33][38][40] while seven[8][21][22][35][36][41][39] carried out CR on all participants. In the remaining three studies, CRs were carried out in addition to or instead of CT[6][34][37]. Seven studies compared CT and CR, with CT frequently identifying findings suggestive of LTBI, which were not identified by CR [6][8][21][22][39][34][35] (Electronic Supplementary Material Table S1). Examples of this can be seen in Fig 3. Tannus-Silva et al. found that CT identified a higher number of positive results (52.9% positivity) than immunological tests demonstrating 36.8% positivity for T.SPOT-TB. and 13.5% for TST[33].

Low-dose CT

Four of the 16 studies used LDCT for the diagnosis of LTBI [34][24][40][25]. The largest of these studies by He et al. provided insufficient detail to confirm whether the acquisition parameters that were used equated to low dose. The exposure parameters

tube voltage and tube current were provided but tube rotation time was omitted[24]. No dose estimations were provided.

Allwood et al. describe using LDCT but provided no detail on imaging parameters[34]. Lee et al.[25] and Yoon et al.[40] did provide a detailed breakdown of parameters and protocols used, and they estimated effective doses per CT scan of 0.19–0.25 mSv and 0.13 ± 0.03 mSv, respectively. Both of these studies were comparable in the dose estimates as they used a similar conversion factor of 0.0145 to estimate effective doses. A limitation of these studies however is the small number of participants ($n=6$)[25] and ($n=17$)[40] who underwent LDCT.

Screening for LTBI

Seven of the studies investigated screening among close contacts and healthcare workers [8][36][24][40][25][41][35]. Yoshiyama et al.[35] followed-up screening with participants for 2 years. In the remaining studies [8][36][24][40][25][41] participants were either not followed up for 2 years or it was unclear.

He et al. recommended replacing CR with CT as a cost-effective option in screening for TB[24]. Allwood et al.[34] felt CT was financially justified in certain circumstances such as TB vaccine research, in advance of commencing immunosuppressive therapy, pre-transplant, and in cases where there was a risk of reactivation; however, Saidenberg-Kermanac'h et al. argued this would not be financially justified due to only one out of 60 patients in their study diagnosed with LTBI using CT[21].

DISCUSSION

This systematic review demonstrates that CT is useful for detecting radiological abnormalities suggestive of LTBI and justifies its use as part of a screening protocol.

There are limited high-quality publications available using LDCT for the detection of LTBI but findings thus far suggest LDCT could be used as an alternative to standard-dose CT for the detection of LTBI. It has also been recognised that there are substantial methodological limitations in this review with some studies limited by including only a small number of participants, and in addition, most studies either did not use a reference standard or CT was itself part of the reference standard. Some of the studies patients were selected to undergo CT based on the results of other tests, which were themselves part of a reference standard. This however is a limitation of the absence of a true reference standard for the diagnosis of LTBI. A more appropriate study design would be to compare LDCT, CR, immunological tests, and the patients' backgrounds.

It is important to recognise that it is not merely a matter of the sensitivity of CT for detecting lung lesions in LTBI, but that complex questions exist around screening for LTBI[8]. The only way to improve the sensitivity is to implement a screening protocol with tests demonstrated to be useful[23]. All studies in this review included a combination of tests or the development of active TB as the reference standard except for Maeda et al.[32], who used CT as the reference standard. As CT is not 100% sensitive[42], this study[32] may have classified TB disease incorrectly, potentially resulting in inaccurate findings. The other studies demonstrated great variation in the accuracy of TST, IGRA, and CR for LTBI diagnosis. Due to the retrospective nature of half of the studies reviewed, selection bias cannot be excluded as CT was generally carried out on individuals with a higher risk of having the disease.

Many CT examinations in the review were interpreted by only one radiologist; however, CT as part of a protocol is prone to reader variability, as highlighted by Yoshiyama et al.[35]. They re-evaluated previous CT findings in 205 contacts, which revealed four contacts who had been judged as having LTBI initially but were later judged as having active TB at re-evaluation. Allwood et al. identified that low inter-reader agreement often limits subjective radiological evaluation[34]. Given the low sensitivity and specificity of CT in some studies, consideration should be given to the interpretation process of CT examinations. Six of the included studies blinded the assessors to the results of other tests [8][21][33][39][34][24] (Electronic Supplementary Material Table S1), which may have disadvantaged them compared to standard practice where they are presented with all of the clinical information to make a report. Seven of the studies did not provide any detail on the blinding of assessors [32][37][6][22][40][25][41].

It was also apparent that radiographic appearances suggestive of LTBI differed between studies (Electronic Supplementary Material Table S1). CT appearances depend on the stage of TB described, with LTBI often detected by another diagnostic method but confirmed by the absence of radiological features indicative of active disease[8][41][35]. Calcified granulomas or lymph nodes indicate LTBI when they are the only radiographic findings. Inactive TB appears radiographically as stable fibronodular changes, scarring, and nodular opacities in apical and upper lung regions [3][24]. Broadly the literature defined LTBI as a blanket term without differentiating latent and inactive TB. Variations in radiographic appearances and reader variability are important to consider as this could modify the accuracy of CT causing a conflation of CT findings for the diagnosis of LTBI.

The Centres for Disease Control and Prevention have recommended that all healthcare workers have baseline TB screening[43] and those with known exposure or with evidence of ongoing TB transmission should undergo serial TB screening[43]. TB granuloma is the greatest indicator of primary and post-primary TB[44]. An indication of progression to active disease is the progression of parenchymal abnormalities on follow-up[25]. Individuals who have been in contact with active TB within 2 years are considered high-risk for the progression from LTBI to active TB[12][39]; however, in six of the studies it was either unclear or participants were not followed up for 2 years[8][24][36][25][40][41]. The stability of imaging findings in a screening protocol is a key consideration, highlighting the benefit of serial CT. Yoon et al. observed radiological changes in their repeated LDCTs. Healthcare workers are at high risk of TB[45], He et al. recommended annual LDCT examinations screening healthcare workers working in TB specialist hospitals[24].

CT has superior sensitivity to CR for detecting minimal changes in lung diseases and helps differentiate active and inactive TB disease [24][46][47][48]. As far back as 1996, Lee et al. demonstrated that CT can effectively detect and determine disease activity in 80% of those with active TB and 89% of those with LTBI[47]. This review revealed many CT lesions associated with past TB that were not identified on CR[6][8][21][22][39][34][35][48][49]. Uzorka et al. identified lesions on CT in 70% of participants compared to 22.7% revealed by CR[6]. Using only CT, Maeda et al. classified past infection and non-infection in a group of rheumatic patients[32]. The addition of CT increases the sensitivity and specificity of TB detection and is a useful additional imaging method for the study of LTBI [24][25]. No diagnostic test used to diagnose

LTBI is 100% sensitive, but sensitivity improves significantly with CT used as part of a screening protocol[35]. Despite these findings, CT is not routinely recommended in international guidelines due to the associated radiation doses.

Radiation doses in CT can be significantly reduced with LDCT or ULDCT, and although LDCT/ULDCT are used to enable the detection of subtle lung disease in common practice they are not ubiquitous. Traditionally, there is a lack of definition between LDCT and ULDCT, which is compounded in this review due to a lack of detail on image acquisition parameters and radiation doses. LDCT or ULDCT can significantly reduce doses compared to conventional-dose CT and, if shown to have equal diagnostic accuracy, could be used for the screening of LTBI. Three of the included studies demonstrated promising results using LDCT to screen for LTBI[24][25][40] despite some lacking detail for reproducible protocols. LDCT/ULDCT appearances suggestive of LTBI identified in corroboration with other screening methods should be implemented to provide an appropriate diagnosis with minimal risk of disease progression.

Like all of the other diagnostic tests used for diagnosing LTBI, CT has suboptimal sensitivity and may fail to identify all patients at risk of reactivation; however, it remains superior to a CR for identifying possible subclinical or latent disease, in order to appropriately refer patients for suitable microbiologic testing. Given its increased sensitivity, the introduction of CT, in particular LDCT, should be considered as a supplementary screening method instead of CR with the possibility of increased sensitivity for the detection of LTBI disease in patients with a positive immunological

test, in particular in patients where the prior probability of asymptomatic TB disease is higher.

In conclusion, given the absence of a reference standard, CT is a corroborative method for diagnosing LTBI. CT will detect many lesions ~~which~~ that may be due to LTBI and should be used as part of the diagnostic work-up in high-risk patients with positive immunological tests in advance of commencing immunosuppressive therapies.

As the sensitivity of CT is much greater than CR, the efficacy of LDCT needs to be considered. This work demonstrates the need for a high-quality prospective clinical trial comparing LDCT/ULDCT to CR to determine its accuracy and associated radiation doses in high-risk individuals. With the recommendation for CT comes the necessity to balance true treatment, overtreatment, and harm of radiation, and further work should investigate this. CT is often considered financially expensive; however, the findings from CT often avoid costly hospital stays or interventional surgeries, both of which increase the risks of iatrogenic errors[50]. Given the differing opinions in relation to cost effectiveness between three of the studies, [21][24][34] cost and risk versus effectiveness also should be monitored and reviewed[51].

REFERENCES

- [1] WHO. Global Tuberculosis Report 2020, vol. XLIX, no. 9-10–11. Available at <https://www.who.int/publications/i/item/9789240013131> (accessed ???).
- [2] Zellweger JP, Sotgiu G, Corradi M, Durando P. The diagnosis of latent tuberculosis infection (Ltbi): currently available tests, future developments, and

perspectives to eliminate tuberculosis (TB). *Med Lav.* 2020;111:170-183. doi: 10.23749/mdl.v111i3.9983.

[3] Nachiappan A, Rahbar K, Shi X et al. Pulmonary tuberculosis: role of radiology in diagnosis and management. *RadioGraphics*, 2017;37:52–72. doi: 10.1148/rg.2017160032.

[4] Ye I, Chapman T, Wen Z et al. Targeted versus universal tuberculosis chemoprophylaxis in 1968 patients with inflammatory bowel disease receiving anti-TNF therapy in a tuberculosis endemic region. *Aliment. Pharmacol. Ther.*, 2021;53:390–399.

[5] Centres for Disease Control and Prevention. Testing for tuberculosis (TB) Fact Sheet. Available at: https://www.cdc.gov/tb/publications/factsheets/testing/tb_testing.htm (accessed ???).

[6] Uzorka J, Duinker D, Kroft L et al. Trends in diagnostic methods and treatment of latent tuberculosis infection in a tertiary care center from 2000 to 2017. *Eur. J. Clin. Microbiol. Infect. Dis.*, 2020;39:1329–1337. doi: 10.1007/s10096-020-03850-7.

[7] Auguste P, Madan J, Tsertsvadze A et al. Identifying latent tuberculosis in high-risk populations: systematic review and meta-analysis of test accuracy. *Int. J. Tuberc. Lung Dis.*, 2019;23:1178–1190, 2019. doi: 10.5588/ijtld.18.0743.

[8] Fujikawa A, Fujii T, Mimura S et al. Tuberculosis contact investigation using interferon-gamma release assay with chest X-ray and computed tomography. *PLoS One* 2014;9:e85612. doi: 10.1371/journal.pone.0085612.

[9] Lalvani A, Pareek M, 'Immigrant screening for TB: A missed opportunity to improve TB control in the United Kingdom', *Pathog. Glob. Health*, 2012;106:5–7. doi: 10.1179/204777312X13305103762501.

[10] Uzorka J, Kroft L, Bakker J et al. Abnormalities suggestive of latent tuberculosis

infection on chest radiography; how specific are they? *J. Clin. Tuberc. Other Mycobact. Dis.*, 2019;15:100089. doi: 10.1016/j.jctube.2019.01.004.

[11] Hasan T, Au E, Chen S, Tong A, Wong G. Screening and prevention for latent tuberculosis in immunosuppressed patients at risk for tuberculosis: a systematic review of clinical practice guidelines. *BMJ Open*, 2018;8:22445. doi: 10.1136/bmjopen-2018-022445.

[12] Lee H, Lee Y, Kim S et al. Comparing tuberculin skin test and interferon γ release assay (T-SPOT.TB) to diagnose latent tuberculosis infection in household contacts', *Korean J. Intern. Med.*, 2017;32:486–496. doi: 10.3904/kjim.2015.344.

[13] World Health Organisation. Latent tuberculosis infection. Updated and consolidated guidelines for programmatic management. 2018. Available at: <https://apps.who.int/iris/handle/10665/260233> (accessed ???).

[14] National TB Advisory Committee. Guidelines on the Prevention and Control of Tuberculosis in Ireland, 2010. Amended 2014. <https://www.hpsc.ie/a-z/vaccinepreventable/tuberculosis/tb/guidance/tbguidelines2010amended2014/File,4349,en.pdf> (accessed ???).

[15] Erkens C, Kamphorst M, Abubakar et al. Tuberculosis contact investigation in low prevalence countries: a European consensus. *Eur. Respir. J.*, 2010;36:925–949. doi: 10.1183/09031936.00201609.

[16] Lenaerts A, Barry C, and Dartois V. Heterogeneity in tuberculosis pathology, microenvironments and therapeutic responses. *Immunol. Rev.*, 2015;264:288–307. doi: 10.1111/imr.12252.

[17] NICE. Tuberculosis. NICE guideline [NG33]. Published: 13 January 2016. Available at: www.nice.org.uk/guidance/ng33 (accessed ???).

[18] World Health Organization. Guidelines on the management of latent

tuberculosis

infection.

Available:

http://www.who.int/tb/publications/lbtbi_document_page/en/ (accessed ???).

- [19] Esmail H, Lai R, Lesosky M et al. Characterization of progressive HIV-associated tuberculosis using 2-deoxy-2-[(18)F]fluoro-D-glucose positron emission and computed tomography. *Nat. Med.*, 2016;22:1090–1093. doi: 10.1038/nm.4161.
- [20] Uzorka J, Wallinga J, Kroft L, Ottenhoff T, Arend S. Radiological signs of latent tuberculosis on chest radiography: a systematic review and meta-analysis. *Open Forum Infect. Dis.*, 2019;6:ofz313. doi: 10.1093/ofid/ofz313.
- [21] Saidenberg-Kermanac'h N, Semerano L, Naccache J et al. Screening for latent tuberculosis in anti-TNF- α candidate patients in a high tuberculosis incidence setting. *Int. J. Tuberc. Lung Dis.*, 2012;16:1307–1314. doi: 10.5588/ijtld.12.0111.
- [22] Guirao-Arrabal E, Santos F, Redel-Montero J et al. Risk of tuberculosis after lung transplantation: the value of pretransplant chest computed tomography and the impact of mTOR inhibitors and azathioprine use. *Transpl. Infect. Dis.*, 2016;18:512–519. doi: 10.1111/tid.12555.
- [23] Piccazzo R, Paparo F, Garlaschi G. Diagnostic accuracy of chest radiography for the diagnosis of tuberculosis (tb) and its role in the detection of latent TB infection: a systematic review. *J. Rheumatol.*, 2014;41(Suppl. 91):32–40. doi: 10.3899/jrheum.140100.
- [24] He W, Chen B, Lv Y et al. Use of low-dose computed tomography to assess pulmonary tuberculosis among healthcare workers in a tuberculosis hospital. *Infect. Dis. Poverty*, 2017;6:68. doi: 10.1186/s40249-017-0274-6.
- [25] Lee S, Yoon S, Goo J, Yim J, and Kim C. Submillisievert computed tomography of the chest in contact investigation for drug-resistant tuberculosis. *J. Korean Med. Sci.*, 2017;32:1779–1783. doi: 10.3346/jkms.2017.32.11.1779.

- [26] Kim Y, Kim Y, Lee B et al. Ultra-low-dose CT of the thorax using iterative reconstruction: evaluation of image quality and radiation dose reduction. *AJR Am. J. Roentgenol.*, 2015;204:1197–1202. doi: 10.2214/AJR.14.13629.
- [27] Huber A, Landau J, Ebner L et al. Performance of ultralow-dose CT with iterative reconstruction in lung cancer screening: limiting radiation exposure to the equivalent of conventional chest radiography imaging. *Eur. Radiol.*, 2016;26:3643–3652. doi: 10.1007/s00330-015-4192-3.
- [28] Moher D, Liberati A, Tetzlaff J, Altman D. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med.*, 2009;6:1000097. doi: 10.1371/journal.pmed.1000097.
- [29] Methley A, Campbell S, Chew-Graham C, McNally R, Cheraghi-Sohi S. PICO, PICOS and SPIDER: A comparison study of specificity and sensitivity in three search tools for qualitative systematic reviews. *BMC Health Serv. Res.*, 2014;14: 579. doi: 10.1186/s12913-014-0579-0.
- [30] CASP. Critical Appraisal Skills Programme (CASP).
- [31] Whiting P, Rutjes A, Westwood M et al. Tailoring QUADAS-2: a revised tool for the Quality Assessment of Diagnostic Accuracy Studies. *Ann Intern Med* 2011;155:529-536
- [32] T. Maeda *et al.*. Comparison of QuantiFERON-TB Gold and the tuberculin skin test for detecting previous tuberculosis infection evaluated by chest CT findings in Japanese rheumatoid arthritis patients. *J. Infect. Chemother.*, 2011;17:842–848. doi: 10.1007/s10156-011-0250-1.
- [33] Tannus Silva DG, Silva BD, Torres PP, Santana PJ Jr, Junqueira-Kipnis AP, Rabahi MF. Latent tuberculosis in rheumatoid arthritis: evaluating cellular response and high-resolution computed tomography. *Arch Bronconeumol.* 2012 May;48(5):144-9. doi:

10.1016/j.arbres.2011.12.013.

[34] Allwood BW, Goldin J, Said-Hartley Q, van Zyl-Smit RN, Calligaro G, Esmail A, Beyers N, Bateman ED. Assessment of previous tuberculosis status using questionnaires, chest X-rays and computed tomography scans. *Int J Tuberc Lung Dis*. 2015 Dec;19(12):1435-40. doi: 10.5588/ijtld.14.0992.

[35] Yoshiyama T, Kurosaki A, Ogata H, Sasaki Y, Okumura M. Limited benefit of CT scans in tuberculosis contact tracing. *J Infect Chemother*. 2019 Oct;25(10):764-768. doi: 10.1016/j.jiac.2019.03.023.

[36] T. Targowski, S. Chelstowska, T. Plusa. IGRA as a predictive factor of silent pulmonary changes in individuals following exposure to tuberculosis. *Lung*, 2014;192:869–874. doi: 10.1007/s00408-014-9637-y.

[37] Agarwal A, Kedia S, Jain S, Gupta V, Bopanna S, Yadav DP, Goyal S, Mouli VP, Dhingra R, Makharia G, Ahuja V. High risk of tuberculosis during infliximab therapy despite tuberculosis screening in inflammatory bowel disease patients in India. *Intest Res*. 2018 Oct;16(4):588-598. doi: 10.5217/ir.2018.00023.

[38] Song D, Tong J, Peng J et al. Tuberculosis screening using IGRA and chest computed tomography in patients with inflammatory bowel disease: a retrospective study. *J. Dig. Dis.*, 2017;18:23–30. doi: 10.1111/1751-2980.12437.

[39] Lyu J, Lee S, Hwang S et al. Chest computed tomography is more likely to show latent tuberculosis foci than simple chest radiography in liver transplant candidates. *Liver Transplant.*, 2011;17:963–968, 2011. doi: 10.1002/lt.22319.

[40] Yoon S, Mihn D, Song J, Kim S, Yim J. Evolution of interferon-gamma release assay results and submillisievert chest CT findings among close contacts of active

pulmonary tuberculosis patients. *Tuberc. Respir. Dis. (Seoul)*, 2020;83:283–288. doi: 10.4046/trd.2020.0038.

[41] Hirama T, Hagiwara K, Kanazawa M. Tuberculosis screening programme using the QuantiFERON®-TB Gold test and chest computed tomography for healthcare workers accidentally exposed to patients with tuberculosis. *J. Hosp. Infect.*, 2011;77:257–262. doi: 10.1016/j.jhin.2010.11.012.

[42] Lew W, Jung Y, Song J et al. Combined use of QuantiFERON®-TB gold assay and chest computed tomography in a tuberculosis outbreak. *Int. J. Tuberc. Lung Dis.*, 2009;13:633–639.

[43] Sosa L, Nije G, Lobato M et al. Tuberculosis screening, testing, treatment of U.S. health care personnel: recommendations from the National Tuberculosis Controllers Association and CDC, 2019. *Am. J. Transplant.*, 2019;19:2383–2387. doi: 10.1111/ajt.15512.

[44] Hunter R. The pathogenesis of tuberculosis: the early infiltrate of post-primary (adult pulmonary) tuberculosis: a distinct disease entity. *Front. Immunol.*, 2018;9:1–9. doi: 10.3389/fimmu.2018.02108.

[45] Baussano J, Nunn P, Williams B, Pivetta E, Bugiani M, Scano F. Tuberculosis among health care workers. *Emerg. Infect. Dis.*, 2011;17:488–494. doi: 10.3201/eid1703.100947.

[46] Pamuk Ö, Yesil Y, Donmez S, Ünlü E, Köker I, Cakir N. The results of purified protein derivative test in ankylosing spondylitis patients: clinical features, HRCT results and relationship with TNF-blocker usage. *Rheumatol. Int.*, 2008;29:179–183. doi: 10.1007/s00296-008-0665-x.

[47] Lee K, Hwang J, Chung M, Kim H, Kwon O. Utility of CT in the evaluation of pulmonary tuberculosis in patients without AIDS. *Chest*, 1996;110:977–984. doi:

10.1378/chest.110.4.977.

[48] Lee S, Jang Y, Park C et al. The role of chest CT scanning in TB outbreak investigation.. *Chest*, 2010;137:1057–1064. doi: 10.1378/chest.09-1513.

[49] Guirao E and Ruiz-cabello M. Diagnostic methods of tuberculosis & Métodos diagnósticos de la tuberculosis. *Med. Clínica* 2020;154:305–307. doi: 10.1016/j.medcle.2019.08.007.

[50] Makary M and Daniel M. Medical error-the third leading cause of death in the US. *BMJ*, 2016;353:1–5. doi: 10.1136/bmj.i2139.

[51] Park J, Kim N, Park H et al. The use of a borderline zone for the interpretation of interferon-gamma release assay results for serial screening of healthcare workers. *PLoS One*, 2020;15:1–12. doi: 10.1371/journal.pone.0235254.

Figure 1. Preferred Reporting Items for Systematic Reviews and Meta-Analyses guidelines for reporting systematic Flowchart

Figure 2. Plain radiographic and CT features of LTBI. (a) Chest radiograph shows right apical fibrotic scarring, fibronodular change, and volume loss, with questionable upper lobe bronchiectasis (black arrow). There are left upper lobe granulomata (white arrows). (b) Conventional-dose chest CT more clearly demonstrates volume loss in the right upper lobe with cicatricial bronchiectasis (black arrow). Calcified granulomata are seen more clearly in the left upper lobe.

Figure 3. LDCT has increased sensitivity for calcified granulomata. This is a non-specific finding but can be useful in the correct clinical context. (a) Normal chest

radiograph]. Calcified granuloma not seen. (b) Unenhanced LDCT of the chest (DLP 6.0 mGy.cm) shows a tiny 1 mm left upper lobe calcified granuloma (white arrow).

Table 1. PICOS elements for eligibility criteria

Population	All studies reporting diagnosis or changes on CT suggestive of LTBI, which include individuals ≥ 16 years of age, with suspected latent TB infection and have had an IGRA/TST and/or a chest radiography performed All studies which include individuals ≥ 16 years of age who are undergoing screening for latent TB and have had a CT thorax and an IGRA/TST and/or a chest radiography performed
Intervention	CT/low-dose CT/ultra-low-dose CT
Comparison	CR and IGRA/TST or IGRA/TST or CR
Outcomes	Detection rate of LTBI using CT or LDCT or ULDCT, detection rate of radiological abnormalities suggestive of LTBI using CT or LDCT or ULDCT
Study Design	All studies with the exception of case reports and case studies

CT, computed tomography; LTBI, latent tuberculosis infection; CR, chest radiography; IGRA, (*M. tuberculosis*-specific) interferon-gamma release assay; TST, tuberculin skin test specific LDCT, low-dose CT; ULDCT, ultra-low dose CT.

Table 2. QUADAS 2 tool, ROB tool and CASP diagnostic checklist and methods of assessment for the detection of latent tuberculosis infection (LTBI)

QUADAS 2: risk of bias					CASP: quality assessment			Methods of assessment
Study name	Patient selection	Index test	Reference standard	Flow and timing	Are the results of the Study Valid?	What are the results?	Will the results help locally?	LTBI diagnostic methods/criteria LTBI
Agarwal et al 2018	✓	?	?	?	?	?	Yes	Patient history + TST/IGRA/CXR/CT
Fujikawa et al 2014	✓	✓	?	?	Yes	Yes	Yes	Patient history + IGRA + CXR ± CT
Allwood et al 2015	✓	✓	X	?	No	Yes	Yes	2 Questionnaires on patient history + CXR + hi-res LDCT
Maeda et al 2011	✓	?	?	✓	Yes	?	Yes	Patient history + CT ± TST/IGRA
Uzorka et al 2020	✓	?	?	X	No	Yes	Yes	Patient history/TST/IGRA ± CXR ± CT
Guirao Arrabal et al 2016	✓	✓	X	X	No	Yes	Yes	Patient history + CT + CXR ± TST
Lyu et al 2011	?	✓	?	X	?	?	Yes	Patient history + CXR ± CT
He et al 2017	✓	✓	?	?	?	?	Yes	Patient history + LDCT
Lee et al 2017	✓	✓	?	✓	Yes	No	Yes	Patient history + CXR + TST ± IGRA + ULDCT
Hirama et al 2011	✓	?	?	?	Yes	No	Yes	Patient history + IGRA + CXR, ± CT
Targowski et al 2014	✓	✓	?	✓	Yes	Yes	Yes	Patient history + CXR + TST + IGRA ± CT


Yoshiyama et al 2019	✓	X	✓	✓	?	No	Yes	Patient history + IGRA + CXR ± CT
Song et al 2017	✓	?	?	?	Yes	Yes	Yes	Patient history + IGRA ± CT
Yoon et al 2020	✓	✓	X	?	Yes	?	Yes	Patient history + TSPOT.TB + QFT-GIT + LDCT chest
Tannus Silva et al 2012	✓	✓	?	?	Yes	?	Yes	Patient history + TST ± IGRA ± hi-res CT chest
Saidenberg-Kermanac'h et al 2012	✓	✓	?	?	Yes	Yes	Yes	Patient history + CXR + TST ± IGRA ± CT


Interferon-gamma release assays (IGRA) are available commercially as QuantiFERON-TB (Gold In tube, QFT-GIT and Gold Plus, QFT-Plus; Qiagen, Hilden, Germany) and T.Spot TB (Oxford Immunotec, Abington, UK)

✓, Low risk; X, high risk;?, unclear; TST, tuberculin skin test; IGRA, *M. tuberculosis*-specific interferon-gamma release assay; CXR, chest radiography; CT, computed tomography; hi-res, high resolution; LDCT, low dose CT.


Table 3. Typical computed tomography (CT) appearances suggestive of latent tuberculosis (TB) infection


CT appearances suggestive of latent TB
Granuloma
Calcified lymph nodes
Stable fibronodular changes
Fibrotic scarring
Calcified nodules in the apical and upper regions
Pleural thickening











Highlights

- Chest x-rays lack sensitivity for the detection of latent TB
- Chest CT can help to discriminate between latent and active TB
- CT is far superior to chest x-ray to identify all patients at risk of latent TB re-activation
- Low dose CT should be considered as a supplementary modality for detecting latent TB

Declaration of interests

☐ The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

☒ The authors declare the following financial interests/personal relationships which may be considered as potential competing interests:

Niamh Moore reports was provided by University College Cork.
--