


|                             |  |
|-----------------------------|--|
| Title                       | Role of the Photorhabdus Dam methyltransferase during interactions with its invertebrate hosts   |
| Authors                     | Payelleville, Amaury;Blackburn, Dana;Lanois, Anne;Pagès, Sylvie;Cambon, Marine C.;Ginibre, Nadege;Clarke, David J.;Givaudan, Alain;Brillard, Julien  |
| Publication date            | 2019-10-09   |
| Original Citation           | Payelleville, A., Blackburn, D., Lanois, A., Pagès, S., Cambon, M. C., Ginibre, N., Clarke, D. J., Givaudan, A. and Brillard, J. [2019] 'Role of the Photorhabdus Dam methyltransferase during interactions with its invertebrate hosts', PLOS ONE, 14(10), e0212655. (14pp.) doi: 10.1371/journal.pone.0212655  |
| Type of publication         | Article (peer-reviewed)  |
| Link to publisher's version | 10.1371/journal.pone.0212655   |
| Rights                      | ©2019 Payelleville et al. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.<br>- <a href="http://creativecommons.org/licenses/by/4.0/">http://creativecommons.org/licenses/by/4.0/</a> |
| Download date               | 2024-04-26 02:27:27  |
| Item downloaded from        | <a href="https://hdl.handle.net/10468/9293">https://hdl.handle.net/10468/9293</a>  |



# UCC

**University College Cork, Ireland**  
Coláiste na hOllscoile Corcaigh

Fig. S1



**Figure S1. Relative expression of the *dam* gene in the *P. luminescens* *dam*-overexpressing strain.** qRT-PCR was carried out with total RNA extracted from 3 independent samples of exponentially growing cells of the *dam*-overexpressing strain and of the control strain. Left panel, Box-plot representing the expression ratio between the two strains is represented for the *dam* gene. Right panel, detailed report of the REST-analysis using with *gyrB* and *fis* used as control genes (see Materials & Methods section for details). The level of expression between the 2 strains was different at  $p<0.001$ .