CORA Cork Open Research Archive
Cartlann Taighde Oscailte Chorcaí

Title	Microplastics in grey seal (Halichoerus grypus) intestines: Are they associated with parasite aggregations?
Authors	Hernandez-Milian, G.;Lusher, A.;MacGabban, S.;Rogan, Emer
Publication date	2019-06-28
Original Citation	Hernandez-Milian, G., Lusher, A., MacGabban, S. and Rogan, E. (2019) 'Microplastics in grey seal (Halichoerus grypus) intestines: are they associated with parasite aggregations?', Marine Pollution Bulletin, 146, pp.349-354. doi: 10.1016/j.marpolbul.2019.06.014
Type of publication	Article (peer-reviewed)
Link to publisher's version	10.1016/j.marpolbul.2019.06.014
Rights	© 2019, Elsevier Ltd. All rights reserved. This manuscript version is made available under the CC BY-NC-ND 4.0 license. - https://creativecommons.org/licenses/by-nc-nd/4.0/
Download date	2026-11-17 14:08:17
Item downloaded from	https://hdl.handle.net/10468/14960

UCC
University College Cork, Ireland
Coláiste na hOllscoile Corcaigh

Accepted Manuscript

This is an Accepted Manuscript of the following article:

G. Hernandez-Milian, A. Lusher, S. MacGabban, E. Rogan.
Microplastics in grey seal (Halichoerus grypus) intestines:
Are they associated with parasite aggregations?
Marine Pollution Bulletin.
Volume 146, 2019, pages 349-354, ISSN 0025-326X.


The article has been published in final form by Elsevier at
http://dx.doi.org/10.1016/j.marpolbul.2019.06.014

© 2019. This manuscript version is made available under the

CC-BY-NC-ND 4.0 license

http://creativecommons.org/licenses/by-nc-nd/4.0/

Graphical Abstract



Highlights

- Knowledge on trophic transfer of microplastics is still scarce
- By caught seals may provide information on microplastic uptake via prey
- All seals had microplastics but no macroplastics were found
- microplastic retention in the intestines may be related to parasite aggregations
- By caught seals are a good source of data to monitor the incidence of microplastic pollution within coastal food webs

*Manuscript (clean - no tracked changes)
Click here to view linked References

Microplastics in grey seal (*Halichoerus grypus*) intestines: are they associated with parasite aggregations?

Hernandez-Milian, G.[1*], Lusher, A[2], MacGabban. S.[1], Rogan, E.[1]

[1*]Aquaculture & Fisheries Development Centre (AFDC), School of Biological, Earth & Environmental Sciences (BEES), University College Cork, Distillery Fields, North Mall, Cork, Ireland

[2]Norwegian Institute of Water Research (NIVA), Gaustadalléen 21, 0349 Oslo, Norway

*corresponding author: ghmilian@gmail.com

ABSTRACT

Between 2012 and 2015, 13 grey seals were recovered from trammel nets targeting monkfish and rays off the south coast of Ireland. Incidence and distribution of microplastics were investigated along the intestines of bycaught seals. No macrodebris items were found, whereas microplastics were detected in all seals. A total of 363 microplastics items were identified (85% fibers, 14% fragments, 1% films). Estimation of microplastic ingestion based on prey ingestion (245 particles) was lower than the observed data. Acantocephala parasites (n=1,543) were found in 12 seals, with an average of 74.5 ± 67.7 parasites per seal. Distribution of microplastics varied between seals, although microplastics tended to accumulate in areas where more parasites were aggregated; however, there was no significant relationship between the number of parasites and microplastics was found. Seals recovered from nets appear to be a good source to monitor the incidence of microplastic pollution within the coastal food webs.

Keywords: grey seals, microplastics, Ireland, parasites, bycatch

INTRODUCTION

Many different forms of marine pollution have been identified, including oil spills, chemicals, and marine debris (e.g. Islam & Tanaka 2004). Over the last few decades, marine debris has emerged as a major threat to the environment and biota. Persistent anthropogenic debris which consists primarily of plastic items is of particular concern (Bergmann et al. 2015). Explosive development of the plastic industry and consumer use has consequently produced a considerable amount of plastic waste (Jambeck et al. 2015). Plastics may ultimately end up in the marine environment, if disposal methods are not appropriate. It is therefore vital to understand the environmental implications of this long-lived pollutant. The importance of this issue is reflected in the implementation of Descriptor 10 of the EU 'Marine Strategy Framework Directive (MSFD)' which monitors marine litter, including plastics, for a 'Good Environmental Status (GES)'. As plastics and other anthropogenic debris move through the marine environment over large areas, they may become dangerous for marine organisms either because debris entering a new ecosystem/habitat could have consequences on the habitat itself, transport species which might behave as invasive organisms or directly impact upon biota themselves. Many species of biota have been reported to interact with, and be affected by, marine debris. Monitoring and detecting marine debris is necessary to understand the implications to different groups of biota (GESAMP 2016; Galgani et al. 2013). Several review publications compiled information on marine debris, and highlighted impacts including entanglement and ingestion by marine mammals which sometimes lead to fatalities (Baulh & Perry 2014, Kühn et al. 2015, CBD 2016). Marine mammals may be affected through direct interaction, or secondary interaction following the ingestion of contaminated prey (e.g. Fossi et al. 2018, Lusher et al. 2018, Panti et al. 2019). It is particularly important to understand the effects of marine debris on megafauna with life histories characterized by slow growth, late maturity, and low fecundity (e.g. Panti et al. 2019).

Microlitter particles are more conspicuous than macrolitter and have many different origins (GESAMP 2016). For instance, plastic items can persist in the environment as large (macrolitter) but over time they are exposed to environmental and mechanical processes which cause them to become brittle and fragment (Andrady 2015). A substantial number of marine species from invertebrates to vertebrates were found to ingest microplastics (e.g. Fossi et al. 2018). Within vertebrates, sea turtles and seabirds are the most commonly studied groups (e.g. Schuyler et al. 2014; van Franeker & Law 2015), although more recently marine mammals have been receiving increased attention (e.g. Eriksson & Bruton 2003, Lusher et al. 2015, Besseling et al. 2015).

It has been hypothesized that microplastics may facilitate the transfer of other pollutants either from the manufacturing process (e.g. plasticers and flame-retardant additives) or from the environment (e.g. organochlorates) (Rochman 2015). Sub-lethal effects associated with microplastics are still unclear (Rochman et al. 2016), and the impacts of microplastics on top predators require further investigation. Therefore, studying the ingestion of microplastics directly through investigation of digestive tracts may be considered as an appropriate method.

Bycatch in commercial fisheries can be an important source of carcasses for different analyses including dietary studies (e.g. Auttila et al., 2014; Lundström et al. 2007) and interactions with marine debris (Bravo-Rebolledo et al. 2013; Lusher et al. 2018). In contrast to stranded animals, where cause of death is often unknown, seals bycaught in fisheries are usually disease free, feeding normally in the marine environment, and therefore should provide a good overview of any effects of pollution in these animals, thus providing indication of the exposure to microplastics for the entire population. In general, the behavior of microplastics and plastics within the digestive tracts of marine mammals are difficult to predict, including how long they may remain in the digestive tract or whether they are egested with feces. The presence of

79 substances in the intestine, such as parasites, may influence the retention of microplastics as
80 they can increase the surface area for adhesion and retention.
81
82 The aims of this study were to (1) report the incidence and characteristics of microplastics
83 within the last part of the digestive tracts (small and large intestines) of grey seals by caught in
84 southern Ireland; (2) investigate the implications of parasite prevalence to the retention of
85 microplastics and their aggregation; and (3) investigate the relationship between microplastic
86 presence and prey items
87
88 **METHODS**
89 **Sample collection**
90 Grey seals by caught in fishing trammel nets targeting monkfish (*Lophius* spp) and rays
91 (Rajaidae) off the Irish south coast (Co. Cork) were recovered for post mortem examination
92 (Table 1). General measurements (body length and maximum girth), sex, age and any marks and
93 external abnormalities were noted. Postmortem examinations were performed following the
94 standard methodology (Dierauf, 1994), including measurement of blubber thickness, and
95 sampling of different organs. Full digestive tracts, from oesophagus to anus, were recovered for
96 all seals. Stomachs were analyzed for dietary purposes (Gosch et al. 2017), and intestines were
97 stored frozen (-20ºC) for parasite infection and microplastic studies.
98
99 **Laboratory analysis**
100 Stomachs were analyzed prior to the development of a microplastics protocol (Lusher et al.
101 2015); consequently, they have not been included in this study. Intestines were thawed,
102 measured and washed with pre-filtered tap water to avoid external contamination. Each intestine
103 was then divided into 20 sections of equal length. Each section was washed with pre-filtered tap
104 water through three nested sieves (from 250 µm on the bottom to 1,000 µm on the top). Food

remains and parasites were transferred into ethanol for further analysis (MacGabban 2015). The remains on the smallest mesh sieve were frozen for microplastic analysis.

A pre-made solution of 10% KOH (Foekema et al. 2013) was used to dissolve organic matter. After the samples were defrosted, the solution was added in a proportion of three times solution to one part sample and left for about three weeks. The remaining solution was filtered using a GF/C microfiber filter paper using a Büchner funnel with a vacuum pump. Identification and measuring of microplastic items were carried out under a microscope (Olympus SZX10) with a mounted camera (Q-imaging Retiga2000R), and classification followed Lusher et al. (2014).
For the purpose of this study we used the term microplastic for both anthropogenic particles, up to five millimeters; FTIR was not used to confirmation synthetic polymers which it was confirmed to be important for particles below one millimeter (Isobe et al. *in press*). Nevertheless, this study sought to investigate the relationship between parasites and anthropogenic particles, therefore the specific polymer identification is not required and visual identification following Lusher et al. (2014) deemed sufficient.
Contamination conditions were avoided during the full process and intestine sections were covered to minimize air exposure, following Lusher and Hernandez-Milian (2018) protocol guidelines. Due to the limited ability to prevent airborne contamination the size limit was set to 200µm, airborne microplastics were monitored with wet filter papers in petri dishes and procedural blanks of replicate filtered distilled water and 10 % KOH were included to monitor procedural contamination.

Data analysis

The number of microplastics and parasites per individual were calculated. In parasite studies, level of aggregation is usually investigated using different indexes (Poulin 2013, MacGabban 2015). These aggregation indexes can also be used to investigate aggregations of microplastics. In this study, two statistical indexes were used to investigate microplastics aggregation; firstly,

we used the Corrected Moment Estimate (1) as is the most common statistical index used in parasite aggregation studies; secondly we used the Poulin's Discrepancy Index, because it provides information when aggregation of parasites varies between hosts.

(1) Corrected Moment Estimate of the negative binomial distribution (k) quantifies aggregations (Sherrard-Smith et al. 2015), where $\bar{x}$ is the average abundance of microplastics, S2 is the variance and 'n' is the sample size (number of seals). The index shows that if 'k' increases, the level of aggregation also increases.

$$k = (\bar{x}^2 - s^2 / n) / (s^2 - \bar{x}) \quad \text{Eq. 1}$$

(2) Poulin's Discrepancy Index (D) measures the degree of inequality between a hypothetical equal distribution and that of the observed distribution, where 'x' is the total number of microplastics in the individual 'j', and 'N' is the total number of individuals investigated (number of seals). D ranges from 0 to 1, where 1 implies all microplastics are aggregated.

$$D=1-2i=1N(j=1ixj)xN(N+1) \quad \text{Eq. 2}$$

In order to investigate the relationship between microplastics and parasites the Kendall's Tau correlation test was used; this test allows parasite concentrations between seals to be compared, using ranks when data is scarce. In addition, Kendall's Tau test was used to investigate the relationship between microplastics and the number of prey ingested by seals. Theoretical ingestion of microplastic of seals was estimated using the average value of 1.9 microplastics ingested per fish given by Lusher et al. (2013). These values were compared with the empirical results.

RESULTS

Sample composition

158 Thirteen grey seals ranging from 107.5 cm to 240.0 cm (Table 1) were recovered dead from
159 trammel nets operating between Crosshaven, Co. Cork (51.79° N, 8.28° W) and Kinsale, Co.
160 Cork (51.66°N, 8.78°W) between 2012 and 2014. Most of the seals were juveniles (77%, n=10),
161 two were subadults (15%) and one an adult (8%). Four of the seals (31%) were juvenile females.
162 Postmortem examinations indicated that seals died due to drowning. Signs of interactions with
163 fisheries (e.g. abrasions, cuts) were also identified.
164 Intestines of the juvenile and sub-adult seals ranged in length from 15.0 m to 22.4 m (with an
165 average of 18.9 m) and the adult seal intestine measured 25.9 m. Average intestine section
166 length ranged from 75.0 cm to 111.7 cm with an average of 94.4 cm (excluding the single adult
167 measurement).
168
169 3.2 Ingested items
170 Food remains in the stomachs were partially digested. Most of the prey found in seals stomachs
171 were demersal prey (96.9% by number), which was composed primarily of gadoids (70.4% by
172 number, Table 2). Comparatively, Low numbers of food remains were found in the intestines
173 which corresponded to prey found in the stomachs.
174
175 There were no macroplastics present in any grey seal stomachs. However, all seal intestines
176 contained microplastics. A total of 363 particles were identified, 85% of which were fibers, 14%
177 fragments and 1% films (Fig 1). The average number of particles per seal was 27.9 ±14.7 (range
178 13 to 71) and the number of microplastics per section ranged from 0 to 19 particles (Fig. 2).
179 Considering an average value of 1.9 (± 0.1 SE) microplastic particles per prey item (see Lusher
180 et al. 2013), a total of 245 particles for all seals in this study were estimated based on the dietary
181 results (Table 1), this was lower than the observed count (363 particles). Most of the seals had a
182 lower estimated microplastic count when compared to the actual count (Table 1). There were
183 four seals where the estimation was higher based on the stomach contents, in one particular seal
184 (HG 3/14) the estimation value was three times greater than the actual value.

185

186 3.3 Aggregation of microplastics and parasites

187 All seals had parasites in their intestines apart from one juvenile male (HG 15/13) Most of the
188 parasites (n=1,543) were identified as Acantocephala (MacGabban 2015), with one unidentified
189 Cestode. Total numbers of acantocephalan parasites ranged from two to 199 per seal (when the
190 adult seal was not included), with an average prevalence of 74.5 ± 67.7 parasites per seal. A
191 total of 606 parasites were counted in the adult seal which accounted for 40% of the overall
192 parasite burden. When the adult seal was included in the analysis, the average prevalence of
193 parasites was 115.0±116.1. (Fig. 3, Table 1).

194

195 Both the corrected moment estimate (k = -0.06) and Poulin's discrepancy index (D = 0.857)
196 detected microplastic aggregation. We hypothesized that aggregations of microplastics may be
197 related to parasite aggregations; however, Kendall's tau correlation test showed that the number
198 of microplastics were not significantly related to parasites (Table 3). Gadoid prey was
199 significantly correlated to microplastic incidence (r_τ= -0.469, p=0.034) however no correlation
200 was found when all the dietary items were considered (Table 3).

201

202 **DISCUSSION**

203 Marine debris is considered as a major threat to the marine environment, and interactions with
204 marine mammals have been reported to reach 56% either due to entanglement or ingestion
205 (Baulch & Perry 2014, Simmonds & Baulch 2016). Recently, Lusher et al. (2018), found that all
206 cetaceans investigated from Irish coasts contained microplastics. In this study, all seal intestines
207 analyzed presented microplastics but none of them contained macroplastics in either their
208 stomachs or their intestines.

209

210 Microplastic presence in cetacean digestive tracts have already been confirmed in a humpback
211 whales (Besseling et al. 2015) and some Odontoceti (Lusher et al. 2018). Plastic items were

found in pinnipeds stomachs, such as in harbor seals (Bravo-Rebolledo et al. 2013), but also microplastics were reported in the scats of fur seals (Eriksson & Burton 2003). Ingestion of microplastics in marine mammals, especially on pinnipeds, is mainly related to consumption, as they target prey by hunting and directly consuming whole prey which might content microplastics. Therefore, it can be assumed that the ingestion of microplastics in pinnipeds may come from prey, as originally suggested by Eriksson & Burton (2003). Estimating microplastic ingestion based on occurrence in prey items is a valuable resource when anthropogenic debris information is not accessible; however, it is necessary to use these calculations with caution as under and overestimations may occur. Interestingly, the estimated number of microplastics based on prey ingestion in one of the seals was 40% less than the number of microplastic found in the intestines. However, the values increased twice (105%) and even three times (213%) when looking at the estimations obtained from two other individual seals. For the remaining individuals, the equation (Lusher et al., 2013) underestimated the observed number of microplastics.

Despite the large amount of microplastics that these top predators may ingest annually, and based on the data presented, most of them, if not all, will be egested following digestion and particles will be released back into the marine environment. Some studies have suggested that plastics, in particular microplastics, might act as vectors of other chemicals such organochlorates, heavy metals and plastic additives (e.g. Rochman 2015; Fossi et al. 2012). These chemicals may be released within the digestive tracts and absorbed into tissues. In fact, some researchers have suggested that if plastics facilitate the transference of chemicals to biota it may be possible to monitor their levels through biopsies (e.g. Fossi et al. 2012). However, this is yet to be verified as any observed levels of contaminants may be related to macrodebris, microplastic or consumed prey items.

Relationships between microplastics presence in prey items, as mentioned above, has been used to estimate levels of ingestion in other studies (Lusher et al. 2016). Prior to this study microplastics or anthropogenic particles smaller than 5mm, have not been studied in relation to other elements of the digestive system, or pathologies, e.g. the presence of parasites. Aggregations of helminth parasites are commonly observed within hosts due to both intrinsic and extrinsic factors, such as sex and site, respectively (e.g. Behnke et al. 2001; Poulin 2013). These aggregations may decrease the intestinal lumen and increase the contact surface within the intestinal lumen, therefore microplastics may have more chances to be retained in these areas. MacGabban (2015) found that most of the Acantocephala parasites in seal intestines tended to aggregate between the 9th and the 15th section. Although no statistical relationship with such aggregations was found in this study, microplastics were found to be more abundant before the 14th section of the intestine. The low sample size (n= 13) and the lack of information regarding microplastic presence in corresponding seal stomachs should may alter the results.

It is notoriously difficult to obtain biopsy samples from marine mammals due to their diving behavior; therefore, stranding and bycaught animals can be a good source of data. In cetaceans, the predicted recovery rate of stranded carcasses ranges from 8 to 30 % of dead dolphins (e.g. Peltier et al. 2012; Carretta et al. 2016**).** With regard to pinnipeds, strandings are rarely reported unless in extreme cases of disease (Bravo-Rebolledo et al. 2013), or deliberate kills or bycatch (Pierce et al. 2011). Post mortem examinations of bycaught animals provide a good source of data for studies of microplastics, pollutant and their trophic transfer within marine mammals. At the time of death these animals were actively feeding and might provide information on healthy organisms that stranded and dead organisms cannot provide, especially when they are found with signs of illness or starvation. This study presents the incidence of microplastic on seals incidentally caught in fisheries activities. Samples were only obtained from south of Ireland, leading to restricted spatial coverage. It will be necessary to investigate stranded and bycaught seals (both adults and juveniles) as well as scats from other areas to obtain a better understanding of the incidence of this type of pollution and the effects at a population level. In

addition, in this research only intestines were available for microplastic analysis, while stomachs were open previously for dietary analysis only. We believe that the use the full digestive tract will provide us a more realistic picture of the incidence of microplastics in the individuals analyzed, however, the result showed us the importance of considering the full digestive tracts and not only the stomach of these species when studying both microplastics and diet.

Nowadays, marine microplastic pollution assessments are primarily carried out analyzing non-biota samples (e.g. sediments and water), algae, invertebrates, or fish (GESAMP 2016). Organisms may act as bioindicators or sentinels (e.g. Fossi et al. 2018), because they provide information on the quality of the environment and changes over time. These species can be use as monitoring tools for ecosystem health; marine litter (macro- and microplastics) is one of the recent threats that these species are confronted with. Management strategies for microplastics are difficult to implement because the information available is still patchy, and the toxicological effects of microplastics are still unclear. Islam & Tanaka (2004) reviewed different issues on marine pollution management including governmental decisions, lack of communication among scientists, and low participation of different sectors; however, these authors found it difficult to address the problems as marine pollution is defined by characteristics such as uncertainty, conflicts, and complicated interactions.

Seals can provide further data on the incidence of microplastic pollution within higher trophic levels of the food web, especially in coastal ecosystems where the highest input occurs (more than 80% of marine debris come from land sources). However, there are three factors to be further considered; i) the effect of microplastic pollution should be studied using different age groups; ii) the stranding rate of seals is low, it is necessary to include long-term sampling, and iii) analysis of additional tissues (e.g. muscle, liver) to investigate the possible effect of chemicals associated with microplastics is required. In addition, the analysis of seal feces may provide scientists with the opportunity to obtain information without the need for carcasses. However, analyzing seal scats needs to be taken with caution because external airborne contamination may occur.

CONCLUSION

Research on the implications of large marine debris has increased during the last decade, including evaluating the interactions between predators (e.g. marine mammals, seabirds, sea turtles, sharks) and ghost nets, plastic bags, and other type of marine debris. More recently, studies on microplastic interactions have highlighted the potential effects on invertebrates and fish. Based on initial results, researchers point out that top predators might also been affected by microplastic pollution; however, incidence and accumulation of microplastics within digestive tracts are important variables needed to understand any adverse effects. Different parameters should be taken into consideration when top predators are being used in studies of pollution. Distribution and accumulation of microplastics, effects of trophic transfer and the presence of parasites are variables to be considered. Trophic transfer has been suggested as one of the main factors of microplastic incidence in top predators, and estimation of microplastic ingestion by prey has been used. The current study shows that these may be underestimates of the incidence of microplastics in top predators, although it is a valuable tool when microplastics analysis cannot be carried out. It is necessary to understand that microplastic distribution varied from one individual to another, however some kind of aggregation may occur. Aggregation can be associated with the retention effect where the reduction of lumen is higher with parasite aggregation, which may also affect the aggregation of microplastics. In this study, aggregation could not be statistically confirmed but there appeared to be some form of aggregation occurring. It is essential for future research to investigate microplastic incidence related to other factors such as prey and parasite prevalence. The knowledge of these associations will be paramount in enabling the use of these predators as monitoring tools in ecosystem management.

ACKNOWLEDGEMENTS

GHM & SMacG were funded under the Beaufort Ecosystem Approach to Fisheries Management award, as part of the Irish Government's National Development Plan (NDP), the

Beaufort Marine Research Award is grant aided by the Department of Communications, Energy and Natural Resources (DCENR) and the Department of Agriculture, Fisheries and Food (DAFF) under the Strategy for Science Technology and Innovation (SSTI) and the Sea Change Strategy. Funding for sampling was also received from Irish Government through BIM Marine Environment Protection Measure and Marine Institute Research Sub-Programme. The authors would like to give special thanks to Kieran Healy for kindly providing us with the seals, Dr Martha Gosch who provided the diet information, and Maria Garagouni and Rosemary Murphy for helping in the laboratory. Finally, authors are grateful to the reviewers and editor for providing comments to improve the quality of the manuscript.

REFERENCES

Andrady, A.L. 2015. Persistence of plastic litter in the oceans. In M. Bergmann, L. Gutow, L. & M. Klages, eds. Marine Anthropogenic Litter, pp. 57-72. Cham, Switzerland, Springer International PublishingAuttila, M., Sinisalo, T., Valtonen, M., Niemi, M., Viljanen, M., Kurkilahti, M. and Kunnasranta, M. 2014. Diet composition and seasonal feeding patterns of a freshwater ringed seal (*Pusa hispida saimensis*). Marine Mammal Science, 31(1), 45-65

Auttila, M., Sinisalo, T., Valtonen, M., Niemi, M., Viljanen, M., Kurkilahti, M. & Kunnasranta, M. 2014. Diet composition and seasonal feeding patterns of a freshwater ringed seal (*Pusa hispida saimensis*). Marine Mammals Science, 31(1), 45-65

Baulch, S. & Perry, C. 2014. Evaluating the impacts of marine debris on cetaceans. Marine Pollution Bulletin, 80(1), 210-221

Behnke, J. M., Bajer, A. Sinski, E. & Wakelin, D. 2001 Interactions involving intestinal nematodes of rodents: experimental field studies. Parasitology, 122, S39-S49.

Bergmann, M., Gutow, L. & Klages, M. (Eds). 2015. Marine Anthropogenic Litter. Springer International Publishing, Heidelberg. B.

Besseling, E., Foekema, E.M., Van Franeker, J.A., Leopold, M.F., Kühn, S., Bravo Rebolledo, E., Heße, E., Mielke, L., IJzer, J., Kamminga, P. & Koelmans, A.A., 2015.

348 Microplastic in a macro filter feeder: humpback whale Megaptera novaeangliae. Marine
349 Pollution Bulletin, 95(1), 248-252
350 Bravo-Rebolledo, E.L., Van Franeker, J.A., Jansen, O.E. & Brasseur, S.M.J.M. 2013.
351 Plastic ingestion by harbour seals (Phoca vitulina) in The Netherlands. Marine Pollution
352 Bulletin, 67(1), 200-202
353 Carretta, J.V., Danil, K., Chivers, S.J., Weller, D.W., Janiger, D.S., Berman- Kowalewski,
354 M., Hernandez, K.M., Harvey, J.T., Dunkin, R.C., Casper, D.R., Stoudt, S., Flannery, M.,
355 Wilkinson, K., Huggins, J. & Lambourn, D.M. 2016. Recovery rates of bottlenose dolphin
356 (Tursiops truncatus) carcasses estimated from stranding and survival rate data. Marine
357 Mammal Science, 32(1), 349-362
358 CBD. 2016. Marine Debris: Understanding, Preventing and Mitigating the Significant
359 Adverse Impacts on Marine and Coastal Biodiversity. Technical Series No.83. Secretariat of
360 the Convention on Biological Diversity, Montreal, 78 pages
361 Cosgrove, R., Gosch, M., Reid, D., Sheridan, M., Chopin, N., Jessopp, M., & Cronin, M.
362 2016. Seal bycatch in gillnet and entangling net fisheries in Irish waters. Fisheries Research
363 183: 192-199
364 Dierauf, L.A. 1994. Pinniped Forensic, Necropsy and Tissue Collection Guide. Wilskinson,
365 D. & Driscoll, C. (Ed.) NOAA Technical Memorandum, USA
366 Eriksson, C. & Burton, H. 2003. Origins and biological accumulation of small plastic
367 particles in fur seals from Macquarie Island. AMBIO: A Journal of the Human Environment,
368 32, 380-384
369 Foekema, E.M., De Gruijter, C., Mergia, M.T., van Franeker, J.A., Murk, A.J. & Koelmans,
370 A.A. 2013. Plastic in North Sea fish. Environ. Sci. Technol., 47(15): 8818-8824
371 Fossi, M.C., Panti, C., Guerranti, C., Coppola, D., Giannetti, M., Marsili, L. & Minutoli, R.
372 2012. Are baleen whales exposed to the threat of microplastics? A case study of the
373 Mediterranean fin whale (Balaenoptera physalus). Marine Pollution Bulletin, 64 (11), 2374-
374 2379

375 Fossi, M.C., Pedà, C., Compa, M., Tsangaris, C., Alomar, C., Claro, F., Loakeimidis, C.,
376 Galgani, F., Hema, T., Seudero, S., Romeo, T., Battaglia, P., Andaloro, F., Caliani, I.,
377 Casini, S., Panti, C. and Baini, M. 2018. Bioindicators for monitoring marine litter ingestión
378 and its impacts on Mediterranean biodiversity. Environmental Pollution, 237, 1023-1040
379 Galgani, F., Hanke, G.,Werner, S., Oosterbaan, L., Nilsson, P., Fleet, D., Kinsey, S.,
380 Thompson, R.C., van Franeker, J., Vlachogianni, T., Scoullos, M., Veiga, J.M., palatinus,
381 A., Matiddi, M., mes, T., Korpinen, S., Budziak, A., Leslie, H., Gago, J. & Liebezeir, G.
382 2013. MSFD GES technical subgroup on marine litter. Guidance on monitoring of marine
383 litter in European Seas. Luxembourg: Joint Research Centre–Institute for Environment and
384 Sustainability, Publications Office of the European Union.
385 GESAMP (2016). Sources, fate and effects of microplastics in the marine environment: part
386 two of a global assessment. (Kershaw, P.J., and Rochman, C.M., eds).
387 (IMO/FAO/UNESCO-IOC/UNIDO/WMO/IAEA/UN/UNEP/UNDP Joint Group of Experts
388 on the Scientific Aspects of Marine Environmental Protection). Rep. Stud. GESAMP No.
389 93, 220 pp
390 Gosch, M. 2017. The diet of the grey seal [*Halichoerus grypus* (Fabricius, 1791)] in Ireland
391 and potential interactions with commercial fisheries. PhD thesis, UCC, pp. 186
392 Islam, M.S. & Tanaka, M. 2004. Impact of pollution on coastal and marine ecosystems
393 including coastal and marine fisheries and approach for management: a review and
394 synthesis, Marine Pollution Bulletin, 48 (7-8): 624-649
395 Isobe, A., Buenaventura, N.T., Chastain, S., Chavanich, S., Cózar, A., DeLorenzo, M.,
396 Hagmann, P., Hinatah, H., Kozlovskii, N., Lusher, A.L., Martí, E., Michidai, Y., Muk, J.,
397 Ohnol, M., Potter, G., Ross, P.S., Shim, W.J., Song, Y.K., Takadan, H., Tokaio, T, Torii, T.,
398 Uchida, K., Vassillenko, K., Vivakarn, V. & Zhang, W. An interlaboratory comparison
399 exercise for the determination of microplastics in standard sample bottles. Marine Pollution
400 Bulletin. *In press*

401 Jambeck, J.R., Geyer, R., Wilcox, C., Siegler, T.R., Perryman, M., Andrady, A., Narayan, R.
402 & Law, K.L. 2015. Plastic waste inputs from land into the ocean. Science, 347(6223): 768-
403 771.
404 Kühn, S., Bravo Rebolledo, E.L. & van Franeker, J.A. 2015. Deleterious effects of litter on
405 marine life. In: Marine Anthropogenic Litter (pp. 75-116). Springer International Publishing
406 Lundström, K., Hjerne, O., Alexandersson, A. and Karlsson O. 2007. Estimation of grey
407 seal (*Halichoerus grypus*) diet composition in the Baltic Sea. NAMMCO Sci. Publ. 6, 177-
408 196
409 Lusher, A.L., Burke, A., O'Connor, I. & Officer, R. 2014. Microplastic pollution in the
410 Northeast Atlantic Ocean: validated and opportunistic sampling. Marine Pollution Bulletin
411 88(1): 325-333.
412 Lusher, A. and Hernandez-Milian, G. 2018. Microplastic Extraction from Marine Vertebrate
413 Digestive Tracts, Regurgitates and Scats: A Protocol for Researchers from All Experience
414 Levels. Bio-Protocol, 8(22), 1-12
415 Lusher, A.L., Hernandez-Milian, G., O'Brien, J., Berrow, S., O'Connor, I. and Officer, R.,
416 (2015). Microplastic and macroplastic ingestion by a deep diving, oceanic cetacean: The
417 True's beaked whale Mesoplodon mirus. Environmental Pollution, 199, pp.185-19
418 Lusher, A.L., McHugh, M. and Thompson, R.C. 2013. Occurrence of microplastics in the
419 gastrointestinal tract of pelagic and demersal fish from the English Channel. Marine
420 Pollution bulletin, 67, 94-99
421 Lusher, A.L., O'Donnell, C., Officer, R. and O'Connor, I. 2016. Microplastic interactions
422 with North Atlantic mesopelagic fish. ICES Journal of Marine Science, 73(4), 1214-1225
423 Lusher, A.L., Hernandez-Milian, G., Berrow, S., Rogan, E. and O'Connor, I. 2018.
424 Incidence of marine litter in cetaceans stranded and bycaught in Ireland: recent findings and
425 a review of historical knowledge. Environmental Pollution, 232, 467-476
426 MacGabban, S. 2015. pH and patterns of distribution and aggregation of the intestinal
427 Acanthocephalan, Corynosoma strumosum in grey seals (Halichoerus grypus): more than a
428 gut feeling?. MSc thesis, UCC, pp. 34

Panti, C., Baini, M., Lusher, A., Hernandez-Milian, G., Bravo Rebolledo, E.L., Unger, B., Syberg, K., Simmonds, M.P.and Fossi, M.C. 2019 Marine litter: one of the major threats for marine mammals. Outcomes from the European Cetacean Society workshop. Environmental Pollution, 247, 72-79

Peltier, H., Dabin, W., Daniel, P., Van Canneyt, O., Dorémus, G., Huon, M. & Ridoux, V., 2012. The significance of stranding data as indicators of cetacean populations at sea: modelling the drift of cetacean carcasses. Ecological Indicators, 18, 278-290

Pierce, G.J., Hernandez-Milian, G., Santos, M.B., Dendrinos, P., Psaradellis, M., Tounta, E., Androukaki, E & Edridge, A. 2011. Diet of monk seal (Monachus monachus) in Greek waters. Aquatic Mammals, 37(3), 284-297

Poulin, R. 2013. Explaining variability in parasite aggregation levels among host samples. Parasitology, 140 (4), 541-546

Rochman, C.M., 2015. The complex mixture, fate and toxicity of chemicals associated with plastic debris in the marine environment. In M. Bergmann, L. Gutow, L. & M. Klages, eds. Marine Anthropogenic Litter, pp. 117-140. Cham, Switzerland, Springer International Publishing

Rochman, C.M., Browne, M.A., Underwood, A.J., van Franeker, J.A., Thompson, R.C. & Amaral- Zettler, L.A. 2016. The ecological impacts of marine debris: unraveling the demonstrated evidence from what is perceived. Ecology, 97(2): 302-312

Schuyler, Q., Hardesty, B.D., Wilcox, C. and Townsend, K. 2014. Global analysis of anthropogenic debris ingestion by sea turtles. Conservation Biology, 28(1), 129-139

Simmonds, M.P. & Baulch, S. 2016. An update on research into marine debris and cetaceans. IWC SC/66a/E/5

van Franeker, J.A. and Law, K.L. 2015. Seabirds, gyres and global trends in plastic pollution. Environmental Pollution, 203, 89-96

TABLES

Table 1. By caught seals in south of Ireland (Co. Cork: 51.79N, 8.28W - 51.66N, 8.78W). M: male, F: female, J: juvenile, SA: subadult, A: adult. Length of seals in cm. Blubber: ventral blubber thickness in mm. T.B.: total burden of parasites. MP: number of microplastics. MPe: number of microplastics estimated using Lusher et al. 2013 value of 1.9 MP/fish.

Table 2. Number of prey ingested by seals

Table 3. Kendall's rank correlation (tau) output. * adult seal was not included. r_τ correlation coefficient tau

Table 1. By caught seals in south of Ireland (Co. Cork: 51.79N, 8.28W - 51.66N, 8.78W). M: male, F: female, J: juvenile, SA: subadult, A: adult. Length of seals in cm. Blubber: ventral blubber thickness in mm. T.B.: total burden of parasites. MP: number of microplastics. MPe: number of microplastics estimated using Lusher et al. 2013 value of 1.9 MP/fish, where (+) and (-) indicates underestimated and overestimated percentages of microplastics when comparing with data obtained from seals.

	Year	Sex	Age	Length	Blubber	T.B.	MP	MPe	Variation
HG1/12	2012	M	J	119	35	2	29	5.7	-80%
HG2/12	2012	M	J	115	20	3	30	1.9	-94%
HG8/13	2013	M	J	130	18	105	13	23	+63%
HG10/13	2013	F	J	145	25	159	19	27	+40%
HG11/13	2013	F	J	147	25	32	32	1.9	-94%
HG12/13	2013	M	SA	156	40	135	71	7.6	-89%
HG13/13	2013	M	SA	162	35	84	30	7.6	-75%
HG14/13	2013	M	J	137	8	199	24	19	-21%
HG15/13	2013	M	J	130	18	0	20	15	-24%
HG16/13	2013	F	J	138	28	8	19	7.6	-60%
HG18/13	2013	M	J	144	21	56	NA	7.6	NA
GAR1/14	2014	M	A	240	25	606	25	-	NA
HG2/14	2014	F	J	124	15	102	36	76	+105%
HG3/14	2014	M	J	140	13	62	15	53	+213%

Table 2. Number of prey ingested by seals.

	HG12/13	HG13/13	HG1/12	HG16/13	HG10/13	HG11/13	HG2/12	HG8/13	HG15/13	HG2/14	HG3/14	HG14/13
Merlangius merlangus				2				2		2	6	
Melanogrammus aeglefinus	1				1			5				6
Gadus morhua					5					12	2	0
Pollachius pollachius					1							
Trisopterus sp.	1	3	1	1		1	3	6	12	7	1	
Gaidropsaurus vulgaris	1											
Molva molva								1	1			
Merluccius merluccius	1	1										1
Trachurus trachurus	1											
Clupea harengus	1											
Belone belone						1						
Ammodytidae	1				5					14	9	1
Callionymus sp.									1		1	
Labrus sp.											1	
Flatfish								1			2	
Unknown fish												1
Total	4	4	3	4	14	1	1	12	8	40	28	10

Table 3. Kendall's rank correlation (tau) output. * adult seal was not included. r_τ correlation coefficient tau

	r_τ	p-value	z
Microplastics vs parasites	0.0260	0.9025	0.11247
Microplastics vs parasites*	0.1378	0.8904	0.13779
Microplastics vs diet	-0.3493	0.1254	-1.5324
Microplastics vs gadoid prey	-0.4688	0.0379	-2.0765
Microplastics vs demersal prey	-0.4252	0.0606	-1.8762
Parasites vs diet	0.3127	0.1655	1.3868
Parasites vs gadoid prey	0.1846	0.4084	0.8268
Parasites vs demersal prey	0.2326	0.2995	1.0376

FIGURES

Figure 1. Photographs of fibers (a, c), films (b), and fragments (d) within the seal intestines.

Figure 2. Average number of microplastics per section of intestines. Error bars show the standard error.

Figure 3. a) number of microplastics (GREY) and parasites (BLACK) in all seals except the adult seal (GAR 1/14)




Figure 1. Photographs of fibers (a, b), films (c), and fragments (d) within the seal intestines.




Figure 2. Average number of microplastics per section of intestines. Error bars show the standard error.




Figure 3. number of microplastics (GREY) and parasites (BLACK) in all seals except the adult seal (GAR 1/14)