CORA Cork Open Research Archive
Cartlann Taighde Oscailte Chorcaí

Title	Appropriate use of elective coronary angiography in patients with suspected stable coronary artery disease
Authors	Hanrahan, Michael T.;O'Flynn, Anne Marie;McHugh, Sheena M.;Kearney, Peter;Kearney, Patricia M.
Publication date	2019-01-05
Original Citation	Hanrahan, M. T., O'Flynn, A. M., McHugh, S. M., Kearney, P., Kearney, P. M. (2019) 'Appropriate use of elective coronary angiography in patients with suspected stable coronary artery disease', Irish Journal of Medical Science, pp. 1-7. doi:10.1007/s11845-018-1953-4
Type of publication	Article (peer-reviewed)
Link to publisher's version	10.1007/s11845-018-1953-4
Rights	© 2019, Royal Academy of Medicine in Ireland. Published by Springer Nature Switzerland AG. This is a post-peer-review, pre-copyedit version of an article published in the Irish Journal of Medical Science. The final authenticated version is available online at: http://dx.doi.org/10.1007/s11845-018-1953-4
Download date	2026-11-16 19:17:07
Item downloaded from	https://hdl.handle.net/10468/7432

UCC
University College Cork, Ireland
Coláiste na hOllscoile Corcaigh

Table 1. Baseline Characteristics		
Total N = 259	Category	Mean ± SD / N (%)
Age		63 ±10
	<65 years	136 (53%)
	≥65 years	123 (47%)
Gender	Male	152 (59%)
	Female	107 (41%)
Ethnicity	White	253 (98%)
	Black/Other	6 (2%)
Marital Status	Married	180 (70%)
	Widow(er)	20 (8%)
	Single	30 (12%)
	Divorced/Separated	24 (9%)
	Unknown	5 (1%)
Medical Care	Public	170 (66%)
	Private	89 (34%)
BMI		29 ±6
BMI Category	Not obese	164 (63%)
	Obese	95 (37%)
Smoking	Never	98 (38%)
	Ex/current	161 (62%)
Family Hx of CAD	Yes	170 (66%)
Hypertension	Yes	137 (53%)
Hyperlipidemia	Yes	162 (63%)
Diabetes mellitus	Yes	38 (15%)
Atrial Fibrillation	Yes	19 (7%)
Total Cholesterol		5.0 +/-1.2
SBP		133 +/-18
DBP		79 +/-11
SCORE Risk	Low	23 (9%)
	Moderate	200 (77%)
	High	25 (10%)
	Very high	11 (4%)
Symptoms	Yes	106 (41%)
Exercise Stress Test	Yes	221 (85%)
Duke Treadmill Score	Low risk	57 (26%)
	Equivocal risk	145 (66%)
	High risk	10 (4%)
	Missing	9 (4%)
Echocardiography	Yes	98 (38%)
CT coronary angiogram	Yes	1 (0.4%)