

|                             |                                                                                                             |
|-----------------------------|-------------------------------------------------------------------------------------------------------------|
| Title                       | Caesarean wound care for midwives                                                                           |
| Authors                     | Murphy, Margaret M.                                                                                         |
| Publication date            | 2013-10                                                                                                     |
| Original Citation           | Murphy, M. (2013) 'Caesarean wound care for midwives', The Practising Midwife, 16 (9), pp. 27-29.           |
| Type of publication         | Article (peer-reviewed)                                                                                     |
| Link to publisher's version | <a href="http://www.ingentaconnect.com/content/mesl/tpm">http://www.ingentaconnect.com/content/mesl/tpm</a> |
| Rights                      | © 2013, Medical Education Solutions Ltd                                                                     |
| Download date               | 2026-09-25 23:51:07                                                                                         |
| Item downloaded from        | <a href="https://hdl.handle.net/10468/2749">https://hdl.handle.net/10468/2749</a>                           |



# UCC

**University College Cork, Ireland**  
 Coláiste na hOllscoile Corcaigh

# Caesarean wound care for midwives

Having investigated the risk factors for surgical site infections, **Margaret Murphy** makes recommendations for midwives to help keep women safe



**SUMMARY** With a rise in caesarean births there is a rise in wound care management issues for midwives and the potential for surgical site infections (SSIs). The burden of SSIs include increases in maternal mortality, morbidity, length of hospital stay and cost. Sepsis is currently the leading cause of maternal mortality, with 50 per cent of the women who die having had a caesarean birth (Centre for Maternal and Child Enquiries (CMACE) 2011). Wound management and the prevention of sepsis are therefore issues of great concern to midwives. This article considers the incidence of wound infections and presents the guidance available to help address this problem.

**Keywords** Wound infection, caesarean, surgical site infection, guidance

**Author** Margaret Murphy, lecturer in the school of nursing and midwifery at University College Cork

## Incidence

The Health Protection Agency (HPA) in the UK cite the most frequent healthcare-associated infections (HCAIs) detected, as being respiratory tract, urinary tract and surgical site infections (HPA 2012). Surgical site infections (SSIs) were third overall and accounted for 15.7 per cent of all HCAs. According to Public Health England (PHE) (2013), rates in England are as diverse as 15.4-34.9 per cent, depending where a woman gives birth. However Leaper et al (2013) would caution that there is non-standardisation of definitions and reporting practices in the UK and Ireland which needs to be addressed to obtain a truly accurate picture of SSIs following caesarean (CS) birth. Wilson et al (2013), in their study of 4,107 women across 14 hospital sites, found a

---

**Independent risk factors for an early wound infection are: obesity, multiple pregnancy, diabetes, hypertension, premature rupture of membranes and emergency CS birth**

---

SSI rate of 404 (10 per cent). Women are more likely to develop SSIs if they laboured prior to the CS birth (Allen et al 2003).

## Risk and compounding factors

Independent risk factors for an early wound infection are: obesity, multiple pregnancy, diabetes, hypertension, premature rupture of membranes and emergency CS birth (Schneid-Kofman et al 2005). Ward et al (2008) identified five risk factors for SSI: body mass index, age, blood loss, method of wound closure and emergency procedures. The main causative organisms for SSIs are *Staphylococcus aureus* (30 per cent), Coagulase-negative staphylococci (13.7 per cent) and *Enterococcus* spp (11.2 per cent) (Hidron et al 2008).

## Prevention of surgical site infections

Current research recommends the use of prophylactic antibiotic before CS birth, closure of subcutaneous space

greater than two centimetres and maintaining normothermia intraoperatively to help reduce the incidence of postoperative wound complications (Tipton et al 2011). In relation to midwifery care, the National Institute for Health and Care Excellence (NICE) (2012: 8-9) recommend the following for routine CS wound care:

- removing the dressing 24 hours after the CS
- specific monitoring for fever
- assessing the wound for signs of infection (such as increasing pain, redness or discharge), separation or dehiscence

## Of paramount importance is the timely recognition and treatment of sepsis in women

- encouraging the woman to wear loose, comfortable clothes and cotton underwear

- gently cleaning and drying the wound daily
- if needed, planning the removal of sutures or clips

In addition, the key recommendations from the surgical infection guideline with particular relevance to post CS wounds include:

- women should be offered clear, consistent information and advice about the risks of surgical site infections;
- the skin should be prepared at the surgical site immediately before incision, using an antiseptic (aqueous or alcohol-based) preparation such as povidone-iodine or chlorhexidine;
- hair removal should not be routinely used, to reduce the risk of surgical site infection. If required, single-use head electric clippers (not razors) should be used on the day of surgery. Surgical incisions anticipated to heal by primary intention should be covered with a film membrane, with or without a central absorbent pad (NICE 2008).

Contributing factors for SSIs following caesarean birth are outlined in *Figure 1*.

Of paramount importance to this is the timely recognition and treatment of sepsis in women (CMACE 2011). Other causes of pyrexia (mastitis, urinary tract infection or upper respiratory tract infection) need to be investigated and excluded (Tharpe 2008). Strict adherence to hand hygiene needs to be used by both women and healthcare staff (CMACE 2011).

Approaches are needed to prevent, or decrease, the risk of SSIs following caesarean birth involving this multidisciplinary team (see *Figure 2*, over page). All women who experience a caesarean birth should have post-discharge surveillance from a multidisciplinary team (Ward et al 2008).

**Figure 1 Contributing factors for SSIs following caesarean birth**

| Pre-existing maternal conditions                                      | Events during labour or birth                             | Procedure/provider related conditions and events |
|-----------------------------------------------------------------------|-----------------------------------------------------------|--------------------------------------------------|
| Extremes of maternal age                                              | Pre-term labour and birth                                 | Pre-operative shaving                            |
| Elevated body mass index                                              | Prolonged rupture of membranes                            | Pre-operative skin preparation technique         |
| Smoking                                                               | Prolonged labour                                          | General anaesthesia                              |
| Primiparity                                                           | Intrapartum fever/pyrexia                                 | Hypothermia                                      |
| Low socioeconomic status                                              | Multiple vaginal examinations                             | Poor aseptic technique                           |
| Poor maternal hygiene                                                 | Post-term pregnancy                                       | Inadequate sterilisation of instruments          |
| Poor nutrition                                                        | Thick meconium staining                                   | Delayed or omitted prophylactic antibiotics      |
| Poor oxygenation                                                      | Internal fetal scalp electrode                            | Suboptimal haemostasis                           |
| Poor tissue perfusion                                                 | Uterine monitoring with an intrauterine pressure catheter | Practitioner skill                               |
| Multiple comorbidities                                                | Operative vaginal birth                                   | Practitioner experience                          |
| Diabetes                                                              | Foley catheter                                            | Length of procedure                              |
| Hypertension                                                          | Caesarean birth                                           | Operative trauma                                 |
| Immune compromise                                                     | Uterine perforation                                       | Contamination of wound or surgical site          |
| Splenectomy                                                           | Manual removal of placenta                                | Residual dead space following wound closure      |
| Severe anaemia                                                        | Retained products of conception                           |                                                  |
| Infection: Bacterial vaginosis, chlamydia, gonorrhoea, trichomoniasis |                                                           |                                                  |

(Andrews et al 2003; Smaill and Hofmeyr 2002; Smaill and Gyte 2010; Mangram et al 1999; Maharaj 2007a; Maharaj 2007b)

## Conclusion

Midwives must be competent in assessing, recognising and caring for women who experience CS birth and who are at risk of developing a surgical site infection. The most recent confidential inquiry (CMACE 2011) has identified the need for midwives to address sepsis prevention and management as a matter of urgency.

## TPM

Margaret Murphy is a lecturer in the school of nursing and midwifery at University College Cork

## Midwives must be competent in assessing, recognising and caring for women who experience CS birth and who are at risk of developing a surgical site infection

## References

Allen VM, O'Connell CM, Liston RM et al (2003).

'Maternal morbidity associated with cesarean delivery without labor compared with

spontaneous onset of labor at term'. *Obst Gyn*, 102(3): 477-482.

Andrews WW, Hauth JC, Cliver SP et al (2003).

'Randomized clinical trial of extended spectrum antibiotic prophylaxis with coverage for *Ureaplasma urealyticum* to reduce post-cesarean delivery endometritis'. *Obst Gyn*, 101(6): 1183-1189.

Greif R, Akca O, Horn EP et al (2000).

'Supplemental perioperative oxygen to reduce the incidence of surgical-wound infection'. *New Eng Jour Med*, 342(3): 161-167.

HPA (2012). *English national point prevalence survey on HAIs and antimicrobial use, 2011*, London: HPA.

Hidron AI, Edwards JR, Patel J et al (2008). 'NHSN annual update: antimicrobial resistant pathogens associated with healthcare associated infections: annual summary of data reported to the National Healthcare Safety Network at the centers for disease control and prevention, 2006–2007'. *Infect Cont and Hosp Epidemiol*, 29(11): 996-1011.

Leaper D, Tanner J and Kiernan M (2013).

'Surveillance of surgical site infection: more accurate definitions and intensive recording needed'. *Jour Hosp Infect*, 83(2): 83-86.

CMACE (2011). 'Saving mothers' lives: reviewing maternal deaths to make motherhood safer: 2006–2008'. *Int Jour of Obst Gyn*, 118: 1-203.

Maharaj D (2007a). 'Puerperal pyrexia: a review. Part 1'. *Obst and Gyn Survey*, 62(6): 393-399.

Maharaj D (2007b). 'Puerperal pyrexia: a review. Part 2'. *Obst and Gyn Survey*, 62(6): 400-406.

Mangram AJ, Horan TC, Pearson ML et al (1999). 'Guideline for prevention of surgical site infection, 1999. Hospital infection control practices advisory committee'. *Infect Cont and Hosp Epidemiol*, 20(4): 250-278.

Melling AC, Ali B, Scott EM et al 2001. 'Effects of preoperative warming on the incidence of wound infection after clean surgery: a randomised controlled trial'. *Lancet*, 358: 876-880.

NICE (2008). *Surgical site infection: prevention and treatment of surgical site infection (CG 74)*, London: NICE.

NICE (2012). *Caesarean section (CG 132)*, London: NICE.

PHE (2013). *Caesarean section rates 2009-2010*, London: PHE.

Schneid-Kofman N, Sheiner E, Levy A et al (2005). 'Risk factors for wound infection

| Figure 2 Team approaches to minimising infection                                    |                                                                        |                                                         |
|-------------------------------------------------------------------------------------|------------------------------------------------------------------------|---------------------------------------------------------|
| Postpartum and postoperative infections: preventive measures                        |                                                                        |                                                         |
| Maternal care                                                                       | Recommended practices                                                  | Surgical technique                                      |
| Screen and treat pre-existing genital tract infection                               | Minimise activity in the procedure room                                | Maintain excellent surgical or delivery technique       |
| Screen and treat medical conditions, particularly those leading to vasoconstriction | Wear clean scrubs and personal protective equipment                    | Limit blood loss, maintain haemostasis                  |
| Pre-procedure antiseptic shower                                                     | Perform recommended surgical/delivery scrub                            | Ensure rigorous adherence to aseptic technique          |
| Minimise number of vaginal examinations in labour                                   | Provide sterile drapes between woman and potential source of infection | Minimise tissue trauma; limit use of ties and cautery   |
| Avoid hair removal or use clippers shortly before procedure                         | Provide supplemental oxygen during and post procedure                  | Remove or debride devitalised tissue                    |
| Provide adequate skin preparation, use occlusive incise drape                       | Pre- and intra-operative warming                                       | Eliminate dead space during closures                    |
| Provide antibiotic prophylaxis as appropriate                                       | Correct volume deficits                                                | Use monofilament suture where possible                  |
| Maintain perineal integrity                                                         | Maintain normoglycaemia                                                | Elevate partial pressure of oxygen (PaO2)               |
| Encourage adequate vitamin C, vitamin A, zinc and omega-3 fatty acid intake         | Provide adequate pain control                                          | Replace fluid losses to maintain peripheral circulation |

(Mangram et al 1999; Greif et al 2000; Melling et al 2001; Ueno et al 2006; Maharaj 2007a; Tharpe 2008)

- following cesarean deliveries'. *Int Jour of Gyn Obst*, 90(1): 10-15.
- Smaill FM and Hofmeyr GJ (2002). 'Antibiotic prophylaxis for cesarean section'. *Cochrane Database of Systematic Reviews (CDSR)*, 3: CD000933.
- Smaill FM and Gyte GM (2010). 'Antibiotic prophylaxis versus no prophylaxis for preventing infection after cesarean section'. *CDSR*, 1: CD007482.
- Tharpe N (2008). 'Postpregnancy genital tract and wound infections'. *Jour Midw Wom Health*, 53(3): 236-246.
- Tipton AM, Cohen SA and Chelmow D (2011). 'Wound infection in the obese pregnant woman'. *Semin in Perinatol*, 35(6): 345-349.
- Ueno C, Hunt TK and Hopf HW (2006). 'Using physiology to improve surgical wound outcomes'. *Plastic Reconstr Surg*, 117(7S) 59S-71S.
- Ward VP, Charlett A, Fagan J et al (2008). 'Enhanced surgical site infection surveillance following caesarean section: experience of a multicentre collaborative post-discharge system'. *Jour Hosp Infect*, 70(2): 166-173.
- Wilson J, Wloch C, Seai A et al (2013). 'Inter-hospital comparison of rates of surgical site infection following caesarean section delivery: evaluation of a multicentre surveillance study'. *Jour Hosp Infect*, 84(1) 44-51.