

Title	Poly(ethylene glycol)-Based Peptidomimetic “PEGtide” of Oligo-Arginine allows for efficient siRNA Transfection and gene inhibition
Authors	Hibbitts, Alan;O'Connor, Aoife M.;McCarthy, Joanna;Forde, Éanna B.;Hessman, Gary;O'Driscoll, Caitríona M.;Cryan, Sally-Ann;Devocelle, Marc
Publication date	2019-06-10
Original Citation	Hibbitts, A., O'Connor, A.M., McCarthy, J., Forde, E.B., Hessman, G., O'Driscoll, C.M., Cryan, S.A. and Devocelle, M., 2019. Poly (ethylene glycol)-Based Peptidomimetic “PEGtide” of Oligo-Arginine Allows for Efficient siRNA Transfection and Gene Inhibition. ACS Omega, 4(6), (10pp). DOI:10.1021/acsomega.9b00265
Link to publisher's version	https://pubs.acs.org/doi/10.1021/acsomega.9b00265 - 10.1021/acsomega.9b00265
Rights	© 2019 American Chemical Society - https://pubs.acs.org/page/policy/authorchoice_termsofuse.html
Download date	2024-05-13 18:32:21
Item downloaded from	https://hdl.handle.net/10468/9139



Supporting Information

Poly(ethylene glycol)-Based Peptidomimetic “PEGtide” of Oligo-Arginine Allows for Efficient siRNA Transfection and Gene Inhibition

Alan Hibbitts,^{†,‡,§,◆} Aoife M. O’Connor,^{‡,◆} Joanna McCarthy,[▼] Éanna B. Forde,[‡] Gary Hessman,[▲] Caitriona M. O’Driscoll,[▼] Sally-Ann Cryan,^{†,§,‡,○} and Marc Devocelle^{*‡}

[†]Tissue Engineering Research Group, Department of Anatomy, [‡]Department of Chemistry, and [§]Drug Delivery & Advanced Materials Team, School of Pharmacy, Royal College of Surgeons in Ireland (RCSI), 123 St. Stephen’s Green, Dublin 2, Ireland

[▲]Trinity Centre for Biomedical Engineering and [○]School of Chemistry, Trinity College Dublin (TCD), The University of Dublin, College Green, Dublin 2, Ireland

[#]Advanced Materials and Bioengineering Research (AMBER) Centre, RCSI & TCD, Dublin 2, Ireland

[▼]Pharmacodelivery Group, School of Pharmacy, University College Cork, Cavanagh Pharmacy Building, College Road, Cork T12 YN60, Ireland

[○]Centre for Research in Medical Devices (CURAM), NUIG & RCSI, Biomedical Sciences, National University of Ireland Galway (NUIG), Newcastle Road, Galway H91 W2TY, Ireland

Table of contents

- Analytical data for octa-D-arginine (Figures S1-S2)
- Analytical data for Poly(glycidylguanidine) (Figures S3-S7)
- Optimisation of pDNA and RNAiFect transfection conditions (Figure S8)
- Settings for In Cell® 1000 Workstation analysis (Tables S1-S2)

Figure S1. RP-HPLC chromatogram of octa-D-arginine peptide




Figure S2. MALDI-TOF MS spectrum of octa-D-arginine peptide





Figure S3. ^1H NMR spectrum of poly(glycidyl *tert*-butylcarbamate) (**1**)



The signal at 3.46 ppm is caused by residual methanol.

Figure S4. ^1H NMR spectrum of poly(glycidylamine) (**2**)





Figure S5. ^1H NMR spectrum of poly(glycidyl *N,N'*-di-Boc-guanidine) (**3**)



Signals at 3.35 ppm and 2.11 ppm correspond to residual methanol and acetone, respectively. Signals at 5.25 and 4.67 ppm are attributed to the secondary and primary alcohols, respectively.

Figure S6. ^1H NMR spectrum of poly(glycidylguanidine)




Figure S7. MALDI-TOF MS spectrum of polyglycidylguanidine.




Figure S8. Optimisation of SuperFect pDNA and RNAiFect siRNA transfection conditions. A) pDNA dose, B) SuperFect-pDNA treatment time with cells prior to washing, C) total incubation time prior to lysis and luciferase analysis, D) SuperFect:pDNA transfection ratios and E) RNAiFect:siRNA transfaction ratios using optimal pDNA transfection conditions.




Table S1. Settings for In Cell® 1000 Workstation analysis of Polymer-siRNA nanoparticle uptake into Calu-3 cells

Feature	Source	Segmentation	Min. Area	Sensitivity	Collar
Nuclei	Wave 1 (Hoechst)	Top Hat	50µm ²	100%	-
Cell	Wave 2 (TRITC)	Collar	-	-	8µm
Organelles	Wave 3 (FITC)	Cytoplasm only	0.05-0.5µm	50%	-

Table S2. Settings for In Cell® 1000 Workstation analysis of Polymer siRNA nanoparticle induced toxicity in Calu-3 cells

Feature	Source	Segmentation	Min. Area	Sensitivity	Collar
Nuclei	Wave 1 (NA, NI, CN)	Top Hat	50µm ²	100%	-
Cell	Wave 2 (PMP)	Collar	-	-	8µm
Reference 1	Wave 3 (Cyt-C)	Pseudo-Cells	-	-	-
Reference 2	Wave 4 (MMP)	Pseudo-Cells	-	-	-
Reference 3	Wave 2 (PMP)	Pseudo-Nuclei	-	-	-