

Title	Suicide and deliberate self harm in older Irish adults
Authors	Corcoran, Paul;Reulbach, Udo;Perry, Ivan J.;Arensman, Ella
Publication date	2010-08-18
Original Citation	Corcoran, P., Reulbach, U., Perry, I. J. and Arensman, E. (2010) 'Suicide and deliberate self harm in older Irish adults', International Psychogeriatrics, 22(08), pp. 1327-1336. http://dx.doi.org/10.1017/S1041610210001377
Type of publication	Article (peer-reviewed)
Link to publisher's version	10.1017/S1041610210001377
Rights	© International Psychogeriatric Association 2010
Download date	2026-08-19 14:09:45
Item downloaded from	https://hdl.handle.net/10468/2916



CAMBRIDGE
UNIVERSITY PRESS

Suicide and deliberate self harm in older Irish adults

Journal:	<i>International Psychogeriatrics</i>
Manuscript ID:	IPG-03-10-047.R1
Manuscript Type:	Original Research Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Corcoran, Paul; National Suicide Research Foundation; University of Oviedo, Psychiatry Reulbach, Udo; National Suicide Research Foundation Perry, Ivan; University College Cork, Epidemiology & Public Health Arensman, Ella; National Suicide Research Foundation
Keyword:	Suicide, Epidemiology, Depression



Suicide and deliberate self harm in older Irish adults

Paul Corcoran^{1,2}, Udo Reulbach¹, Ivan J Perry³, Ella Arensman¹

Affiliations

¹ National Suicide Research Foundation, 1 Perrott Avenue, College Road, Cork, Ireland

² Department of Psychiatry, School of Medicine, University of Oviedo, Centro de Investigación Biomédica en Red de Salud Mental (CIBERSAM), Julian Claveria 6, 33006 Oviedo, Spain

³ Department of Epidemiology and Public Health, University College, Cork, Ireland

Corresponding author: Paul Corcoran, National Suicide Research Foundation, 1 Perrott Avenue, College Road, Cork, Ireland. Email: paul.nsr@iol.ie Telephone: +353 21 4277499 Fax: +353 21 4277545

Suicidal behavior in older Irish adults

ABSTRACT

Background Hospital-treated deliberate self harm and suicide among older adults have rarely been examined at a national level.

Methods The Irish Central Statistics Office provided suicide and undetermined death data for 1980-2006. The National Registry of Deliberate Self Harm collected data relating to deliberate self harm presentations made in 2006-2008 to all forty Irish hospital emergency departments.

Results Rates of female suicide among older adults (over 55 years) were relatively stable in Ireland during 1980-2006 whereas male rates increased in the 1980s and decreased in more recent decades. Respectively, the annual male and female suicide and undetermined death rate was 22.1 and 7.6 per 100,000 in 1997-2006. Male and female deliberate self harm was 3.0 and 11.0 times higher at 67.4 and 83.4 per 100,000, respectively. Deliberate self harm and suicide decreased in incidence with increasing age. Deliberate self harm generally involved drug overdose (male: 72%; female 85%) or self-cutting (male: 15%; female 9%). The most common methods of suicide were hanging (41%) and drowning (29%) for men and drowning (39%) and drug overdose (24%) for women. City and urban district populations had the highest rates of hospital-treated self harm. The highest suicide rates were in urban districts.

Conclusions Older Irish adults have high rates of hospital-treated deliberate self harm but below average rates of suicide. Drowning was relatively common as a method of suicide. Restricting availability of specific medications may reduce both forms of suicidal behavior.

Abstract word count: 237

Keywords: Suicide, undetermined death, deliberate self harm, suicidal behavior, older adults

Running title: Suicidal behavior in older Irish adults

Manuscript word count: 3,596

Introduction

Suicide is among the 10 leading causes of death worldwide and with approaching 1.5 million people dying from suicide each year, it is expected to represent 2.4% of global mortality by 2020 (Bertolote and Fleischmann, 2002). In most industrialized countries, suicide in older adults increased since the 1980s (Kaplan *et al.*, 1998), while older people also constitute the fastest growing population in most **developed** countries (Conwell *et al.*, 2002). Therefore, the number of suicides in older people is expected to increase even more in the future. Moreover, in many European countries, older men are at highest risk of suicide (Cattell and Jolley, 1995) although the pattern of increasing suicide rates with increasing age was not evident in Ireland when previously examined (Kelleher *et al.*, 1997).

It has been suggested that deliberate self harm in older people can frequently be regarded as a failed suicide attempt (Dennis *et al.*, 2007). This suggestion is supported by the lower case fatality rate observed in older people. In a WHO/EURO multicenter study on suicidal behavior (De Leo *et al.*, 2001), the ratio between fatal (suicide) and non-fatal (deliberate self harm) behaviors was 1:2. More recent studies from the UK (Hawton and Harriss, 2008b) and Ireland (Corcoran *et al.*, 2003) showed the rate ratio of deliberate self harm to suicide to decrease markedly with increasing age. The former also showed there to be a higher proportion of high suicidal intent cases of deliberate self harm in the older adults. Self harm is the strongest risk factor for suicide (Hawton and van Heeringen, 2009) and particularly so in older people (Cooper *et al.*, 2005; Hawton and Harriss, 2006; Hawton *et al.*, 2003). This has significant implications for clinicians involved in assessment and management of patients who present to hospital due to deliberate self harm (Kapur, 2006; Reulbach and Bleich, 2008).

A recent review reflecting on deliberate self harm in older adults (Chan *et al.*, 2007), underlined the requirement for more population-based studies with adequate sample size. The aim of the present population-based study was to examine the incidence and nature of

Suicidal behavior in older Irish adults

hospital-treated deliberate self harm and suicide mortality among people aged 55 years and older in Ireland.

Methods*Suicide data*

The Irish Central Statistics Office (CSO) provided data relating to all deaths by suicide and deaths of undetermined intent (respectively, codes E950-959 and E980-989 of the Ninth Revision of the International Classification of Diseases, Injuries and Causes of Death (ICD-9)) registered as occurring in 1980-2006, inclusively (2006 being the most recent year for which complete data were available).

Deliberate self harm data

The Irish National Registry of Deliberate Self Harm collected data on deliberate self harm presentations to all forty hospital emergency departments (EDs) operating in Ireland in 2006-2008. The definition of deliberate self harm used was that developed by the former WHO/Euro Multicentre Study on Parasuicide (Platt *et al.*, 1992):

‘An act with a non-fatal outcome, in which an individual deliberately initiates a non-habitual behavior that, without intervention from others, will cause self-harm, or deliberately ingests a substance in excess of the prescribed or generally recognised therapeutic dosage, and which is aimed at realising changes which the subject desired via the actual or expected physical consequences.’

While the definition was associated with the term parasuicide, the Registry utilises the term deliberate self harm. All data were collected **and coded** by **trained data** registration officers following the Registry’s standardised methodology, described in detail in its Annual Reports (National Suicide Research Foundation, 2009). This involves the classification of the method of self harm according to the ICD-10 codes for intentional injury (X60-X84). **Quality control measures include regular team meetings to reinforce the standardised application of**

Suicidal behavior in older Irish adults

case-definition and ascertainment criteria as well as cross-checking exercises which have shown high levels of agreement (Kappa statistic > 0.9) and reliability across registration officers. The Registry was granted ethical approval by the National Research Ethics Committee of the Faculty of Public Health Medicine.

Setting and population data

The Republic of Ireland, population 4.24 million according to the 2006 census, is made up of 26 counties and five cities. Dublin is by far the largest city and its expansion has urbanised all of Dublin county where 28% (1.19 million) of the country resides. The four other cities (Cork, Galway, Limerick and Waterford) account for 8% (0.29 million) of the country's population. There are 54 urban districts across the country accounting for almost 10% (0.40 million) of the population and the remainder of the country consists of rural districts where more than half of the country's population live (2.36 million, 56%).

Population data from the 1981, 1986, 1991, 1996, 2002 and 2006 national censuses and population estimates for intercensal years were obtained from the CSO website (<http://www.cso.ie/>). Small-area level population data were obtained from the 2002 and 2006 censuses and aggregated to provide figures for the population of Dublin, the other four Irish cities, the urban districts and the rural districts. Population estimates were extrapolated and interpolated for the intercensal years in the period 1997-2008 by assuming that the population change between the two censuses reflected a trend that applied across the twelve years.

Calculation of rates

Annual rates per 100,000 of male and female (aged over 55 years) suicide, suicide plus undetermined deaths and deliberate self harm were calculated and age-adjusted using the European standard population (Waterhouse *et al.*, 1976). Annual age-sex-specific rates were

Suicidal behavior in older Irish adults

calculated for each five-year age group. The calculation of incidence rates for deliberate self harm only considered one presentation per person per calendar year. For 3.5% of all deliberate self harm presentations by adults over 55 years of age, no address was recorded. A further 5.5% of presentations by residents of institutions such as prisons and hospitals, by homeless people and non-residents of the country were also not included in the rate calculations by geographic area.

To help assess rate differences, 95% confidence intervals (CIs) were calculated. Assuming that the number of deaths or persons presenting with deliberate self harm (x) followed a Poisson distribution, 95% CIs for the rates were calculated using the Normal approximation, i.e. $CI = (x \pm 2 \cdot \sqrt{x}) \cdot 100,000 / \text{population}$. The 95% CIs are presented as error bars in some charts.

Statistical analysis

Poisson regression analysis assessed the magnitude and statistical significance of rate differences and associations with sex, age group and area type. Whether associations differed due to an interaction factor was tested by fitting Poisson regression models with and then without the relevant interaction term and performing a likelihood ratio test (LRT). Rate differences assessed by Poisson regression were reported as incidence rate ratios (IRRs) with their 95% CIs and p-values. Poisson regression was carried out using Stata version 6.0 (StataCorp, 1999).

The ICD-9 external cause of death codes recorded in the mortality data for suicide and undetermined death and the ICD-10 intentional injury codes recorded in the Registry data for deliberate self harm were reclassified into comparable categories as follows: Drug overdose (ICD-9: E950.0-950.5, E980.0-980.5; ICD-10: X60-64); Poisoning, excluding alcohol (ICD-9: E950.6-950.9, E951-952, E980.6-980.9, E981-982; ICD-10: X66-69); Self-cutting or piercing (ICD-9: E956, E986; ICD-10: X78); Hanging or strangulation (ICD-9: E953, E983; ICD-10:

Suicidal behavior in older Irish adults

X70); Drowning (ICD-9: E954, E984; ICD-10: X71); Firearms (ICD-9: E955, E985; ICD-10: X72-74); Other (ICD-9: E957-959, E987-989; ICD-10: X75-77, X79-84). Chi-square tests were used to assess whether method of self harm differed in deliberate self harm and suicide and by gender. Whereas only one method of self harm was recorded in the mortality data, multiple methods were sometimes recorded for deliberate self harm.

All analyses were carried out for suicide deaths only and for suicide plus undetermined deaths. If it was not possible to present findings for both clearly, results from the latter were presented.

Results

Since 1980, the female suicide rate among over 55 year-olds in Ireland has been relatively stable whereas the male rate has fluctuated (Figure 1). It increased during most of the 1980s with a decrease of smaller magnitude in the 1990s and some evidence of a further decrease in recent years. The addition of deaths of undetermined intent made a significant difference to the male and female rates in the 1980s. During the 1990s, rates of undetermined death (UD) diminished greatly but were notable again by the end of that decade.

Figure 1 here

During the 10 years 1997-2006, there were 921 suicides (male: 691 (75.0%); female: 230 (25.0%)) and 175 deaths of undetermined intent (male: 110 (62.9%); female: 65 (37.1%)) among the population aged over 55 years in Ireland. The annual age-adjusted male suicide rate was 19.1 per 100,000, approximately three times the female rate (5.9 per 100,000; ratio = 3.2). The three-fold difference remained when UD's were added (male rate: 22.1 per 100,000; female rate: 7.6 per 100,000; ratio = 2.9).

Suicidal behavior in older Irish adults

In 2006-2008, 2,352 deliberate self harm presentations were made to hospital by over 55 year-olds, the majority by women (1,019 (43.3%) by men; 1,332 (56.7%) by women; sex unknown in one case). Respectively, 715, 813 and 824 deliberate self harm presentations were made in 2006, 2007 and 2008 by 592, 640 and 699 individuals. **The average number of presentations per person per year was 1.22 (maximum = 13). Only one presentation per year was made by 85.4% of patients, two presentations per year were made by 11.0%, three were made by 2.2% and more than three presentations were made by 1.4% of patients.** The annual age-adjusted male deliberate self harm rate was 67.4 per 100,000 (95% CI = 62.8-72.1 per 100,000). At 83.4 per 100,000 (95% CI = 78.8-88.1 per 100,000), the female rate was significantly higher (+24%).

The incidence of older male deliberate self harm was therefore 3.5 times the 19.1 per 100,000 rate of suicide and 3.0 times the 22.1 per 100,000 rate of suicide + UD. For older women, the incidence of deliberate self harm was 14.0 times the 5.9 per 100,000 rate of suicide and 11.0 times the 7.6 per 100,000 rate of suicide + UD.

With increasing age, there was a stepped decrease in the incidence of both suicide + UD and deliberate self harm (Figure 2). The age association with suicide + UD differed significantly from that with deliberate self harm (LRT chi-square = 24.14, df = 6, $p < 0.001$) because older age was associated with greater decrease in the incidence of deliberate self harm. The male rate of suicide + UD was significantly higher than the female rate in each five-year age group by a ratio of 2.6-4.7:1 (Table 1). **The male rate of deliberate self harm was 19-29% lower than the female rate in 55-59, 60-64 and 65-69 year-olds whereas among over 85 year-olds, the male rate was almost three times higher.**

Figure 2 here

Table 1 here

Suicidal behavior in older Irish adults

The method of self harm causing death differed significantly between men and women (Chi-square = 130.2, df = 5, p < 0.001; Table 2). Hanging was the most common method used by men (40.6%), followed by drowning (28.7%) which was the most common method used by women (39.3%). Drug overdose was the external cause of almost one in four female deaths compared to just 7.6% of male deaths. While relatively uncommon, 9.0% of the male deaths involved firearms as opposed to just one female death. A similarly significant gender difference was evident when only officially-classified suicides were considered (Chi-square = 123.2, df = 5, p < 0.001) with the most common methods used - hanging by men (46.5%) and drowning by women (45.2%) – even more pronounced.

Drug overdose was by far the most common method involved in hospital-treated self harm and especially so among women (Table 2; Chi-square = 56.18, df = 1, p < 0.001). The only other method that was relatively common was self-cutting and this was used by men more than women (Chi-square = 18.50, df = 1, p < 0.001). More lethal methods of self harm were rare but two were more commonly used by men (Hanging: chi-square = 6.26, df = 1, p = 0.012; Drowning: chi-square = 19.37, df = 1, p < 0.001). Other methods were also more often associated with male deliberate self harm (Chi-square = 13.46, df = 1, p < 0.001). In addition, alcohol was more often involved in male acts of deliberate self harm (Male: 45.9%; Female: 34.4%; Chi-square = 32.22, df = 1, p < 0.001).

Table 2 here

The methods of self harm involved in suicide + UD differed markedly from those involved in hospital-treated deliberate self harm. The latter involved a higher proportion of drug overdose and self-cutting whereas hanging, drowning, firearms and poisoning were significantly more common in acts of suicide + UD (all p < 0.001).

Suicidal behavior in older Irish adults

Figure 3 here

The incidence of suicide + UD varied markedly by geographic area (Figure 3). For both sexes, there were particularly high rates in the urban district populations, about twice the rate in rural districts (Table 3). The rate of suicide + UD was lower in Dublin for men only. Among women, the rate suicide + UD was significantly elevated in cities other than Dublin.

Table 3 here

There was also a marked urban-rural difference in the male and female incidence of hospital-treated deliberate self harm. Compared to rural districts, the male rate of self harm was twice as high in Dublin and 2.5 times higher in other cities and urban districts. Relative to rural districts, the female rate of deliberate self harm was almost twice as high in cities other than Dublin and about 2.5 times higher in Dublin and urban districts.

Discussion

The significance of suicidal behavior has become well recognised with many countries implementing national prevention strategies. Studies have found deliberate self harm in older adults generally involves greater suicidal intent (Hawton and Harriss, 2008b) than in the young and that the link between deliberate self harm and suicide is also stronger in older adults (Cooper *et al.*, 2005; Hawton and Harriss, 2006; Hawton *et al.*, 2003). However, the incidence of both forms of suicidal behavior have rarely been compared. To our knowledge, only one previous study has compared the incidence of deliberate self harm and suicide at a national level in older adults (Shah, 2009b).

Methodological issues

Suicidal behavior in older Irish adults

The national perspective of this study is one of its strengths **although there was limited overlap between the years with available suicide mortality data and deliberate self harm data.** Suicide and deliberate self harm in older adults have previously been compared in 10 urban centers across Europe (De Leo *et al.*, 2001) and in Western Australia (Lawrence *et al.*, 2000). The incidence of deliberate self harm reported in this paper was based on persons presenting to hospital emergency departments. This is an improvement on studies based on deliberate self harm that results in admission to a hospital ward (Shah, 2009a) because admission rates following self harm presentation vary greatly between hospitals (Bennewith *et al.*, 2004) thereby leading to significant bias. Our self harm data suffered some bias, however, as untreated deliberate self harm and that treated in other settings were unavailable.

The geographic comparisons made in this paper are an added strength and a rare feature of study of older adult suicidal behavior (Shah *et al.*, 2009). We distinguished between Dublin, other cities, urban districts and rural districts. While this independent stratification has justification, it classifies some Irish towns and suburban areas as rural which is a limitation. Furthermore, small-area postcodes do not exist in Ireland thereby hampering the geocoding of addresses. The mortality data, in particular, were geocoded with more limited resources. As a result, unreliable geocoding must be borne in mind as a potential contributor to the geographic differences in suicide rates illustrated by this study.

Suicide research is greatly hampered by the question of data reliability due to the difficulty inherent in establishing suicide as a cause of death (Claassen *et al.*, 2010). Misclassification of suicide deaths has most often been associated with the cause of death category ‘deaths of undetermined intent’ (Cooper and Milroy, 1995; Jouglu *et al.*, 2002; Ohberg and Lonnqvist, 1998). We carried out all analyses using data related to officially-classified suicides and suicides plus undetermined deaths in order to show that the findings were robust to the effects of misclassification. **However, adjustment was not made for misclassification to**

Suicidal behavior in older Irish adults

other causes of death including natural causes. This may be more common among older persons with severe and/or terminal illness who bring about their deaths through the cessation of vital medication.

Incidence rates

The incidence of deliberate self harm among older adults in Ireland is higher than that reported for a range of states in the USA (Center for Disease Control, 2007). Other comparable studies have had predominantly urban catchment areas. The Irish rates of deliberate self harm among older adults in urban districts and cities appear similar to those of Oxford, England (Hawton and Harriss, 2008a) which had previously ranked third highest of 15 European centers (De Leo *et al.*, 2001). Thus, hospital-treated deliberate self harm among older adults in Ireland is likely to be high in the European context.

Ireland's suicide rate among older adults is lower than most European countries. Older Irish suicide also has a different age association because the Irish rate decreases with age whereas the opposite is seen in many other European countries (http://www.who.int/mental_health/prevention/suicide/country_reports/en/index.html). Ireland and the UK are similar in terms of the age pattern in suicide rates but the peak in young men is more pronounced in Ireland. The rapid social change that Ireland has experienced in recent decades seems relevant. Older Irish adults lived most of their lives in an Ireland where the Catholic Church was dominant and suicide was strongly sanctioned religiously and legally. This may have created a cohort who associate such stigma and shame with suicide that their suicide rate is relatively low (Cleary and Brannick, 2007). Younger Irish adults have grown up in a secular Ireland where the Church has little or no (positive or negative) influence on their lives and suicide is far less stigmatised. In times of despair when older generations may have turned to the Church or their religion for support they do not and seem far more likely to resort to suicide.

Suicidal behavior in older Irish adults

The incidence of deliberate self harm among older adults also decreased with increasing age and at a greater rate than for suicide. Therefore, the ratio of non-fatal to fatal suicidal behavior was lowest among the very oldest as has been shown elsewhere (Hawton and Harriss, 2008b).

There was a 3:1 male:female rate ratio in relation to suicide deaths whereas the older female rate of deliberate self harm was 24% higher than the male rate. These gender differences are similar to those found previously across a range of European centers (De Leo *et al.*, 2001) although an absence of a gender difference in older adult deliberate self harm has recently been reported from Oxford, England (Hawton and Harriss, 2008a).

Geographic differences

We have shown striking geographic differences in suicide and deliberate self harm rates among older adults in Ireland. The latter had 2-2.5 times higher rates in the Irish city and urban district populations than in rural districts. Given that hospital emergency departments are situated in urban centers, proximity may partly explain the higher rate of self harm presentations by urban populations. However, we have previously shown evidence that the distribution of socioeconomic deprivation is strongly associated with the geographic differences in rates of deliberate self harm (Corcoran *et al.*, 2007). In relation to suicide, the highest rates were in urban districts and especially so for men. As mentioned above, unreliable geocoding of mortality data cannot be discounted as a potential confounder of these findings. However, the extent of the differences warrants further investigation. We observed lower incidence of suicide among older men in Dublin. An artefactual reason may be behind this finding as we previously showed the lower Dublin suicide rate to be due to deficits in the operation of the suicide recording system in Dublin (National Suicide Research Foundation, 2007).

*Suicidal behavior in older Irish adults**Methods of self harm*

Expectedly, the more lethal methods of self harm (hanging and drowning) were commonly involved in suicide deaths rather than in hospital-treated deliberate self harm. Hanging is the commonest method of suicide in most European countries but drowning is relatively rare (Biermann *et al.*, 2009; Chen *et al.*, 2009; Varnik *et al.*, 2008), whereas it was involved in about 30% and 40% of the older male and female suicide deaths in this study, respectively. Drug overdose was involved in the vast majority of deliberate self harm presentations and especially so in female acts. It was also involved in a significant minority of suicide deaths. Drug overdose is consistently found to be the predominant method used by older adults who self harm (Chan *et al.*, 2007; De Leo *et al.*, 2001; Shah, 2009a). While less common, self-cutting and hanging were more often associated with male than female deliberate self harm, as has been previously shown for older adults in Europe (De Leo *et al.*, 2001).

Conclusions

Older adults in Ireland have high rates of hospital-treated deliberate self harm but below average rates of suicide. Both forms of suicidal behavior decrease in incidence with increasing age, especially deliberate self harm. There are striking geographic differences with particularly high rates among urban populations that are worthy of further investigation. Drug overdose was involved in the vast majority of older adult deliberate self harm and a significant minority of suicides. Previous success in suicide prevention through means restriction (Daigle, 2005; Hawton *et al.*, 2009; Hawton *et al.*, 2004) suggests that restriction of specific medications may be an effective approach to the prevention of suicidal behavior in older Irish adults.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Conflicts of interest: None

Description of authors’ roles: All authors contributed to formulating the research questions. IJ Perry and P Corcoran designed the Registry upon which much of the study is based. P Corcoran carried out the statistical analysis and drafted the manuscript. U Reulbach drafted the Introduction. All authors contributed to completing the paper.

Acknowledgements

We would like to acknowledge the staff of the Irish Central Statistics Office for their work in compiling the mortality and population data used in this paper. We would like to thank all who contribute to the Irish National Registry of Deliberate Self Harm, in particular Ms Eileen Williamson, Executive Director, the data managers and data registration officers. The Registry is funded by the Irish Health Service Executive’s National Office for Suicide Prevention.

*Suicidal behavior in older Irish adults***References**

- Bennewith, O., Gunnell, D., Peters, T., Hawton, K. and House, A.** (2004). Variations in the hospital management of self harm in adults in England: observational study. *British Medical Journal*, 328, 1108-1109.
- Bertolote, J. M. and Fleischmann, A.** (2002). Suicide and psychiatric diagnosis: a worldwide perspective. *World Psychiatry*, 1, 181-185.
- Biermann, T., Sperling, W., Bleich, S., Kornhuber, J. and Reulbach, U.** (2009). Particularities of suicide in the elderly. A population-based study. *Aging Clinical Experimental Research*, 21, 470-474.
- Cattell, H. and Jolley, D. J.** (1995). One hundred cases of suicide in elderly people. *British Journal of Psychiatry*, 166, 451-457.
- Center for Disease Control** (2007). Nonfatal self-inflicted injuries among adults aged > or = 65 years--United States, 2005. *Morbidity Mortality Weekly Report*, 56, 989-993.
- Chan, J., Draper, B. and Banerjee, S.** (2007). Deliberate self-harm in older adults: a review of the literature from 1995 to 2004. *International Journal of Geriatric Psychiatry*, 22, 720-732.
- Chen, Y. Y., Park, N. S. and Lu, T. H.** (2009). Suicide methods used by women in Korea, sweden, taiwan and the United States. *Journal of the Formosan Medical Association*, 108, 452-459.
- Claassen, C. A., Yip, P. S., Corcoran, P., Bossarte, R. M., Lawrence, B. A. and Currier, G. W.** (2010). National suicide rates a century after Durkheim: A staged model to estimate mortality rate error. *Suicide and Life-Threatening Behavior*, (in press).
- Cleary, A. and Brannick, T.** (2007). Suicide and changing values and beliefs in Ireland. *Crisis*, 28, 82-88.
- Conwell, Y., Duberstein, P. R. and Caine, E. D.** (2002). Risk factors for suicide in later life. *Biological Psychiatry*, 52, 193-204.
- Cooper, J., et al.** (2005). Suicide after deliberate self-harm: a 4-year cohort study. *American Journal of Psychiatry*, 162, 297-303.

Suicidal behavior in older Irish adults

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Cooper, P. N. and Milroy, C. M. (1995). The coroner's system and under-reporting of suicide. *Medicine, Science and Law*, 35, 319-326.

Corcoran, P., Arensman, E. and Perry, I. J. (2007). The area-level association between hospital-treated deliberate self-harm, deprivation and social fragmentation in Ireland. *Journal of Epidemiology and Community Health*, 61, 1050-1055.

Corcoran, P., Keeley, H. S., O'Sullivan, M. and Perry, I. J. (2003). Parasuicide and suicide in the south-west of Ireland. *Irish Journal of Medical Science*, 172, 107-111.

Daigle, M. S. (2005). Suicide prevention through means restriction: assessing the risk of substitution. A critical review and synthesis. *Accident; Analysis and Prevention*, 37, 625-632.

De Leo, D., et al. (2001). Attempted and completed suicide in older subjects: results from the WHO/EURO Multicentre Study of Suicidal Behaviour. *Internation Journal of Geriatric Psychiatry*, 16, 300-310.

Dennis, M. S., Wakefield, P., Molloy, C., Andrews, H. and Friedman, T. (2007). A study of self-harm in older people: mental disorder, social factors and motives. *Aging and Mental Health*, 11, 520-525.

Hawton, K., et al. (2009). Effect of withdrawal of co-proxamol on prescribing and deaths from drug poisoning in England and Wales: time series analysis. *British Medical Journal*, 338, b2270.

Hawton, K. and Harriss, L. (2006). Deliberate self-harm in people aged 60 years and over: characteristics and outcome of a 20-year cohort. *International Journal of Geriatric Psychiatry*, 21, 572-581.

Hawton, K. and Harriss, L. (2008a). The changing gender ratio in occurrence of deliberate self-harm across the lifecycle. *Crisis*, 29, 4-10.

Hawton, K. and Harriss, L. (2008b). How often does deliberate self-harm occur relative to each suicide? A study of variations by gender and age. *Suicide and Life-Threatening Behavior*, 38, 650-660.

Hawton, K., et al. (2004). UK legislation on analgesic packs: before and after study of long term effect on poisonings. *British Medical Journal*, 329, 1076.

Suicidal behavior in older Irish adults

- Hawton, K. and van Heeringen, K.** (2009). Suicide. *Lancet*, 373, 1372-1381.
- Hawton, K., Zahl, D. and Weatherall, R.** (2003). Suicide following deliberate self-harm: long-term follow-up of patients who presented to a general hospital. *British Journal of Psychiatry*, 182, 537-542.
- Jougla, E., Pequignot, F., Chappert, J., Rossollin, F., Le Toullec, A. and Pavillon, G.** (2002). [Quality of suicide mortality data]. *Revue d'Epidemiologie et de Sante Publique*, 50, 49-62.
- Kaplan, M. S., Adamek, M. E. and Rhoades, J. A.** (1998). Prevention of elderly suicide. Physicians' assessment of firearm availability. *American Journal of Preventive Medicine*, 15, 60-64.
- Kapur, N.** (2006). Self-harm in the general hospital. *Clinical Medicine*, 6, 529-532.
- Kelleher, M. J., Keohane, B., Corcoran, P. and Keeley, H. S.** (1997). Elderly suicides in Ireland. *Irish Medical Journal*, 90, 72, 74.
- Lawrence, D., Almeida, O. P., Hulse, G. K., Jablensky, A. V. and Holman, C. D.** (2000). Suicide and attempted suicide among older adults in Western Australia. *Psychological Medicine*, 30, 813-821.
- National Suicide Research Foundation** (2007). Inquested deaths in Ireland: a study of routine data and recording practices. Cork: National Suicide Research Foundation.
- National Suicide Research Foundation** (2009). National Registry of Deliberate Self Harm Ireland Annual Report 2008. Cork: National Suicide Research Foundation.
- Ohberg, A. and Lonnqvist, J.** (1998). Suicides hidden among undetermined deaths. *Acta Psychiatrica Scandinavica*, 98, 214-218.
- Platt, S., et al.** (1992). Parasuicide in Europe: the WHO/EURO multicentre study on parasuicide. I. Introduction and preliminary analysis for 1989. *Acta Psychiatrica Scandinavica*, 85, 97-104.
- Reulbach, U. and Bleich, S.** (2008). Suicide risk after a suicide attempt. *British Medical Journal*, 337, a2512.

Suicidal behavior in older Irish adults

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Shah, A. (2009a). Attempted suicide in the elderly in England: age-associated rates, time trends and methods. *International Psychogeriatrics*, 1-7.

Shah, A. (2009b). Does improvement in the treatment of those who attempt suicide contribute to a reduction in elderly suicide rates in England? *International Psychogeriatrics*, 21, 768-773.

Shah, A., Sinha, T. and Makena, R. (2009). The relationship between elderly suicide rates, population density and room density. *International Psychogeriatrics*, 1-2.

StataCorp (1999). Stata Statistical Software: Release 6.0. College Station: Stata Corporation.

Varnik, A., et al. (2008). Suicide methods in Europe: a gender-specific analysis of countries participating in the "European Alliance Against Depression". *Journal of Epidemiology and Community Health*, 62, 545-551.

Waterhouse, J., Muir, C., Correa, P. and Powell, J. (1976). *Cancer incidence in five continents*. Lyon: IARC.


Suicidal behavior in older Irish adults

Figure 1. Trends in male and female suicide and suicide plus undetermined death (UD) among over 55 year-olds in Ireland, 1980-2006

Suicidal behavior in older Irish adults




Figure 2. Annual age-sex-specific rate of suicide and undetermined death and hospital-treated deliberate self harm among over 55 year-olds in Ireland

Suicidal behavior in older Irish adults

Table 1. Gender difference by age in the incidence of suicide and undetermined death and hospital-treated deliberate self harm among over 55 year-olds in Ireland

	Suicide and undetermined death, 1997-2006		Deliberate self harm, 2006-2008	
	IRR ¹	(95% CI)	IRR ¹	(95% CI)
55-59 years	2.84***	(2.23-3.61)	0.81**	(0.70-0.93)
60-64 years	2.91***	(2.22-3.81)	0.71***	(0.60-0.85)
65-69 years	2.59***	(1.88-3.55)	0.78*	(0.61-0.99)
70-74 years	3.23***	(2.17-4.82)	1.05	(0.76-1.44)
75-79 years	3.26***	(2.09-5.07)	0.70	(0.47-1.06)
80-84 years	4.68***	(2.42-9.03)	1.27	(0.73-2.22)
85 years+	3.49**	(1.51-8.07)	2.82**	(1.43-5.55)

¹ Incidence rate ratio of the male rate to the female rate.

Suicidal behavior in older Irish adults

Table 2. Methods of self harm involved in suicide and undetermined death and hospital-treated deliberate self harm among over 55 year-olds in Ireland

	Suicide and undetermined death				Deliberate self harm			
	Male		Female		Male		Female	
	n	(%)	n	(%)	n	(%)	n	(%)
Drug overdose	61	(7.6)	71	(24.1)	732	(71.8)	1126	(84.5)
Poisoning	65	(8.1)	17	(5.8)	29	(2.8)	36	(2.7)
Self-cutting	0	(0)	0	(0)	149	(14.6)	119	(8.9)
Hanging	325	(40.6)	51	(17.3)	36	(3.5)	25	(1.9)
Drowning	230	(28.7)	116	(39.3)	62	(6.1)	33	(2.5)
Firearms	72	(9.0)	1	(0.3)	4	(0.4)	1	(0.1)
Other	48	(6.0)	39	(13.2)	48	(4.7)	27	(2.0)

Note: Multiple methods of self harm were involved in some cases of deliberate self harm and therefore the percentages sum to more than 100%.


Suicidal behavior in older Irish adults

Figure 3. Annual age-adjusted rate of suicide and undetermined death and hospital-treated deliberate self harm among over 55 year-olds by area type in Ireland

Suicidal behavior in older Irish adults

Table 3. Geographic variation in the incidence of suicide and undetermined death and hospital-treated deliberate self harm among over 55 year-olds in Ireland.

	Suicide and undetermined death, 1997-2006				Deliberate self harm, 2006-2008			
	Male		Female		Male		Female	
	IRR ¹	(95% CI)	IRR ¹	(95% CI)	IRR ¹	(95% CI)	IRR ¹	(95% CI)
Dublin	0.75**	(0.62-0.90)	1.19	(0.90-1.58)	1.96***	(1.66-2.31)	2.32***	(2.01-2.68)
Other cities	1.07	(0.81-1.40)	1.61*	(1.08-2.40)	2.54***	(1.99-3.22)	1.83***	(1.43-2.32)
Urban districts	1.85***	(1.55-2.22)	2.11***	(1.55-2.86)	2.46***	(1.98-3.04)	2.57***	(2.13-3.10)
Rural districts	1.00	---	1.00	---	1.00	---	1.00	---

¹ Incidence rate ratio of age-adjusted rates with rural districts as the reference group