

Title	Enhancing the stress responses of probiotics for a lifestyle from gut to product and back again
Authors	Mills, S.;Stanton, Catherine;Fitzgerald, Gerald F.;Ross, R. Paul
Publication date	2011-08
Original Citation	Mills, S., Stanton, C., Fitzgerald, G. F. and Ross, R.P. (2011) Enhancing the Stress Responses of Probiotics for a Lifestyle from Gut to Product and Back Again. Microbial Cell Factories 10(Suppl 1):S19
Type of publication	Article (peer-reviewed)
Link to publisher's version	http://www.microbialcellfactories.com/content/10/S1/S19 - 10.1186/1475-2859-10-S1-S19
Rights	© 2011 Mills et al; licensee BioMed Central Ltd. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/2.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. - http://creativecommons.org/licenses/by/2.0
Download date	2026-08-29 13:24:15
Item downloaded from	https://hdl.handle.net/10468/1031



UCC

University College Cork, Ireland
 Coláiste na hOllscoile Corcaigh

PROCEEDINGS

Open Access

Enhancing the stress responses of probiotics for a lifestyle from gut to product and back again

Susan Mills¹, Catherine Stanton^{1,3}, Gerald F Fitzgerald^{2,3}, R Paul Ross^{1,3*}

From 10th Symposium on Lactic Acid Bacterium

Egmond aan Zee, the Netherlands. 28 August - 1 September 2011

Abstract

Before a probiotic bacterium can even begin to fulfill its biological role, it must survive a battery of environmental stresses imposed during food processing and passage through the gastrointestinal tract (GIT). Food processing stresses include extremes in temperature, as well as osmotic, oxidative and food matrix stresses. Passage through the GIT is a hazardous journey for any bacteria with deleterious lows in pH encountered in the stomach to the detergent-like properties of bile in the duodenum. However, bacteria are equipped with an array of defense mechanisms to counteract intracellular damage or to enhance the robustness of the cell to withstand lethal external environments. Understanding these mechanisms in probiotic bacteria and indeed other bacterial groups has resulted in the development of a molecular toolbox to augment the technological and gastrointestinal performance of probiotics. This has been greatly aided by studies which examine the global cellular responses to stress highlighting distinct regulatory networks and which also identify novel mechanisms used by cells to cope with hazardous environments. This review highlights the latest studies which have exploited the bacterial stress response with a view to producing next-generation probiotic cultures and highlights the significance of studies which view the global bacterial stress response from an integrative systems biology perspective.

Introduction

To suggest that probiotic bacteria have entered the realm of the highly exploited bacterial groups is an understatement. Moreover, man's expectations of probiotic bacteria have perhaps become the most demanding for any bacterial group to date. With numerous health-promoting attributes, many of which have been scientifically validated in animal and human clinical trials, probiotic bacteria have become an integral element to everyday healthy living. Indeed, the global market for probiotics is estimated to exceed US\$28.8 billion by 2015 [1]. Probiotic bacteria generally execute their biological role in the gut, which can involve a plethora of effects from immunomodulation to production of bioactive metabolites, all of which have dramatic consequences for disease evasion [2-4]. The present target of any probiotic food product in terms of probiotic cell numbers is to have up to 10^7 colony forming units

(cfu)/g at the end of its shelf life [5], although it should be emphasized that this will probably become strain and application specific as new clinical evidence emerges supporting health claims. Thus, before probiotic bacteria can even begin to fulfill their physiological role in the gut, the bacteria themselves must endure a number of stresses to ensure they reach the target site in adequate numbers to elicit an effect. These stresses can be considered in two distinct contexts; firstly, probiotic bacteria must be processed in suitable form to enable oral consumption and secondly the bacteria must be able to withstand the harsh conditions imposed during passage through the gastrointestinal tract (GIT).

From gut...

Potential probiotic cultures have been isolated from a variety of sources including animal, human and food sources. However, there is now growing evidence that strains are host specific [6] and for that reason it is generally recommended that human isolates should be used for human applications as a starting point. In this

* Correspondence: paul.ross@teagasc.ie

¹Teagasc Food Research Centre, Moorepark, Fermoy, Co. Cork, Ireland
Full list of author information is available at the end of the article

respect human strains have been isolated from various sites from oral to stool samples.

To product...

In terms of processing, probiotics are commonly grown to high numbers before undergoing a drying process to produce a high-cell density probiotic powder. This can then be added to a specific carrier, such as a dairy product, which is stored under suitable conditions. Probiotic robustness can be compromised even in the initial growing stages. Indeed, Muller et al. [7] recently demonstrated that the nutrient composition of the growth medium can have a significant effect on the technological properties of probiotics by altering membrane composition and morphology. The presence of linoleic and linolenic acids in a minimal growth medium reduced the acid and heat tolerance of *Lactobacillus johnsonii* NCC533 by 6- and 20-fold, respectively [7]. The stresses encountered during drying include extremes in temperatures, from very high temperatures in spray-drying, to very low-temperatures during freeze drying and storage, as well as osmotic and oxidative stresses. During spray-drying temperatures can reach as high as 200°C [8] and while the exposure time for the cells is extremely short, the integrity of viable bacterial cells can be severely compromised. The cytoplasmic membrane is the primary target for heat damage, resulting in damage to fatty acids and aggregation of proteins, however, intracellular proteins, ribosomes and RNA can also be duly impaired [9,10]. In addition to heat stress, spray-drying also exposes cells to osmotic stress, dehydration and oxidative stress [10]. At the other end of the scale, cells are typically frozen at -196°C during freeze-drying, and then dried by sublimation under high vacuum [11-13]. Freeze-drying is less harsh on cells than spray-drying resulting in higher cell viability [14]. However, the low temperature still compromises cellular integrity where the main consequences include reduction in membrane fluidity, increases in the rate of DNA strand breakage, stabilization of RNA and DNA secondary structures which in turn alters transcription, translation and replication [15]. Ribosome functions, protein folding and enzyme activity are also disturbed at low temperatures [16,17]. Oxygen stress can affect probiotic bacteria at various stages of their production, from fermentation to drying and even storage etc. While oxygen itself does not damage the cells, its partial reduction to water generates reactive oxygen species (the superoxide anion radical O_2^- , the hydroxyl radical $OH^\bullet$, and hydrogen peroxide H_2O_2) which can lethally damage proteins, lipids and DNA [5,18]. Cysteine in the active sites of enzymes can also be a target for oxidation [19]. Oxygen sensitivity can be particularly problematic for bifidobacteria, although variations have been observed across this

species in terms of oxygen tolerance [20,21]. Indeed, two species isolated from porcine caecum, *Bifidobacterium aerophilum* and *Bifidobacterium psychroaerophilum*, were found to be aerotolerant with the ability to grow on agar-medium under aerobic conditions [22]. Certain strategies have proven successful in terms of reducing the effects of oxygen toxicity. For example, electro-reduction, de-aeration or addition of reducing agents to non-fermented pasteurized milk was shown to enhance the viability of bifidobacteria during extended storage [23]. In fermented milk, co-cultivation of *Bifidobacterium longum* BB536 with *Lactococcus lactis* ssp. *lactis* MCC866, improved the survival of the bifidobacterial strain [24]. In this case, *L. lactis* ssp. *lactis* MCC866 was found to express several genes involved in protection against active oxygen species at higher levels during refrigerated storage, compared to non-effective lactococcal strains. Moreover, the food matrix itself can impose multiple stresses on bacterial survival. For example, several genes contributing to stress-related functions were induced in the commercial meat starter *Lactobacillus sakei* 23K during meat fermentation [25]. The global transcriptional response of *Lactobacillus reuteri* ATCC 55730 to the sourdough environment also revealed the up-regulation of various stress-related genes during sourdough fermentation [26]. Probiotic viability in Cheddar cheese was found to be markedly reduced when cells were added just before cheddaring but cell losses were halved when the probiotic was added to milk before renneting [27]. Moreover, in the same study salt was found to negatively compromise cell viability. However, food matrix stresses may also contribute to the overall survival of probiotic bacteria. Leverrier et al. [28] demonstrated that freeze-dried propionibacteria included at the beginning of milk fermentation were much more robust when exposed to acid and bile salt stresses as opposed to reconstituted freeze-dried cells which were dramatically injured under the same stress conditions.

And back again...

Passage of probiotics through the mammalian GIT is a hazardous journey, with the initial stages designed to jeopardize the survival of pathogenic microorganisms. The principal stresses include shifting pH encountered in the stomach resulting from gastric acid, as well as bile, a digestive secretion of the hepatic system which serves to emulsify and solubilize lipids and lipid soluble vitamins for metabolism, and steep gradients of oxygen. Exposure to acid negatively affects the proton motive force (PMF) across the membrane as a result of the accumulation of protons inside the cell [29]. In addition to cell membrane structural damage, acid stress causes damage to DNA and proteins [15]. Exposure to bile

disrupts the integrity of the cell membrane, affects DNA, RNA structure formation as well as protein folding and exposes the cell to oxidative stress and low intracellular pH [30,31].

The probiotic stress response

Bacterial cells are naturally equipped with a plethora of defense mechanisms to enhance survival in hostile environments [15,29,32-34]. These include chaperone proteins which assist the folding of misfolded proteins, proteases which degrade proteins which are irreversibly damaged, transport systems to maintain correct osmolarity, catalases and superoxide dismutases to tackle reactive oxygen species, as well as proton pumps, decarboxylases and transporters to combat decreases in intracellular pH [29] (Table 1, Figure 1). Understanding the intricacies of these systems provides scientists with a plethora of molecular tools to improve bacterial endurance. For example, a recent patent was filed describing numerous stress-related proteins of the probiotic strain *Lactobacillus acidophilus* and genetic engineering approaches which may be used to improve the stress response of the strain [35].

The aim of this review is to highlight the latest studies which take advantage of the bacterial stress response to produce 'super-fit' bacteria through genetic manipulation. Moreover, the upsurge in genome sequencing alongside systems biology approaches is now enabling scientists to view the global cellular response to the stresses encountered during processing and transit in the gut. As well as providing a deeper understanding of the probiotic stress response, this information is generating a wealth of novel molecular tools, which may find themselves as central players in the science of probiotic enhancement in the future.

Taking advantage of the bacterial stress response - genetic manipulation

Exploiting probiotic stress responses

It is well accepted that adaptation to a sub-lethal dose of a specific physical or chemical stress can dramatically improve subsequent performance in compromising environments and as a consequence is a popular strategy to increase both the technological and gastrointestinal robustness of a strain. The success of this strategy is owed to the induction of the bacterial stress response during the adaptation process whereby a specific system may be induced with the result that the cells survive a previously lethal dose of the same stress, or a more general system may be targeted enabling cross-protection against a range of stresses. For example, in order to improve the viability of the probiotic strain *Lactobacillus paracasei* NFBC 338 during spray-drying Desmond et al. [36] demonstrated that pre-stressing the culture by

exposure to 52°C for 15 minutes improved survival of the strain 700-fold (in reconstituted skim milk) during heat stress and 18 fold during spray-drying compared to un-adapted cells. Exposure to salt stress also afforded a level of thermotolerance. Indeed, following exposure to 0.3 M NaCl survival of the strain improved by 16-fold during spray-drying [36]. A proteomics analysis of the biological response of *Lb. paracasei* NFBC 338 to heat adaptation revealed an altered level of expression of at least 12 proteins, where expression of the chaperone GroEL was most dramatically induced in the test culture [37]. GroEL works in concert with its co-chaperone GroES where the complex ensures the correct folding of proteins in an ATP-regulated manner under normal growth conditions and under conditions of stress [29]. The genes encoding GroESL are negatively regulated by HrcA, which binds to a palindromic operator sequence called CIRCE (controlling inverted repeat of chaperone expression). Subsequently, overexpression of the *groESL* operon of *Lb. paracasei* NFBC 338 using the nisin controlled expression system in *Lb. paracasei* resulted in increased survival during exposure to stressful levels of heat, salt and butanol [37]. Moreover, *Lb. paracasei* over-expressing GroESL exhibited 10-fold better survival during spray-drying and 2-fold better survival during freeze-drying demonstrating that the GroESL chaperone is an effective molecular target for enhancing the technological performance of probiotic lactobacilli during spray-drying [5] (Figure 2). Interestingly, mRNA transcripts of the chaperones GroEL and DnaK were recently up-regulated 2-fold by down-regulating the negative regulator HrcA using anti-sense RNA technology [Kearney et al. unpublished]. However, although the anti-sense strain exhibited an increased exponential growth rate compared to the control, the anti-sense strain remained as sensitive as the control strain to heat, acid, bile solvent and osmotic stresses.

The ATP-independent chaperones, the small heat shock proteins (sHsps), have also been associated with enhanced bacterial survival during stress. While these proteins are not involved in protein re-folding they are necessary for normal cellular functions including growth, stability of DNA and RNA and they prevent the formation of inclusion bodies [38-40]. Heat, cold and ethanol stresses were previously shown to enhance the expression of the heat shock genes *hsp18.5*, *hsp19.3* and *hsp18.55* encoded on the genome of *Lactobacillus plantarum* [41,42]. Interestingly, *hsp18.5* and *hsp19.3* genes were preceded by an inverted repeat at the 5' end with homology to the operator sequence CIRCE [41] whereas the 5' region preceding *hsp18.55* revealed the presence of putative *cis* elements able to bind alternative sigma factor σ^B [42]. Inactivation of *hsp18.55* revealed that the protein encoded by the gene may be involved in the

Table 1 Examples of Proteins and Genes Involved in the Stress Responses of Probiotics

Stress	Protein/Gene/System (General Role/Description)	Fold Induction	Strain	Ref
Heat	GroEL (chaperone protein)	+49.1	<i>Lactobacillus paracasei</i> NFBC338	[37]
Heat	GroEL (chaperone protein)	+15	<i>Lactobacillus rhamnosus</i> HN001	[118]
Osmotic		+3		
Heat	<i>clpL1</i> (Clp ATPase family, members act as chaperones and regulators of proteolysis)	+20	<i>Lactobacillus rhamnosus</i> E800	[119]
Heat, Cold, Ethanol	<i>hsp18.5</i> (heat shock protein) <i>hsp19.3</i> (heat shock protein)	nd	<i>Lactobacillus plantarum</i> Lp90	[41]
Heat, Cold, Ethanol	<i>hsp18.55</i> (heat shock protein)	nd	<i>Lactobacillus plantarum</i> Lp90	[42]
Heat	DnaK (chaperone protein)	+4.4	<i>Lactobacillus gasseri</i> ATCC 33323	[120]
	GroEL (chaperone protein)	+3.8		
	ClpE (Clp ATPase family, members act as chaperones and regulators of proteolysis)	+1.7		
Heat	FtsH (protease and chaperone activity)	+8	<i>Lactobacillus plantarum</i> WCFS1	[121]
Heat	HtrA (protease and chaperone activity)	+10 +15	<i>Bifidobacterium longum</i> 3A	[122]
	GroEL (chaperone protein)	+5 +10		
	DnaK (chaperone protein)	+10 +15		
	GrpE (chaperone protein)	+2 +5		
Heat	<i>groEL-cspA</i> , (chaperone protein-cold shock protein) <i>groES</i> (chaperone protein)	+12 +8	<i>Bifidobacterium breve</i> UCC 2003	[123]
Heat	<i>clpC</i> (Clp ATPase family, members act as chaperones and regulators of proteolysis)	+15	<i>Bifidobacterium breve</i> UCC 2003	[124]
Heat	<i>clpP</i> operon: <i>clpP1</i> , <i>clpP2</i> (Clp ATPase family, members act as chaperones and regulators of proteolysis)	+15	<i>Bifidobacterium breve</i> UCC 2003	[125]
Heat	<i>hsp20</i> (heat shock protein)	+28	<i>Bifidobacterium breve</i> UCC 2003	[126]
Osmotic		+25		
Temperature downshift	<i>cspA</i> (cold shock protein)	+20	<i>Lactobacillus bulgaricus</i> V1104	[127]
Cold	ClpP (Clp ATPase family, members act as chaperones and regulators of proteolysis) Pyruvate kinase Glycoprotein endopeptidase	≥2	<i>Lactobacillus acidophilus</i> RD758	[128]
Temperature downshift	<i>cspL</i> (cold shock protein)	+17	<i>Lactobacillus plantarum</i> NC8	[46]
Osmotic	<i>dnaK</i> operon: <i>dnaK</i> , <i>grpE</i> , <i>dnaJ</i> (chaperones), ORF1 (similarity to predicted heat-controlled transcriptional regulators of Mer family)	+15, +14	<i>Bifidobacterium breve</i> UCC 2003	[129]
Osmotic	<i>dnaJ₂</i> (chaperone)	+4.5	<i>Bifidobacterium breve</i> UCC 2003	[130]
Oxygen	NADH oxidase & NADH peroxidase		<i>Lactobacillus acidophilus</i> 2400 <i>Lactobacillus acidophilus</i> 2409 <i>Bifidobacterium lactis</i> 1941 <i>Bifidobacterium pseudolongum</i> 1944 <i>Bifidobacterium longum</i> 1944	[103]
Oxygen	AhpC (alkyl hydroperoxide reductase C22) PNDR (Pyridine nucleotide-disulfide reductase) Dps (DNA-binding ferritin-like protein) NrdA (ribonucleotide reductase) MutT1 (NTP phosphohydrolases)	+10 +2 +1.7 +2 +1.8	<i>Bifidobacterium longum</i> BBMN68	[104]
Bile	HtrA (protease and chaperone activity) DnaK (chaperone protein) GroEL (chaperone protein)	+1.5+2 +1.5+2 +1.5+2	<i>Bifidobacterium longum</i> 3A	[122]
Bile	Bsh1 (bile salt hydrolase)		<i>Lactobacillus plantarum</i> WCFS1	[131]
Bile	Ctr (cholate transporter)		<i>Bifidobacterium longum</i> NCIMB 702259T	[132]
Acid	<i>clpL</i> (Clp ATPase family, members act as chaperones and regulators of proteolysis) Ir1516 (putative cell wall-altering esterase)	+2+3 >5	<i>Lactobacillus reuteri</i> ATCC 55730	[133]
Acid	GrpE (chaperone protein)	>3.5	<i>Lactobacillus sanfranciscensis</i> CB1	[134]
Acid	<i>LBA1524HK</i> (histidine kinase)		<i>Lactobacillus acidophilus</i> NCFM	[135]
Acid	<i>F₀F₁</i> -ATPase operon (involved in ATP synthesis and proton extrusion)	+2	<i>Lactobacillus acidophilus</i> NCFM/N2	[136]
Acid	<i>F₀F₁</i> -ATPase operon (involved in ATP synthesis and proton extrusion)	+15	<i>Bifidobacterium lactis</i> DSM10140	[137]




Figure 1 Diagrammatic representation of the main stress responses to acid pH and bile salt in bifidobacteria. (1) Conjugated bile acids enter the bacterial cytoplasm and are cleaved by BSH (bile salt hydrolase) (2) releasing one amino acid and one de-conjugated bile acid moiety. (3) De-conjugated bile acid can also enter the cell by passive diffusion and becomes deprotonated (4). (5) Ionized bile salts are non-permeable and are excreted by the action of certain transporters e.g. Ctr (cholate transporter) of *Bifidobacterium longum*. (6) Synthesis of molecular chaperones is also increased and a shift in fatty acid composition of cell membrane can occur (7). Exposure to acid pH or bile salt deprotonation results in acidification of the cytoplasm (8). This can be counteracted by the production of ammonia from glutamine deamination (9) or pumping of protons from the cell by the F_1F_0 -ATPase (10). ATP required for driving these systems is generated through glycolysis (11) (reproduced from Sanchez et al. [138]).




Figure 2 Survival of *Lb. paracasei* NFBC 338 (control) and GroESL-overproducing *Lb. paracasei* NFBC 338 (GroESL) following spray-drying (a) and freeze-drying (b). White bars represent powders plated on MRS and shaded bars represent powders plated on MRS containing NaCl (5% wt/vol). Insets show transmission electron micrographs of *Lb. paracasei* NFBC 338 (1) and GroESL-overproducing *Lb. paracasei* NFBC 338 (2) following spray-drying, and *Lb. paracasei* NFBC 338 (3) and GroESL-overproducing *Lb. paracasei* NFBC 338 (4) following freeze-drying. GroESL-overproducing *Lb. paracasei* NFBC 338 appeared to be more robust following both drying procedures compared to the control strain (adapted from Corcoran et al. [5]).

recovery of stressed cells during the early stage of high temperature stress [43]. Indeed, the $\Delta hsp18.55$ mutant strain displayed a longer lag phase under conditions of short intense heat stress (50°C). Physical parameters were also disturbed in the mutant strain including membrane fluidity and surface properties. Overproduction of each of the three heat shock proteins in *Lb. plantarum* improved the growth of the genetically-modified cultures during heat shock (37°C or 40°C) and cold shock (12°C) and overproduction of Hsp18.55 and Hsp19.3 also improved survival during solvent stress (1% v/v butanol or 12% v/v ethanol) [44].

Cold shock proteins (Csps) are induced as a result of cold shock pre-treatment where they have been associated with the stabilisation of mRNA [45]. Three induced Csps were previously identified in *Lb. plantarum* following cold shock treatment, namely CspL, CspP, CspC [46]. Overproducing each of the three proteins in *Lb. plantarum* resulted in improved performance during treatments involving a down-shift in temperature and under conditions of nutritional deprivation [47]. For example, overproduction of CspC led to faster growth when stationary phase cells were added to fresh medium when compared to controls, suggesting a role for CspC in nutrient adaptation. Cells overproducing CspL did not suffer a reduction in growth rate following exposure of exponentially growing cells to cold shock (8°C), while overproduction of CspP resulted in enhanced survival during 6 cycles of freeze-thawing at -80°C resulting in a 4-fold survival level.

Exploiting stress responses of pathogenic microorganisms

Patho-biotechnology is a relatively new concept which aims to exploit stress response systems of pathogenic bacteria for biotechnological and biomedical purposes [48]. Indeed, pathogenic bacteria can be a rich reservoir of sophisticated mechanisms which ensure bacterial endurance in some of the most stressful environments including the host immune system or those encountered during food processing. For example, certain bacterial species accumulate protective compounds called compatible solutes as part of the stress response [49]. These compounds are generally highly soluble with no net charge at physiological pH [50]. Within the cells they serve as osmotic balancers [51] but also serve to protect enzyme function against the stresses of high temperature, salinity, freeze-thawing and drying [52,53]. The trimethylammonium compound glycine betaine is used as a compatible solute by the majority of bacteria including *Listeria monocytogenes* [54]. In this respect, the secondary glycine betaine transporter of *L. monocytogenes*, BetL, has been shown to improve survival of the strain in certain foods [55], as well as improve its osmotolerance, cryotolerance and barotolerance [56-58].

Interestingly, heterologous expression of BetL from *L. monocytogenes* in the probiotic strain *Lactobacillus salivarius* UCC118 using the nisin controlled expression system resulted in a genetically modified strain with improved resistance to numerous stresses [54]. The BetL⁺ strain accumulated 65 times more betaine than the BetL⁻ strain and was capable of growth in 7% NaCl, unlike the control which was unable to grow efficiently at the same concentration. Repeated freeze-thaw cycles at -20°C or -70°C demonstrated that the BetL⁺ strain was two logs greater than the control at -20°C and 0.5 logs greater at -70°C. The barotolerance of the test strain also improved yielding a 2-log increase over the control strain following exposure to 350 MPa. Moreover, the BetL⁺ strain survived freeze-drying and spray-drying much more efficiently than the control with a 36% survival level for test versus 18% for control following freeze-drying and a 1.4% survival level for test versus 0.3% for control after spray-drying. Cloning the *L. monocytogenes* BetL system into the probiotic strain *Bifidobacterium breve* UCC2003 significantly improved the tolerance of the strain to gastric juice and elevated osmolarity [59]. In addition, BetL⁺*B. breve* was recovered at much higher levels from the faeces, intestines and caecum of BetL⁺ inoculated mice compared to control mice inoculated with the BetL⁻ strain, and *Listeria*-challenge experiments revealed that BetL⁺ mice exhibited reduced levels of *Listeria* infection in the spleen compared to controls [59].

The marked ability of *L. monocytogenes* to survive the stresses encountered in the upper small intestine has been linked to its capacity to endure high bile concentrations. Indeed, *L. monocytogenes* has been isolated from the human gallbladder [60,61]. A novel bile resistance mechanism, termed BilE was identified in *L. monocytogenes* and operates by excluding bile from the cell [62]. Heterologous expression of BilE in *L. lactis* NZ9800 and *B. breve* UCC2003 resulted in modified strains with enhanced capacities to survive in the presence of bile [63]. For example, the *bilE*-containing strain of *L. lactis* exhibited a 2.5 log enhanced resistance (compared to control) following a 20 minute exposure to 1% porcine bile, a similar result was observed for the *bilE*-containing strain of *B. breve*. In addition, both genetically modified strains exhibited an enhanced ability to survive *in vivo* conditions using mouse models. This was particularly noteworthy for the *bilE*-containing *L. lactis* strain considering that it is generally not considered to be a gut inhabitant due to its poor survival rates in the GIT. Indeed, the BilE⁺*L. lactis* strain was detected at high levels in murine faeces 3 days after inoculation whereas the BilE⁻*L. lactis* strain was undetectable in the faecal samples on day 2 after inoculation. While both BilE⁺ and BilE⁻ strains of *B. breve* colonised

the murine GI tract at relatively similar rates (10^7 cfu/g of faeces), the persistence of the BiE⁺ strain began to differ significantly 12 days after inoculation. Indeed, by day 19 the BiE⁻ strain had reached a level of 1×10^5 cfu/g of faeces, whereas the BiE⁺ strain persisted at a level of 4.5×10^7 cfu/g of faeces. Moreover, by day 19 the levels of the BiE⁺ strain in the small intestine, caecum and large intestine were significantly higher compared to the control strain. This was a particularly important finding for the small intestine, a region of the GIT which has the highest levels of bile [64]. Indeed, the persistence levels of the genetically modified strain were 100 fold greater than the persistence levels of the control. The BiE⁺ strain also enhanced clearance of *L. monocytogenes* from the liver at significant levels when compared to the control strain.

Exploiting generic microbial mechanisms

Improving probiotic robustness can also be achieved by looking to other microbial mechanisms which may not be necessarily associated with pathogenic survival. For example, various studies have demonstrated that probiotic robustness can be significantly enhanced by adding a protectant to the growth medium prior to exposure of the strain to environmental stress. An exudate gum from trees, gum acacia, enhanced survival of *Lb. paracasei* NFBC 338 from the stresses of heat, bile and H₂O₂ when added to the growth medium and also enhanced survival during spray-drying [65]. We proposed that the protective effect of gum acacia may be related to cellular encapsulation by the polysaccharide thus protecting the cells from the harsh environmental conditions. Many microorganisms produce exopolysaccharides (EPS) which can be either excreted into the medium or form a capsule around the cell, the capsular polysaccharides (CPS). Recently, a positive correlation was observed between EPS production and resistance to bile salt and low pH stress in *Bifidobacterium* species isolated from breast milk and infant faeces [66]. Interestingly, both the technological and gastrointestinal durability of *Lb. paracasei* NFBC 338 was significantly improved by equipping the strain with an inherent ability to produce the EPS beta-glucan [67]. The gene encoding this particular EPS, the membrane-associated glycosyltransferase enzyme (*gtf*) comes from *Pediococcus parvulus* 2.6. Indeed, heterologous expression of the gene in *Lb. paracasei* increased its heat tolerance 60-fold, its acid-tolerance 20-fold, its ability to survive in simulated gastric juice by 15-fold and its ability to survive in bile by 5.5 fold, compared to the control strain (Figure 3). Moreover, beta-glucan has been linked with many health-promoting properties including an ability to lower serum cholesterol levels [68,69], immunomodulation [70,71], anticancer properties [72] and it behaves as a prebiotic [73].

Thus, as well as enhancing the gastrointestinal robustness of the strain Stack et al., [67] simultaneously improved its health-promoting properties.

Other microbial mechanisms have also proven worthwhile as tools for probiotic enhancement. Heterologous expression of the manganese superoxide dismutase gene (*sodA*) from *Streptococcus thermophilus* in the intestinal strains *Lactobacillus gasseri* and *Lb. acidophilus* enabled the cells to tolerate greater concentrations of H₂O₂ (up to 1.6 mM) compared to the control which could not adequately tolerate concentrations greater than 1.2 mM [74]. Integration of *sodA* into the chromosome of *Lb. gasseri* also dramatically improved its oxidative tolerance, whereby the modified strain was capable of surviving up to 45 mM H₂O₂ after 90 minutes of exposure, unlike the control strain which was unable to tolerate concentrations greater than 25 mM [75]. Moreover, cloning the catalase gene *kata* from *Lb. sakei* SR911 into catalase-deficient *Lb. plantarum* TISTR850 under a strong lactococcal promoter improved the oxidative tolerance of the modified strain markedly [76]. Indeed, the modified strain survived 6 logs better than the control following 60 hours of growth under oxidative stress. Likewise, cloning *kata* from *Lb. sakei* into *Lactobacillus rhamnosus* AS 1.2466 significantly improved the oxidative resistance of the recombinant strain [77]. The exponential-phase culture of the modified strain increased 600-fold following H₂O₂ challenge and the stationary-phase culture increased 1000-fold under the same conditions when compared to the control. Endowing the probiotic strain *Lactobacillus casei* with both *kata* (from *Lb. sakei*) and the bile salt hydrolase gene *bsh1* (from *Lb. plantarum*) dramatically enhanced both the oxidative tolerance and the bile salt resistance of the strain [78]. The survival ratio of the recombinant strain was 40-fold greater than the control after 8 hours of incubation in the presence of 8 mM H₂O₂, while the recombinant strain survived in the presence of 0.5% glycodeoxycholate reaching 10^5 cfu/ml, a concentration which killed the control cells.

New players involved in the probiotic stress response

Proteomics and whole genome DNA microarrays alongside heterologous expression studies and the generation of deletion mutants continue to provide important insights into the response and adaptation of probiotic bacteria to environmental stresses. Indeed, studies which employ such techniques continually highlight the importance of the main molecular defense mechanisms, reveal stress-associated regulatory networks, but also provide insight into novel systems which serve to protect the stressed cell providing new molecular tools for probiotic enhancement. A global analysis of the transcriptomes of




Figure 3 Survival of *Lb. paracasei* NFBC 338 expressing the *gtf* gene (GTF⁺) and control strain in the presence of acid, gastric juice and elevated temperature. Inset displays the loop-touch test of *Lb. paracasei* NFBC 338 producing EPS demonstrating the 'ropy' phenotype of the modified strain (adapted from Stack et al. [67]).

two heat-shock tolerant strains of *B. longum* (isolated following serial passages at heat-shock temperatures) demonstrated that overexpression of the *dnaK* operon and *clpB* protease-encoding gene was linked to point mutations in the gene encoding the negative regulator, HspR [79]. Indeed, in one mutant a tyrosine residue was replaced by a histidine in the helix-turn-helix domain of the regulator, while in the second mutant a tyrosine residue was replaced with a cysteine in the winged 1 motif of the protein. Sensitivity to heat was restored by complementing the mutant strains with the wild-type *hspR* gene. Interestingly, the arginine deiminase (ADI) pathway of *Lactobacillus fermentum* IMDO 130101, a strain isolated from sourdough, was shown to respond to salt and temperature stresses resulting in an increase in ornithine production [80]. The ADI pathway functions to produce extra ATP and aid with acid stress

[80]. The authors suggest that the main function of the ADI pathway at high temperature is to provide energy to stationary phase cells and improve energy generation for growing cells under osmotic stress.

Heterologous expression of myosin-cross reactive antigen (MCRA), from *B. breve* NCIMB 702258, in *Lactococcus* and *Corynebacterium* improved sensitivity of both recombinant strains to heat and solvent stresses suggesting that this FAD-dependent fatty acid hydratase is involved in protecting the cell from environmental stress [81]. In addition, deletion of MCRA in *Lb. acidophilus* NCFM resulted in reduced growth of the deletion mutant in the presence of acids, bile and salt and significantly reduced the ability of the mutant strain to adhere to Caco-2 cells suggesting a role for the gene in the stress response, cell division and Caco-2 cell adherence [82]. Poly P granules are polyanionic inorganic biopolymers of

orthophosphate residues which have been associated with stress responses in bacteria [83]. The putative polyphosphate kinase gene of *Bifidobacterium scardovii* responsible for poly P synthesis was recently linked to the oxidative stress response as well as providing protection against other environmental stresses such as low pH [83].

Highest survival of *Lb. casei* ATCC 334 to acid stress was achieved by exposing the cells to pH 4.5 for 10 or 20 minutes [84]. Whole genome DNA microarrays revealed the up-regulation of 104 genes and the down-regulation of 216 genes after 20 minutes at pH 4.5. Malolactic fermentation and histidine accumulation were also revealed as important features of acid adaptation in *Lb. casei*. Malolactic enzyme was up-regulated 16-fold and 7 fold following 5 and 20 minutes of acid exposure, respectively. This enzyme functions to decarboxylate L-malate to L-lactate and CO₂, thus contributing to alkalinization of the cytoplasm and enabling the production of ATP through H⁺-ATPase [85,86]. The transcriptome of cells exposed to acid for 20 minutes also revealed the up-regulation of an eight gene cluster involved in histidine biosynthesis. The authors postulated that histidine accumulation may contribute to the buffering capacity within the cell and is the first report that histidine accumulation may enhance acid resistance in bacteria. Interestingly, addition of either 30 mM histidine or 30 mM malate during acid exposure (pH 2.5) for 60 minutes or 2 hours improved cell survival more than 100-fold or more than 10⁷-fold, respectively. The activity of the LuxS-mediated quorum sensing system, which is responsible for generating the universal signaling molecule called autoinducer-2 (AI-2) was found to be significantly increased following exposure to acid shock in *Lb. acidophilus* NCFM and *Lb. rhamnosus* GG suggesting a role for the *luxS* gene in the acid stress response of lactobacilli [87]. The gene encoding the universal stress protein Usp1 was also recently suggested to be an important mediator in the acid stress response of *Lb. plantarum* [88]. Indeed, studies in *Escherichia coli* demonstrated that Usp1 from *Lb. plantarum* inactivates the negative regulator PadR which is involved in the phenolic acid stress response by negatively regulating *padA*, a gene which encodes the phenolic acid decarboxylase enzyme.

A whole genome DNA microarray was employed to determine the transcriptional response of *Lb. reuteri* ATCC 55730 to bile [89]. Differential expression of a wide variety of genes involved in cell envelope stress, protein denaturation and DNA damage was observed. Interestingly, survival of *Lb. reuteri* in the presence of bile was decreased by mutating a Clp chaperone, a putative esterase and a gene of unknown function, whereas the ability of the strain to restart growth in the presence of bile was hampered by mutating two operon genes, a

multidrug resistance (MDR) transporter and a gene of unknown function, suggesting their importance in the bile stress response. The importance of MDR transporters for the bile stress response of *Lb. acidophilus* was also revealed by Pfeiler and Klaenhammer [90]. Indeed, of the ten most highly induced genes in *Lb. acidophilus* in the presence of bile, two were found to encode MDR transporters. These transporters function by extruding structurally unrelated compounds from the cell including antibiotics and bile salts [90]. MDR transporters were also identified in *B. longum* and *B. breve* following exposure to sub-inhibitory concentrations of bile [91]. Expression of the MDR transporter from *B. longum* in *E. coli* conferred bile resistance on the heterologous host when exposed to 3% bile [91]. However, the modified strain exhibited a reduced growth rate; hence the authors suggest that production of the MDR transporter is toxic for the cells. A gene responsible for S-layer production in *Lb. acidophilus* ATCC 4356, *slpA*, was found to be induced in bile concentrations ranging from 0.01-0.05% [92]. S-layer proteins are external bacterial structures which have been associated with protection against hostile environmental elements and the establishment of *Lb. acidophilus* in the GIT [93-95]. However, the expression of *slpA* in 0.1% bile was lower than that recorded in 0.02 and 0.05% bile although the level of S-layer protein on the cell surface increased in concentration. The authors therefore suggest that *slpB*, rather than *slpA*, may be expressed during unfavorable growth conditions [92]. A putative aggregation-promoting factor (Apf) of *Lb. acidophilus* NCFM has been linked to survival of the strain during passage through the GIT and may even be involved in bacterium-host interactions [96]. Indeed, a Δ *apf* deletion mutant was much more susceptible to bile and detergent, and survival rates of the mutant strain were decreased in simulated gastric juice compared to the control. Moreover, adherence of stationary phase mutant cells to an intestinal epithelial cell line was reduced. Overall, the results of the study suggest that Apf-like proteins are important for the gastrointestinal robustness of *Lb. acidophilus*.

Holistic approaches to understanding and exploiting the probiotic stress response

The general global cellular response of probiotic bacteria to environmental stresses can probably be grouped into six broad biological categories based on differential expression of the associated genes and on the encountered stress(es): (i) stress response genes, (ii) genes involved in energy metabolism, (iii) transcription and translation associated genes, (iv) genes involved in nucleotide metabolism and amino acid biosynthesis genes, (v) cell envelope and cell wall-associated genes, (vi) genes which have no assigned function. For

example, whole genome DNA microarrays exploited to study the complete cellular response of *B. longum* NCC2705 to heat treatment of 50°C for 3, 7 and 12 minutes revealed that genes involved in cell growth (ribosomal proteins, aminoacyl-tRNA synthetases, genes involved in cell division etc.) were markedly repressed, which as the authors suggest may represent a strategy to conserve energy which can then be directed towards protection mechanisms in the cell [97]. The most highly and rapidly induced genes included *dnaK*, *grpE*, *dnaJ* and the transcriptional repressor HspR. Genes encoding *groEL* and *groES* and the transcriptional regulator HrcA were induced but at lower levels. Numerous transcriptional regulators were induced which included 2 two-component response regulators and the corresponding sensor histidine kinases. Ten LacI-type sugar-responsive-repressors were up-regulated and genes involved in the SOS response were also induced including the transcriptional repressor LexA and genes encoding RecA, RecX and RecN as well as the *trans*-translation machinery. This latter mechanism involves a ribonucleoprotein complex made up of tmRNA (with properties of a tRNA and mRNA-encoded by *ssrA*) and the protein SmpB. Stalled translational complexes serve as the target for tmRNA-SmpB which 'rescues' the ribosome and tags the nascent polypeptide and mRNAs for degradation [98]. While *ssrA* was constitutively expressed, the gene encoding SmpB was highly induced following heat-treatment in *B. longum*.

An investigation into the global stress response of *B. breve* UCC2003 to moderate and severe heat treatment as well as osmotic, solvent and oxidative stress revealed that an interactive regulatory network controls the stress response in this strain [99,100]. Interestingly, exposure to moderate temperatures of 42°C and 44°C for 1 hour resulted in the induction of 5 and 17 genes, respectively and down-regulation of 11 and 92 genes, respectively. In contrast, 267 genes were induced during severe heat treatment (47°C) while 266 genes were down-regulated. Under conditions of severe heat shock, genes belonging to carbohydrate transport and metabolism, energy production and conversion and nucleotide transport and metabolism were negatively regulated. Of the up-regulated genes only a fraction were involved in protein misfolding and DNA damage. Based on the overall results, the authors proposed a model interactive regulatory gene network for the bifidobacterial stress response whereby the negative regulator HspR controls the SOS response and the ClgR regulon, which in turn is regulated by and regulates the negative regulator HrcA [99,100].

Previous studies have suggested that different bifidobacteria use NADH peroxidase to prevent the accumulation of H₂O₂ (101,102,103), however, none of the

sequenced genomes to date contain gene analogs for this enzyme. Interestingly, a recent proteomic study of the oxidative stress-related responses of *B. longum* BBMN68 revealed a change in the expression of 51 protein spots following oxygen exposure revealing a distinctively different set of detoxification proteins compared to other anaerobes (104). Proteins involved in protecting proteins, DNA and RNA were identified including alkyl hydroperoxide reductase (AhpC), pyridine nucleotide-disulfide reductase (PNDR), and DNA oxidative damage-protective proteins including DNA-binding ferritin-like protein (Dps), ribonucleotide reductase (NrdA) and NTP phosphohydrolases (Mut1). The activity of Dps in oxidative stress protection was confirmed by *in vitro* and *in vivo* studies. Indeed, *in vitro* studies revealed that Dps binds DNA to protect it from oxidative degradation and over-expression of the protein in *E. coli* increased survival of the cells under oxidative challenge.

Adapting *Lb. casei* ATCC 334 to a broad range of acid stresses to improve the acid tolerance of the strain demonstrated that exposure to pH 4.5 for 10 or 20 minutes resulted in the highest survival [84]. Assessment of the transcriptional responses of the strain following 5 and 20 minutes of exposure to this pH revealed a dramatic increase in the number of responsive genes following the 20 minute treatment (320 genes with altered expression levels). The majority of genes were down-regulated. This was particularly apparent for genes involved in information storage and processing including translation, ribosomal structure and biogenesis, transcription, DNA replication, recombination and repair as well as genes involved in cellular processes such as protein turnover and remarkably stress response genes and those involved in cell secretion and cell envelope biogenesis. Up-regulated genes included the malolactic enzyme (as discussed earlier) and genes involved in amino acid transport and metabolism, including those involved in the transport of histidine as previously discussed. Interestingly, genes involved in the mobile DNA elements category were also up-regulated following acid adaptation for 20 minutes. Of the poorly characterized up-regulated genes, three were associated with phospholipid turnover: an acetyl transferase, an esterase, and a putative membrane-associated phospholipid phosphatase. Acid-adapted cells had higher total percentages of saturated and cyclopropane fatty acids in the cytoplasmic membrane than control cells which may be linked to the up-regulation of the three phospholipid-associated genes. Other studies have also demonstrated that acid stress induces changes in the cytoplasmic membrane fatty acid content in *Lactobacillus* [105,106]. Moreover, several stresses have been shown to cause an increase in the concentration of cyclopropane fatty acids

in the cell membrane including heat stress [107], osmotic stress [108], and bile stress [109,110]. Indeed, using both transcriptomics and strategic proteomics approaches which also enabled a study of cell surface properties (surfome) Koskenniemi et al., [109] demonstrated the up-regulation of a gene encoding cyclopropane-fatty-acyl-phospholipid synthase in *Lb. rhamnosus* GG in response to bile stress, although the level of up-regulation was not deemed to be statistically significant. This study also demonstrated that bile shock resulted in the repression of EPS-encoding genes. The authors postulated that EPS serves to protect *Lb. rhamnosus* GG cells from the harsh environmental conditions of the stomach. However, the presence of bile serves as a signal of gut entrance and hence down-regulation in EPS production to enable better adherence of the bacterium to intestinal cells. Genes involved in the D-alanylation of the negatively charged lipoteichoic acids were also up-regulated in response to bile stress which was also observed for *Lb. plantarum* [111]. Such a strategy serves to increase the positive surface charge, and as the authors suggest possibly serves to repulse the cationic compounds in bile. Indeed, alteration of surface charge has also been associated with resistance to cationic peptides [112,113] and we recently observed the same phenomenon in response to bacteriophage challenge in *L. lactis* (unpublished). Several two-component systems, multi-drug transporters, the F_1F_0 -ATP synthase and a bile salt hydrolase were also up-regulated in response to bile stress as well as several chaperones and proteases directly involved in the stress response.

Discussion

The ability of probiotic bacteria to survive the harsh environments encountered during processing and gastrointestinal transit has been a major factor in their selection criteria. Indeed, induction of the probiotic stress response through pre-adaptation strategies may not always ensure the improved performance of a strain in compromising environments. For example, exposing *B. longum* DJO10A to sub-lethal stresses of acid, cold and centrifugation before addition to Cheddar cheese as a starter adjunct did not influence the viability of the culture during Cheddar cheese ripening [114]. Moreover, exposure of *Lb. acidophilus* La-5, *Lb. rhamnosus* GG and *Lb. fermentum* ME-3 to temperature, acid or bile stress did not positively affect survival of the strains in a gastrointestinal tract simulator, with a marked decrease in cell numbers for all three strains after the bile phase [115]. In addition, the biological efficacy of probiotic cells may be compromised following exposure to stress. Indeed, spray drying was recently shown to negatively influence the adhesion capacity of *Lactobacillus kefir* 8348 but not *Lb. plantarum* 83114 or *Lb. kefir* 8321 to intestinal cells [116].

Developing a molecular toolbox, whether through patho-biotechnology, targeting indigenous defense strategies of lactic acid bacteria or indeed other bacterial groups, should ensure that the most biologically/functionally active strains can be confidently selected for probiotic development. Indeed, we have already seen how probiotic robustness can be dramatically improved by targeting even a single mechanism. By targeting several cellular defense mechanisms in one strain we should be able to develop designer organisms with the capacity to overcome the plethora of stresses presented during processing and *in vivo* survival.

However, despite the advantages of using such approaches this field of science is not without its limitations. Probiotics which have been enhanced in this way are genetically modified organisms and with the exception of the United States and Canada, there is still uncertainty in the public arena towards the use of genetic manipulation. Moreover, the use of pathogen derived genes in genetically modified probiotics through patho-biotechnology is a concept that undoubtedly consumer and regulatory groups may find hard to accept. Yet despite this, designer probiotics offer huge potential for both technological and clinical applications. For example, Sleator and Hill [117] suggest that 'bioengineered probiotics' may provide a safer alternative to attenuated pathogens which are currently used as vaccine delivery platforms or may be used as novel drug delivery vehicles. Moreover, one of the major obstacles associated with genetic manipulation to date is that the benefits of the technology rarely benefit the consumer but rather serve to maximize corporate profit. Genetically modified probiotics, on the other hand, should directly benefit the consumer. However, studies which evaluate the safety of engineered probiotics are crucial if the technology is to gain acceptance. For this reason continued investigation and understanding of the bacterial stress response is a highly worthwhile endeavor in probiotic science providing strategies for scientists to manipulate probiotics to their full potential. In this way next-generation probiotic cultures will be better equipped to face technological and gastrointestinal challenges, (which at the moment can be a rate-limiting step in probiotic selection) as well as meeting medical demands.

Acknowledgements

This article has been published as part of *Microbial Cell Factories* Volume 10 Supplement 1, 2011: Proceedings of the 10th Symposium on Lactic Acid Bacterium. The full contents of the supplement are available online at <http://www.microbialcellfactories.com/supplements/10/S1>.

Author details

¹Teagasc Food Research Centre, Moorepark, Fermoy, Co. Cork, Ireland.

²Department of Microbiology, University College Cork, Ireland. ³Alimentary Pharmabiotic Centre, Cork, Ireland.

Competing interests

The authors declare that they have no competing interests.

Published: 30 August 2011

References

- Global Industry Analysts Inc: **Probiotics: A Global Strategic Business Report**. 2010 [http://www.researchandmarkets.com].
- Mills S, Stanton C, Ross RP: **Microbial production of bioactives: from fermented Functional Foods to probiotic mechanisms**. *Austral. J. Dairy Technol* 2009, **64**:41-49.
- O'Flaherty S, Saulnier DM, Pot B, Versalovic J: **How can probiotics and prebiotics impact mucosal immunity?** *Gut Microbes* 2010, **1**:293-300.
- Ross RP, Mills S, Hill C, Fitzgerald GF, Stanton C: **Specific metabolite production by gut microbiota as a basis for probiotic function**. *Int Dairy J* 2010, **20**:269-276.
- Corcoran BM, Ross RP, Fitzgerald GF, Dockery P, Stanton C: **Enhanced survival of GroESL-overproducing *Lactobacillus paracasei* NFBC 338 under stressful conditions induced by drying**. *Appl Environ Microbiol* 2006, **72**:5104-5107.
- Guinane CM, Kent RM, Norberg S, Hill C, Fitzgerald GF, Stanton C, Ross RP: **Host specific diversity in *Lactobacillus johnsonii* as evidenced by a major chromosomal inversion and phage resistance mechanisms**. *Plos One* 2011, **6**:e18740.
- Muller JA, Ross RP, Sybesma WFH, Fitzgerald GF, Stanton C: **Modification of technical properties of *Lactobacillus johnsonii* NCC 533 by supplementing growth medium with unsaturated fatty acids**. *Appl Environ Microbiol* 2011, doi:10.1128/AEM.05213-11.
- Meng XC, Stanton C, Fitzgerald GF, Daly C, Ross RP: **Anhydrobiotics: the challenges of drying probiotic cultures**. *Food Chemistry* 2008, **106**:1406-1416.
- Somero GN: **Proteins and temperature**. *Annu Rev Physiol* 1995, **57**:43-68.
- Teixeira P, Castro H, Mohacs-Farkas C, Kirby R: **Identification of sites of injury in *Lactobacillus bulgaricus* during heat stress**. *J Appl Microbiol* 1997, **83**:219-226.
- Santivarangkna C, Kulozik U, Foerst P: **Alternative drying processes for the industrial preservation of lactic acid starter cultures**. *Biotechnol Prog* 2007, **23**:302-315.
- Jennings TA: **Lyophilisation - introduction and basic principles**. Boca Raton: CRC Press; 1999.
- Oetjen GW: **Freeze-drying**. Weinheim: Wiley-VCH; 1999.
- Wang YC, Yu RC, Chou CC: **Viability of lactic acid bacteria and bifidobacteria in fermented soymilk after drying, subsequent rehydration and storage**. *Int J Food Microbiol* 2004, **93**:209-217.
- van de Guchte M, Serror P, Chervaux C, Smokvina T, Ehrlich SD, Maguin E: **Stress responses in lactic acid bacteria**. *Antonie Van Leeuwenhoek* 2002, **82**:187-216.
- Bayles DO, Tunick MH, Foglia TA, Miller AJ: **Cold shock and its effect on ribosomes and thermal tolerance in *Listeria monocytogenes***. *Appl Environ Microbiol* 2000, **66**:4351-4355.
- Graumann PL, Marahiel MA: **Cold shock response in *Bacillus subtilis***. *J Mol Microbiol Biotechnol* 1999, **1**:203-209.
- Miyoshi A, Rochat T, Gratadoux JJ, Le Loir Y, Oliveira SC, Langella P, Azevedo V: **Oxidative stress in *Lactococcus lactis***. *Genet Mol Res* 2003, **2**:348-359.
- Vido K, Diemer H, Van Dorsselaer A, Leize E, Juillard V, Gruss A, Gaudu P: **Roles of thioredoxin reductase during the aerobic life of *Lactococcus lactis***. *J Bacteriol* 2005, **187**:601-610.
- Lee JH, O'Sullivan DJ: **Genomic insights into bifidobacteria**. *Microbiol Mol Biol Rev* 2010, **74**:378-416.
- Cronin M, Ventura M, Fitzgerald GF, van Sinderen D: **Progress in genomics, metabolism and biotechnology of bifidobacteria**. *Int J Food Microbiol* 2011, **149**:4-18.
- Simpson PJ, Ross RP, Fitzgerald GF, Stanton C: ***Bifidobacterium psychrophilum* sp. nov. and *Aeriscardovia aeriphila* gen. nov., sp. nov., isolated from a porcine caecum**. *Int J Syst Evol Microbiol* 2004, **54**:401-406.
- Bolduc MP, Raymond Y, Fustier P, Champagne CP, Vuilleumard JC: **Sensitivity of bifidobacteria to oxygen and redox potential in non-fermented pasteurised milk**. *Int Dairy J* 2006, **16**:1038-1048.
- Odumaki T, Xiao JZ, Yonezawa S, Yaeshima T, Iwatsuki K: **Improved viability of bifidobacteria in fermented milk by cocultivation with *Lactococcus lactis* subspecies *lactis***. *J Dairy Sci* 2011, **94**:1112-1121.
- Hufner E, Markieton T, Chaillou S, Crutz-Le Coq AM, Zagorec M, Hertel C: **Identification of *Lactobacillus sakei* genes induced during meat fermentation and their role in survival and growth**. *Appl Environ Microbiol* 2007, **73**:2522-2531.
- Hufner E, Britton RA, Roos S, Jonsson H, Hertel C: **Global transcriptional response of *Lactobacillus reuteri* to the sourdough environment**. *Syst Appl Microbiol* 2008, **31**:323-338.
- Fortin MH, Champagne CP, St-Gelais D, Britten M, Fustier P, Lacroix M: **Effect of time of inoculation, starter addition, oxygen level and salting on the viability of probiotic cultures during Cheddar cheese manufacture**. *Int Dairy J* 2011, **21**:75-82.
- Leverrier P, Fremont Y, Rouault A, Boyaval P, Jan G: **In vitro tolerance to digestive stresses of propionibacteria: influence of food matrices**. *Food Microbiol* 2005, **22**:11-18.
- Corcoran BM, Stanton C, Fitzgerald G, Ross RP: **Life under stress: the probiotic stress response and how it may be manipulated**. *Curr Pharm Des* 2008, **14**:1382-1399.
- Begley M, Gahan CG, Hill C: **The interaction between bacteria and bile**. *FEMS Microbiol Rev* 2005, **29**:625-651.
- Leverrier P, Dimova D, Pichereau V, Auffray Y, Boyaval P, Jan G: **Susceptibility and adaptive response to bile salts in *Propionibacterium freudenreichii*: physiological and proteomic analysis**. *Appl Environ Microbiol* 2003, **69**:3809-3818.
- De Angelis M, Gobbetti M: **Environmental stress responses in *Lactobacillus*: a review**. *Proteomics* 2004, **4**:106-122.
- Sugimoto S, Abdullah AI M, Sonomoto K: **Molecular chaperones in lactic acid bacteria: physiological consequences and biochemical properties**. *J Biosci Bioeng* 2008, **106**:324-336.
- van Schaik W, Abee T: **The role of σB in the stress response of Gram-positive bacteria - targets for food preservation and safety**. *Curr Opin Biotechnol* 2005, **16**:218-224.
- Klaenhammer TR, Altermann E, Azcarate-Peril MA, McAuliffe O, Russell M: ***Lactobacillus acidophilus* nucleic acid sequences encoding stress-related proteins and uses thereof**. 2010, Patent No. WO/2005/086794 (USA).
- Desmond C, Stanton C, Fitzgerald GF, Collins K, Ross RP: **Environmental adaptation of probiotic lactobacilli towards improvement of performance during spray drying**. *Int Dairy J* 2001, **11**:801-808.
- Desmond C, Fitzgerald GF, Stanton C, Ross RP: **Improved stress tolerance of GroESL-overproducing *Lactococcus lactis* and probiotic *Lactobacillus paracasei* NFBC 338**. *Appl Environ Microbiol* 2004, **70**:5929-5936.
- Jakob U, Gaestel M, Engel K, Buchner J: **Small heat shock proteins are molecular chaperones**. *J Biol Chem* 1993, **268**:1517-1520.
- Narberhaus F: **Alpha-crystallin-type heat shock proteins: socializing minichaperones in the context of a multichaperone network**. *Microbiol Mol Biol Rev* 2002, **66**:64-93.
- Veinger L, Diamant S, Buchner J, Goloubinoff P: **The small heat-shock protein *IbpB* from *Escherichia coli* stabilizes stress-denatured proteins for subsequent refolding by a multichaperone network**. *J Biol Chem* 1998, **273**:11032-11037.
- Spano G, Capozzi V, Vernile A, Massa S: **Cloning, molecular characterization and expression analysis of two small heat shock genes isolated from wine *Lactobacillus plantarum***. *J Appl Microbiol* 2004, **97**:774-782.
- Spano G, Beneduce L, Perrotta C, Massa S: **Cloning and characterization of the *hsp 18.55* gene, a new member of the small heat shock gene family isolated from wine *Lactobacillus plantarum***. *Res Microbiol* 2005, **156**:219-224.
- Capozzi V, Weidmann S, Fiocco D, Rieu A, Hols P, Guzzo J, Spano G: **Inactivation of a small heat shock protein affects cell morphology and membrane fluidity in *Lactobacillus plantarum* WCFS1**. *Res Microbiol* 2011.
- Fiocco D, Capozzi V, Goffin P, Hols P, Spano G: **Improved adaptation to heat, cold, and solvent tolerance in *Lactobacillus plantarum***. *Appl Microbiol Biotechnol* 2007, **77**:909-915.
- Phadtare S, Inouye M: **Role of CspC and CspE in regulation of expression of RpoS and UspA, the stress response proteins in *Escherichia coli***. *J Bacteriol* 2001, **183**:1205-1214.
- Derzelle S, Hallet B, Francis KP, Ferain T, Delcour J, Hols P: **Changes in *cspL*, *cspP*, and *cspC* mRNA abundance as a function of cold shock and growth phase in *Lactobacillus plantarum***. *J Bacteriol* 2000, **182**:5105-5113.
- Derzelle S, Hallet B, Ferain T, Delcour J, Hols P: **Improved adaptation to cold-shock, stationary-phase, and freezing stresses in *Lactobacillus***

- plantarum* overproducing cold-shock proteins. *Appl Environ Microbiol* 2003, **69**:4285-4290.
48. Sleator RD, Hill C: Patho-biotechnology: using bad bugs to do good things. *Curr Opin Biotechnol* 2006, **17**:211-216.
 49. Sleator RD, Hill C: Bacterial osmoadaptation: the role of osmolytes in bacterial stress and virulence. *FEMS Microbiol Rev* 2001, **26**:49-71.
 50. Galinski EA: Osmoadaptation in bacteria. *Adv Microb Physiol* 1995, **37**:272-328.
 51. Brown AD: Microbial water stress. *Bacteriol Rev* 1976, **40**:803-846.
 52. Lippert K, Galinski EA: Enzyme stabilisation by ectoine-type compatible solutes: protection against heating, freezing and drying. *Appl Microbiol Biotechnol* 1992, **37**:61-65.
 53. Welsh DT: Ecological significance of compatible solute accumulation by micro-organisms: from single cells to global climate. *FEMS Microbiol Rev* 2000, **24**:263-290.
 54. Sheehan VM, Sleator RD, Fitzgerald GF, Hill C: Heterologous expression of BetL, a betaine uptake system, enhances the stress tolerance of *Lactobacillus salivarius* UCC118. *Appl Environ Microbiol* 2006, **72**:2170-2177.
 55. Sleator RD, Francis GA, O'Beirne D, Gahan CG, Hill C: Betaine and carnitine uptake systems in *Listeria monocytogenes* affect growth and survival in foods and during infection. *J Appl Microbiol* 2003, **95**:839-846.
 56. Sleator RD, Gahan CG, Abbe T, Hill C: Identification and disruption of BetL, a secondary glycine betaine transport system linked to the salt tolerance of *Listeria monocytogenes* LO28. *Appl Environ Microbiol* 1999, **65**:2078-2083.
 57. Sleator RD, Gahan CG, Hill C: A postgenomic appraisal of osmotolerance in *Listeria monocytogenes*. *Appl Environ Microbiol* 2003, **69**:1-9.
 58. Smiddy M, Sleator RD, Patterson MF, Hill C, Kelly AL: Role for compatible solutes glycine betaine and L-carnitine in listerial barotolerance. *Appl Environ Microbiol* 2004, **70**:7555-7557.
 59. Sheehan VM, Sleator RD, Hill C, Fitzgerald GF: Improving gastric transit, gastrointestinal persistence and therapeutic efficacy of the probiotic strain *Bifidobacterium breve* UCC2003. *Microbiology* 2007, **153**:3563-3571.
 60. Briones V, Blanco MM, Marco A, Prats N, Fernandez-Garayzabal JF, Suarez G, Domingo M, Dominguez L: Biliary excretion as possible origin of *Listeria monocytogenes* in fecal carriers. *Am J Vet Res* 1992, **53**:191-193.
 61. Hardy J, Francis KP, DeBoer M, Chu P, Gibbs K, Contag CH: Extracellular replication of *Listeria monocytogenes* in the murine gall bladder. *Science* 2004, **303**:851-853.
 62. Sleator RD, Wemekamp-Kamphuis HH, Gahan CG, Abbe T, Hill C: A PrfA-regulated bile exclusion system (BilE) is a novel virulence factor in *Listeria monocytogenes*. *Mol Microbiol* 2005, **55**:1183-1195.
 63. Watson D, Sleator RD, Hill C, Gahan CG: Enhancing bile tolerance improves survival and persistence of *Bifidobacterium* and *Lactococcus* in the murine gastrointestinal tract. *BMC Microbiol* 2008, **8**:176.
 64. Begley M, Sleator RD, Gahan CG, Hill C: Contribution of three bile-associated loci, *bsh*, *pva*, and *btbB*, to gastrointestinal persistence and bile tolerance of *Listeria monocytogenes*. *Infect Immun* 2005, **73**:894-904.
 65. Desmond C, Ross RP, O'Callaghan E, Fitzgerald G, Stanton C: Improved survival of *Lactobacillus paracasei* NFB3 338 in spray-dried powders containing gum acacia. *J Appl Microbiol* 2002, **93**:1003-1011.
 66. Alp G, Aslim B: Relationship between the resistance to bile salts and low pH with exopolysaccharide (EPS) production of *Bifidobacterium* spp. isolated from infants feces and breast milk. *Anaerobe* 2010, **16**:101-105.
 67. Stack HM, Kearney N, Stanton C, Fitzgerald GF, Ross RP: Association of beta-glucan endogenous production with increased stress tolerance of intestinal lactobacilli. *Appl Environ Microbiol* 2010, **76**:500-507.
 68. Theuwissen E, Mensink RP: Water-soluble dietary fibers and cardiovascular disease. *Physiol Behav* 2008, **94**:285-292.
 69. Wilson TA, Nicolosi RJ, Delaney B, et al: Reduced and high molecular weight barley beta-glucans decrease plasma total and non-HDL-cholesterol in hypercholesterolemic Syrian golden hamsters. *J Nutr* 2004, **134**:2617-2622.
 70. Akramiene D, Kondrotas A, Didziapetriene J, Kevelaitis E: Effects of beta-glucans on the immune system. *Medicina (Kaunas)* 2007, **43**:597-606.
 71. Volman JJ, Ramakers JD, Plat J: Dietary modulation of immune function by beta-glucans. *Physiol Behav* 2008, **94**:276-284.
 72. Mantovani MS, Bellini MF, Angeli JP, Oliveira RJ, Silva AF, Ribeiro LR: beta-Glucans in promoting health: prevention against mutation and cancer. *Mutat Res* 2008, **658**:154-161.
 73. Snart J, Bibiloni R, Grayson T, et al: Supplementation of the diet with high-viscosity beta-glucan results in enrichment for lactobacilli in the rat cecum. *Appl Environ Microbiol* 2006, **72**:1925-1931.
 74. Bruno-Barcena JM, Andrus JM, Libby SL, Klaenhammer TR, Hassan HM: Expression of a heterologous manganese superoxide dismutase gene in intestinal lactobacilli provides protection against hydrogen peroxide toxicity. *Appl Environ Microbiol* 2004, **70**:4702-4710.
 75. Bruno-Barcena JM, Azcarate-Peril MA, Klaenhammer TR, Hassan HM: Marker-free chromosomal integration of the manganese superoxide dismutase gene (*sodA*) from *Streptococcus thermophilus* into *Lactobacillus gasserii*. *FEMS Microbiol Lett* 2005, **246**:91-101.
 76. Noonpakdee W, Sitthimonchai S, Panyim S, Lertsiri S: Expression of the catalase gene *katA* in starter culture *Lactobacillus plantarum* TISTR850 tolerates oxidative stress and reduces lipid oxidation in fermented meat product. *Int J Food Microbiol* 2004, **95**:127-135.
 77. An H, Zhou H, Huang Y, Wang G, Luan C, Mou J, Luo Y, Hao Y: High-level expression of heme-dependent catalase gene *katA* from *Lactobacillus sakei* protects *Lactobacillus rhamnosus* from oxidative stress. *Mol Biotechnol* 2010, **45**:155-160.
 78. Wang G, Yin S, An H, Chen S, Hao Y: Coexpression of bile salt hydrolase gene and catalase gene remarkably improves oxidative stress and bile salt resistance in *Lactobacillus casei*. *J Ind Microbiol Biotechnol* 2010.
 79. Berger B, Moine D, Mansourian R, Arigoni F: HspR mutations are naturally selected in *Bifidobacterium longum* when successive heat shock treatments are applied. *J Bacteriol* 2010, **192**:256-263.
 80. Vrancken G, Rimaux T, Wouters D, Leroy F, De Vuyst L: The arginine deiminase pathway of *Lactobacillus fermentum* IMDO 130101 responds to growth under stress conditions of both temperature and salt. *Food Microbiol* 2009, **26**:720-727.
 81. Rosberg-Cody E, Liavonchanka A, Gobel C, Ross RP, O'Sullivan O, Fitzgerald GF, Feussner I, Stanton C: Myosin-cross-reactive antigen (MCRA) protein from *Bifidobacterium breve* is a FAD-dependent fatty acid hydratase which has a function in stress protection. *BMC Biochem* 2011, **12**:9.
 82. O'Flaherty SJ, Klaenhammer TR: Functional and phenotypic characterization of a protein from *Lactobacillus acidophilus* involved in cell morphology, stress tolerance and adherence to intestinal cells. *Microbiology* 2010, **156**:3360-3367.
 83. Qian Y, Borowski WJ, Calhoon WD: Intracellular granule formation in response to oxidative stress in *Bifidobacterium*. *Int J Food Microbiol* 2011, **145**:320-325.
 84. Broadbent JR, Larsen RL, Deibel V, Steele JL: Physiological and transcriptional response of *Lactobacillus casei* ATCC 334 to acid stress. *J Bacteriol* 2010, **192**:2445-2458.
 85. Poolman B, Molenaar D, Smid EJ, Ubbink T, Abbe T, Renault PP, Konings WN: Malolactic fermentation: electrogenic malate uptake and malate/lactate antiport generate metabolic energy. *J Bacteriol* 1991, **173**:6030-6037.
 86. Renault P, Gaillardin C, Heslot H: Role of malolactic fermentation in lactic acid bacteria. *Biochimie* 1988, **70**:375-379.
 87. Moslehi-Jenabian S, Gori K, Jespersen L: Al-2 signalling is induced by acidic shock in probiotic strains of *Lactobacillus* spp. *Int J Food Microbiol* 2009, **135**:295-302.
 88. Gury J, Seraut H, Tran NP, Barthelmebs L, Weidmann S, Gervais P, Cavin JF: Inactivation of PadR, the repressor of the phenolic acid stress response, by molecular interaction with Usp1, a universal stress protein from *Lactobacillus plantarum*, in *Escherichia coli*. *Appl Environ Microbiol* 2009, **75**:5273-5283.
 89. Whitehead K, Versalovic J, Roos S, Britton RA: Genomic and genetic characterization of the bile stress response of probiotic *Lactobacillus reuteri* ATCC 55730. *Appl Environ Microbiol* 2008, **74**:1812-1819.
 90. Pfeiler EA, Klaenhammer TR: Role of transporter proteins in bile tolerance of *Lactobacillus acidophilus*. *Appl Environ Microbiol* 2009, **75**:6013-6016.
 91. Gueimonde M, Garrigues C, van Sinderen D, de los Reyes-Gavilan CG, Margolles A: Bile-inducible efflux transporter from *Bifidobacterium longum* NCC2705, conferring bile resistance. *Appl Environ Microbiol* 2009, **75**:3153-3160.
 92. Khaleghi M, Kermanshahi RK, Yaghoobi MM, Zarkesh-Esfahani SH, Baghizadeh A: Assessment of bile salt effects on s-layer production, *slp* gene expression and some physicochemical properties of *Lactobacillus acidophilus* ATCC 4356. *J Microbiol Biotechnol* 2010, **20**:749-756.
 93. Avall-Jaaskelainen S, Palva A: *Lactobacillus* surface layers and their applications. *FEMS Microbiol Rev* 2005, **29**:511-529.
 94. Frece J, Kos B, Svetec IK, Zgaga Z, Mrsa V, Suskovic J: Importance of S-layer proteins in probiotic activity of *Lactobacillus acidophilus* M92. *J Appl Microbiol* 2005, **98**:285-292.

95. Kos B, Suskovic J, Vukovic S, Simpraga M, Frece J, Matosic S: **Adhesion and aggregation ability of probiotic strain *Lactobacillus acidophilus* M92.** *J Appl Microbiol* 2003, **94**:981-987.
96. Goh YJ, Klaenhammer TR: **Functional roles of aggregation-promoting-like factor in stress tolerance and adherence of *Lactobacillus acidophilus* NCFM.** *Appl Environ Microbiol* 2010, **76**:5005-5012.
97. Rezzonico E, Lariani S, Barretto C, Cuanoud G, Giliberti G, Delley M, Arigoni F, Pessi G: **Global transcriptome analysis of the heat shock response of *Bifidobacterium longum*.** *FEMS Microbiol Lett* 2007, **271**:136-145.
98. Keiler KC: **Biology of trans-translation.** *Annu Rev Microbiol* 2008, **62**:133-151.
99. Zomer A, Fernandez M, Kearney B, Fitzgerald GF, Ventura M, van Sinderen D: **An interactive regulatory network controls stress response in *Bifidobacterium breve* UCC2003.** *J Bacteriol* 2009, **191**:7039-7049.
100. Zomer A, van Sinderen D: **Intertwinement of stress response regulons in *Bifidobacterium breve* UCC2003.** *Gut Microbes* 2010, **1**:100-102.
101. Shimamura S, Abe F, Ishibashi N, Miyakawa H, Yaeshima T, Araya T, Tomita M: **Relationship between oxygen sensitivity and oxygen metabolism of *Bifidobacterium* species.** *J Dairy Sci* 1992, **75**:3296-3306.
102. Shin SY, Park JH: **Activities of oxidative enzymes related with oxygen tolerance in *Bifidobacterium* sp.** *J Microbiol Biotechnol* 1997, **7**:356-359.
103. Talwalkar A, Kailasapathy K: **Metabolic and biochemical responses of probiotic bacteria to oxygen.** *J Dairy Sci* 2003, **86**:2537-2546.
104. Xiao M, Xu P, Zhao J, Wang Z, Zuo F, Zhang J, Ren F, Li P, Chen S, Ma H: **Oxidative stress-related responses of *Bifidobacterium longum* subsp. *longum* BBMN68 at the proteomic level after exposure to oxygen.** *Microbiology* 2011, **157**:1573-1588.
105. Montanari C, Sado Kamdem SL, Serrazanetti DI, Etoa FX, Guerzoni ME: **Synthesis of cyclopropane fatty acids in *Lactobacillus helveticus* and *Lactobacillus sanfranciscensis* and their cellular fatty acids changes following short term acid and cold stresses.** *Food Microbiol* 2010, **27**:493-502.
106. Streit F, Delettre J, Corrieu G, Beal C: **Acid adaptation of *Lactobacillus delbrueckii* subsp. *bulgaricus* induces physiological responses at membrane and cytosolic levels that improves cryotolerance.** *J Appl Microbiol* 2008, **105**:1071-1080.
107. Broadbent JR, Lin C: **Effect of heat shock or cold shock treatment on the resistance of *Lactococcus lactis* to freezing and lyophilization.** *Cryobiology* 1999, **39**:88-102.
108. Guillot A, Obis D, Mistou MY: **Fatty acid membrane composition and activation of glycine-betaine transport in *Lactococcus lactis* subjected to osmotic stress.** *Int J Food Microbiol* 2000, **55**:47-51.
109. Koskeniemi K, Laakso K, Koponen J, et al: **Proteomics and transcriptomics characterization of bile stress response in probiotic *Lactobacillus rhamnosus* GG.** *Mol Cell Proteomics* 2011, **10**:M1110 002741.
110. Taranto MP, Fernandez Murga ML, Lorca G, de Valdez GF: **Bile salts and cholesterol induce changes in the lipid cell membrane of *Lactobacillus reuteri*.** *J Appl Microbiol* 2003, **95**:86-91.
111. Bron PA, Molenaar D, de Vos WM, Kleerebezem M: **DNA micro-array-based identification of bile-responsive genes in *Lactobacillus plantarum*.** *J Appl Microbiol* 2006, **100**:728-738.
112. Kovacs M, Halfmann A, Fedtke I, Heintz M, Peschel A, Vollmer W, Hakenbeck R, Bruckner R: **A functional *dlt* operon, encoding proteins required for incorporation of d-alanine in teichoic acids in gram-positive bacteria, confers resistance to cationic antimicrobial peptides in *Streptococcus pneumoniae*.** *J Bacteriol* 2006, **188**:5797-5805.
113. Peschel A, Otto M, Jack RW, Kalbacher H, Jung G, Gotz F: **Inactivation of the *dlt* operon in *Staphylococcus aureus* confers sensitivity to defensins, protegrins, and other antimicrobial peptides.** *J Biol Chem* 1999, **274**:8405-8410.
114. Scheller M, O'Sullivan DJ: **Comparative analysis of an intestinal strain of *Bifidobacterium longum* and a strain of *Bifidobacterium animalis* subspecies *lactis* in Cheddar cheese.** *J Dairy Sci* 2011, **94**:1122-1131.
115. Sumeri I, Arike L, Stekolstikova J, Uusna R, Adamberg S, Adamberg K, Paalme T: **Effect of stress pretreatment on survival of probiotic bacteria in gastrointestinal tract simulator.** *Appl Microbiol Biotechnol* 2010, **86**:1925-1931.
116. Golowczyc MA, Silva J, Teixeira P, De Antoni GL, Abraham AG: **Cellular injuries of spray-dried *Lactobacillus* spp. isolated from kefir and their impact on probiotic properties.** *Int J Food Microbiol* 2011, **144**:556-560.
117. Sleanor RD, Hill C: **Bioengineered bugs - a patho-biotechnology approach to probiotic research and applications.** *Med Hypotheses* 2008, **70**:167-169.
118. Prasad J, McJarrow P, Gopal P: **Heat and osmotic stress responses of probiotic *Lactobacillus rhamnosus* HN001 (DR20) in relation to viability after drying.** *Appl Environ Microbiol* 2003, **69**:917-925.
119. Suokko A, Savijoki K, Malinen E, Palva A, Varmanen P: **Characterization of a mobile *clpL* gene from *Lactobacillus rhamnosus*.** *Appl Environ Microbiol* 2005, **71**:2061-2069.
120. Suokko A, Poutanen M, Savijoki K, Kalkkinen N, Varmanen P: **ClpL is essential for induction of thermotolerance and is potentially part of the HrcA regulon in *Lactobacillus gasserii*.** *Proteomics* 2008, **8**:1029-1041.
121. Fiocco D, Collins M, Muscarelli L, Hols P, Kleerebezem M, Msadek T, Spano G: **The *Lactobacillus plantarum* *ftsH* gene is a novel member of the CtsR stress response regulon.** *J Bacteriol* 2009, **191**:1688-1694.
122. Savijoki K, Suokko A, Palva A, Valmu L, Kalkkinen N, Varmanen P: **Effect of heat-shock and bile salts on protein synthesis of *Bifidobacterium longum* revealed by [35S]methionine labelling and two-dimensional gel electrophoresis.** *FEMS Microbiol Lett* 2005, **248**:207-215.
123. Ventura M, Canchaya C, Zink R, Fitzgerald GF, van Sinderen D: **Characterization of the *groEL* and *groES* loci in *Bifidobacterium breve* UCC 2003: genetic, transcriptional, and phylogenetic analyses.** *Appl Environ Microbiol* 2004, **70**:6197-6209.
124. Ventura M, Fitzgerald GF, van Sinderen D: **Genetic and transcriptional organization of the *clpC* locus in *Bifidobacterium breve* UCC 2003.** *Appl Environ Microbiol* 2005, **71**:6282-6291.
125. Ventura M, Zhang Z, Cronin M, Canchaya C, Kenny JG, Fitzgerald GF, van Sinderen D: **The CigR protein regulates transcription of the *clpP* operon in *Bifidobacterium breve* UCC 2003.** *J Bacteriol* 2005, **187**:8411-8426.
126. Ventura M, Canchaya C, Zhang Z, Fitzgerald GF, van Sinderen D: **Molecular characterization of *hsp20*, encoding a small heat shock protein of *Bifidobacterium breve* UCC2003.** *Appl Environ Microbiol* 2007, **73**:4695-4703.
127. Serror P, Dervyn R, Ehrlich SD, Maguin E: ***csp*-like genes of *Lactobacillus delbrueckii* ssp. *bulgaricus* and their response to cold shock.** *FEMS Microbiol Lett* 2003, **226**:323-330.
128. Wang Y, Delettre J, Guillot A, Corrieu G, Beal C: **Influence of cooling temperature and duration on cold adaptation of *Lactobacillus acidophilus* RD758.** *Cryobiology* 2005, **50**:294-307.
129. Ventura M, Zink R, Fitzgerald GF, van Sinderen D: **Gene structure and transcriptional organization of the *dnaK* operon of *Bifidobacterium breve* UCC 2003 and application of the operon in bifidobacterial tracing.** *Appl Environ Microbiol* 2005, **71**:487-500.
130. Ventura M, Canchaya C, Bernini V, Del Casale A, Dellaglio F, Neviani E, Fitzgerald GF, van Sinderen D: **Genetic characterization of the *Bifidobacterium breve* UCC 2003 *hrcA* locus.** *Appl Environ Microbiol* 2005, **71**:8998-9007.
131. Lambert JM, Bongers RS, de Vos WM, Kleerebezem M: **Functional analysis of four bile salt hydrolase and penicillin acylase family members in *Lactobacillus plantarum* WCFS1.** *Appl Environ Microbiol* 2008, **74**:4719-4726.
132. Price CE, Reid SJ, Driessen AJ, Abratt VR: **The *Bifidobacterium longum* NCIMB 702259T *ctr* gene codes for a novel cholate transporter.** *Appl Environ Microbiol* 2006, **72**:923-926.
133. Wall T, Bath K, Britton RA, Jonsson H, Versalovic J, Roos S: **The early response to acid shock in *Lactobacillus reuteri* involves the ClpL chaperone and a putative cell wall-altering esterase.** *Appl Environ Microbiol* 2007, **73**:3924-3935.
134. De Angelis M, Bini L, Pallini V, Cocconcelli PS, Gobetti M: **The acid-stress response in *Lactobacillus sanfranciscensis* CB1.** *Microbiology* 2001, **147**:1863-1873.
135. Azcarate-Peril MA, McAuliffe O, Altermann E, Lick S, Russell WM, Klaenhammer TR: **Microarray analysis of a two-component regulatory system involved in acid resistance and proteolytic activity in *Lactobacillus acidophilus*.** *Appl Environ Microbiol* 2005, **71**:5794-5804.
136. Kullen MJ, Klaenhammer TR: **Identification of the pH-inducible, proton-translocating F1F0-ATPase (*atpBEFHAGDC*) operon of *Lactobacillus acidophilus* by differential display: gene structure, cloning and characterization.** *Mol Microbiol* 1999, **33**:1152-1161.

137. Ventura M, Canchaya C, van Sinderen D, Fitzgerald GF, Zink R: *Bifidobacterium lactis* DSM 10140: identification of the *atp* (*atpBEFHAGDC*) operon and analysis of its genetic structure, characteristics, and phylogeny. *Appl Environ Microbiol* 2004, **70**:3110-3121.
138. Sanchez B, Ruiz L, de los Reyes-Gavilan CG, Margolles A: Proteomics of stress response in *Bifidobacterium*. *Front Bioscience* 2008, **13**:6905-6919.

doi:10.1186/1475-2859-10-S1-S19

Cite this article as: Mills et al.: Enhancing the stress responses of probiotics for a lifestyle from gut to product and back again. *Microbial Cell Factories* 2011 **10**(Suppl 1):S19.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

