

Title	Emergence of winner-takes-all connectivity paths in random nanowire networks
Authors	Manning, Hugh G.;Niosi, Fabio;de Rocha, Claudia Gomes;Bellew, Allen T.;O'Callaghan, Colin;Biswas, Subhajit;Flowers, Patrick F.;Wiley, Benjamin J.;Holmes, Justin D.;Ferreira, Mauro S.;Bolan, John J.
Publication date	2018-08-13
Original Citation	Manning, H. G., Niosi, F., da Rocha, C. G., Bellew, A. T., O'Callaghan, C., Biswas, S., Flowers, P. F., Wiley, B. J., Holmes, J. D., Ferreira, M. S. and Boland, J. J. (2018) 'Emergence of winner-takes-all connectivity paths in random nanowire networks', Nature Communications, 9(1), 3219 (9 pp). doi: 10.1038/s41467-018-05517-6
Type of publication	Article (peer-reviewed)
Link to publisher's version	https://www.nature.com/articles/s41467-018-05517-6 - 10.1038/ s41467-018-05517-6
Rights	© The Author(s) 2018. Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit http://creativecommons.org/licenses/by/4.0/ - http:// creativecommons.org/licenses/by/4.0/
Download date	2025-07-05 13:05:37



University College Cork, Ireland Coláiste na hOllscoile Corcaigh

Description of Additional Supplementary Files

File Name: Supplementary Movie 1

Description: Animation for the Ag nanowire network shown in Supplementary Figure 7 with $A_j=0.05$ and $\alpha_j=1$. One can see the formation of two superimposed conductive paths in the first power-law regime. Subsequent paths are formed as current increases causing the observed changes in slopes in the conductance curve.

File Name: Supplementary Movie 2

Description: Animation for the Ag nanowire network shown in Supplementary Figure 7 with $A_j=0.05$ and $\alpha_j=1.1$. One can see the formation of a single conductive path in the power-law regime. Once all the junctions in this path are fully optimized, the network becomes temporarily Ohmic, i.e. its conductance does not change within a certain current window. Further paths are formed in a quantized manner as current is loaded onto the electrodes with the conductance curve depicting a stepwise increase.

File Name: Supplementary Movie 3

Description: Animation for the Ag nanowire network used in Supplementary Figure 8 with $A_j=0.05$ and $\alpha_j=1$. One can see the formation of multiple conductive paths in the first power-law regime. Subsequent paths are formed as current increases causing the observed changes in slopes in the conductance curve.

File Name: Supplementary Movie 4

Description: Animation for the Ag nanowire network used in Supplementary Figure 8 with A_j =0.05 and α_j =1.1. One can see the formation of a single conductive path practically slicing the network at half in the power-law regime. Once all the junctions in this path are fully optimized, the network becomes temporarily Ohmic, i.e. its conductance does not change within a certain current window. After the first conductance plateau, one can observe the formation of two independent conductive paths. As more current is loaded onto the terminals, additional paths are formed in a quantized manner with the conductance curve depicting a stepwise increase.